

## UNIT-01

### # Taylor's Series

$$y' = \frac{dy}{dx} = f(x, y)$$

$$y = y_0 + xy'_0 + \frac{x^2 y''_0}{2!} + \frac{x^3 y'''_0}{3!} + \dots$$

### # Rolle's Theorem

- +(i)  $f(x)$  is continuous in  $[a, b]$
- +(ii)  $f(x)$  is differentiable in  $(a, b)$
- +(iii)  $f(a) = f(b)$

then  $\exists$  atleast one point  $c \in (a, b)$  so that  $f'(c) = 0$

### # Mean Value theorem

- +(i)
- +(ii) same Rolle's theorem
- +(iii)  $f(a) \neq f(b)$

then  $\exists c \in (a, b)$  so that

$$f'(c) = \frac{f(b) - f(a)}{b - a}$$

### # Bisection Method

let  $f(x) = 0$

Suppose,  $f(0) = -ve$   
 $f(1) = -ve$   
 $f(2) = +ve$  } root lies b/w them

1st Approx.

$$x_1 = \frac{1+2}{2} = 1.5 ; f(1.5) = -ve$$

2nd Approx.

$$x_2 = \frac{1.5+2}{2} = 1.75 ; f(1.75) = +ve$$

3rd Approx.

$$x_3 = \dots$$

do on till required accuracy.

### # Secant Method

let  $f(x) = 0$

Suppose,  $f(0) = -ve$   
 $f(1) = -ve \rightarrow x_0$   
 $f(2) = +ve \rightarrow x_1$

$$x_{n+1} = \frac{x_{n-1} f(x_n) - x_n f(x_{n-1})}{f(x_n) - f(x_{n-1})}$$

$$x_2 = \frac{x_0 f(x_1) - x_1 f(x_0)}{f(x_1) - f(x_0)} ; f(x_2) =$$

$x_3 =$  similarly -

### # Newton Raphson Method

let  $f(x) = 0$

Suppose,  $f(0) = -ve$

$f(1) = -ve$   
 $f(2) = +ve$  } root lies b/w them

let  $x_0 = 1$  or  $2$  or  $\frac{1+2}{2} = 1.5$

also find  $f'(x)$

$$x_n = x_{n-1} - \frac{f(x_{n-1})}{f'(x_{n-1})}$$

$$n=1 \rightarrow x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}$$

# Absolute Error: If  $x$  is true value &  $x'$  its Approximate value.  
 $E_a = |x - x'|$

# Relative Error:  $E_R = \left| \frac{x - x'}{x} \right|$

# % Error:  $E_p = \left| \frac{x - x'}{x} \right| \times 100$

# Inherent errors: Errors which are already present in the statement of a problem before its solution.

# Rounding errors: Errors which arise from the process of rounding off the numbers.

# Truncation errors: Errors caused by using Approximate results.

! Do it Yourself 😊

- + Fibonacci
- + Golden Search
- + Newton's Method
- + Steepest
- + Nelder-Mead Algo.

## # Interpolation

→ Newton forward :

$$f(x+hu) = f(x) + \frac{u}{1!} \Delta f(x) + \frac{u(u-1)}{2!} \Delta^2 f(x) + \frac{u(u-1)(u-2)}{3!} \Delta^3 f(x) + \dots$$

$\nwarrow$  initial / Base  
 $\uparrow$  Interval

→ Newton Backward :

$$f(x+hu) = f(x) + u \nabla f(x) + \frac{u(u+1)}{2!} \nabla^2 f(x) + \frac{u(u+1)(u+2)}{3!} \nabla^3 f(x) + \dots$$

$u = \frac{x - x_0}{h}$  where  $x_0$  is initial / Base

# Lagrange's interpolation for unequal interval :

Ex:-

x	5	6	9	11
y	12	13	14	16

$$f(x) = \frac{(x-6)(x-9)(x-11)}{(5-6)(5-9)(5-11)} \times 12 + \frac{(x-5)(x-9)(x-11)}{(6-5)(6-9)(6-11)} \times 13 + \frac{(x-5)(x-6)(x-11)}{(9-5)(9-6)(9-11)} \times 14 + \frac{(x-5)(x-6)(x-9)}{(11-5)(11-6)(11-9)} \times 16$$

# Newton Divided Difference :

x	f(x)	$\Delta f(x)$	$\Delta^2 f(x)$	$\Delta^3 f(x)$
x <sub>0</sub> 5	12	$\frac{13-12}{6-5} = 1$	$\frac{\frac{1}{3}-1}{9-5} = -\frac{1}{6}$	$\frac{\frac{2}{15}-(-\frac{1}{6})}{11-5} = \frac{1}{20}$
x <sub>1</sub> 6	13	$\frac{14-13}{9-6} = \frac{1}{3}$		
x <sub>2</sub> 9	14	$\frac{16-14}{11-9} = 1$		
x <sub>3</sub> 11	16			

given : x=10

$$f(x) = f(x_0) + (x-x_0) \Delta f(x_0) + (x-x_0)(x-x_1) \Delta^2 f(x_0) + (x-x_0)(x-x_1)(x-x_2) \Delta^3 f(x_0) + \dots$$

# Numerical integration :

$$I = \int_a^b f(x) dx ; h = \frac{b-a}{n}$$

① Trapezoidal Rule :-

$$\int_a^b f(x) dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})]$$

∴ for any no. of interval.

② Simpson  $\frac{1}{3}$  Rule : (Even interval)

$$\int_a^b f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots) + 2(y_2 + y_4 + y_6 + \dots)]$$

③ Simpson  $\frac{3}{8}$  Rule : (for multiple of 3)

$$\int_a^b f(x) dx = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_4 + \dots) + 2(y_3 + y_6 + y_9 + \dots)]$$

# Gauss quadrature formula

Ques →  $I = \int_a^b f(x) dx$   
 step ①: change interval [a,b] into [-1,1] using  $x = \frac{b-a}{2}t + \frac{b+a}{2}$

step ②: Substitute x in f(x)  
 step ③: formula - integrand.

one-point :  $\int_{-1}^1 f(x) dx = 2f(0)$  (n=0)

two-point :  $\int_{-1}^1 f(x) dx = f(-\frac{1}{\sqrt{3}}) + f(\frac{1}{\sqrt{3}})$

three-point :  $\int_{-1}^1 f(x) dx = \frac{1}{9} [5f(-\frac{\sqrt{3}}{5}) + 8f(0) + 5f(\frac{\sqrt{3}}{5})]$



## UNIT - 03

### # Gauss Elimination Method.

Consider the system of eq<sup>s</sup>.

$$a_{11}x + a_{12}y + a_{13}z = b_1$$

$$a_{21}x + a_{22}y + a_{23}z = b_2$$

$$a_{31}x + a_{32}y + a_{33}z = b_3$$

Convert in Matrix form  $AX=B$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$$

make Augmented matrix,  $C=[A:B]$

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & : & b_1 \\ a_{21} & a_{22} & a_{23} & : & b_2 \\ a_{31} & a_{32} & a_{33} & : & b_3 \end{bmatrix}$$

∴ use Row transformations,

$$C = \begin{bmatrix} - & - & - & : & - \\ \bullet & - & - & : & - \\ \bullet & \bullet & - & : & - \end{bmatrix}$$

Convert back to system of eqs.

$$c_{11}x + c_{12}y + c_{13}z = d_1$$

$$c_{22}y + c_{23}z = d_2$$

$$c_{33}z = d_3$$

Now, solve these to get the Answers.

### # Gauss Jordan

Similar to Gauss Elimination

$$\begin{bmatrix} 1 & \bullet & \bullet & : & - \\ \bullet & 1 & \bullet & : & - \\ \bullet & \bullet & 1 & : & - \end{bmatrix}$$

$$x = d_1$$

$$y = d_2$$

$$z = d_3$$

### # Eigen Value Problem & Power Method.

Ex:  $AX_0 = \begin{bmatrix} 3 & -5 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -2 \\ 2 \end{bmatrix}$  Eigen Value  $\lambda = -2$  Eigen Vector  $\begin{bmatrix} -1 \\ 1 \end{bmatrix} \Rightarrow 2x_1$

$AX_1 = \begin{bmatrix} 3 & -5 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} +8 \\ -6 \end{bmatrix}$  bigger one  $\begin{bmatrix} 1 \\ -0.75 \end{bmatrix} = 8x_2$

$AX_2 = \begin{bmatrix} 3 & -5 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ -0.75 \end{bmatrix} = -x_3$

So on till we get same value of successive iteration

### # LU Decomposition Methods.

#### # DoLittle :

①  $AX=B$

②  $A=LU$

$$L = \begin{bmatrix} 1 & 0 & 0 \\ l_{21} & 1 & 0 \\ l_{31} & l_{32} & 1 \end{bmatrix} \quad U = \begin{bmatrix} u_{11} & u_{12} & u_{13} \\ 0 & u_{22} & u_{23} \\ 0 & 0 & u_{33} \end{bmatrix}$$

③  $LY=B$  ∴  $LU\underbrace{x}_y = B$

where,  $y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$  Solve to find  $y_1, y_2, y_3$

④  $UX=y$

from this we can find  $X(x, y, z)$  req. sol. ✓

#### # Crout's :

Same Procedure as DoLittle

$$L = \begin{bmatrix} a & 0 & 0 \\ b & c & 0 \\ d & e & f \end{bmatrix} \quad U = \begin{bmatrix} 1 & g & h \\ 0 & 1 & i \\ 0 & 0 & 1 \end{bmatrix}$$

#### # Cholesky's : $A \rightarrow \text{Matrix.}$

Conditions  $\left\{ \begin{array}{l} \rightarrow A \text{ is Symmetric i.e. } A^T = A \\ \rightarrow A \text{ is positive definite.} \end{array} \right.$

①  $AX=B$

②  $A=LL^T$

$$A = \begin{bmatrix} a & 0 & 0 \\ b & c & 0 \\ d & e & f \end{bmatrix} \begin{bmatrix} a & b & d \\ 0 & c & e \\ 0 & 0 & f \end{bmatrix} = \begin{bmatrix} a^2 & ab & ad \\ ab & b^2+c^2 & bd+ce \\ ad & bd+ce & d^2+e^2+f^2 \end{bmatrix}$$

③  $LY=B$ , where  $y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}$

④  $L^T X = y$



## # Picard's Method.

$$y = y_0 + \int_{x_0}^x f(x, y) dx$$

1st Approx. -

$$y_1 = y_0 + \int_{x_0}^x f(x, y_0) dx$$

2nd Approx. -

$$y_2 = y_0 + \int_{x_0}^x f(x, y_1) dx$$

Doing this till the desired degree of accuracy obtained.

## # Taylor's Series ( $f(x, y) = y = \frac{dy}{dx}$ )

$$y_1 = y_0 + h y_0' + \frac{h^2}{2!} y_0'' + \frac{h^3}{3!} y_0''' + \dots$$

$$h = (x - x_0)$$

$$x = x_0 + h$$

$$y_2 = y_1 + h y_1' + \frac{h^2}{2!} y_1'' + \frac{h^3}{3!} y_1''' + \dots$$

and so on - - -

## # Euler's Method

Consider,  $\frac{dy}{dx} = f(x, y)$ ;  $y(x_0) = y_0$

To find:  $y(x_n) = y_n$  (Initial)

Acc. to Euler's Method,

$$y_n = y_{n-1} + h \cdot f(x_{n-1}, y_{n-1}), n = 1, 2, 3, \dots$$

width of differencing

$$h = \frac{x_n - x_0}{n} = x_n - x_{n-1} = x_1 - x_0$$

Put  $n=1$ ,  $y_1 = y_0 + h f(x_0, y_0)$

Put  $n=2$ ,  $y_2 = y_1 + h f(x_1, y_1)$

and so on.

from Euler's Method, starting from  $y_0$  and using eqn ①, we can get  $y_1, y_2, y_3, \dots, y_n$ .

## # Modified Euler's Method. ( $x = x_0 + h$ )

Step ①:-  $y_1^{(0)} = y_0 + h f(x_0, y_0)$  (Euler Method)

$$y_1^{(n+1)} = y_0 + \frac{h}{2} [f(x_0, y_0) + f(x_1, y_1^{(n)})] \text{ (Modified)}$$

Put  $n = 0, 1, 2, 3, \dots$

Step ②: 1st Approx.

$$y_1^{(1)} = y_0 + \frac{h}{2} [f(x_0, y_0) + f(x_1, y_1^{(0)})]$$

Similarly.

Step ③: 2nd Approx.

$$y_1^{(2)} = y_0 + \frac{h}{2} [f(x_0, y_0) + f(x_1, y_1^{(1)})]$$

and so on. till required accuracy.

• if two consecutive values of  $y_1^{(k)}$  &  $y_1^{(k+1)}$  are same then  $y_1 = y_1^{(k)}$

preceding in the same manner,  $y_2, y_3, \dots$  are calculated.

( $x = x_0 + h$ )

## # Runge - Kutta (Second Order)

$$\frac{dy}{dx} = f(x, y); y(x_0) = y_0$$

To find  $y(x_1)$

$$k_1 = h f(x_n, y_n)$$

$$k_2 = h f(x_n + h, y_n + k_1)$$

$$y_{n+1} = y_n + \frac{1}{2} (k_1 + k_2)$$

## # Runge - Kutta (4th Order)

$$k_1 = h f(x_n, y_n)$$

$$k_2 = h f(x_n + \frac{h}{2}, y_n + \frac{k_1}{2})$$

$$k_3 = h f(x_n + \frac{h}{2}, y_n + \frac{k_2}{2})$$

$$k_4 = h f(x_n + h, y_n + k_3)$$

then compute;

$$k = \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$

$$y_{n+1} = y_n + k$$

## # Adams-Bashforth Predictor Corrector Method:

Consider,  $\frac{dy}{dx} = f(x, y)$  with initial  $y(x_0) = y_0$

To find:  $y(x_n)$  must be 4 starting values of  $y$ . find  $y(x_1), y(x_2), y(x_3)$  using any methods.

Then calculate -

$$f_0 = f(x_0, y_0), f_1, f_2, f_3$$

$$\text{Predictor} \rightarrow y_4 = y_3 + \frac{h}{24} (55f_3 - 59f_2 + 37f_1 - 9f_0)$$

$$\text{find } f_4 = f(x_4, y_4)$$

$$\text{Corrector} \rightarrow y_4 = y_3 + \frac{h}{24} (9f_4 + 19f_3 - 5f_2 + f_1)$$

again find  $f_4$  & again corrector.



## # Milne Predictor-Corrector Method:

$$\frac{dy}{dx} = f(x, y) \text{ with initial Cond}^n y(x_0) = y_0$$

To find  $y(x_4)$

find  $y(x_1), y(x_2), y(x_3)$  using any method!

Then Calculate:

$$f_0 = f(x_0, y_0) \quad f_1 = f(x_1, y_1)$$

$$f_2 = f(x_2, y_2) \quad f_3 = f(x_3, y_3)$$

Milne's Corrector formulae:

$$y_4 = y_2 + \frac{h}{3} (f_2 + 4f_3 + f_4)$$

find  $f_4$  and again Apply Milne method to find better value of  $y_4$ , we repeat this step until  $y_4$  remains unchanged.

## # Numerical Solution of Partial Diff. eq<sup>n</sup>. (Parabolic, Hyperbolic, elliptical)

$$a \frac{\partial^2 \phi}{\partial x^2} + b \frac{\partial^2 \phi}{\partial x \partial y} + c \frac{\partial^2 \phi}{\partial y^2}$$

$$D = b^2 - 4ac$$

$D > 0$  Hyperbola, 2 charac<sup>n</sup> sol<sup>n</sup>  
 $D = 0$  Parabola, "  
 $D < 0$  Ellipse, No Sol<sup>n</sup>.