


Comenius University

Faculty of Mathematics, Physics and Informatics

DIPLOMA THESIS




Jakub Bahyl

Quantum Turbulence Measurements in Superfluid Helium and Numerical Simulations

Bratislava, 2018

Comenius University
Faculty of Mathematics, Physics and Informatics

DIPLOMA THESIS



Jakub Bahyl

Quantum Turbulence Measurements in Superfluid Helium and Numerical Simulations

Department of Low Temperature Physics, Charles University in Prague

Study programme: Condensed Matter Physics

Study programme number: mFTL

Supervisor of the thesis: RNDr. David Schmoranzer, PhD.

Consultant of the thesis: Mgr. Emil Varga

Bratislava, 2018



Comenius University in Bratislava
Faculty of Mathematics, Physics and Informatics

THESIS ASSIGNMENT

Name and Surname:	Jakub Bahyl
Study programme:	Physics (Single degree study, bachelor I. deg., full time form)
Field of Study:	Physics
Type of Thesis:	Bachelor's thesis
Language of Thesis:	English
Secondary language:	Slovak
Title:	Measurement of Quantum Turbulence in Superfluid 4He Using Second Sound Attenuation
Aim:	During the work on this Thesis, the student will become familiar with the theoretical foundations of the description of quantum fluids and superfluidity. The student will learn about several experimental methods used for characterization of the flows of superfluid helium, such as second sound attenuation, which differs significantly from any techniques used in classical fluids. Given the focus of the Thesis, classical hydrodynamics will be used as a stepping-stone to help in the understanding of quantum fluids. On the practical side, the student will learn to design and run complex cryogenic experiments. This work will cover a wide range of skills, starting with the understanding of safety procedures for handling cryoliquids, of specific measures needed for operation of experiments with liquid helium at temperatures between 1.2 K and 4.2 K, and including work with sensitive electronic detectors such as quartz tuning forks, as well as a fundamental understanding of modern automated data acquisition and processing.
Literature:	<p>[1] L.D. Landau, E.M. Lifshitz, Fluid Mechanics, Pergamon Books, 1987</p> <p>[2] L. Skrbek a kol., Fyzika nízkých teplot, Matfyzpress, 2011</p> <p>[3] D.R. Tilley, J. Tilley, Superfluidity and Superconductivity, Adam Hilger, 1986</p> <p>[4] R.J. Donnelly: Quantized vortices in helium II, Cambridge University Press, 2005</p> <p>[5] E. Varga, S. Babuin, L. Skrbek, Second-sound studies of coflow and counterflow of superfluid He-4 in channels, Phys. Fluids 27(6), 065101 (2015)</p> <p>[6] Vinen, W. F. Mutual friction in a heat current in liquid helium II. II. Experiments on transient effects. Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences. 240, 1220 (1957).</p> <p>[7] R. Blaauwgeers, M. Blažková, M. Človečko, V.B. Eltsov, R. de Graaf, J. Hosio, M. Krusius, D. Schmoranzer, W. Schoepe, L. Skrbek, P. Skyba, R.E. Solntsev, D.E. Zmeev: Quartz Tuning Fork: Thermometer, Pressure- and Viscometer for Helium Liquids, J. Low Temp. Phys. 146, 537-562 (2007)</p> <p>[8] S.L. Ahlstrom, D.I. Bradley, M. Človečko, S.N. Fisher, A.M. Guénault, E.A. Guise, R.P. Haley, O. Kolosov, P.V.E. McClintock, G.R. Pickett, M. Poole, V. Tsepelin, and A.J. Woods, Phys. Rev. B 89, 014515 (2014).</p>
Annotation:	The main focus of this Thesis will be study of oscillatory flows of superfluid helium at low temperatures. Specifically, we will characterize the transition to



Univerzita Komenského v Bratislave
Fakulta matematiky, fyziky a informatiky

ZADANIE ZÁVEREČNEJ PRÁCE

- Meno a priezvisko študenta:** Jakub Bahyl
Študijný program: fyzika (Jednoodborové štúdium, bakalársky I. st., denná forma)
Študijný odbor: 4.1.1. fyzika
Typ záverečnej práce: bakalárska
Jazyk záverečnej práce: slovenský
Sekundárny jazyk: anglický
- Názov:** Meranie intenzity kvantovej turbulencie v supratekutom 4He pomocou tlmenia druhého zvuku
Measurement of Quantum Turbulence in Superfluid 4He Using Second Sound Attenuation
- Cieľ:** V rámci tejto práce sa študent po teoretickej stránke zoznámí so základmi popisu kvantových kvapalín, ich supratekutosti a spôsobmi, akými sa dajú charakterizovať supratekuté prúdenia, ktoré sa veľakrát odlišujú od správania sa bežných kvapalín. S ohľadom na zvolenú problematiku budú predmetom štúdia aj vybrané kapitoly klasickej dynamiky tekutín. Prakticky sa študent naučí vykonávať a obsluhovať komplexný kryogénny experiment počínajúc prácou s kryokvapalinami cez nastavovanie špecifických opatrení pre experimenty s kvapalným héliom pri teplotách 1,2 až 4,2 K až po počítačové riadenie experimentu a jeho vyhodnotenie.
- Literatúra:** [1] L.D. Landau, E.M. Lifshitz, Fluid Mechanics, Pergamon Books, 1987
[2] L. Skrbek a kol., Fyzika nízkých teplot, Matfyzpress, 2011
[3] D.R. Tilley, J. Tilley, Superfluidity and Superconductivity, Adam Hilger, 1986
[4] R.J. Donnelly: Quantized vortices in helium II, Cambridge University Press, 2005
[5] E. Varga, S. Babuin, L. Skrbek, Second-sound studies of coflow and counterflow of superfluid He-4 in channels, Phys. Fluids 27(6), 065101 (2015)
[6] Vinen, W. F. Mutual friction in a heat current in liquid helium II. II. Experiments on transient effects. Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences. 240, 1220 (1957).
- Anotácia:** Náplňou tejto bakalárskej práce bude štúdium prúdenia supratekutého hélia pri nízkych teplotách. Konkrétne sa budeme zaujímať o meranie hustoty kvantových vírov v supratekutom héliu metódou tlmenia druhého zvuku. Budú študované rôzne druhy prúdenia supratekutého hélia vrátane tzv. tepelného protiprúdu, v ktorom sa študent zameria najmä na vznik turbulentného prúdenia v prúdovom kanáli. Jedným z hlavných cieľov bude súbežným meraním na viacerých senzoroch druhého zvuku zistiť, či kvantové víry vznikajú v celom objeme tekutiny súčasne, alebo či sa šíri postupne z miesta prvých nestabilít prúdení. Táto práca bude teda priamo nadväzovať na výskum tejto problematiky, ktorý sa robí v Laboratóriách supratekutosti na MFF UK v Prahe [5].

Acknowledgment

!!TODO!!

BACHELOR:

My special thanks go to my supervisor, David Schmoranzer, who gave me a lot of experiences, knowledge and much more of his time, without which I definitely could not write this Thesis. Together with consultant Martin Jackson, they guided me during the whole experiment and helped me to understand all the important aspects included in this work. Moreover, I am thankful to Ladislav Skrbek and Patrik Švančara for giving me such a great opportunity to find out how astounding the experimental physics can be and to work with Superfluidity group within Charles University in Prague.

In the end, I would like to hugely thank my friends and family for all their „*Good luck*” and „*Keep going*” motivational quotes and other sort of supports, which I deeply appreciated.

DIPLOMA:

Thanks.

I declare that I carried out this master thesis independently, and only with the cited sources, literature and other professional sources.

In date

Signature of the author:

!!TODO!!

Názov práce

Kvantová turbulencia v supratekutom héliu v teplotnej limite absolútnej nuly

Autor

Jakub Bahyl

Školiteľ bakalárskej práce

RNDr. David Schmoranzer, Ph.D.

Abstrakt

V tejto práci prezentujeme meranie kvantovej turbulencie generovanej oscilujúcim 6.5 kHz kremenným oscilátorom, ponoreným v supratekutom ^4He pri viacerých teplotách pod kritickou $T_\lambda = 2.17\text{ K}$. Pozorovaná nelineárna odporová sila pôsobiaca na oscilátor je kvalitatívne spôsobená prítomnosťou klasickej turbulencie a kvantovaných vírov. Odporové sily a množstvo vytvorených kvantovaných vírov (hustota vírov čiar L) sú nepriamo určené z útlmu druhého zvuku a mechanicko-elektrických vlastností oscilátora. Výsledky, ktoré prezentujeme, kvantitatívne charakterizujú tvorbu klasickej aj kvantovej turbulencie. Ako prví ukazujeme, že obidve turbulencie vedia vzniknúť samostatne a nezávisle. Tento fakt je prezentovaný, v prvom priblížení, pomocou *prúdového fázového diagramu*, v ktorom pre danú teplotu vieme predpokladať, kedy a ktorá turbulencia vzniká.

Kľúčové slová

supratekutosť ^4He • kvantový vír a turbulencia

Title

Quantum turbulence in superfluid helium down to the zero temperature limit

Author

Jakub Bahyl

Supervisor

RNDr. David Schmoranzer, Ph.D.

Abstract

In this work we present measurements of quantum turbulence generated by a 6.5 kHz oscillating quartz tuning fork submerged in superfluid ^4He at various temperatures below $T_\lambda = 2.17\text{ K}$. The observed non-linear drag acting on the tuning fork is qualitatively described by a presence of classical turbulence and quantized vortices. Drag forces and the amount of produced quantized vortices (the vortex line density L) are determined indirectly by the attenuation of second sound and by the measurement of mechanical and electrical properties of the tuning fork. We present results which characterize quantitatively the formation of classical and quantum turbulence. For the first time, we show that both turbulences can arise separately. This is presented, in first approximation, via a *flow phase diagram*, where for a given temperature, one can predict when each type of turbulence forms.

Keywords

superfluidity of ^4He • quantum vortex and turbulence

Contents

Introduction	3
1 Theoretical Background (15 pgs)	4
Microscopic model	5
1.1 Bose-Einstein condensate	5
1.2 Quantum vortex properties	5
Mesoscopic model	5
1.3 Filaments theory	5
1.4 Filaments dynamics	5
Macroscopic model	5
1.5 Two-fluid model	5
1.6 Oscillatory motion	5
1.7 Quantum turbulence	5
1.8 Quantum rings	5
2 Experimental Setup (20 pgs)	6
2.1 Apparatus	6
2.2 QT Generators	6
2.3 Measurements and Processing	6
2.4 Scaling, Second Sound	6
3 Simulations (15 pgs)	7
3.1 Codebase	7
3.2 Finite differences	7
3.3 Stepping	7
3.4 Resegmentation	7

3.5	Vortex ring	7
4	Results (10 pgs)	8

Introduction / Motivation (3 pgs)

- helium background, phase diagram, superfluid transition, interesting properties
- quantum turbulence, comparison with classical turbulence
- overview on experiments with QT
- overview on numerical simulations of QT
- motivations, goals of this work

1. Theoretical Background (15 pgs)

The theoretical part of this Thesis is composed of three chapters:

1. Microscopic view - wave function, collapse, quantum vortices
2. Mesoscopic view - filaments, drag and magnus force, dynamics of vortex, Schwarz equation, Kelvin waves (?)
3. Macroscopic view - hydrodynamics, two-fluid model,

Microscopic model

Bose-Einstein condensate

Quantum vortex properties

Mesoscopic model

Filaments theory

Filaments dynamics

Macroscopic model

Two-fluid model

Oscillatory motion

Quantum turbulence

Quantum rings

2. Experimental Setup (20 pgs)

Apparatus

QT Generators

Quartz Tuning Fork

Measurements and Processing

Scaling, Second Sound

3. Simulations (15 pgs)

Codebase

Finite differences

Stepping

Resegmentation

Vortex ring

4. Results (10 pgs)

Experiments

Simulations

Bibliography

- [1] KAPITSA, P. *The Super-Fluidity of Liquid Helium II*. Nature **141**, 74 (1938)
- [2] TONG, D. *Statistical physics*. University of Cambridge Part II
- [3] TISZA, L. *The viscosity of liquid helium and the Bose-Einstein statistics*. Comptes Rendus Acad. Sciences, **207**:1186-1189 (1952)
- [4] LANDAU, L.D. and LIFSHITZ, E.M. *Fluid Mechanics*, Second English Edition. Pergamon Books Ltd., (1987). ISBN 0-08-033933-6
- [5] LANDAU, L.D. *The theory of superfluidity of helium II*. J. Phys. USSR, Vol. **11**, 91 (1947)
- [6] ANDRONIKASHVILI, E.L. J. Phys. USSR, **10**, 201 (1946)
- [7] SKRBK, L. et al. *Fyzika nízkých teplot*, 1. vydání. Praha: MatfyzPress, (2011). ISBN 978-80-7378-168-2.