MHF4U Advanced Functions - Unit 1 Practice Test

Version 3 | Name: _____ Date: _____ 75 minutes | 50 marks

Show calculations in the lined space and use the graph grids. A calculator may be used; formal differentiation rules are not required.

1. Piecewise functions [9 marks]

The function S is defined by

```
            2x + 4      if -3 <= x < 0
S(x) =      4 - x^2     if 0 <= x <= 2
            x + 1       if 2 < x <= 4
```

a. Sketch S on one set of axes. Label all open and closed endpoints. [K2 C2]

b. State its domain and range in interval notation. [K2]

c. State the intervals where S is increasing. [K1]

d. Would the inverse relation be a function? Explain using your graph. [T1 C1]

Q1 - piecewise graph (choose and label a suitable scale)



Q1 - domain, range, intervals and inverse reasoning

2. Absolute value graph [6 marks]

Sketch `g(x) = 3|x - 2| - 6`. Label the vertex and both x-intercepts. [K2 C1] State the domain and range in interval notation. [K2] Explain how the coefficient 3 changes the graph of `y = |x|`. [T1]

Q2 - absolute value graph (choose and label a suitable scale)



Q2 - domain, range and transformation

3. Transforming a function [5 marks]

The point `(9, 3)` lies on `y = f(x)`, where `f(x) = sqrt(x)`. Define `h(x) = -2f(4(x + 2)) + 5`.

a. Find the corresponding point on h, showing the input equation. [T2]

b. State the domain and range of h. [K2]

c. Describe the horizontal transformation. [C1]

Q3 - point mapping, domain and range

4. Inverse relations [5 marks]

Let `q(x) = (x - 3)^2 + 2`, initially with domain all real numbers.

a. State the domain and range of its inverse relation. [K2]

b. Explain why this inverse relation is not a function. [T1]

c. Restrict the domain of q so that its inverse becomes a function, then write that inverse function. [A2]

Q4 - inverse reasoning

5. Observation wheel diagram [7 marks]

An observation wheel has radius 5 m. Its lowest seat is 1 m above the ground. It makes exactly one rotation in 480 s at constant speed. A rider begins at the lowest point at $t = 0$.



radius = 5 m

centre height = ?

greatest height = ?

lowest seat: 1 m above ground

one rotation: 480 s

a. From the diagram, determine the centre height and greatest seat height. [K2]

b. State the domain and range of the rider's height function for this ride, in interval notation. [K2]

c. A student says the greatest height is 6 m because $1 + 5 = 6$. Explain the error. [T1]

d. Sketch a qualitative height-versus-time graph for one rotation. Label the height at $t = 0$, $t = 240$, and $t = 480$. [C2]

Q5 a-c - height calculations and explanation

Q5 d - height versus time (label t in seconds and h in metres)



6. Average and instantaneous rates [8 marks]

During its first 6 seconds, a ball's height above the ground is modelled by `H(t) = -3t^2 + 18t + 2` metres.

a. Calculate the average rate of change of height over `[0, 2]`. [A2]

b. Estimate the instantaneous rate at `t = 2` using the centred interval `[1.9, 2.1]`. Show the two heights used. [A2 T1]

c. At what time does the ball reach its maximum modelled height, and what is that height? [K2]

d. Explain why your answers in a and b differ. [C1]

Q6 - calculations and interpretation

7. Graph properties and interval notation [5 marks]

Let `r(x) = x^4 - 3x^2`.

a. Show algebraically whether `r` is even, odd, or neither. [T2]

b. If `(2, 4)` lies on its graph, use symmetry to name one more point. [K1]

c. Write the solution to `|x + 2| < 4` in interval notation. [K2]

Q7 - symmetry and interval notation

8. Secant and tangent diagram [5 marks]

The graph shows `y = (x - 2)^2 + 2`, a secant through `A(2,2)` and `B(4,6)`, and the tangent at A.



a. Which line represents the average rate of change on `[2,4]`? Calculate its slope. [K1 A1]

b. Which line represents the instantaneous rate at `x = 2`? State its slope. [K1 A1]

c. Explain why the two rates differ even though both lines touch the graph at A. [T1]

Q8 - secant and tangent explanations