

Automated classification of *Chandra* X-ray point sources using machine learning methods

Shivam Kumaran¹ ¹★, Samir Mandal,¹ Sudip Bhattacharyya² ² and Deepak Mishra³ ³

¹Department of Earth and Space Sciences, Indian Institute of Space Science and Technology, Thiruvananthapuram 695547, India

²Department of Astronomy and Astrophysics, Tata Institute of Fundamental Research, Mumbai 400005, Maharashtra, India

³Department of Avionics, Indian Institute of Space Science and Technology, Thiruvananthapuram 695547, India

Accepted 2023 January 31. Received 2022 December 28; in original form 2022 September 14

ABSTRACT

A large number of unidentified sources found by astronomical surveys and other observations necessitate the use of an automated classification technique based on machine learning (ML) methods. The aim of this paper is to find a suitable automated classifier to identify the point X-ray sources in the *Chandra* Source Catalogue (CSC) 2.0 in the categories of active galactic nuclei (AGN), X-ray emitting stars, young stellar objects (YSOs), high-mass X-ray binaries (HMXBs), low-mass X-ray binaries (LMXBs), ultra luminous X-ray sources (ULXs), cataclysmic variables (CVs), and pulsars. The catalogue consists of $\approx 317\,000$ sources, out of which we select 277 069 point sources based on the quality flags available in CSC 2.0. In order to identify unknown sources of CSC 2.0, we use MW features, such as magnitudes in optical/ultraviolet bands from Gaia-EDR3, Sloan Digital Sky Survey, and GALEX, and magnitudes in infrared bands from 2MASS, WISE, and MIPS-Spitzer, in addition to X-ray features (flux and variability) from CSC 2.0. We find the Light Gradient Boosted Machine, an advanced decision tree-based ML classification algorithm, suitable for our purpose and achieve 93 per cent precision, 93 per cent recall score, and 0.91 Mathew's Correlation coefficient score. With the trained classifier, we identified 54 770 (14 066) sources with more than 3σ (4σ) confidence, out of which there are 32 600 (8574) AGNs, 16,148 (5,166) stars, 5,184 (208) YSOs, 439 (46) HMXBs, 197 (71) LMXBs, 50 (0) ULXs, 89 (1) CVs, and 63 (0) pulsars. This method can also be useful for identifying sources of other catalogues reliably.

Key words: methods: statistical – astronomical data bases: miscellaneous – catalogues – surveys – X-rays: general.

1 INTRODUCTION

Huge amounts of high-quality data from many astronomical sources are becoming available due to large-scale surveys and an open data access policy. Many of these sources are unidentified. These data have a great potential for discoveries of novel classes of sources, new sources of known classes, new observational phenomena, and even perhaps new physics. However, the sheer volume of data necessitates taking an automated approach to source classification. Such an automated classifier can be designed using machine learning (ML) methods (Ball & Brunner 2010). ML algorithms are capable of learning patterns in big data and can identify decision boundaries based on the already identified examples. Unlike the manual methods of identifying sources, for example, based on the colour–colour diagram clustering limited to three-dimension, ML methods can create decision boundaries in very high-dimensional feature space.

In optical/infrared (IR) astronomy, several works have been done for source identification using ML (Krakowski et al. 2016; Tang, Scaife & Leahy 2019; Tous, Solanes & Perea 2020; Ćiprijanović et al. 2021). However, in X-ray astronomy, the use of ML is relatively less. Farrell, Murphy & Lo (2015) did classification of variable sources in the third XMM-Newton Serendipitous Source Catalogue (3XMM) using a RF classifier with timing properties. Zhang, Zhao & Wu (2021) used RF and LogiBoost to classify the sources in XMM-

Newton's 4XMM-DR9 using multiwavelength (MW) properties from GAIA, WISE, and 2MASS. Classification of X-ray binaries based on whether the compact object is a black hole or a neutron star was done by De Beurs et al. (2022) using MAXI/GSC lightcurve. Falocco, Carrera & Larsson (2022) used RF and AdaBoost to develop an automated classifier for the identification of active galactic nuclei (AGN) and to classify them as Type-I or Type-II AGN further with the data from XMM-Newton and Sloan Digital Sky Survey (SDSS). Tranin et al. (2022) used MW data to classify sources in Swift-XRT and XMM-Newton serendipitous Source Catalogues using Naive Bayes classifier. Pattnaik et al. (2021) used the spectrum in the energy range of 5–25 keV from the Rossi X-ray Timing Explorer to identify the nature of the compact object in the Low Mass X-ray Binary objects using RF classifier.

In this paper, we aim to classify the point X-ray sources in the *Chandra* Source Catalogue (CSC) 2.0, which is the second source catalogue (Evans et al. 2020) of the CSC Observatory. A unique strength of *Chandra* is its angular resolution (≤ 1 arcsec), which is substantially better than the full-width-at-half-maximum (FWHM) ($6''$; Jansen et al. (2001)) of XMM-NEWTON and on-axis angular resolution [$5''$; Aschenbach, Bräuninger & Kettenring (1982)] of ROSAT. CSC 2.0 contains properties of about 317 000 sources from the observations till the end of 2014 and total sky coverage of 558.65 deg^2 . Most of these sources remain unidentified.

To the best of our knowledge, an approach adopted in this paper on automated classification of unidentified sources in CSC 2.0 has not been published yet. Due to the sub-arcsec spatial resolution and high

* E-mail: kumaranshivam57@gmail.com

Table 1. Quality flags used to filter sources in CSC 2.0 and their description.

Flag code	Description
<code>pileup_flag</code>	ACIS pile-up fraction exceeds $\sim 10\%$
<code>sat_src_flag</code>	Saturated source in all observations
<code>conf_flag</code>	Source confused (source and/or background regions in different stacks may overlap)
<code>streak_src_flag</code>	Source located on ACIS CCD read-out streak

sensitivity of Chandra, the source spatial population density in the CSC 2.0 is the highest among X-ray source catalogues. Thus, CSC 2.0 offers an excellent opportunity for the serendipitous discovery of objects of known classes and new exotic objects (Martinez Galarza et al. 2019).

In this paper, we discuss the development of an automated classifier based on supervised machine-learning algorithms for the point sources in CSC 2.0. The classifier primarily uses the features available in CSC 2.0, which are flux in five different bands of Chandra’s ACIS instrument and variability properties. In addition to the X-ray features, the source identification can be improved with the use of features available in other wavelengths. We obtain MW features from 2MASS, WISE, Gaia-EDR3, MIPS-Spitzer, SDSS, and GALEX. We explore decision tree based supervised ML classification algorithms. We find that the Light Gradient Boosted Machine (LightGBM) gives the best classification performance.

In the Section 2, we describe the details of the data, standardizing the data, and the method for identifying the training set. In Section 3, we present various classifier models which we explore and the methodology for selection and validation of the classifier. In Section 4, we present the result of the model validation and performance evolution. In Section 5, we give the summary and conclusions.

2 THE DATA

The objective of a ML classification model is to learn the relation between the features of a sample and its class label. In supervised ML methods, this relation is learnt using already labelled samples. For astrophysical objects, the features can be observed properties like magnitudes or flux in various wavelength bands. CSC 2.0 provides tabulated values of various observed properties. Apart from X-ray features, the source properties in other MW bands can be used to improve the classification further.

2.1 X-ray data

Chandra have two focal plane instruments: Advanced CCD Imaging Spectrometer (ACIS) and High Resolution Camera (HRC). The ACIS instrument observes in broad (b): 0.5–7.0 keV, ultrasoft (u): 0.2–0.5 keV, soft (s): 0.5–1.2 keV, medium (m): 1.2–2.0 keV, and hard (h): 2.0–7.0 keV bands. The HRC instrument observes in 0.1–10 keV energy band and is designated as ‘W’ band. CSC 2.0 was prepared with the observations from ACIS and HRC till the end of 2014. It contains the information of 317 167 sources, out of which 296 473 are point sources, selected by the parameter `extent_flag == 0` in the catalogue. We further filter the sources based on the quality flags available in CSC 2.0, which are: `pileup_flag`, `sat_src_flag`, `conf_flag`, `streak_src_flag` (Table 1).

For CSC 2.0, the energy flux in each band is determined using aperture photometry. The source count is derived from an elliptical source region and subtracted by the background count in the surrounding region. To convert the count rate to energy flux, the

total count rate is summed up and then scaled by the local ancillary response function. In this work, we use aperture-corrected average net-flux in b, u, s, m, and h bands, which are named as b-csc, u-csc, s-csc, m-csc, h-csc (Table 2), respectively.

The CSC also include various features indicating the short-term temporal variability (intra-observation) and long-term temporal variability (inter-observation) in each energy band. The intra-observation variability features are calculated using three methods:

- (i) Gregory–Loredo (GL) variability probability
- (ii) Kolmogorov–Smirnov (KS) test
- (iii) Kuiper’s test

The GL (Gregory & Loredo 1992) analysis provides the probability (*var_prob*) that the count rate in the source region varies over a flat distribution for each observation. It also calculates the odds ratio, time-integrated average count rate, and standard deviation (σ). The inverse of the odds ratio represents the significance of the observed distribution. GL algorithm defines a variability index *var_index* (an integer between 0 and 10)¹ using *var_prob*, odds ratio and the fraction of light curve within 3σ and 5σ . Any source having *var_index* > 6 is considered a variable. The highest *var_prob* and *var_index* across the observations are recorded as *var_intra_prob* and *var_intra_index* in the master table for the given source.

KS test calculates the probability that the arrival times of the events within the source region are inconsistent with a constant source count rate. It compares the cumulative photon distribution in the source region with a constant count rate. The highest probability value across the observations is recorded as *ks_intra_prob* in the master table for the given source. Similarly, a variability index, *kp_intra_index*, is defined using Kuiper’s test. While the KS test is sensitive around the median value, Kuiper’s test is more effective towards the tail of a probability distribution.

The inter-observations variability probability (*var_inter_prob*) defines the variation of the photon flux between the contributing observations. The maximum log-likelihood functions of the observed photon flux distribution are calculated assuming a different flux for each observation and constant photon flux. The difference between these two maximum log-likelihood functions follows a χ^2 distribution. The *var_inter_prob* represents the cumulative probability of this χ^2 distribution. The inter-observation variability index, *var_inter_index* (an integer between 0 and 8), is defined based on *var_inter_prob* and the number of degrees of freedom. The variance of the individual observation fluxes is indicated by the feature *var_inter_sigma* parameter.

In this work, we use four intra-observation variability properties in b-band (*var_intra_prob_b*, *var_intra_index_b*, *ks_intra_prob_b*, *kp_intra_prob_b*) and three inter-observation variability properties in b-band (*var_inter_prob_b*, *var_inter_index_b* and *var_inter_sigma_b*).

We use Chandra Interactive Analysis of Observations (CIAO-4.14) (Fruscione et al. 2006) to download the data from CSC 2.0 using Astronomical Data Query Language (ADQL).²

2.2 Multiwavelength data

AllWISE catalogue (Cutri et al. 2021) is the all IR survey catalogue built by combining the data from Wide-field Infrared Survey Explorer

¹ See variability index criterion table: https://cxc.cfa.harvard.edu/csc/why/gregory_loredo.html#varindex

² <https://www.ivoa.net/documents/ADQL/20180112/PR-ADQL-2.1-20180112.html>

Table 2. MW features from various catalogues used in this work.

Feature source	Feature name	Feature description
CSC 2.0	gal_l2	Galactic longitude
	gal_b2	Galactic Latitude
	b-csc	Flux in ACIS broad (b) band (0.5–7.0 keV)
	u-csc	Flux in ACIS ultrasoft (u) band (0.2–0.5 keV)
	s-csc	Flux in ACIS soft (s) band (0.5–1.2 keV)
	m-csc	Flux in ACIS medium (m) band (1.2–2.0 keV)
	h-csc	Flux in ACIS hard (h) band (2.0–7.0 keV)
	var_inter_prob_b	Inter-observation variability probability in ACIS b band
	var_inter_sigma_b	Standard deviation in Inter-observation flux variability
	var_inter_index_b	Inter-observation variability index
	var_intra_prob_b	Intra-observation Gregory-Loredo variability probability in b band
	ks_intra_prob	Kolmogorov–Smirnov Intra-observation variability probability b-band
GAIA-EDR3	var_intra_prob_b	Intra-observation Kupier’s test variability probability in b band
	var_intra_index_b	Intra-observation variability index
	G	Gaia Green (G) pass-band magnitude
GALEX	Bp	Gaia Blue (G_BP) pass-band magnitude
	Rp	Gaia Red (G_RP) pass-band magnitude
	FUV	Magnitude in GALEX FUV band
SDSS	NUV	Magnitude in GALEX NUV band
	u-sdss	SDSS u band magnitude
	g-sdss	SDSS g band magnitude
WISE	r-sdss	SDSS r band magnitude
	i-sdss	SDSS i band magnitude
	z-sdss	SDSS z band magnitude
MIPS-Spitzer	W1	WISE W1(3.4 micron) band magnitude
	W2	WISE W2(4.6 micron) band magnitude
	W3	WISE W3 (12 micron) band magnitude
	W4	WISE W4 (22 micron) band magnitude
2MASS	24_microns_MIPS	Magnitude in 24 micron band of MIPS on Spitzer
	J	J-band (1.235 micron) band magnitude
	H	H-band (1.662 micron) band magnitude
	K_s	Ks-band (2.159 micron) band magnitude
Colour*	B-R	Magnitude in Gaia Bp—magnitude in Gaia Rp
	G-J	Magnitude in Gaia G band—magnitude in 2MASS J band
	G-W2	Magnitude in Gaia G band—magnitude in WISE W2 band
	Bp-H	Magnitude in Gaia G_BP band—magnitude in 2MASS H band
	Bp-W3	Magnitude in Gaia G_BP band—magnitude in WISE W3 band
	Rp-K	Magnitude in Gaia G_RP band—magnitude in 2MASS K band
	J-H	Magnitude in 2MASS H—magnitude in 2MASS H band
	J-W1	Magnitude in 2MASS J—magnitude in WISE W1 band
	W1-W2	Magnitude in WISE W1—magnitude in WISE W2 band

Note. *The ‘colour’ features are computed using available magnitude values.

(WISE) mission’s two all-sky survey projects: WISE cryogenic phase (Wright et al. 2010) and the post cryogenic NEOWISE survey (Mainzer et al. 2011). The AllWISE is an all-sky infra-red survey at the wavelength bands 3.4 μm , 4.6 μm , 12 μm , and 22 μm named W1, W2, W3, and W4 bands, respectively. The AllWISE catalogue contains 747 634 026 sources with limiting sensitivities $W1 < 17.1$, $W2 < 15.7$, $W3 < 11.5$, and $W4 < 7.7$ magnitude. In our work, we use W1, W2, W3, and W4 magnitude from the AllWISE catalogue.

The Gaia (Prusti et al. 2016) is an optical telescope launched and operated by European Space Agency. The Gaia Early Data Release-3 (Gaia-EDR3) (Forveille & Kotak 2021) contains 1,811,709,771 sources and gives magnitudes in three broadband optical passbands, green (G), blue (G_{BP}) and red (G_{RP}) passbands. In this work, we use Gaia-EDR3 G, G_{BP} and G_{RP} band magnitudes. We find the association with Gaia using CDS X-match positional cross-match service (Boch, Pineau & Derriere 2014) such that the source must be within 3σ positional error of CSC 2.0 and Gaia EDR-3 error circle.

The 2 Micron All Sky Survey (2MASS) (Skrutskie et al. 2006) is the survey of the entire celestial sphere with a 99.99 per cent sky

coverage in the infra-red domain at 1.25 μm (J), 1.65 μm (H), and 2.16 μm (K_s) bands. The survey data were taken by two identical telescopes of diameter 1.3 m at Arizona and Chile in the northern and southern hemispheres, respectively. The Survey catalogue contains 470,992,970 point sources.

The Multiband Imaging Photometer (MIPS) onboard *Spitzer* (Rieke et al. 2004; Capak et al. 2013) covers the IR spectrum in the wavebands of 24 μm , 70 μm , and 160 μm . In this work, we use the 24 μm band (bandwidth $\sim 5 \mu\text{m}$) data due to its highest photometric accuracy of 6".

The telescope *Galaxy Evolution Explorer* (GALEX) takes observation in the far ultraviolet (FUV: 1344–1786 Å) and near ultraviolet (NUV: 1771–2831 Å) wavelengths with a resolution of $\sim 4.5''$ (FWHM) and $\sim 6.0''$ (FWHM), respectively (Morrissey et al. 2005).

The SDSS (York et al. 2000) is an extensive photometric and spectroscopic survey with an astrometric accuracy of the order of 0.1 arcsec. The SDSS obtains the data in five optical bands: u, g, r, i, and z with the central wavelengths of 3560 Å, 4680 Å, 6180 Å, 7500 Å, and 8870 Å, respectively. The limiting magnitudes of u, g, r, i, and

Table 3. Published catalogues used to identify sources in various classes.

Class	Catalogue source	Catalogue details	Reference
AGN	VERONCAT	Veron Catalogue of Quasars & AGN, 13th Edition	(Véron-Cetty & Véron 2010)
STAR	SKIFF	Catalogue of Stellar Spectral Classifications	Skiff (2013)
YSO		The Spitzer Space Telescope Survey...	Megeath et al. (2012)
		The Spitzer/IRAC Candidate YSO Catalogue...	Kuhn et al. (2021)
HMXB	HEASARC	SMCPSCXMM	Sturm et al. (2013)
		High-Mass X-Ray Binaries Catalogue	Liu, Van Paradijs & Van Den Heuvel (2006)
		INTEGRAL Reference Catalogue	Ebisawa et al. (2003)
		Magellanic Clouds High-Mass X-Ray Binaries Catalogue	Liu, van Paradijs & van den Heuvel (2005)
		IBIS/ISGRI Soft Gamma-Ray Survey Catalogue	Bird et al. (2016)
LMXB	HEASARC	INTEGRAL/ISGRI Catalogue of Variable X-Ray Sources	Tezhinsky et al. (2010)
		NGC 3115 Chandra X-Ray Point Source Catalogue	Lin et al. (2015)
		Ritter Low-Mass X-Ray Binaries Catalogue	Ritter & Kolb (2003)
		Low-Mass X-Ray Binaries Catalogue	Liu, van Paradijs & van den Heuvel (2007)
		INTEGRAL Reference Catalogue	Ebisawa et al. (2003)
		XMM-Newton M 31 Survey Catalogue	Pietsch, Freyberg & Haberl (2005)
		M 31... Point Source Catalogue	Hofmann et al. (2013)
		ROSAT All-Sky Survey	Haakonsen & Rutledge (2009)
		INTEGRAL IBIS Hard X-Ray Survey	Krivonos et al. (2015)
		INTEGRAL IBIS 9-Year Galactic Hard X-Ray Survey Catalog	Krivonos et al. (2012)
		IBIS/ISGRI Soft Gamma-Ray Survey Catalogue	Bird et al. (2016)
		M 31 XMM-Newton... X-Ray Point Source Catalogue	Shaw Greening et al. (2009)
ULX	ULXRBCAT		Liu & Mirabel (2005)
CV	The Open CV Cat.	The Open Cataclysmic Variable Catalogue	Jackim et al. (2020)
PULSAR	ATNF		Manchester et al. (2005)
	FERMI LAT (4FGL)	Fermi LAT Second Catalogue of Gamma-Ray Pulsars (2PC)	Abdo et al. (2013)

z are 21.6, 22.2, 22.2, 21.3, and 20.7, respectively. This work uses the 16th data release of SDSS (SDSS-DR16; Ahumada et al. 2020). This release includes the data from the previous release combined with the Apache Point Observatory Galactic Evolution Experiment 2 (APOGEE-2) survey and from the Extended Baryon Oscillation Spectroscopic Survey (eBOSS).

We use NASA/IPAC Extragalactic Database (NED) to obtain MW information for the CSC sources. With the November 2021 release, NED integrated the CSC. With the help of the cross-match algorithm Match Expert (MatchEx; Ogle et al. 2015), 80 per cent of the CSC sources were found to have associations with already existing objects, and 20 per cent became new objects in the NED database. We use CSC 2.0 names as the object identifier in NED to obtain the MW property with identifier-based query using `astroquery` package. We obtain MW data for 277 069 sources from NED. However, the NED server responded with error messages and the data could not be retrieved for 648 sources. Out of 277 069 objects in CSC 2.0, we could find an association for 60 per cent sources in Gaia-EDR3, 55 per cent in 2MASS, 43 per cent for MIPS-Spitzer, 41 per cent for WISE, 24 per cent for SDSS, and 17 per cent for GALEX.

The MW features used from these catalogues are given in Table 2. Besides these features, we compute the colours from the magnitude available in different bands and use them as additional features. We use an online MW visualization tool developed by Yang et al. (2021) to identify the colours that show the best class-wise clustering in a colour-colour diagram.

2.3 The training set

We aim to classify the point sources in CSC 2.0 into the following classes: AGN, X-ray emitting stars (STAR), Young Stellar Objects (YSO), High Mass X-ray binaries (HMXB), Low Mass X-ray binaries (LMXB), Ultra Luminous X-ray sources (ULX), Cataclysmic variables (CVs) and pulsars. First, we prepare a list of already

Table 4. Number of sources under various classes in the training set.

Class	% of training set	Number of sources
AGN	31 %	2395
STAR	36 %	2790
YSO	15 %	1149
HMXB	10 %	748
LMXB	2 %	143
ULX	3 %	211
CV	2 %	166
PULSAR	1 %	101
Total training set		7703
Unidentified Sources		269 366
Total		277069

identified sources belonging to these classes. Various published catalogues that we use to identify known sources are given in Table 3.

We cross-match the coordinates of the known sources with all the 277 069 sources in our list. We use `ASTROPY`³, which is a PYTHON package to perform cross-matching. We select a cross-match radius of 1". In case there is more than one source in the cross-match radius, we consider the source with the least angular separation from the target source. Using this, we identify a total of 7703 sources, of which there are 2395 AGNs, 2790 stars, 1149 YSOs, 748 HMXBs, 143 LMXBs, 211 ULXs, 166 CVs, and 101 pulsars. The class-wise percentages of identified sources are given in Table 4. These sources are used to train the supervised machine learning algorithm. In our training set, we have a large fraction of

³This work made use of Astropy: <http://www.astropy.org>, a community-developed core Python package and an ecosystem of tools and resources for astronomy

AGNs, stars, and YSOs, which comprise a total of about 80 per cent of the entire training set. The classes LMXB, ULX, CV, and pulsar are minorities with populations of only 1–3 per cent of the training set.

We create a data-table with 41 MW features (Table 2) from CSC-2.0, GAIA-EDR3, 2MASS, SDSS, WISE, GALEX, and MIPS-Spitzer for 277 069 point sources in CSC 2.0. Out of this, we keep a separate data table of 7703 known sources as the training data set. We attempt to identify the rest (269 366) of the sources.

3 METHODOLOGY

We prepare the MW data for a set of already identified sources. We use this set to train a supervised ML model. The model learns the pattern of the features in the training set and identifies the best possible decision boundary in the feature space. For designing the ML classification model, we use the python package *Scikit-Learn* (Pedregosa et al. 2011). In *Scikit-Learn*, various ML models are defined as Python classes with several options to customize the model. These models also implement general routine functions like ‘fit’ to train the model and use the trained model to ‘predict’ the class of a new sample. From *Scikit-Learn*, we test Multi-Layer Perceptron, K-Nearest Neighbour, RF, and Gradient Boosted Decision Trees (GBDTs). We find that decision tree-based models – RF and GBDT – perform better than other models. We also explore the LightGBM (Ke et al. 2017), which is an advanced development over GBDT, in this work.

3.1 Classifier models

3.1.1 Random forest

RF (Breiman 2001) is an ensemble of decision trees. Each decision tree is built from a randomly selected bootstrapped sample from the training set. Each tree, thus built is unique in nature and acts as a parallel weak learner. For a given source, each tree votes for it belonging to one of the eight classes. The fraction of trees out of the entire ensemble voting for a particular class is treated as the class membership probability (CMP) of the given source. For example, if 8 out of 10 trees vote for a particular sample to belong to class AGN, then the sample is said to be AGN with a membership probability of 0.8.

3.1.2 Gradient boosted decision tree

GBDT is an ensemble of weak learners (Friedman 2001). Compared to a decision tree, where each tree is built independently, in GBDT the trees are built sequentially based on the error of the previous tree. For each newly constructed tree, a loss is calculated based on the error between the predicted and the true values. In the case of the classification algorithm, categorical cross-entropy is used as the loss function and is defined as:

$$Loss = - \sum y_i \times \log \hat{y}_i \quad (1)$$

where y_i is a vector of length 8 (number of classes) with 1 for the true value and 0 otherwise. Here, \hat{y}_i is also a vector of length 8 representing the probability of the object belonging to each class. The gradient of this loss function at $(m - 1)^{th}$ tree is used to construct a new tree. It is then combined with the previous trees after multiplying it with a weight factor called learning rate η , which varies from 0 to 1. Essentially, each new tree is built to minimize the error from the previous tree. The main advantage of GBDT over RF is that

each newly constructed tree uses the loss from the previous tree and thus tries hard to better classify the previously incorrectly classified sources. The GBDT model can also learn more complex decision boundaries than the RF.

3.1.3 Light gradient boosted machine

LightGBM was developed by Ke et al. (2017). LightGBM is an advanced and efficient version of Gradient Boosted algorithms. Compared to GBDT, where each feature value is compared at the decision nodes of a tree, LightGBM first discretizes the value of the input features and then uses these values to construct the decision trees. To make learning more efficient, LightGBM implements two novel techniques, namely, Gradient-based One-Side Sampling (GOSS) and Exclusive Feature Bundling (EFB). With GOSS, LightGBM downsamples the low-gradient examples and upsamples high-gradient examples, which are more difficult to learn. Using EFB, LightGBM bundles the mutually exclusive features (the features that rarely take zero simultaneously) to reduce the dimension of the feature space. Another major capability of LightGBM is that it can handle missing values. It is very useful as we have a large number of missing values in our data set. It uses the Block Propagation method (Josse et al. 2019). In this method for splitting the nodes of the tree, only the available features are used. Wherever a missing value in a sample is found, it is sent to the side that would minimize the final loss.

3.2 Data normalization

The feature value in the data table varies in order of magnitude. Any ML model, in this scenario, would artificially tend to give more importance to the feature with higher magnitude. Thus it is a general practice to normalize the data set such that the magnitude variation across features remains uniform before feeding the data to the model. In our case, we normalize the data in such a way that the values lie between 0 and 10 using the following equation,

$$X_{norm} = 10 \times \frac{X - \min(X)}{\max(X) - \min(X)}, \quad (2)$$

where X_{norm} is normalized values of the feature X , $\min(X)$ and $\max(X)$ is the minimum and the maximum value of the feature X .

3.3 Missing value imputation

In our data table, we compile features from different MW catalogues. Due to the difference in coverage of these catalogues and the differences in limiting sensitivity, objects may not be available in all the catalogues. For example, the SDSS survey coverage is limited to the northern hemisphere. Similarly, we may have missing values in the data table due to differences in the intrinsic luminosity of the source across different wavelengths.

For example, X-ray binaries in the quiescent stage have lower luminosity in optical-UV and IR but are prominent in X-rays. In the X-ray domain, based on the variability timescales of the objects, the variability features are not available for some of the objects. Fig. 1 shows the fraction of sources for which the given set of features are available. In the figure, different features from the same instruments are grouped together. For *Chandra*, the fluxes in different bands are mentioned separately. The four intra-observation and three inter-observation variability properties are grouped together. We can see that 2MASS, MIPS, and WISE are




Figure 1. Plot showing the percentage of availability for different features group (see Section 3.3 for details).

mostly available only for AGNs, stars, and YSOs. The availability of X-ray variability features is significantly higher for X-ray binaries.

Most ML classification models need an input of a fixed size and are incompatible with missing values. One of the methods to avoid missing values is to remove the instances with missing features. We have hardly any source with 100 per cent features’ availability. For RF and GBDT models, we must have a method to handle these missing values. Imputation refers to the method of filling in these missing values with a suitable guess. Imputation can be done using statistics on the available data. In a tabular data format where rows represent the sources and columns represent the features, the missing values can be filled using the mean/mode/median of the feature column. For RF and GBDT models, we select to impute the missing values using column mode. However, the imputation method may not give satisfactory results for a high percentage of missing values, and in some cases, it is counter-productive to the final output. The RF and GBDT require missing value imputation and, therefore, may not perform well, particularly for minority classes where the percentage of missing values is high. Moreover, in some cases, the missing values themselves may be important features. For example, X-ray binaries (in the quiescent stage) are less likely to be observed in optical wavelengths. Therefore, we have tried to avoid imputation altogether using the model LightGBM as our final classifier. This model is capable of handling the missing values in the data table and also provides better results in every aspect compared to RF and GBDT.

3.4 Class imbalance problem

Table 4 shows that the numbers of AGNs, stars, and YSOs are typically an order of magnitude higher than those of LMXBs,

ULXs, CVs, and pulsars. Therefore, it is obvious that there is a vast imbalance in the number of training sources, with the majority classes being AGN, star, YSO and the minority classes being LMXB, ULX, CV, and pulsar. Any classifier model can achieve higher accuracy by biasing itself towards the majority class and thus would fail to perform on the new data.

To tackle the class imbalance problem, we use Synthetic Minority Oversampling Technique (SMOTE; Chawla et al. (2011)). In the feature space, it performs linear interpolations between k -neighbouring points (which represent a source in the feature space) and synthetic sources are sampled. Using this technique, each class is sampled such that the number of sources in the minority class becomes equal to the same in the most populous class. To keep our result insensitive to the oversampling, we perform SMOTE only on the training set and not on the validation set. SMOTE is used only for the RF and GBDT models. In LightGBM, we are working with missing data, and SMOTE cannot be performed with missing values. In LightGBM, we use the class-weight technique, which assigns higher weightage to the samples belonging to the minority class in calculating the loss function [equation (1)] during training. Essentially in the loss function, we make sure that equal contribution comes from each class.

3.5 Strategy for model performance validation

We compare the performance of the classifiers using a custom version of k -fold cross validation, which we call cumulative k -fold cross validation (CCV).

The flowchart in the Fig. 3 shows the cumulative cross validation method. Here, we divide the training set into k -fold, and we train the classifier using $k - 1$ folds and keeping aside k^{th} fold as the validation set in each iteration. After training, we make a prediction on the k^{th} fold (represented by the dashed box in Fig. 3). The predictions of classes of these samples are then stored in a prediction table. In the next iteration, when a different set of samples are in the validation set, their predictions are stored in the prediction table. In this manner, after k iterations, in the prediction table, we have the predictions for each training source coming only from the iterations when the source was in the validation set. Finally, the elements in the matrix are calculated using this prediction table.

We use precision, recall and F1-score for comparing classifier performance. The precision score is the probability that the predicted class is actually the true class for the sample and is defined as:

$$precision_A = \frac{TP}{TP + FP}, \quad (3)$$

where $precision_A$ represents the precision score for class A , TP represents the true positive, i.e. the number predicted as class A actually belonging to class A . Here, FP measures the false positive, i.e. the number of samples for which prediction are class A , while they belong to some other classes.

In probabilistic terms, the recall score is the probability of identifying the samples truly belonging to that particular class. The recall score for class A is defined as:

$$recall_A = \frac{TP}{TP + FN}, \quad (4)$$

where FN represents the number of samples belonging to class A but predicted to be in a class other than A .

F1 score is the harmonic mean of the precision and the recall score:

$$F1_A = 2 \times \frac{precision_A \times recall_A}{precision_A + recall_A}. \quad (5)$$




Figure 2. Confusion matrix for RF classifier showing the comparison between no upsampling (top) and upsampling using SMOTE (bottom). The confusion matrix is normalized by the true number of sources available in each class. See Section 3.4 for details.

We also use Mathew’s correlation coefficient (MCC), first introduced by Matthews (1975), which is supposed to be a better representation of classification score, particularly for imbalanced data set (Boughorbel, Jarray & El-Anbari 2017). MCC is defined as

$$MCC = \frac{TP \times TN - FP \times FN}{\sqrt{(TP + FP)(TP + FN)(TN + FP)(TN + FN)}} \quad (6)$$

where TN is the number of true negatives. MCC score is comparable to the correlation between the output predictions and the true labels whose values vary between +1, 0, and -1. The perfect prediction means +1, 0 means the model is as good as random predictions, and -1 shows a complete disagreement between the predicted and the true labels.

We use a confusion matrix for summarizing the model performance. An element of the matrix shows the percentage of sources which truly belong to the class given on the y-axis being classified by the model to a class given on the x-axis. The diagonal elements essentially are the recall score for the individual classes. In the CCV, for each iteration, after the data set is split into training and test set, SMOTE is applied to the training set for the models: RF and GBDT.

Fig.2 shows the confusion matrix for 20-fold cumulative cross-validation without upsampling (top) and with SMOTE upsampling (bottom) for the RF model. The numbers in the matrix are normalized by the number of sources belonging to the class shown on the vertical axis (true label). The diagonal elements in the figure represent the correct classification, and the rest are wrongly classified. We also notice from the top figure that most of the sources belonging to minority classes, mainly CV and pulsar, are getting classified as




Figure 3. Flowchart showing the cumulative cross validation algorithm. The components inside the dashed box represent one fold of validation. The algorithm is discussed in detail in Section 3.5.

AGN, STAR, and YSO if no upsampling is done. However, the correctly classified pulsars improve from 18 per cent to 45 per cent using SMOTE, and ~10% improvement happens for other minority classes. In general, the reduced contribution at the lower left corner of the bottom figure, as compared to the same in the top figure, shows the effectiveness of SMOTE in reducing the bias of the classifier towards the majority class.

Nevertheless, any class balancing method is not capable of reproducing an equal class distribution. The minority classes: LMXB, CV, ULX, and pulsars, together constitute less than 10 per cent of the total sample. In such a case, for SMOTE, where we are essentially interpolating source points in the feature space, the number of data points available may not capture the true distribution of the sources and may be affected by the boundary cases. For these minority classes, however, the upsampling shows significant improvement but does not take the score to a satisfactory level.

4 RESULT AND DISCUSSION

We train the classifier models: RF, GBDT, and LightGBM and evaluate the performance using the cumulative k -fold cross-validation. We implement the models using Scikit-Learn Python library. For

Table 5. Precision, recall, and F1 score for different classes for RF, GBDT, and LightGBM models. The scores are calculated by 20-fold cumulative cross-validation.

Score→ class↓ Model→	RF	Precision GBDT	LightGBM	RF	Recall GBDT	LightGBM	RF	1 score GBDT	LightGBM
AGN	96.7 ± 0.1	97.8 ± 0.2	96.8 ± 0.2	93.3 ± 0.2	89.6 ± 0.3	97.6 ± 0.2	95.0 ± 0.1	93.5 ± 0.2	97.2 ± 0.1
STAR	95.6 ± 0.1	97.1 ± 0.2	96.0 ± 0.2	94.1 ± 0.2	91.4 ± 0.2	95.7 ± 0.2	95.0 ± 0.1	94.1 ± 0.1	95.9 ± 0.1
YSO	91.6 ± 0.2	91.2 ± 0.3	92.7 ± 0.3	92.4 ± 0.3	93.7 ± 0.3	95.4 ± 0.3	92.0 ± 0.2	92.4 ± 0.2	94.1 ± 0.2
HMXB	79.2 ± 0.7	83.4 ± 0.8	91.6 ± 0.5	83.4 ± 0.5	87.4 ± 0.7	90.7 ± 0.6	81.2 ± 0.5	85.3 ± 0.5	91.2 ± 0.4
LMXB	84.8 ± 1.3	77.4 ± 2.8	94.8 ± 1.6	80.4 ± 1.1	80.8 ± 0.9	80.9 ± 1.6	82.5 ± 0.8	79.1 ± 1.5	87.2 ± 0.9
ULX	52.4 ± 1.0	47.4 ± 1.1	72.2 ± 1.4	63.5 ± 1.1	75.0 ± 1.1	71.1 ± 1.4	57.4 ± 0.9	57.9 ± 1.0	71.5 ± 1.2
CV	49.1 ± 1.1	42.4 ± 1.0	61.5 ± 1.6	53.8 ± 1.2	60.1 ± 1.6	55.3 ± 1.6	51.4 ± 0.3	49.7 ± 1.1	57.4 ± 1.5
PULSAR	35.8 ± 1.9	28.3 ± 1.2	42.1 ± 1.8	47.1 ± 2.6	55.8 ± 1.8	44.2 ± 1.8	40.7 ± 2.1	37.6 ± 1.4	43.7 ± 2.0

Table 6. MCC score and weighted average precision, recall, and F1 score for RF, GBDT, and LightGBM models.


Score	RF	Model GBDT	LightGBM
Precision	90.7 ± 0.1	91.3 ± 0.1	93.2 ± 0.1
Recall	90.0 ± 0.1	89.0 ± 0.2	93.2 ± 0.1
F1 score	90.3 ± 0.1	89.9 ± 0.2	93.2 ± 0.1
MCC	0.87 ± 0.00	0.86 ± 0.00	0.91 ± 0.00

the final selected models, we tune the hyperparameters using FLAML (A Fast Library for Automated Machine Learning & Tuning) by Wang et al. (2019). To compare the model performance, using the CCV method, we compute the precision, recall and F1 score for each class.

We perform several iterations of the 20-fold CCV, each time starting with a new random seed. These iterations are performed to compute the spread in the scoring metrics till the errors stabilize, which occurred after about 15 iterations. For each model, 15 iterations of CCV are performed, and the reported mean and the standard deviations are estimated over these iterations. The precision, recall, and F1 score for 20-fold cumulative cross validation are given in Table 5. The performance for majority classes (AGN, star, and YSO) with precision, recall, and F1 scores greater than 92 per cent are significantly higher than ULX, CV, and pulsar, for which scores are between 40 and 60 per cent. Performance is moderately good for LMXB and HMXB. We observe a similar trend across all three models. However, when models are compared, LightGBM performs the best for each class, whereas RF and GBDT have similarly lower performances.

The LightGBM model performs marginally better than RF for AGN, STAR, and YSO, but the scores are significantly higher for LMXB, HMXB, and ULX. For minority classes, there is a large difference between precision and recall for RF and GBDT models, whereas these two scores are comparable for LightGBM. For example, GBDT tries more aggressively to increase the prediction chances for the minority class and hence improves the recall score by about 7 per cent for ULX, CV, and pulsars, but this results in a drop in the precision score for the minority classes. Similarly, the RF model precision score for pulsars is 35 per cent, whereas the recall score is 47 per cent. It means that the model is trying hard to predict more pulsars to achieve a higher recall score, and therefore precision score for pulsars is reduced. This factor is well balanced for LightGBM, resulting highest F1 score across all classes.

Table 6 shows the MCC score and the weighted average of precision, recall, and F1 score for RF, GBDT, and LightGBM classifier, with the weighting factor being the proportion of sample

**Figure 4.** Cumulative histogram of class membership probability for RF, GBDT and LightGBM models, for the training set calculated during cumulative cross validation.

available in each class. The weighted average is taken to account for the class imbalance. LightGBM has the highest F1 score of 93.3 per cent, whereas the same for RF and GBDT are about 90 per cent. Once again, it shows that the LightGBM classifier performs the best. The MCC score for LightGBM is the highest (0.91), which means that the prediction made by LightGBM is more correlated with the true value than other models.

Apart from high precision, recall, and F1 score, a good classifier should be capable of predicting the class membership with high confidence. Fig. 4 shows the cumulative histogram of the class membership probabilities (CMP) assigned to the sources in the training set during CCV. The plot shows the histogram for all three models: RF, GBDT, and LightGBM. Any point on the plot shows the number of sources with CMP below the value given on the horizontal axis. The more sharply peaked the plot towards unity, the higher the predicted CMP and the more confident the model is. The figure reveals that the LightGBM is the most confident classifier model among the three, with only about 500 out of 7703 sources below CMP of 0.8, while RF is the least with more than 2000 sources below CMP of 0.8. Therefore, the cumulative cross validation of the models shows that LightGBM is the best classifier model of choice.

To study the importance of MW features, apart from the base sample, which includes all 41 features mentioned in Table 2, we create another sample keeping only the X-ray features obtained from CSC 2.0. We perform 15 iterations of the 20-fold CCV for the X-ray sample with the LightGBM model. Then we compare the model performance with the base sample score to see the class-wise

Table 7. Class-wise precision, recall, and F1 score for CCV using LightGBM model. The validation is done for various sets of ‘sample type’ indicated in the second column: ‘all features’ where all the 41 features in Table 2 are used and ‘X-ray’, the sample with no optical, UV and IR features.

Class	Sample type	Precision	Recall	F1 Score
AGN	All-features	96.8 ± 0.2	97.6 ± 0.2	97.2 ± 0.1
	X-ray	91.0 ± 0.2	95.0 ± 0.2	93.0 ± 0.1
STAR	All-features	96.0 ± 0.1	95.7 ± 0.1	95.8 ± 0.1
	X-ray	89.1 ± 0.3	88.2 ± 0.2	88.6 ± 0.2
YSO	All-features	92.7 ± 0.3	95.4 ± 0.3	94.0 ± 0.2
	X-ray only	82.9 ± 0.2	89.5 ± 0.5	86.1 ± 0.3
HMXB	All-features	91.6 ± 0.5	90.5 ± 0.5	91.1 ± 0.4
	X-ray	92.0 ± 0.5	89.9 ± 0.4	90.9 ± 0.4
LMXB	All-features	94.7 ± 1.6	80.9 ± 0.7	87.3 ± 0.9
	X-ray	95.0 ± 1.7	82.1 ± 0.5	88.0 ± 0.8
ULX	All-features	72.2 ± 1.4	71.1 ± 1.5	71.6 ± 1.2
	X-ray only	61.3 ± 2.0	42.7 ± 2.0	50.3 ± 1.9
CV	All-features	61.5 ± 1.6	55.3 ± 1.7	58.2 ± 1.5
	X-ray	56.0 ± 2.0	44.9 ± 1.8	49.8 ± 1.7
PULSAR	All-features	42.1 ± 1.8	44.2 ± 2.7	43.1 ± 2.0
	X-ray	28.2 ± 1.9	19.0 ± 1.4	22.7 ± 1.4

dependency on the MW features which are presented in Table 7 and Fig. 5.

Table 7 shows the mean and standard deviation of the precision, recall and F1 score over 15 times of the 20-fold cumulative cross validation. For each class, the table shows the scores for the two cases: one with X-ray features combined with MW features (all-features) and the other with only X-ray features. We can see a clear performance gain for all the classes when the MW features are used combined with X-ray features. The gain is most pronounced for pulsars with an enhancement of F1 score from 22 per cent to 43 per cent. Considering the recall score, 44 per cent of the pulsars can be retrieved using all features. However, only 19 per cent of the pulsars are classified properly with only X-ray features. For AGN, if all the MW features are simultaneously removed, the precision, recall and F1 score drop by 5 per cent, 2 per cent and 4 per cent, respectively. Even with only X-ray features, AGNs have a remarkably high F1 score of 93 per cent. Stars and YSOs, too, follow a similar trend but the dependency on MW features is higher than AGN, with typically 10 per cent drop in F1 score if MW features are removed. HMXB and LMXB are insensitive to optical/UV and IR features, and there is hardly any drop in performance without MW features. This can be attributed to the fact that X-ray binaries are, in general, faint in optical/UV/IR wavelengths.

For the ULXs, if MW features are removed, the F1 score drops about 20 per cent, and it is seen that a large percentage of ULXs are classified as AGNs (Fig. 5, bottom panel). It results in a slight drop in the F1 score for AGN, but it translates to a very high drop in the F1 score of ULX due to a very small population. For CVs, Optical/UV and IR features are important, and the F1 score drops by 10 per cent when only X-ray features are used.

Fig. 5 shows the confusion matrix for the two cases: one with all the features (top) and the other one with only X-ray features (bottom). Looking at the ULX row, with MW features, only 9 per cent (top figure) of ULXs are identified as AGNs, but the same increases to 41 per cent if only X-ray features are used. Using MW features, the classifier can better separate ULXs from AGNs. Similarly, we see pulsars are most likely to be identified as stars without MW features, and the incorrect prediction increases from 11 per cent to 41 per cent. From the confusion matrix, it is evident that the network becomes




Figure 5. Confusion matrix for the two sample sets: all-features (top) and only X-ray feature (bottom). The vertical axis represents the percentage of sources belonging to a class, and that being classified into various classes are shown on the horizontal axis.

more biased towards AGN, star and YSO without MW features and the fraction of ULX, CV, and pulsar being classified as AGN, star and YSO increases.

To find the impact of different MW bands, we did a similar analysis adding only optical/UV features with X-ray features, and then only IR features with X-ray features. For AGN and Stars, both optical/UV and IR features result in identical improvement in the F1 score, but for YSOs addition of IR features has a higher impact than optical/UV features. For ULX and pulsar, each additional MW band (Optical/UV or IR) provide 7–10 per cent improvement in the F1 scores.

Even with the MW features and class weight balancing, the model does not provide a satisfactory result for the minority classes: LMXB, ULX, CV, and pulsars. We attempt to explore to what extent the presence of minority classes affects the classification of majority classes. We train another model using only the four majority classes: AGN, YSO, Stars, and HMXB with and without MW features. Table 8 shows the scores for the model trained only on the majority classes. Comparing Table 7 and Table 8, the later model has shown a slightly improved performance for these classes. For the all-features case, the F1 scores of AGN, Star and YSO show ~ 1 per cent, and HMXB shows ~ 2 per cent improvement. Considering only the X-ray feature, AGN, Star, YSO, and HMXB have 3 per cent, 3 per cent, 1 per cent, and 2 per cent improved F1 scores, respectively. The inclusion of minority classes does not significantly affect the majority class scores if MW features and the X-ray features are used together. We keep the minority classes in our classification

Table 8. Same as Table 7, but the classifier is trained and validated only for the majority class.

Class	Sample type	Precision	Recall	F1 score
AGN	All-features	98.3 ± 0.1	98.4 ± 0.1	98.3 ± 0.1
	X-ray	95.5 ± 0.2	96.5 ± 0.2	96.0 ± 0.1
STAR	All-features	97.3 ± 0.1	96.3 ± 0.1	96.8 ± 0.1
	X-ray	92.7 ± 0.2	89.6 ± 0.2	91.1 ± 0.2
YSO	All-features	93.7 ± 0.2	96.5 ± 0.2	95.1 ± 0.2
	X-ray	84.4 ± 0.3	91.2 ± 0.4	87.7 ± 0.3
HMXB	All-features	93.9 ± 0.4	93.0 ± 0.3	93.4 ± 0.2
	X-ray	93.8 ± 0.4	90.7 ± 0.4	92.2 ± 0.3

Table 9. The number of sources identified in various classes with the LightGBM. The column name ‘all’ represents the sources classified based on maximum CPM of the class.

Class	All	Number of sources CMP>3 σ	CMP>4 σ
AGN	114 642	32 600	8574
STAR	63 967	16 148	5166
YSO	40 524	5184	208
HMXB	8321	439	46
LMXB	1688	197	71
ULX	6083	50	0
CV	10 999	89	1
PULSAR	23 142	63	0
Total	269 366	54 770	14 066

scheme; otherwise, these minority class objects will be classified as the majority class.

From the results discussed above, we select LightGBM as the final classifier with all the 41 features in Table 2 for training and classification of objects into eight classes.

The trained LightGBM model is applied to all the 269 366 unidentified sources. For each source, the trained model gives a probability of belonging to each of the eight classes. For a given source, the class having the highest probability is assigned to it, and the corresponding class probability is called the class membership

probability (CMP). Thus we get the class identification and the corresponding CMPs for each of the 269 366 sources.

To identify the class-wise confidence of the classifier, we dwell deeper into the corresponding predicted probabilities. After the classification is done, we compute the class-wise probability density function (PDF) of these CMP. For example, 114 642 objects out of 269 366 are identified as AGNs (Table 9). Using the CMP of these 114 642 objects, we calculate the PDF of AGN’s CMP. The probability of getting an object of class A with a CMP = x , lying between the values a and b is given by the area under the curve of the PDF and is given as

$$P(a < x < b) = \int_a^b f_A(x) dx, \quad (7)$$


where the function $f_A(x)$ is the PDF of the class A . It is calculated from the CMP histogram with N bins and counts (n_i) in i^{th} bin using the following equation

$$f_A(x_i) = \frac{n_i}{\sum_0^N n_i \times \Delta x}, \quad (8)$$

where Δx is the bin size. We plot the probability density function in Fig. 6 to present the class-wise distribution of the CMP. The peak of the PDF is marked by red dashed lines, which is the highest CMP value for a given class. The PDF curves show the overall classification confidence for each class. The PDFs are more sharply peaked close to 1.0 for the majority classes (e.g. AGN, STAR, YSO). It means the LightGBM model can predict these classes with high confidence.

In contrast, for some other classes (e.g. ULX, CV), the peak appears at lower CMP values, implying a good fraction of sources with lower classification confidence. We find double-peak profiles for some classes (e.g. LMXB). This implies that the nature of some sources of such a class is predicted with low confidence values, while that of other sources of the same class is predicted with high confidence values. The reason could be the availability or unavailability of one or more observational features for certain sources of that class. For example, we find that the inter-observation variability feature is not available for a larger fraction of low-confidence LMXBs.

To pick the sources which are identified with a very high membership probability, we set a higher threshold to select only those

**Figure 6.** PDF of the distribution of CMP of all the unidentified sources predicted using the LightGBM model for different classes (marked on the plots). The most probable value of the class membership probability is shown in the plot with a vertical dashed line.

sources above the CMP threshold. Table 9 shows the number of newly identified sources in various classes. The number of sources in every class based on the most probable values of CMP (red line in Fig. 6) is given in the second column (all). Similarly, the number of sources above the probability confidence threshold of 3σ (CMP >0.997) and 4σ (CMP >0.9999) are shown in the third and fourth column of Table 9 respectively. We identify 54 770 new sources in the existing classes with 3σ confidence and 14 066 new sources with 4σ confidence. This significantly increases the number of sources in various classes.

5 SUMMARY AND CONCLUSIONS

The Chandra Source Catalogue CSC 2.0 contains $\approx 317\,000$ sources, including $\approx 277\,000$ point sources, with a majority of them unidentified. In this work, we implement the decision tree based classifier Light Gradient Boosted Machine to identify the CSC 2.0 objects in the classes of AGN, Star, YSO, HMXB, LMXB, ULX, CV, and pulsar. For the classification, we use X-ray properties from *Chandra*, optical/UV properties from *Gaia*, *SDSS*, and *GALEX*, and IR properties from 2MASS, WISE, and MIPS-Spitzer. We train the classifier and applied the trained classifier to the unidentified sources to estimate the class membership probabilities of these sources. We achieve a classification weighted precision score of 93 per cent, recall score of 93 per cent, F1 score of 93 per cent, and Mathew's Correlation coefficient of 0.91.

We identify 54 770 new point sources out of which there are 32 600 AGNs, 16 148 stars, 5184 YSOs, 197 LMXBs, 439 HMXBs, 50 ULXs, 89 CVs and 63 pulsars with a confidence of more than 3σ . Even at a higher confidence (more than 4σ), we get 8574 AGNs, 5166 stars, 208 YSOs, 46 HMXBs, 71 LMXBs, and 1 CV but not ULXs and pulsars.

The identification of an unknown source as a member of a known class is equivalent to the discovery of a new source of that class. While the main aim of this paper is to find a suitable classifier and apply it to CSC 2.0 point sources, in a subsequent paper we will list those sources which could be assigned to various classes with high significance values and discuss their properties in detail. Finally, we believe that our method can be reliable and promising for other catalogues as well.

Very recently, we came across the work by Yang et al. (2022) which aims to develop a pipeline called MUWCLASS using a MW approach to classify sources in CSC 2.0. They applied the pipeline to a sample of 66 369 CSC 2.0 sources. The training data set, peer-reviewed catalogues, the choice of classifier, the validation methodology and the overall list of unidentified sources make our work unique.

ACKNOWLEDGEMENTS

This research has used the *Chandra* Data Archive and the *Chandra* Source Catalogue, and software provided by the *Chandra* X-ray Center (CXC) in the application packages CIAO and Sherpa; NASA/IPAC Extragalactic Database (NED), which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration; data from the European Space Agency (ESA) mission *Gaia* (<https://www.cosmos.esa.int/gaia>), processed by the *Gaia* Data Processing and Analysis Consortium (DPAC, <https://www.cosmos.esa.int/web/gaia/dpac/consortium>); the cross-match service provided by CDS, Strasbourg; Multiwavelength visualisation tool by Yang et al. (2021) to identify the colour-colour properties to be used.

This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, and NEOWISE, which is a project of the Jet Propulsion Laboratory/California Institute of Technology. WISE and NEOWISE are funded by the National Aeronautics and Space Administration.

DATA AVAILABILITY

All the data accumulated/generated in this work will be made available on a publicly accessible portal which is in the development phase. The portal will include the MW data for the 277 069 sources. The portal will also allow the user to perform a cone search based on the coordinates. The classification table and the class membership probabilities generated in this work will be integrated into the portal such that the user will be able to filter the search for their class of interest with their selected confidence threshold. The sources identified with very high confidence will be released as a machine-readable data table in a subsequent paper. The training data set with the corresponding reference will be shared as a CSV table at the reader's request.

REFERENCES

- Abdo A. A. et al., 2013, *ApJS*, 208, 17
- Ahumada R. et al., 2020, *ApJS*, 249, 3
- Aschenbach B., Bräuninger H., Kettenring G., 1982, *Adv. Space Res.*, 2, 251
- Ball N. M., Brunner R. J., 2010, *Int. J. Mod. Phys. D*, 19, 1049
- Bird A. J. et al., 2016, *ApJS*, 223, 15
- Boch T., Pineau F., Derriere S., Astronomical Society of the Pacific Conference Series 461 2012, CDS xMatch service documentation, <http://cdsxmatch.u-strasbg.fr/xmatch/doc/CDSXMatchDoc.pdf>
- Boughorbel S., Jarray F., El-Anbari M., 2017, *PloS One*, 12, e017767828574989
- Breiman L., 2001, *Machine Learning*, 45, 5
- Capak P. L., Teplitz H. I., Brooke T. Y., Laher R., Science Center S., 2013, in American Astronomical Society Meeting Abstracts #221. p. 340.06
- Chawla N. V., Bowyer K. W., Hall L. O., Kegelmeyer W. P., *Journal Of Artificial Intelligence Research* 16 321–357 2002,
- Ćiprijanović A. et al., 2021, *MNRAS*, 506, 677
- Cutri R. M. et al., 2014, AllWISE Data Release Centre de Données astronomique de Strasbourg (CDS) Vizier Online Data Catalog, p. II/328
- De Beurs Z. L., Islam N., Gopalan G., Vrilek S. D., The Astrophysical Journal 2022 933 116
- Ebisawa K., Bourban G., Bodaghee A., Mowlavi N., 2003, *Astron. Astrophys.*, 411, L59
- Evans I. N. et al., 2020, in American Astronomical Society Meeting Abstracts #235. p. 154.05
- Falocco S., Carrera F. J., Larsson J., 2022, *MNRAS*, 510, 161
- Farrell S. A., Murphy T., Lo K. K., 2015, *ApJ*, 813, 28
- Forveille T., Kotak R., 2021, *A&A*, 649, E1
- Friedman J. H., 2001, *Ann. Stat.*, pp 1189
- Fruscione A. et al., 2006, in Silva D. R., Duxsey R. E. eds, Society of Photo-Optical Instrumentation Engineers (SPIE) Conf. Ser. Vol. 6270, Society of Photo-Optical Instrumentation Engineers (SPIE) Conf. Ser. p. 62701V,
- Gregory P. C., Loredó T. J., 1992, *ApJ*, 398, 146
- Haakonsen C. B., Rutledge R. E., 2009, *ApJS*, 184, 138
- Hofmann F., Pietsch W., Henze M., Haberl F., Sturm R., Della Valle M., Hartmann D. H., Hatzidimitriou D., 2013, *A&A*, 555, A65
- Jackim R., Szkody P., Hazelton B., Benson N. C., 2020, *Res. Notes Am. Astron. Soc.*, 4, 219
- Jansen F. et al., 2001, *A&A*, 365, L1

- Josse J., Prost N., Scornet E., Varoquaux G., 2019, arXiv preprint [arXiv:1902.06931](https://arxiv.org/abs/1902.06931), p. 18
- Ke G., Meng Q., Finley T., Wang T., Chen W., Ma W., Ye Q., Liu T.-Y., 2017, *Adv. Neural Inf. Processing Syst.*, 30
- Krakowski T., Małek K., Bilicki M., Pollo A., Kurcz A., Krupa M., 2016, *A&A*, 596, A39
- Krivosos R., Tsygankov S., Lutovinov A., Revnivitsev M., Churazov E., Sunyaev R., 2012, *A&A*, 545, A27
- Krivosos R., Tsygankov S., Lutovinov A., Revnivitsev M., Churazov E., Sunyaev R., 2015, *MNRAS*, 448, 3766
- Kuhn M. A., de Souza R. S., Krone-Martins A., Castro-Ginard A., Ishida E. O., Povich M. S., Hillenbrand L. A., COIN Collaboration, 2021, *ApJS*, 254, 33
- Lin D. et al., 2015, *ApJ*, 808, 19
- Liu Q. Z., Mirabel I. F., 2005, *A&A*, 429, 1125
- Liu Q. Z., van Paradijs J., van den Heuvel E. P. J., 2005, *A&A*, 442, 1135
- Liu Q., Van Paradijs J., Van Den Heuvel E., 2006, *Astron. Astrophys.*, 455, 1165
- Liu Q. Z., van Paradijs J., van den Heuvel E. P. J., 2007, *A&A*, 469, 807
- Mainzer A. et al., 2011, *ApJ*, 731, 53
- Manchester R. N., Hobbs G. B., Teoh A., Hobbs M., 2005, *AJ*, 129, 1993
- Martinez Galarza J., D’Abrusco R., Civano F., Evans I., 2019, in *AAS/High Energy Astrophysics Division*. p. 109.29
- Matthews B., 1975, *Biochimica et Biophysica Acta (BBA)—Protein Structure*, 405, 442
- Megeath S. T. et al., 2012, *AJ*, 144, 192
- Morrissey P. et al., 2005, *ApJ*, 619, L7
- Ogle P. M., Mazzarella J., Ebert R., Fadda D., Lo T., Terek S., Schmitz M., NED Team, 2015, in *Taylor A. R., Rosolowsky E. eds, Astron. Soc. Pacific Conf. Ser. Vol. 495, Astronomical Data Analysis Software and Systems XXIV (ADASS XXIV)*. p. 25([arXiv:1503.01184](https://arxiv.org/abs/1503.01184))
- Pattanaik R., Sharma K., Alabarta K., Altamirano D., Chakraborty M., Kembhavi A., Méndez M., Orwat-Kapola J. K., 2021, *MNRAS*, 501, 3457
- Pedregosa F. et al., 2011, *J. Machine Learning Res.*, 12, 2825
- Pietsch W., Freyberg M., Haberl F., 2005, *A&A*, 434, 483
- Prusti T. et al., 2016, *A&A*, 595, A1
- Rieke G. H. et al., 2004, *ApJS*, 154, 25
- Ritter H., Kolb U., 2003, *A&A*, 404, 301
- Shaw Greening L., Barnard R., Kolb U., Tonkin C., Osborne J. P., 2009, *A&A*, 495, 733
- Skiff B. A., VizieR Online Data Catalog: Catalogue of Stellar Spectral Classifications (Skiff, 2009-2013) 2013, VizieR Online Data Catalog, p. B/mk [ADS](https://arxiv.org/abs/1305.0001)
- Skrutskie M. F. et al., 2006, *AJ*, 131, 1163
- Sturm R. et al., 2013, *Astron. Astrophys.*, 558, A3
- Tang H., Scaife A. M. M., Leahy J. P., 2019, *MNRAS*, 488, 3358
- Teleszhinsky I., Eckert D., Savchenko V., Neronov A., Produit N., Courvoisier T. J. L., 2010, *A&A*, 522, A68
- Tous J. L., Solanes J. M., Perea J. D., 2020, *MNRAS*, 495, 4135
- Tranin H., Godet O., Webb N., Primorac D., 2022, *A&A*, 657, A138
- Véron-Cetty M.-P., Véron P., 2010, *Astron. Astrophys.*, 518, A10
- Wang C., Wu Q., Weimer M., Zhu E., 2021, *Proceedings of Machine Learning and Systems* 3 434–447
- Wright E. L. et al., 2010, *AJ*, 140, 1868
- Yang H., Hare J., Volkov I., Kargaltsev O., 2021, *Res. Notes Am. Astron. Soc.*, 5, 102
- Yang H., Hare J., Kargaltsev O., Volkov I., Chen S., Rangelov B., 2022, *ApJ*, 941, 104
- York D. G. et al., 2000, *AJ*, 120, 1579
- Zhang Y., Zhao Y., Wu X.-B., 2021, *MNRAS*, 503, 5263

This paper has been typeset from a \LaTeX file prepared by the author.

List of astronomical key words (Updated on 2020 January)

This list is common to *Monthly Notices of the Royal Astronomical Society*, *Astronomy and Astrophysics*, and *The Astrophysical Journal*. In order to ease the search, the key words are subdivided into broad categories. No more than *six* subcategories altogether should be listed for a paper.

The subcategories in boldface containing the word ‘individual’ are intended for use with specific astronomical objects; these should never be used alone, but always in combination with the most common names for the astronomical objects in question. Note that each object counts as one subcategory within the allowed limit of six.

The parts of the key words in *italics* are for reference only and should be omitted when the keywords are entered on the manuscript.

General

editorials, notices
errata, addenda
extraterrestrial intelligence
history and philosophy of astronomy
miscellaneous
obituaries, biographies
publications, bibliography
sociology of astronomy
standards

Physical data and processes

acceleration of particles
accretion, accretion discs
asteroseismology
astrobiology
astrochemistry
astroparticle physics
atomic data
atomic processes
black hole physics
chaos
conduction
convection
dense matter
diffusion
dynamo
elementary particles
equation of state
gravitation
gravitational lensing: micro
gravitational lensing: strong
gravitational lensing: weak
gravitational waves
hydrodynamics
instabilities
line: formation
line: identification
line: profiles
magnetic fields
magnetic reconnection
(*magnetohydrodynamics*) MHD
masers
molecular data
molecular processes
neutrinos
nuclear reactions, nucleosynthesis, abundances
opacity
plasmas
polarization

radiation: dynamics
radiation mechanisms: general
radiation mechanisms: non-thermal
radiation mechanisms: thermal
radiative transfer
relativistic processes
scattering
shock waves
solid state: refractory
solid state: volatile
turbulence
waves

Astronomical instrumentation, methods and techniques

atmospheric effects
balloons
instrumentation: adaptive optics
instrumentation: detectors
instrumentation: high angular resolution
instrumentation: interferometers
instrumentation: miscellaneous
instrumentation: photometers
instrumentation: polarimeters
instrumentation: spectrographs
light pollution
methods: analytical
methods: data analysis
methods: laboratory: atomic
methods: laboratory: molecular
methods: laboratory: solid state
methods: miscellaneous
methods: numerical
methods: observational
methods: statistical
site testing
space vehicles
space vehicles: instruments
techniques: high angular resolution
techniques: image processing
techniques: imaging spectroscopy
techniques: interferometric
techniques: miscellaneous
techniques: photometric
techniques: polarimetric
techniques: radar astronomy
techniques: radial velocities
techniques: spectroscopic
telescopes

Astronomical data bases

astronomical data bases: miscellaneous
atlases
catalogues
surveys
virtual observatory tools

Software

software: data analysis
software: development
software: documentation
software: public release
software: simulations

Astrometry and celestial mechanics

astrometry
celestial mechanics
eclipses
ephemerides
occultations
parallaxes
proper motions
reference systems
time

The Sun

Sun: abundances
Sun: activity
Sun: atmosphere
Sun: chromosphere
Sun: corona
Sun: coronal mass ejections (CMEs)
Sun: evolution
Sun: faculae, plages
Sun: filaments, prominences
Sun: flares
Sun: fundamental parameters
Sun: general
Sun: granulation
Sun: helioseismology
Sun: heliosphere
Sun: infrared
Sun: interior
Sun: magnetic fields
Sun: oscillations
Sun: particle emission
Sun: photosphere
Sun: radio radiation
Sun: rotation
(*Sun*:) solar–terrestrial relations
(*Sun*:) solar wind
(*Sun*:) sunspots
Sun: transition region
Sun: UV radiation
Sun: X-rays, gamma-rays

Planetary systems

comets: general

comets: individual: . . .

Earth
interplanetary medium
Kuiper belt: general

Kuiper belt objects: individual: . . .

meteorites, meteors, meteoroids

minor planets, asteroids: general

minor planets, asteroids: individual: . . .

Moon

Oort Cloud

planets and satellites: atmospheres
planets and satellites: aurorae
planets and satellites: composition
planets and satellites: detection
planets and satellites: dynamical evolution and stability
planets and satellites: formation
planets and satellites: fundamental parameters
planets and satellites: gaseous planets
planets and satellites: general

planets and satellites: individual: . . .

planets and satellites: interiors
planets and satellites: magnetic fields
planets and satellites: oceans
planets and satellites: physical evolution
planets and satellites: rings
planets and satellites: surfaces
planets and satellites: tectonics
planets and satellites: terrestrial planets
planet–disc interactions
planet–star interactions
protoplanetary discs
zodiacal dust

Stars

stars: abundances
stars: activity
stars: AGB and post-AGB
stars: atmospheres
(*stars*:) binaries (*including multiple*): close
(*stars*:) binaries: eclipsing
(*stars*:) binaries: general
(*stars*:) binaries: spectroscopic
(*stars*:) binaries: symbiotic
(*stars*:) binaries: visual
stars: black holes
(*stars*:) blue stragglers
(*stars*:) brown dwarfs
stars: carbon
stars: chemically peculiar
stars: chromospheres
(*stars*:) circumstellar matter
stars: coronae
stars: distances
stars: dwarf novae
stars: early-type
stars: emission-line, Be
stars: evolution
stars: flare
stars: formation
stars: fundamental parameters
(*stars*:) gamma-ray burst: general
(*stars*:) **gamma-ray burst: individual: . . .**
stars: general
(*stars*:) Hertzsprung–Russell and colour–magnitude diagrams
stars: horizontal branch
stars: imaging
stars: individual: . . .
stars: interiors

- stars: jets
- stars: kinematics and dynamics
- stars: late-type
- stars: low-mass
- stars: luminosity function, mass function
- stars: magnetars
- stars: magnetic fields
- stars: massive
- stars: mass-loss
- stars: neutron
- (stars:) novae, cataclysmic variables
- stars: oscillations (*including pulsations*)
- stars: peculiar (*except chemically peculiar*)
- (stars:) planetary systems
- stars: Population II
- stars: Population III
- stars: pre-main-sequence
- stars: protostars
- (stars:) pulsars: general
- (stars:) **pulsars: individual: . . .**
- stars: rotation
- stars: solar-type
- (stars:) starspots
- stars: statistics
- (stars:) subdwarfs
- (stars:) supergiants
- (stars:) supernovae: general
- (stars:) **supernovae: individual: . . .**
- stars: variables: Cepheids
- stars: variables: Scuti
- stars: variables: general
- stars: variables: RR Lyrae
- stars: variables: S Doradus
- stars: variables: T Tauri, Herbig Ae/Be
- (stars:) white dwarfs
- stars: winds, outflows
- stars: Wolf–Rayet

Interstellar medium (ISM), nebulae

- ISM: abundances
- ISM: atoms
- ISM: bubbles
- ISM: clouds
- (ISM:) cosmic rays
- (ISM:) dust, extinction
- ISM: evolution
- ISM: general
- (ISM:) HII regions
- (ISM:) Herbig–Haro objects

ISM: individual objects: . . .

- (*except planetary nebulae*)
- ISM: jets and outflows
- ISM: kinematics and dynamics
- ISM: lines and bands
- ISM: magnetic fields
- ISM: molecules
- (ISM:) photodissociation region (PDR)
- (ISM:) planetary nebulae: general
- (ISM:) **planetary nebulae: individual: . . .**
- ISM: structure
- ISM: supernova remnants

The Galaxy

- Galaxy: abundances
- Galaxy: bulge
- Galaxy: centre
- Galaxy: disc
- Galaxy: evolution
- Galaxy: formation
- Galaxy: fundamental parameters
- Galaxy: general
- (Galaxy:) globular clusters: general
- (Galaxy:) **globular clusters: individual: . . .**
- Galaxy: halo
- Galaxy: kinematics and dynamics
- (Galaxy:) local interstellar matter
- Galaxy: nucleus
- (Galaxy:) open clusters and associations: general
- (Galaxy:) **open clusters and associations: individual: . . .**
- (Galaxy:) solar neighbourhood
- Galaxy: stellar content
- Galaxy: structure

Galaxies

- galaxies: abundances
- galaxies: active
- galaxies: bar
- (galaxies:) BL Lacertae objects: general
- (galaxies:) **BL Lacertae objects: individual: . . .**
- galaxies: bulges
- galaxies: clusters: general

galaxies: clusters: individual: . . .

- galaxies: clusters: intracluster medium
- galaxies: disc
- galaxies: distances and redshifts
- galaxies: dwarf
- galaxies: elliptical and lenticular, cD
- galaxies: evolution
- galaxies: formation
- galaxies: fundamental parameters
- galaxies: general
- galaxies: groups: general

galaxies: groups: individual: . . .

- galaxies: haloes
- galaxies: high-redshift

galaxies: individual: . . .

- galaxies: interactions
- (galaxies:) intergalactic medium
- galaxies: irregular
- galaxies: ISM
- galaxies: jets
- galaxies: kinematics and dynamics
- (galaxies:) Local Group
- galaxies: luminosity function, mass function
- (galaxies:) Magellanic Clouds
- galaxies: magnetic fields
- galaxies: nuclei
- galaxies: peculiar
- galaxies: photometry
- (galaxies:) quasars: absorption lines
- (galaxies:) quasars: emission lines
- (galaxies:) quasars: general

(*galaxies:*) **quasars: individual: . . .**
(*galaxies:*) quasars: supermassive black holes
galaxies: Seyfert
galaxies: spiral
galaxies: starburst
galaxies: star clusters: general

galaxies: star clusters: individual: . . .
galaxies: star formation
galaxies: statistics
galaxies: stellar content
galaxies: structure

Cosmology

(*cosmology:*) cosmic background radiation
(*cosmology:*) cosmological parameters
(*cosmology:*) dark ages, reionization, first stars
(*cosmology:*) dark energy
(*cosmology:*) dark matter
(*cosmology:*) diffuse radiation
(*cosmology:*) distance scale
(*cosmology:*) early Universe
(*cosmology:*) inflation
(*cosmology:*) large-scale structure of Universe
cosmology: miscellaneous
cosmology: observations
(*cosmology:*) primordial nucleosynthesis
cosmology: theory

Resolved and unresolved sources as a function of wavelength

gamma-rays: diffuse background
gamma-rays: galaxies
gamma-rays: galaxies: clusters
gamma-rays: general
gamma-rays: ISM
gamma-rays: stars
infrared: diffuse background
infrared: galaxies
infrared: general
infrared: ISM
infrared: planetary systems
infrared: stars
radio continuum: galaxies
radio continuum: general
radio continuum: ISM
radio continuum: planetary systems
radio continuum: stars
radio continuum: transients
radio lines: galaxies
radio lines: general
radio lines: ISM
radio lines: planetary systems
radio lines: stars
submillimetre: diffuse background
submillimetre: galaxies
submillimetre: general
submillimetre: ISM
submillimetre: planetary systems
submillimetre: stars
ultraviolet: galaxies

ultraviolet: general
ultraviolet: ISM
ultraviolet: planetary systems
ultraviolet: stars
X-rays: binaries
X-rays: bursts
X-rays: diffuse background
X-rays: galaxies
X-rays: galaxies: clusters
X-rays: general
X-rays: individual: . . .
X-rays: ISM
X-rays: stars

Transients

(*transients:*) black hole mergers
(*transients:*) black hole - neutron star mergers
(*transients:*) fast radio bursts
(*transients:*) gamma-ray bursts
(*transients:*) neutron star mergers
transients: novae
transients: supernovae
transients: tidal disruption events