CAP-Lab-7 Estimation of parameter using Maximum Likelihood

The angular distribution of electron scattering experiments show a probability distribution function of the form: $p(\cos \theta) = N(1 + \alpha \cos^2 \theta)$, where θ is the angle between incident and scattered direction.

An experiment measures scattering events and finds the following values of $\cos\theta$:

```
-.999, -.983, -.956, -.946, -.933, -.925, -.916, -.910, -.881, -.739, -.734, -.717, -.715, -.675, -.665, -.649, -.621, -.537, -.522, -.508, -.499, -.471, -.460, -.419, -.403, -.311, -.305, -.281, -.170, -.162, -.063, 0.214, 0.438, .444, .508, .586, .638, .677, .721, .730, 0.438, .444, .508, .586, .638, .677, .721, .730, 0.768, .785, .790, .793, .877, .896, .931, .938, 0.948, 0.999
```

We wish to calculate the value of α using the Maximum Likelihood Method.

- (a) Calculate N from normalized probability (notice that it depends on α)
- (b) Write down the Likelihood Function (\mathcal{L}) for this problem.
- (c) Make a plot of the relative Likelihood Function ($\mathcal{L}/\mathcal{L}_{max}$) vs α . Use this plot to find a guess value of α . Maximizes the Likelihood Function to estimate α . (Note: You can use log scale if you wish. You are encouraged to be innovative for the maximization method)
- (d) Make a histogram of the observed data and over plot the probability distribution for the best estimated parameters.