

# $\infty$ -Cosmoi for Lean

Emily Riehl

Dominic Verity



# Contents

Chapter 1. $\infty$ -Cosmoi	5
1.1. Overview	5
1.2. Simplicial sets	6
1.3. Enriched limits	14
1.4. $\infty$ -Cosmoi	16
1.5. Change of base	20
1.6. The homotopy 2-category	24
Bibliography	31



## CHAPTER 1

# $\infty$ -Cosmoi

### 1.1. Overview

Following [RV22], from which this document was excerpted, we aim to develop the basic theory of  $\infty$ -categories in a model independent fashion using a common axiomatic framework that is satisfied by a variety of models. In contrast with prior “analytic” treatments of the theory of  $\infty$ -categories — in which the central categorical notions are defined in reference to the coordinates of a particular model — our approach is “synthetic,” proceeding from definitions that can be interpreted simultaneously in many models to which our proofs then apply.

To achieve this, our strategy is not to axiomatize what infinite-dimensional categories *are*, but rather to axiomatize the categorical “universe” in which they *live*. This motivates the notion of an  $\infty$ -cosmos, which axiomatizes the universe in which  $\infty$ -categories live as objects.<sup>1</sup> So that theorem statement about  $\infty$ -cosmoi suggest their natural interpretation, we recast  $\infty$ -category as a technical term, to mean an object in some (typically fixed)  $\infty$ -cosmos. Several common models of  $(\infty, 1)$ -categories<sup>2</sup> are  $\infty$ -categories in this sense, but our  $\infty$ -categories also include certain models of  $(\infty, n)$ -categories<sup>3</sup> as well as fibered versions of all of the above. Thus each of these objects are  $\infty$ -categories in our sense and our theorems apply to all of them.<sup>4</sup> This usage of the term “ $\infty$ -categories” is meant to interpolate between the classical one, which refers to any variety of weak infinite-dimensional categories, and the common one, which is often taken to mean quasi-categories or complete Segal spaces.

Much of the development of the theory of  $\infty$ -categories takes place not in the full  $\infty$ -cosmos but in a quotient that we call the *homotopy 2-category*, the name chosen because an  $\infty$ -cosmos is something like a category of fibrant objects in an enriched model category and the homotopy 2-category is then a categorification of its homotopy category. The homotopy 2-category is a strict 2-category — like the 2-category of categories, functors, and natural transformations<sup>5</sup> — and in this way the foundational proofs in the theory of  $\infty$ -categories closely resemble the classical foundations of ordinary category theory except that the universal properties they characterize, e.g., when a

---

<sup>1</sup>Metaphorical allusions aside, our  $\infty$ -cosmoi resemble the fibrational cosmoi of Street [Str74].

<sup>2</sup>Quasi-categories, complete Segal spaces, Segal categories, and 1-complicial sets (naturally marked quasi-categories) all define the  $\infty$ -categories in an  $\infty$ -cosmos.

<sup>3</sup> $n$ -quasi-categories,  $\Theta_n$ -spaces, iterated complete Segal spaces, and  $n$ -complicial sets also define the  $\infty$ -categories in an  $\infty$ -cosmos, as do saturated (née weak) complicial sets, a model for  $(\infty, \infty)$ -categories.

<sup>4</sup>There is a sense, however, in which many of our definitions are optimized for those  $\infty$ -cosmoi whose objects are  $(\infty, 1)$ -categories. A good illustration is provided by the notion of *discrete  $\infty$ -category*. In the  $\infty$ -cosmoi of  $(\infty, 1)$ -categories, the discrete  $\infty$ -categories are the  $\infty$ -groupoids, but this is not true for the  $\infty$ -cosmoi of  $(\infty, n)$ -categories.

<sup>5</sup>In fact this is another special case: there is an  $\infty$ -cosmos whose objects are ordinary categories and its homotopy 2-category is the usual category of categories, functors, and natural transformations. This 2-category is as old as category theory itself, introduced in Eilenberg and Mac Lane’s foundational paper [EML45].

functor between  $\infty$ -categories defines a cartesian fibration, are slightly weaker than in the familiar case of strict 1-categories.

There are many alternate choices we could have made in selecting the axioms of an  $\infty$ -cosmos. One of our guiding principles, admittedly somewhat contrary to the setting of homotopical higher category theory, was to allow us to work as strictly as possible, with the aim of shortening and simplifying proofs. As a consequence of these choices, the  $\infty$ -categories in an  $\infty$ -cosmos and the functors and natural transformations between them assemble into a 2-category rather than a bicategory. To help us achieve this counterintuitive strictness, each  $\infty$ -cosmos comes with a specified class of maps between  $\infty$ -categories called *isofibrations*. The isofibrations have no homotopy-theoretic meaning, as any functor between  $\infty$ -categories is equivalent to an isofibration with the same codomain. However, isofibrations permit us to consider strictly commutative diagrams between  $\infty$ -categories and allow us to require that the limits of such diagrams satisfy a universal property up to simplicially enriched isomorphism. Neither feature is essential for the development of  $\infty$ -category theory. Similar proofs carry through to a weaker setting, at the cost of more time spent considering coherence of higher cells.

An  $\infty$ -cosmos is a particular sort of *simplicially enriched category* with certain *simplicially enriched limits*. In §1.2, we first review some prerequisites from the theory of simplicial sets, most of which are either currently in Mathlib or on their way. While the notion of simplicially enriched category currently exists in Mathlib, simplicially enriched limits do not, so in §1.3 we first introduce the prerequisite notions of simplicially enriched limits that will be required to state the definition of an  $\infty$ -cosmos in §1.4. The homotopy 2-category of an  $\infty$ -cosmos is then obtained by applying the general theory of change-of-base from enriched category, which is also currently missing from Mathlib. This theory is described in §1.5 and then used to define the homotopy 2-category in §1.6.

Additional chapters will be added to this blueprint in the future, containing excerpts of the material that can be found in [RV22, Chapters 2-5]. The broader aim of this project is to formalize the core basic theory of  $\infty$ -categories, covering those aspects that can be defined in the homotopy 2-category of an  $\infty$ -cosmos.

The authors of this blueprint are particularly indebted to:

- Mario Carneiro, who contributed greatly to the original Lean formalization of  $\infty$ -cosmoi and prerequisite results about the homotopy category functor;
- Johan Commelin, who suggested restructuring this as a blueprint project; and
- Pietro Monticone, who created a template for blueprint-driven formalization projects in Lean, from which this repository was forked.

Special thanks are also due to the Hausdorff Research Institute for Mathematics and the organizers of the Trimester Program “Prospects of Formal Mathematics,” where the genesis of this project took place.

## 1.2. Simplicial sets

Before introducing an axiomatic framework that allows us to develop  $\infty$ -category theory in general, we first consider one model in particular: *quasi-categories*, which were introduced in 1973 by Boardman and Vogt [BV73] in their study of homotopy coherent diagrams. Ordinary 1-categories give examples of quasi-categories via the construction of Definition 1.2.15. Joyal first undertook the task of extending 1-category theory to quasi-category theory in [Joy02] and [Joy08] and in several unpublished draft book manuscripts. The majority of the results in this section are due to him.

DEFINITION 1.2.1 (the simplex category). Let  $\Delta$  denote the **simplex category** of finite nonempty ordinals  $[n] = \{0 < 1 < \dots < n\}$  and order-preserving maps.

The maps in the simplex category include in particular:

DEFINITION 1.2.2 (elementary face maps). The **elementary face operators** are the maps

$$[n-1] \xrightarrow{\delta^i} [n] \quad 0 \leq i \leq n$$

whose images omit the element  $i \in [n]$ .

DEFINITION 1.2.3 (elementary degeneracy maps). The **elementary degeneracy operators** are the maps

$$[n+1] \xrightarrow{\sigma^i} [n] \quad 0 \leq i \leq n$$

whose images double up on the element  $i \in [n]$ .

The following decomposition result is yet to be proven, though there are related results on the image factorization in the simplex category:

PROPOSITION 1.2.4. *Every morphism in  $\Delta$  factors uniquely as an epimorphism followed by a monomorphism; these epimorphisms, the **degeneracy operators**, decompose as composites of elementary degeneracy operators, while the monomorphisms, the **face operators**, decompose as composites of elementary face operators.*

PROOF. The image factorizations have been formalized but the canonical decompositions into elementary face and degeneracy operators remain to be done.  $\square$

DEFINITION 1.2.5 (simplicial set). A **simplicial set** is a presheaf on the simplex category.

DEFINITION 1.2.6 (the category of simplicial sets). The category of **simplicial sets** is the category  $sSet := Set^{\Delta^{op}}$  of presheaves on the simplex category.

Standard examples of simplicial sets include:

DEFINITION 1.2.7 (standard simplex). We write  $\Delta[n]$  for the **standard  $n$ -simplex** the simplicial set represented by  $[n] \in \Delta$ .

DEFINITION 1.2.8 (simplex boundary). We write  $\partial\Delta[n] \subset \Delta[n]$  for the **boundary sphere** of the  $n$ -simplex. The sphere  $\partial\Delta[n]$  is the simplicial subset generated by the codimension-one faces of the  $n$ -simplex.

DEFINITION 1.2.9 (simplicial horn). We write  $\Lambda^k[n] \subset \Delta[n]$  for the  **$k$ -horn** in the  $n$ -simplex. The horn  $\Lambda^k[n]$  is the further simplicial subset of  $\partial\Delta[n]$  that omits the face opposite the vertex  $k$ , but it is defined as a subset of  $\Delta[n]$ .

Given a simplicial set  $X$ , it is conventional to write  $X_n$  for the set of  **$n$ -simplices**, defined by evaluating at  $[n] \in \Delta$ . This is implemented as a scoped notation “ $\_ [n]$ ” accessible with “open Simplicial” or “open scoped Simplicial.”

By the Yoneda lemma:

LEMMA 1.2.10. *Each  $n$ -simplex  $x \in X_n$  corresponds to a map of simplicial sets  $x: \Delta[n] \rightarrow X$ . Accordingly, we write  $x \cdot \delta^i$  for the  $i$ th face of the  $n$ -simplex, an  $(n-1)$ -simplex classified by the composite map*

$$\begin{array}{ccc}
\Lambda^k[n] & \longrightarrow & A \\
\downarrow & \nearrow & \\
\Delta[n] & & 
\end{array}
\quad \text{for } n \geq 2, 0 < k < n.$$

$$\Delta[n-1] \xrightarrow{\delta^i} \Delta[n] \xrightarrow{x} X.$$

PROOF. This is a special case of the Yoneda lemma.  $\square$

The right action of the face operator defines a map  $X_n \xrightarrow{\cdot \delta^i} X_{n-1}$ . Geometrically,  $x \cdot \delta^i$  is the “face opposite the vertex  $i$ ” in the  $n$ -simplex  $x$ .

The category of simplicial sets, as a presheaf category, is very well-behaved:

COROLLARY 1.2.11. *The category of simplicial sets is complete.*

PROOF. Presheaf categories are complete.  $\square$

COROLLARY 1.2.12. *The category of simplicial sets is cocomplete.*

PROOF. Presheaf categories are cocomplete.  $\square$

Instances of these facts currently appear in Mathlib, which likely also knows that the category of simplicial sets is cartesian closed.

The definition of a quasi-category can be found in Mathlib as well.

DEFINITION 1.2.13 (quasi-category). A **quasi-category** is a simplicial set  $A$  in which any **inner horn** can be extended to a simplex, solving the displayed lifting problem:

Quasi-categories were first introduced by Boardman and Vogt [BV73] under the name “weak Kan complexes,” as they generalize the following notion:

DEFINITION 1.2.14 (Kan complex). A **Kan complex** is a simplicial set admitting extensions as in (1.2.13) along all horn inclusions  $n \geq 1, 0 \leq k \leq n$ .

Since any topological space can be encoded as a Kan complex,<sup>6</sup> in this way spaces provide examples of quasi-categories.

Categories also provide examples of quasi-categories via the nerve construction.

DEFINITION 1.2.15 (nerve). The category  $\mathcal{Cat}$  of 1-categories embeds fully faithfully into the category of simplicial sets via the **nerve** functor. An  $n$ -simplex in the nerve of a 1-category  $C$  is a sequence of  $n$  composable arrows in  $C$ , or equally a functor  $\mathfrak{n}+1 \rightarrow C$  from the ordinal category  $\mathfrak{n}+1$  with objects  $0, \dots, n$  and a unique arrow  $i \rightarrow j$  just when  $i \leq j$ .

DEFINITION 1.2.16 (nerve functor). The map  $[n] \mapsto \mathfrak{n}+1$  defines a fully faithful embedding  $\Delta \hookrightarrow \mathcal{Cat}$ . From this point of view, the nerve functor can be described as a “restricted Yoneda embedding” which carries a category  $C$  to the restriction of the representable functor  $\text{hom}(-, C)$  to the image of this inclusion.

---

<sup>6</sup>The total singular complex construction defines a functor from topological spaces to simplicial sets that is an equivalence on their respective homotopy categories — weak homotopy types of spaces correspond to homotopy equivalence classes of Kan complexes [Qui67, §II.2]. The left adjoint “geometrically realizes” a simplicial set as a topological space.



This is an instance of a more general family of “nerve-type constructions.”

**PROPOSITION 1.2.17.** *The nerve of a category  $C$  is **2-coskeletal** as a simplicial set, meaning that every sphere  $\partial\Delta[n] \rightarrow C$  with  $n \geq 3$  is filled uniquely by an  $n$ -simplex in  $C$ , or equivalently that the nerve is canonically isomorphic to the right Kan extension of its restriction to 2-truncated simplicial sets.<sup>7</sup>*

**PROOF.** Note a sphere  $\partial\Delta[2] \rightarrow C$  extends to a 2-simplex if and only if that arrow along its diagonal edge is the composite of the arrows along the edges in the inner horn  $\Lambda^1[2] \subset \partial\Delta[2] \rightarrow C$ . The simplices in dimension 3 and above witness the associativity of the composition of the path of composable arrows found along their **spine**, the 1-skeletal simplicial subset formed by the edges connecting adjacent vertices. In fact, as suggested by the proof of Proposition 1.2.18, any simplicial set in which inner horns admit *unique* fillers is isomorphic to the nerve of a 1-category. This characterization of nerves is not yet in Mathlib, however, we have proven the one-way result, namely that nerves of categories satisfy the “strict Segal condition” and this is used in the proof of 2-coskeletality.  $\square$

In the book that is the primary source this text [RV22], as in much of the  $\infty$ -categories literature, we decline to introduce explicit notation for the nerve functor, preferring instead to identify 1-categories with their nerves. As we shall discover the theory of 1-categories extends to  $\infty$ -categories modeled as quasi-categories in such a way that the restriction of each  $\infty$ -categorical concept along the nerve embedding recovers the corresponding 1-categorical concept. For instance, the standard simplex  $\Delta[n]$  is isomorphic to the nerve of the ordinal category  $n+1$ , and we frequently adopt the latter notation — writing  $1 := \Delta[0]$ ,  $2 := \Delta[1]$ ,  $3 := \Delta[2]$ , and so on — to suggest the correct categorical intuition. However, Mathlib notates nerves explicitly, so at some point this document should be adapted to follow that convention.

To begin down this path, we must first verify the implicit assertion that has just been made. A proof of the following result, due to Johan Commelin, will appear in Mathlib soon (see `Wombat.lean` for now).

**PROPOSITION 1.2.18** (nerves are quasi-categories). *Nerves of categories are quasi-categories.*

This is not the proof that was formalized but we include it for fun:

**PROOF.** Via the isomorphism  $C \cong \text{cosk}_2 C$  from Proposition 1.2.17 and the associated adjunction  $\text{sk}_2 \dashv \text{cosk}_2$  of, the required lifting problem displayed below-left transposes to the one displayed below-right:

$$\begin{array}{ccc} \Lambda^k[n] & \longrightarrow & C \cong \text{cosk}_2 C \\ \downarrow & \nearrow & \\ \Delta[n] & & \end{array} \quad \Leftrightarrow \quad \begin{array}{ccc} \text{sk}_2 \Lambda^k[n] & \longrightarrow & C \\ \downarrow & \nearrow & \\ \text{sk}_2 \Delta[n] & & \end{array}$$

The functor  $\text{sk}_2$  replaces a simplicial set by its **2-skeleton**, the simplicial subset generated by the simplices of dimension at most two. For  $n \geq 4$ , the inclusion  $\text{sk}_2 \Lambda^k[n] \hookrightarrow \text{sk}_2 \Delta[n]$  is an isomorphism, in which case the lifting problems on the right admit (unique) solutions. So it remains only to solve the lifting problems on the left in the cases  $n = 2$  and  $n = 3$ .

To that end consider

---

<sup>7</sup>The equivalence between these two perspectives is non-obvious and makes use of Reedy category theory (see [RV22, §C.4-5]), which does not currently exist in Mathlib.

$$\begin{array}{ccc}
\Lambda^1[2] \longrightarrow C & \Lambda^1[3] \longrightarrow C & \Lambda^2[3] \longrightarrow C \\
\downarrow & \downarrow & \downarrow \\
\Delta[2] & \Delta[3] & \Delta[3]
\end{array}$$

An inner horn  $\Lambda^1[2] \rightarrow C$  defines a composable pair of arrows in  $C$ ; an extension to a 2-simplex exists precisely because any composable pair of arrows admits a (unique) composite.

An inner horn  $\Lambda^1[3] \rightarrow C$  specifies the data of three composable arrows in  $C$ , as displayed in the following diagram, together with the composites  $gf$ ,  $hg$ , and  $(hg)f$ .

$$\begin{array}{ccccc}
& & c_1 & & \\
& f \nearrow & \downarrow & \nwarrow hg & \\
c_0 & \xrightarrow{\quad} & & \xrightarrow{(hg)f} & c_3 \\
& g \searrow & \downarrow & \nearrow h & \\
& & c_2 & & 
\end{array}$$

Because composition is associative, the arrow  $(hg)f$  is also the composite of  $gf$  followed by  $h$ , which proves that the 2-simplex opposite the vertex  $c_1$  is present in  $C$ ; by 2-coskeletality, the 3-simplex filling this boundary sphere is also present in  $C$ . The filler for a horn  $\Lambda^2[3] \rightarrow C$  is constructed similarly.  $\square$

We now turn to the homotopy category functor. The following definitions and results are not currently in Mathlib.

**DEFINITION 1.2.19** (homotopy relation on 1-simplices). A parallel pair of 1-simplices  $f, g$  in a simplicial set  $X$  are **homotopic** if there exists a 2-simplex whose boundary takes either of the following forms<sup>8</sup>

$$\begin{array}{ccc}
& y & \\
f \nearrow & \parallel & \\
x & \xrightarrow{g} & y
\end{array}
\quad
\begin{array}{ccc}
& x & \\
\parallel & \searrow f & \\
x & \xrightarrow{g} & y
\end{array}$$

or if  $f$  and  $g$  are in the same equivalence class generated by this relation.

In a quasi-category, the relation witnessed by either of the types of 2-simplex on display in Definition 1.2.19 is an equivalence relation and these equivalence relations coincide.

**LEMMA 1.2.20** (homotopic 1-simplices in a quasi-category). *Parallel 1-simplices  $f$  and  $g$  in a quasi-category are homotopic if and only if there exists a 2-simplex of any or equivalently all of the forms displayed in Definition 1.2.19.*

**PROOF.** A lengthy exercise in low-dimensional horn filling.  $\square$

**DEFINITION 1.2.21.** By 1-truncating, any simplicial set  $X$  has an underlying **reflexive quiver** or **reflexive directed graph** with the 0-simplices of  $X$  defining the objects and the 1-simplices defining the arrows:

$$X_1 \begin{array}{c} \xrightarrow{\cdot\delta^1} \\ \xleftarrow{\cdot\sigma^0} \\ \xrightarrow{\cdot\delta^0} \end{array} X_0,$$

<sup>8</sup>The symbol “=” is used in diagrams to denote a degenerate simplex or an identity arrow.

By convention, the source of an arrow  $f \in X_1$  is its 0th face  $f \cdot \delta^1$  (the face opposite 1) while the target is its 1st face  $f \cdot \delta^0$  (the face opposite 0).

PROPOSITION 1.2.22. *The functor that carries a category to its underlying reflexive quiver has a left adjoint, defining the free category on a reflexive quiver:*

$$\begin{array}{ccc} & F & \\ \mathcal{C}at & \xleftarrow{\quad} & rQuiv \\ & \perp & \\ & U & \end{array}$$

PROOF. This has been formalized and is currently in the [CatHasColimits branch of Mathlib](#).  $\square$

DEFINITION 1.2.23 (the homotopy category [GZ67, §2.4]). The **free category** on this reflexive directed graph has  $X_0$  as its object set, degenerate 1-simplices serving as identity morphisms, and nonidentity morphisms defined to be finite directed paths of nondegenerate 1-simplices. The **homotopy category**  $hX$  of  $X$  is the quotient of the free category on its underlying reflexive directed graph by the congruence<sup>9</sup> generated by imposing a composition relation  $h = g \circ f$  witnessed by 2-simplices

$$\begin{array}{ccc} & x_1 & \\ f \nearrow & & \searrow g \\ x_0 & \xrightarrow{h} & x_2 \end{array}$$

By soundness of the quotient construction:

LEMMA 1.2.24. *Homotopic 1-simplices in a simplicial set represent the same arrow in the homotopy category.*

PROOF. This should be relatively straightforward.  $\square$

PROPOSITION 1.2.25. *The homotopy category of the nerve of a 1-category is isomorphic to the original category, as the 2-simplices in the nerve witness all of the composition relations satisfied by the arrows in the underlying reflexive directed graph.*

PROOF. This has been formalized and is currently in the [CatHasColimits branch of Mathlib](#).  $\square$

Indeed, the natural isomorphism  $hC \cong C$  forms the counit of an adjunction, embedding  $\mathcal{C}at$  as a reflective subcategory of  $sSet$ .

PROPOSITION 1.2.26. *The nerve embedding admits a left adjoint, namely the functor which sends a simplicial set to its homotopy category:*

$$\begin{array}{ccc} & h & \\ \mathcal{C}at & \xleftarrow{\quad} & sSet \\ & \perp & \end{array}$$

The adjunction of Proposition 1.2.26 exists for formal reasons, via results which have already been formalized in Mathlib, once the category  $\mathcal{C}at$  is known to be cocomplete. A proof of this fact did not exist in Mathlib, however, and in fact the adjunction between the homotopy category and the nerve can be used to construct colimits of categories, as it embeds  $\mathcal{C}at$  as a reflective subcategory of a cocomplete category (see [Rie16, 4.5.16]). Thus, we instead formalized a direct proof.

<sup>9</sup>A binary relation  $\sim$  on parallel arrows of a 1-category is a **congruence** if it is an equivalence relation that is closed under pre- and post-composition: if  $f \sim g$  then  $hfk \sim hgk$ .

PROOF. This has been formalized and is currently in the [CatHasColimits branch of Mathlib](#).

For any simplicial set  $X$ , there is a natural map from  $X$  to the nerve of its homotopy category  $\mathbf{h}X$ ; since nerves are 2-coskeletal, it suffices to define the map  $\mathrm{sk}_2 X \rightarrow \mathbf{h}X$ , and this is given immediately by the construction of Definition 1.2.23. Note that the quotient map  $X \rightarrow \mathbf{h}X$  becomes an isomorphism upon applying the homotopy category functor and is already an isomorphism whenever  $X$  is the nerve of a category. Thus the adjointness follows by direct verification of the triangle equalities.  $\square$

By inspection:

PROPOSITION 1.2.27. *The nerve functor is fully faithful.*

PROOF. This has been formalized and is currently in the [CatHasColimits branch of Mathlib](#).  $\square$

As a corollary, it follows that  $\mathcal{C}at$  has colimits.

The homotopy category of a quasi-category admits a simplified description.

LEMMA 1.2.28 (the homotopy category of a quasi-category). *If  $A$  is a quasi-category then its homotopy category  $\mathbf{h}A$  has*

- the set of 0-simplices  $A_0$  as its objects
- the set of homotopy classes of 1-simplices  $A_1$  as its arrows
- the identity arrow at  $a \in A_0$  represented by the degenerate 1-simplex  $a \cdot \sigma^0 \in A_1$
- a composition relation  $h = g \circ f$  in  $\mathbf{h}A$  between the homotopy classes of arrows represented by any given 1-simplices  $f, g, h \in A_1$  if and only if there exists a 2-simplex with boundary

$$\begin{array}{ccc} & a_1 & \\ f \nearrow & & \searrow g \\ a_0 & \xrightarrow{h} & a_2 \end{array}$$

PROOF. Another lengthy exercise in low-dimensional horn filling.  $\square$

Later we will require either of the following results:

LEMMA 1.2.29.

- (i) *The functor  $\mathbf{h}: s\mathcal{S}et \rightarrow \mathcal{C}at$  preserves finite products.*
- (ii) *The functor  $\mathbf{h}: \mathcal{Q}Cat \rightarrow \mathcal{C}at$  preserves small products.*

PROOF. For the first statement, preservation of the terminal object is by direct calculation. By Proposition 1.2.25, preservation of binary products is equivalent to the statement that the canonical map  $N(\mathcal{D}^c) \rightarrow N(\mathcal{D})^{Nc}$  involving nerves of categories is an isomorphism. On  $n$ -simplices, this is defined by uncurrying, which is bijection since  $\mathcal{C}at$  is cartesian closed.

For the second statement, we have a canonical comparison functor from the homotopy category of the products to the product of the homotopy categories. It follows from Lemma 1.2.28 that this is an isomorphism on underlying quivers, which suffices.  $\square$

DEFINITION 1.2.30 (isomorphism in a quasi-category). A 1-simplex in a quasi-category is an **isomorphism**<sup>10</sup> just when it represents an isomorphism in the homotopy category. By Lemma 1.2.28 this means that  $f: a \rightarrow b$  is an isomorphism if and only if there exists a 1-simplex  $f^{-1}: b \rightarrow a$  together with a pair of 2-simplices

<sup>10</sup>Joyal refers to these maps as “isomorphisms” while Lurie refers to them as “equivalences.” We prefer, wherever possible, to use the same term for  $\infty$ -categorical concepts as for the analogous 1-categorical ones.

$$\begin{array}{ccc}
& b & \\
f \nearrow & & \searrow f^{-1} \\
a & \xlongequal{\quad} & a
\end{array}
\qquad
\begin{array}{ccc}
& a & \\
f^{-1} \nearrow & & \searrow f \\
b & \xlongequal{\quad} & b
\end{array}$$

The properties of the isomorphisms in a quasi-category are somewhat technical to prove and will likely be a pain to formalize (see [RV22, §D]). Here we focus on a few essential results, which are more easily obtainable.

DEFINITION 1.2.31. The **homotopy coherent isomorphism**  $\mathbb{I}$ , is the nerve of the free-living isomorphism.

Just as the arrows in a quasi-category  $A$  are represented by simplicial maps  $2 \rightarrow A$  whose domain is the nerve of the free-living arrow, the isomorphisms in a quasi-category can be represented by diagrams  $\mathbb{I} \rightarrow A$  whose domain is the homotopy coherent isomorphism:

PROPOSITION 1.2.32. *An arrow  $f$  in a quasi-category  $A$  is an isomorphism if and only if it extends to a homotopy coherent isomorphism*

$$\begin{array}{ccc}
2 & \xrightarrow{f} & A \\
\downarrow & \nearrow & \uparrow \\
\mathbb{I} & & 
\end{array}$$

REMARK 1.2.33. *If this result proves too annoying to formalize without the general theory of “special-outer horn filling,” we might instead substitute a finite model of the homotopy coherent isomorphism for  $\mathbb{I}$ .*

Quasi-categories define the fibrant objects in a model structure due to Joyal. We use the term *isofibration* to refer to the fibrations between fibrant objects in this model structure, which admit the following concrete description.

DEFINITION 1.2.34 (isofibration). A simplicial map  $f: A \rightarrow B$  between quasi-categories is an **isofibration** if it lifts against the inner horn inclusions, as displayed below-left, and also against the inclusion of either vertex into the free-living isomorphism  $\mathbb{I}$ .

$$\begin{array}{ccc}
\Lambda^k[n] & \longrightarrow & A \\
\downarrow & \nearrow & \downarrow f \\
\Delta[n] & \longrightarrow & B
\end{array}
\qquad
\begin{array}{ccc}
\mathbb{I} & \longrightarrow & A \\
\downarrow & \nearrow & \downarrow f \\
\mathbb{I} & \longrightarrow & B
\end{array}$$

To notationally distinguish the isofibrations, we depict them as arrows “ $\twoheadrightarrow$ ” with two heads.

We now introduce the weak equivalences and trivial fibrations between fibrant objects in the Joyal model structure.

DEFINITION 1.2.35 (equivalences of quasi-categories). A map  $f: A \rightarrow B$  between quasi-categories is an **equivalence** if it extends to the data of a “homotopy equivalence” with the free-living isomorphism  $\mathbb{I}$  serving as the interval: that is, if there exist maps  $g: B \rightarrow A$ ,

$$\begin{array}{ccc}
& A & \\
& \parallel & \uparrow \text{ev}_0 \\
A & \xrightarrow{\alpha} & A^{\mathbb{I}} \\
& \searrow gf & \downarrow \text{ev}_1 \\
& A & 
\end{array}
\quad \text{and} \quad
\begin{array}{ccc}
& B & \\
& \parallel & \uparrow \text{ev}_0 \\
B & \xrightarrow{\beta} & B^{\mathbb{I}} \\
& \searrow fg & \downarrow \text{ev}_1 \\
& B & 
\end{array}$$

We write “ $\xrightarrow{\sim}$ ” to decorate equivalences and  $A \simeq B$  to indicate the presence of an equivalence  $A \xrightarrow{\sim} B$ .

LEMMA 1.2.36. *If  $f: A \rightarrow B$  is an equivalence of quasi-categories, then the functor  $hf: hA \rightarrow hB$  is an equivalence of categories, where the data displayed above defines an equivalence inverse  $hg: hB \rightarrow hA$  and natural isomorphisms encoded by the composite<sup>11</sup> functors*

$$hA \xrightarrow{h\alpha} h(A^\flat) \longrightarrow (hA)^\flat \qquad hB \xrightarrow{h\beta} h(B^\flat) \longrightarrow (hB)^\flat$$

DEFINITION 1.2.37. A map  $f: X \rightarrow Y$  between simplicial sets is a **trivial fibration** if it admits lifts against the boundary inclusions for all simplices

$$\begin{array}{ccc} \partial\Delta[n] & \longrightarrow & X \\ \downarrow & \nearrow \sim & \downarrow f \\ \Delta[n] & \longrightarrow & Y \end{array} \quad \text{for } n \geq 0$$

We write “ $\xrightarrow{\sim}$ ” to decorate trivial fibrations.<sup>12</sup>

The notation “ $\xrightarrow{\sim}$ ” is suggestive: the trivial fibrations between quasi-categories are exactly those maps that are both isofibrations and equivalences. This can be proven by a relatively standard although rather technical argument in simplicial homotopy theory [RV22, D.5.6].

### 1.3. Enriched limits

A simplicially enriched category—commonly called a “simplicial category” for short—is a category that is enriched over the cartesian monoidal category of simplicial sets. We recall the definition, which already exists in Mathlib.

DEFINITION 1.3.1 (simplicial categories as enriched categories).

The data of a **simplicial category** is a **simplicially enriched category** with a set of objects and a simplicial set  $\mathcal{A}(x, y)$  of morphisms between each ordered pair of objects. Each endo-hom space contains a distinguished 0-simplex  $\text{id}_x \in \mathcal{A}(x, x)_0$ , and composition is required to define a simplicial map

$$\mathcal{A}(y, z) \times \mathcal{A}(x, y) \xrightarrow{\circ} \mathcal{A}(x, z)$$

The composition is required to be associative and unital, in a sense expressed by the commutative diagrams of simplicial sets

$$\begin{array}{ccc} \mathcal{A}(y, z) \times \mathcal{A}(x, y) \times \mathcal{A}(w, x) & \xrightarrow{\circ \times \text{id}} & \mathcal{A}(x, z) \times \mathcal{A}(w, x) \\ \text{id} \times \circ \downarrow & & \downarrow \circ \\ \mathcal{A}(y, z) \times \mathcal{A}(w, y) & \xrightarrow{\circ} & \mathcal{A}(w, z) \end{array} \qquad \begin{array}{ccc} \mathcal{A}(x, y) & \xrightarrow{\text{id}_y \times \text{id}} & \mathcal{A}(y, y) \times \mathcal{A}(x, y) \\ \text{id} \times \text{id}_x \downarrow & \searrow \text{id} & \downarrow \circ \\ \mathcal{A}(x, y) \times \mathcal{A}(x, x) & \xrightarrow{\circ} & \mathcal{A}(x, y) \end{array}$$

DEFINITION 1.3.2. For each  $n \geq 0$ , an  $n$ -simplex in  $\mathcal{A}(x, y)$  is referred to as an  $n$ -**arrow** from  $x$  to  $y$ .

<sup>11</sup>Note that  $h(A^\flat) \not\cong (hA)^\flat$  in general. Objects in the latter are homotopy classes of isomorphisms in  $A$ , while objects in the former are homotopy coherent isomorphisms, given by a specified 1-simplex in  $A$ , a specified inverse 1-simplex, together with an infinite tower of coherence data indexed by the nondegenerate simplices in  $\mathbb{I}$ .

<sup>12</sup>Please help us find an html friendly version of this symbol.

LEMMA 1.3.3. *For any simplicial category  $\mathcal{A}$  and  $n \geq 0$ , the  $n$ -arrows assemble into the arrows of an ordinary category  $\mathcal{A}_n$  with the same set of objects as  $\mathcal{A}$ .*

PROOF. The category of  $n$ -arrows is easy to construct directly. Alternatively, this result can be proven by applying the theory of change-of-base of §1.5 to the functor  $\text{ev}_n: s\text{Set} \rightarrow \text{Set}$ .  $\square$

In particular, the underlying category of a simplicial category can be identified with the category of 0-arrows. Below we link the general Mathlib construction of the underlying category of an enriched category, though it might be useful to formalize a lemma characterizing it in the way stated here.

DEFINITION 1.3.4. The category  $\mathcal{A}_0$  of 0-arrows is the **underlying category** of the simplicial category  $\mathcal{A}$ , which forgets the higher dimensional simplicial structure.

There is alternate presentation of the data of a simplicial category as a simplicial object in the category of categories and identity-on-objects functors.<sup>13</sup>

DIGRESSION 1.3.5 (simplicial categories as simplicial objects). A **simplicial category**  $\mathcal{A}$  is equivalently given by categories  $\mathcal{A}_n$ , with a common set of objects and whose arrows are called  **$n$ -arrows**, that assemble into a diagram  $\Delta^{\text{op}} \rightarrow \text{Cat}$  of identity-on-objects functors

$$\begin{array}{ccccc} & \xrightarrow{\cdot\delta^3} & & \xrightarrow{\cdot\delta^2} & \\ \xleftarrow{\cdot\sigma^2} & & \xleftarrow{\cdot\sigma^1} & & \xleftarrow{\cdot\sigma^0} \\ \xrightarrow{\cdot\delta^2} & & \xrightarrow{\cdot\delta^1} & & \xrightarrow{\cdot\delta^0} \\ \cdots \mathcal{A}_3 & \xleftarrow{\cdot\sigma^1} & \mathcal{A}_2 & \xleftarrow{\cdot\sigma^0} & \mathcal{A}_1 & \xleftarrow{\cdot\sigma^0} & \mathcal{A}_0 =: \mathcal{A} \\ \xrightarrow{\cdot\delta^1} & & \xrightarrow{\cdot\delta^0} & & & & \\ \xleftarrow{\cdot\sigma^0} & & \xleftarrow{\cdot\sigma^0} & & & & \\ \xrightarrow{\cdot\delta^0} & & \xrightarrow{\cdot\delta^0} & & & & \end{array}$$

By contrast, the notion of simplicially enriched limit remains to be formalized. Fortunately, we do not (immediately) require the general notion of *weighted limits*, as the notion of an  $\infty$ -cosmos only requires two special cases: *cotensors* and *conical limits*.

DEFINITION 1.3.6 (simplicial cotensors). Let  $\mathcal{A}$  be a simplicial category. The **cotensor** of an object  $A \in \mathcal{A}$  by a simplicial set  $U$  is given by the data of an object  $A^U \in \mathcal{A}$  together with a cone  $U \rightarrow \mathcal{A}(A^U, A)$  so that the induced map defines an isomorphism of simplicial sets:

$$(1.3.7) \quad \mathcal{A}(X, A^U) \cong \mathcal{A}(X, A)^U$$

Note by construction the isomorphism (1.3.7) is automatically simplicially natural in  $X$ . This simplicial naturality is an important aspect of the enriched universal property.

DEFINITION 1.3.8 (simplicial cotensors). A simplicial category  $\mathcal{A}$  **has cotensors** when all cotensors exist.

LEMMA 1.3.9. *When a simplicial category has cotensors, cotensors are associative: given  $A \in \mathcal{A}$  and simplicial sets  $U$  and  $V$  there are canonical isomorphisms*

$$(A^U)^V \cong A^{U \times V} \cong (A^V)^U.$$

PROOF. By the enriched Yoneda lemma, these objects represent the same simplicial functors  $\mathcal{A}^{\text{op}} \rightarrow s\text{Set}$ .  $\square$

LEMMA 1.3.10. *Assuming such objects exist, the simplicial cotensor defines a bifunctor*

<sup>13</sup>The phrase “simplicial object in  $\text{Cat}$ ” is reserved for the more general yet less common notion of a diagram  $\Delta^{\text{op}} \rightarrow \text{Cat}$  that is not necessarily comprised of identity-on-objects functors.

$$\begin{aligned} sSet^{\text{op}} \times \mathcal{A} &\longrightarrow \mathcal{A} \\ (U, A) &\longmapsto A^U \end{aligned}$$

in a unique way making the isomorphism (1.3.7) natural in  $U$  and  $A$  as well.

PROOF. Functoriality in each variable follows from the universal property.  $\square$

The other simplicial limit notions postulated by axiom 1.4.1i are **conical**, which is the term used for ordinary 1-categorical limit shapes that satisfy an enriched analog of the usual universal property. Such limits also define limits in the underlying category, but the usual universal property is strengthened.

DEFINITION 1.3.11 (simplicial conical limits). Consider a limit cone  $(\lim_{j \in J} A_j \rightarrow A_j)_{j \in J}$  in the underlying category  $\mathcal{A}_0$  of a simplicially-enriched category  $\mathcal{A}$ . By applying the covariant representable functor  $\mathcal{A}(X, -): \mathcal{A}_0 \rightarrow sSet$  to a limit cone  $(\lim_{j \in J} A_j \rightarrow A_j)_{j \in J}$  in  $\mathcal{A}_0$ , we obtain a natural comparison map

$$(1.3.12) \quad \mathcal{A}(X, \lim_{j \in J} A_j) \rightarrow \lim_{j \in J} \mathcal{A}(X, A_j).$$

We say that  $\lim_{j \in J} A_j$  defines a **simplicially enriched limit** if and only if (1.3.12) is an isomorphism of simplicial sets for all  $X \in \mathcal{A}$ .

REMARK 1.3.13. *The notion of cotensors and conical limits could be introduced for categories enriched over arbitrary cartesian monoidal categories or more generally for categories enriched over symmetric monoidal categories. This might be worth doing as a service to the broader Mathlib.*

#### 1.4. $\infty$ -Cosmoi

There are a variety of models of infinite-dimensional categories for which the category of “ $\infty$ -categories,” as we call them, and “ $\infty$ -functors” between them is enriched over quasi-categories and admits classes of isofibrations, equivalences, and trivial fibrations satisfying certain properties that are familiar from abstract homotopy theory.<sup>14</sup> In particular, the use of isofibrations in diagrams guarantees that their strict limits are equivalence invariant, so we can take advantage of up-to-isomorphism universal properties and strict functoriality of these constructions while still working “homotopically.” This motivates the following axiomatization:

DEFINITION 1.4.1 ( $\infty$ -cosmos). An  $\infty$ -**cosmos**  $\mathcal{K}$  is a category that is enriched over quasi-categories,<sup>15</sup> meaning in particular that

- its morphisms  $f: A \rightarrow B$  define the vertices of a quasi-category denoted  $\text{Fun}(A, B)$  and referred to as a **functor space**,

that is also equipped with a specified collection of maps that we call **isofibrations** and denote by “ $\twoheadrightarrow$ ” satisfying the following two axioms:

- (i) (completeness) The quasi-categorically enriched category  $\mathcal{K}$  possesses a terminal object, small products, pullbacks of isofibrations, limits of countable towers of isofibrations, and

<sup>14</sup>More specifically, these classes form a *category of fibrant objects* à la Brown [Bro73].

<sup>15</sup>This is to say  $\mathcal{K}$  is a simplicially enriched category (see Definition 1.3.1) whose hom spaces are all quasi-categories.



cotensors with simplicial sets, each of these limit notions satisfying a universal property that is enriched over simplicial sets.<sup>16</sup>

- (ii) (isofibrations) The isofibrations contain all isomorphisms and any map whose codomain is the terminal object; are closed under composition, product, pullback, forming inverse limits of towers, and Leibniz cotensors with monomorphisms of simplicial sets; and have the property that if  $f: A \twoheadrightarrow B$  is an isofibration and  $X$  is any object then  $\mathrm{Fun}(X, A) \twoheadrightarrow \mathrm{Fun}(X, B)$  is an isofibration of quasi-categories.

For ease of reference, we refer to the simplicially enriched limits of diagrams of isofibrations enumerated in [i](#) as the **cosmological limit notions**.

DEFINITION 1.4.2. In an  $\infty$ -cosmos  $\mathcal{K}$ , a morphism  $f: A \rightarrow B$  is an **equivalence** just when the induced map  $f_*: \mathrm{Fun}(X, A) \xrightarrow{\sim} \mathrm{Fun}(X, B)$  on functor spaces is an equivalence of quasi-categories for all  $X \in \mathcal{K}$ .

DEFINITION 1.4.3. In an  $\infty$ -cosmos  $\mathcal{K}$ , a morphism  $f: A \rightarrow B$  is a **trivial fibration** just when  $f$  is both an isofibration and an equivalence.

These classes are denoted by “ $\xrightarrow{\sim}$ ” and “ $\xrightarrow{\sim}\twoheadrightarrow$ ”, respectively.<sup>17</sup>

Put more concisely, one might say that an  $\infty$ -cosmos is a “quasi-categorically enriched category of fibrant objects.”

CONVENTION 1.4.4 ( $\infty$ -category, as a technical term). Henceforth, we recast  $\infty$ -**category** as a technical term to refer to an object in an arbitrary ambient  $\infty$ -cosmos. Similarly, we use the term  $\infty$ -**functor** — or more commonly the elision “**functor**” — to refer to a morphism  $f: A \rightarrow B$  in an  $\infty$ -cosmos. This explains why we refer to the quasi-category  $\mathrm{Fun}(A, B)$  between two  $\infty$ -categories in an  $\infty$ -cosmos as a “functor space”: its vertices are the  $(\infty)$ -functors from  $A$  to  $B$ .

DEFINITION 1.4.5. The underlying category  $\mathcal{K}_0$  of an  $\infty$ -cosmos  $\mathcal{K}$  is the category whose objects are the  $\infty$ -categories in  $\mathcal{K}$  and whose morphisms are the 0-arrows, i.e., the vertices in the functor spaces.

In all of the examples to appear in what follows, this recovers the expected category of  $\infty$ -categories in a particular model and functors between them.

The following theorem should be quite difficult to formalize:

PROPOSITION 1.4.6 (the  $\infty$ -cosmos of quasi-categories). *The full subcategory  $\mathcal{QCat} \subset sSet$  of quasi-categories defines an  $\infty$ -cosmos in which the isofibrations, equivalences, and trivial fibrations coincide with the classes already bearing these names.*

PROOF. The proof requires myriad combinatorial results about the class of isofibrations between quasi-categories. See [\[RV22, §D\]](#). □

Two further examples fit into a common paradigm: both arise as full subcategories of the  $\infty$ -cosmos of quasi-categories and inherit their  $\infty$ -cosmos structures from this inclusion (see Lemma [\[RV22, 6.1.4\]](#)), but it is also instructive, and ultimately takes less work, to describe the resulting  $\infty$ -cosmos structures directly.

<sup>16</sup>This is to say, these are simplicially enriched limit notions, in the sense described in Definitions [1.3.6](#) and [1.3.11](#).

<sup>17</sup>Please help us find an html friendly version of the trivial fibration symbol.

PROPOSITION 1.4.7 (the  $\infty$ -cosmos of categories). *The category  $\mathcal{Cat}$  of 1-categories defines an  $\infty$ -cosmos whose isofibrations are the **isofibrations**: functors satisfying the displayed right lifting property:*

$$\begin{array}{ccc} \mathbb{1} & \longrightarrow & A \\ \downarrow & \nearrow \text{dashed} & \downarrow f \\ \mathbb{1} & \longrightarrow & B \end{array}$$

*The equivalences are the equivalences of categories and the trivial fibrations are **surjective equivalences**: equivalences of categories that are also surjective on objects.*

PROOF. It is well-known that the 2-category of categories is complete (and in fact also cocomplete) as a  $\mathcal{Cat}$ -enriched category (see [Kel89]). The categorically enriched category of categories becomes a quasi-categorically enriched category by applying the nerve functor to the hom-categories (see §1.5). Since the nerve functor is a right adjoint, it follows formally that these 2-categorical limits become simplicially enriched limits. In particular, as proscribed in Proposition 1.5.9, the cotensor of a category  $A$  by a simplicial set  $U$  is defined to be the functor category  $A^{\mathbf{h}U}$ . This completes the verification of axiom i.

Since the class of isofibrations is characterized by a right lifting property, the isofibrations are closed under all of the limit constructions of 1.4.1ii except for the last two. For these, the Leibniz closure subsumes the closure under exponentiation.

To verify that isofibrations of categories  $f: A \twoheadrightarrow B$  are stable under forming Leibniz cotensors with monomorphisms of simplicial sets  $i: U \hookrightarrow V$ , we must solve the lifting problem below-left

$$\begin{array}{ccc} \mathbb{1} & \xrightarrow{s} & A^{\mathbf{h}V} \\ j \downarrow & \nearrow \gamma & \downarrow \mathbf{h}i^* f \\ \mathbb{1} & \xrightarrow{\langle \beta, \alpha \rangle} & B^{\mathbf{h}V} \times_{B^{\mathbf{h}U}} A^{\mathbf{h}U} \end{array} \quad \Leftrightarrow \quad \begin{array}{ccc} \mathbf{h}U \times \mathbb{1} \cup_{\mathbf{h}U} \mathbf{h}V & \xrightarrow{\langle \alpha, s \rangle} & A \\ \mathbf{h}i \times j \downarrow & \nearrow \gamma & \downarrow f \\ \mathbf{h}V \times \mathbb{1} & \xrightarrow{\beta} & B \end{array}$$

which transposes to the lifting problem above-right, which we can solve by hand. Here the map  $\beta$  defines a natural isomorphism between  $f s: \mathbf{h}V \rightarrow B$  and a second functor. Our task is to lift this to a natural isomorphism  $\gamma$  from  $s$  to another functor that extends the natural isomorphism  $\alpha$  along  $\mathbf{h}i: \mathbf{h}U \rightarrow \mathbf{h}V$ . Note this functor  $\mathbf{h}i$  need not be an inclusion, but it is injective on objects, which is enough.

We define the components of  $\gamma$  by cases. If an object  $v \in \mathbf{h}V$  is equal to  $i(u)$  for some  $u \in \mathbf{h}U$  define  $\gamma_{i(u)} := \alpha_u$ ; otherwise, use the fact that  $f$  is an isofibration to define  $\gamma_v$  to be any lift of the isomorphism  $\beta_v$  to an isomorphism in  $A$  with domain  $s(v)$ . The data of the map  $\gamma: \mathbf{h}V \times \mathbb{1} \rightarrow A$  also entails the specification of the functor  $\mathbf{h}V \rightarrow A$  that is the codomain of the natural isomorphism  $\gamma$ . On objects, this functor is given by  $v \mapsto \text{cod}(\gamma_v)$ . On morphisms, this functor defined in the unique way that makes  $\gamma$  into a natural transformation:

$$(k: v \rightarrow v') \mapsto \gamma_{v'} \circ s(k) \circ \gamma_v^{-1}.$$

This completes the proof that  $\mathcal{Cat}$  defines an  $\infty$ -cosmos. Since the nerve of a functor category, such as  $A^{\mathbb{1}}$ , is isomorphic to the exponential between their nerves, the equivalences of categories coincide with the equivalences of Definition 1.2.35. It follows that the equivalences in the  $\infty$ -cosmos of categories coincide with equivalences of categories, and since the surjective equivalences are the intersection of the equivalences and the isofibrations, this completes the proof.  $\square$

Similarly:

PROPOSITION 1.4.8 (the  $\infty$ -cosmos of Kan complexes). *The category  $\mathcal{Kan}$  of Kan complexes defines an  $\infty$ -cosmos whose isofibrations are the **Kan fibrations**: maps that lift against all horn inclusions  $\Lambda^k[n] \hookrightarrow \Delta[n]$  for  $n \geq 1$  and  $0 \leq k \leq n$ .*

Several consequences of the  $\infty$ -cosmos axioms are mentioned in [RV22, §1.2]. For now, we focus on just one.

By a Yoneda-style argument, the “homotopy equivalence” characterization of the equivalences in the  $\infty$ -cosmos of quasi-categories of Definition 1.2.35 extends to an analogous characterization of the equivalences in any  $\infty$ -cosmos:

LEMMA 1.4.9 (equivalences are homotopy equivalences). *A map  $f: A \rightarrow B$  between  $\infty$ -categories in an  $\infty$ -cosmos  $\mathcal{K}$  is an equivalence if and only if it extends to the data of a “homotopy equivalence” with the free-living isomorphism  $\mathbb{I}$  serving as the interval: that is, if there exist maps  $g: B \rightarrow A$*

$$\begin{array}{ccc}
 & A & \\
 \nearrow & \uparrow \sim \text{ev}_0 & \\
 A & \xrightarrow{\alpha} A^{\mathbb{I}} & \\
 \searrow & \downarrow \sim \text{ev}_1 & \\
 & A &
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 & B & \\
 \nearrow fg & \uparrow \sim \text{ev}_0 & \\
 B & \xrightarrow{\beta} B^{\mathbb{I}} & \\
 \searrow & \downarrow \sim \text{ev}_1 & \\
 & B &
 \end{array}$$

in the  $\infty$ -cosmos.

PROOF. By hypothesis, if  $f: A \rightarrow B$  defines an equivalence in the  $\infty$ -cosmos  $\mathcal{K}$  then the induced map on post-composition  $f_*: \text{Fun}(B, A) \xrightarrow{\sim} \text{Fun}(B, B)$  is an equivalence of quasi-categories in the sense of Definition 1.2.35. Evaluating the inverse equivalence  $\tilde{g}: \text{Fun}(B, B) \xrightarrow{\sim} \text{Fun}(B, A)$  and homotopy  $\tilde{\beta}: \text{Fun}(B, B) \rightarrow \text{Fun}(B, B)^{\mathbb{I}}$  at the 0-arrow  $\text{id}_B \in \text{Fun}(B, B)$ , we obtain a 0-arrow  $g: B \rightarrow A$  together with an isomorphism  $\beta: \mathbb{I} \rightarrow \text{Fun}(B, B)$  from the composite  $fg$  to  $\text{id}_B$ . By the defining universal property of the cotensor (1.3.7), this isomorphism internalizes to define the map  $\beta: B \rightarrow B^{\mathbb{I}}$  in  $\mathcal{K}$  displayed on the right of the displayed equation in the statement.

Now the hypothesis that  $f$  is an equivalence also provides an equivalence of quasi-categories  $f_*: \text{Fun}(A, A) \xrightarrow{\sim} \text{Fun}(A, B)$ , and the map  $\beta f: A \rightarrow B^{\mathbb{I}}$  represents an isomorphism in  $\text{Fun}(A, B)$  from  $fgf$  to  $f$ . Since  $f_*$  is an equivalence, we conclude from Lemma 1.2.36 that  $\text{id}_A$  and  $gf$  are isomorphic in the quasi-category  $\text{Fun}(A, A)$ : explicitly, such an isomorphism may be defined by applying the inverse equivalence  $\tilde{h}: \text{Fun}(A, B) \rightarrow \text{Fun}(A, A)$  and composing with the components at  $\text{id}_A, gf \in \text{Fun}(A, A)$  of the isomorphism  $\tilde{\alpha}: \text{Fun}(A, A) \rightarrow \text{Fun}(A, A)^{\mathbb{I}}$  from  $\text{id}_{\text{Fun}(A, A)}$  to  $\tilde{h}f_*$ . Now by Proposition 1.2.32 this isomorphism is represented by a map  $\mathbb{I} \rightarrow \text{Fun}(A, A)$  from  $\text{id}_A$  to  $gf$ , which internalizes to a map  $\alpha: A \rightarrow A^{\mathbb{I}}$  in  $\mathcal{K}$  displayed on the left of the displayed equation in the statement.

The converse is easy: the simplicial cotensor construction commutes with  $\text{Fun}(X, -)$ , so a homotopy equivalence induces a homotopy equivalence of quasi-categories as in Definition 1.2.35.  $\square$

Many, though not all, of the  $\infty$ -cosmoi we encounter “in the wild” satisfy an additional axiom:

DEFINITION 1.4.10 (cartesian closed  $\infty$ -cosmoi). An  $\infty$ -cosmos  $\mathcal{K}$  is **cartesian closed** if the product bifunctor  $- \times -: \mathcal{K} \times \mathcal{K} \rightarrow \mathcal{K}$  extends to a simplicially enriched two-variable adjunction

$$\text{Fun}(A \times B, C) \cong \text{Fun}(A, C^B) \cong \text{Fun}(B, C^A)$$

in which the right adjoints  $(-)^A: \mathcal{K} \rightarrow \mathcal{K}$  preserve isofibrations for all  $A \in \mathcal{K}$ .

For instance, the  $\infty$ -cosmos of quasi-categories is cartesian closed, with the exponentials defined as (special cases of) simplicial cotensors. This is one of the reasons that we use the same notation for cotensor and for exponential. Note in this case the functor spaces and the exponentials coincide. The same is true for the cartesian closed  $\infty$ -cosmoi of categories and of Kan complexes. In general, the functor space from  $A$  to  $B$  is the “underlying quasi-category” of the exponential  $B^A$  whenever it exists.

### 1.5. Change of base

“Change of base,” first considered by Eilenberg and Kelly in [EK66], refers to a systematic procedure by which enrichment over one category  $\mathcal{V}$  is converted into enrichment over another category  $\mathcal{W}$ . This will be applied in §1.6 to convert an  $\infty$ -cosmos into a simpler structure. For a cartesian closed category  $\mathcal{V}$ , there is a 2-category  $\mathcal{V}\text{-Cat}$  of  $\mathcal{V}$ -categories,  $\mathcal{V}$ -functors, and  $\mathcal{V}$ -natural transformations. The first main result, appearing as Proposition 1.5.4, gives conditions under which a functor  $T: \mathcal{V} \rightarrow \mathcal{W}$  between cartesian closed categories induces a change-of-base 2-functor  $T_*: \mathcal{V}\text{-Cat} \rightarrow \mathcal{W}\text{-Cat}$ .

As the context we are working in here is less general than the one considered by Eilenberg and Kelly — our base categories are cartesian closed while theirs are closed symmetric monoidal — we take a shortcut which covers all of our examples and is easier to explain. In general, all that is needed to produce a change of base 2-functor is a *lax monoidal* functor between symmetric monoidal categories, but the lax monoidal functors we encounter between cartesian closed categories are in fact finite-product-preserving, so we content ourselves with explicating the results in that case instead.

However, lax monoidal functors exist in Mathlib already, so we briefly recall the definition.

**DEFINITION 1.5.1.** A **(lax) monoidal functor** between cartesian closed categories  $\mathcal{V}$  and  $\mathcal{W}$  is a functor  $T: \mathcal{V} \rightarrow \mathcal{W}$  equipped with natural transformations

$$\begin{array}{ccc} \mathcal{V} \times \mathcal{V} & \xrightarrow{T \times T} & \mathcal{W} \times \mathcal{W} \\ \times \downarrow & \Downarrow \phi & \downarrow \times \\ \mathcal{V} & \xrightarrow{T} & \mathcal{W} \end{array} \quad \begin{array}{ccc} \mathbb{1} & \xrightarrow{1} & \mathcal{V} \\ & \nearrow \uparrow \phi_0 & \downarrow T \\ & 1 & \mathcal{W} \end{array}$$

so that the evident associativity and unit diagrams commute.

Except in a special case that we now introduce, the maps  $\phi$  and  $\phi_0$  are to be regarded as part of the structure of a lax monoidal functor, rather than a property the functor  $T$  enjoys.

Recall that a functor  $T: \mathcal{V} \rightarrow \mathcal{W}$  between cartesian closed categories **preserves finite products** just when the natural maps defined for any  $u, v \in \mathcal{V}$

$$T(u \times v) \xrightarrow{\cong} Tu \times Tv \quad \text{and} \quad T1 \xrightarrow{\cong} 1$$

are isomorphisms. These maps satisfy the duals of the coherence conditions mentioned in Definition 1.5.1 and make  $T$  into a **strong monoidal functor** between the cartesian closed categories  $\mathcal{V}$  and  $\mathcal{W}$ . The inverse isomorphisms then provide the structure maps of Definition 1.5.1.

For example:

**EXAMPLE 1.5.2.** Since representable functors preserve products, for any cartesian closed category  $\mathcal{V}$ , the underlying set functor  $(-)_0: \mathcal{V} \rightarrow \mathbf{Set}$  is product-preserving

EXAMPLE 1.5.3. In a cartesian closed category  $\mathcal{V}$ , finite products distribute over arbitrary coproducts. In particular, for any sets  $X$  and  $Y$  there is an isomorphism

$$\coprod_{X \times Y} 1 \cong (\coprod_X 1) \times (\coprod_Y 1)$$

between coproducts of the terminal object  $1$ , which proves that the functor

$$\mathcal{Set} \xrightarrow{\coprod - 1} \mathcal{V}$$

is finite-product-preserving.

A finite-product-preserving functor may be used to change the base as follows:

PROPOSITION 1.5.4. *A finite-product-preserving functor  $T: \mathcal{V} \rightarrow \mathcal{W}$  between cartesian closed categories induces a change-of-base 2-functor*

$$\mathcal{V}\text{-Cat} \xrightarrow{T_*} \mathcal{W}\text{-Cat}.$$

An early observation along these lines was first stated as [EK66, II.6.3], with the proof left to the reader. We adopt the same tactic and leave the diagram chases to the reader or to [Cru08, 4.2.4] and instead just give the construction of the change-of-base 2-functor, which is the important thing. The construction of a  $\mathcal{W}$ -category  $T_*\mathcal{C}$  from a  $\mathcal{V}$ -category  $\mathcal{C}$  exists in Mathlib in the more general setting of a lax monoidal functor  $T$ , but change of base for enriched functors or natural transformations has not been formalized.

PROOF. Let  $\mathcal{C}$  be a  $\mathcal{V}$ -category and define a  $\mathcal{W}$ -category  $T_*\mathcal{C}$  to have the same objects and to have mapping objects  $T_*\mathcal{C}(x, y) := T\mathcal{C}(x, y)$ . The composition and identity maps are given by the composites

$$T\mathcal{C}(y, z) \times T\mathcal{C}(x, y) \xrightarrow{\cong} T(\mathcal{C}(y, z) \times \mathcal{C}(x, y)) \xrightarrow{T\circ} T\mathcal{C}(x, z) \quad 1 \xrightarrow{\cong} T1 \xrightarrow{T\text{id}_x} T\mathcal{C}(x, x)$$

which make use of the inverses of the natural maps that arise when a finite-product-preserving functor is applied to a finite product. A straightforward diagram chase verifies that  $T_*\mathcal{C}$  is a  $\mathcal{W}$ -category.

If  $F: \mathcal{C} \rightarrow \mathcal{D}$  is a  $\mathcal{V}$ -functor, then we define a  $\mathcal{W}$ -functor  $T_*F: T_*\mathcal{C} \rightarrow T_*\mathcal{D}$  to act on objects by  $c \in \mathcal{C} \mapsto Fc \in \mathcal{D}$  and with internal action on arrows defined by

$$T\mathcal{C}(x, y) \xrightarrow{TF_{x,y}} T\mathcal{D}(Fx, Fy)$$

Again, a straightforward diagram chase verifies that  $T_*F$  is  $\mathcal{W}$ -functorial. It is evident from this definition that  $T_*(GF) = T_*G \cdot T_*F$ .

Finally, let  $\alpha: F \Rightarrow G$  be a  $\mathcal{V}$ -natural transformation between  $\mathcal{V}$ -functors  $F, G: \mathcal{C} \rightarrow \mathcal{D}$  and define a  $\mathcal{W}$ -natural transformation  $T_*\alpha: T_*F \Rightarrow T_*G$  to have components

$$1 \xrightarrow{\cong} T1 \xrightarrow{T\alpha_c} T\mathcal{D}(Fc, Gc)$$

Another straightforward diagram chase verifies that  $T_*\alpha$  is  $\mathcal{W}$ -natural.

It remains to verify this assignment is functorial for both horizontal and vertical composition of enriched natural transformations. The component of  $T_*(\beta \cdot \alpha)$  is defined by the top-horizontal composite below while the component of the vertical composite of  $T_*\alpha$  with  $T_*\beta: T_*G \Rightarrow T_*H$  is defined by the bottom composite:

$$\begin{array}{ccccc}
1 & \xrightarrow{\cong} & T1 & \xrightarrow{T(\beta_c \times \alpha_c)} & T(\mathcal{D}(Gc, Hc) \times \mathcal{D}(Fc, Gc)) \xrightarrow{T\circ} T\mathcal{D}(Fc, Hc) \\
& \searrow \cong & \uparrow \cong & & \uparrow \cong \\
& & T1 \times T1 & \xrightarrow{T\beta_c \times T\alpha_c} & T\mathcal{D}(Gc, Hc) \times T\mathcal{D}(Fc, Gc)
\end{array}$$

The square commutes by the naturality of the isomorphism  $T(u \times v) \cong Tu \times Tv$ , while the triangle commutes because 1 is terminal, so the inverses of the displayed isomorphisms form a commutative triangle. The argument for functoriality of horizontal composites is similar.  $\square$

REMARK 1.5.5. *In fact, the “change of base” procedure  $\mathcal{V} \mapsto \mathcal{V}\text{-Cat}$  is itself a 2-functor from the 2-category of cartesian closed categories, finite-product-preserving functors, and natural transformations to the 2-category of 2-categories, 2-functors, and 2-natural transformations. See [Cru08, §4.3] for a discussion and proof.*

As an immediate consequence of the 2-functoriality of Remark 1.5.5:

PROPOSITION 1.5.6. *Any adjunction between cartesian closed categories whose left adjoint preserves finite products induces a change-of-base 2-adjunction*

$$\mathcal{V} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathcal{W} \quad \rightsquigarrow \quad \mathcal{V}\text{-Cat} \begin{array}{c} \xrightarrow{F_*} \\ \perp \\ \xleftarrow{U_*} \end{array} \mathcal{W}\text{-Cat}$$

PROOF. Of course right adjoints always preserve products, so the adjoint pair of functors  $F \dashv U$  defines an adjunction in the 2-category of cartesian closed categories and finite-product-preserving functors described in Remark 1.5.5. The 2-functor  $\mathcal{V} \mapsto \mathcal{V}\text{-Cat}$  then carries the adjunction displayed on the left to the adjunction displayed on the right.  $\square$

As a special case we have a free-forgetful adjunction between  $\mathcal{C}at$  and  $\mathcal{V}\text{-Cat}$ . Some pieces of this are in the enriched categories folder of Mathlib.

COROLLARY 1.5.7. *For any cartesian closed category  $\mathcal{V}$  with coproducts, the underlying category construction and free category construction define adjoint 2-functors*

$$\mathcal{C}at \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{(-)_0} \end{array} \mathcal{V}\text{-Cat}$$

In light of Proposition 1.5.6 and results to follow, an adjunction between cartesian closed categories whose left adjoint preserves finite products provides a **change-of-base adjunction**. While Proposition 1.5.6 permits the change of base along either adjoint of a finite-product-preserving adjunction, the next series of results reveal that change of base along the right adjoint is somewhat better behaved.

LEMMA 1.5.8. *Any adjunction comprised of finite-product-preserving functors between cartesian closed categories*

$$\mathcal{V} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathcal{W} \quad \rightsquigarrow \quad \mathcal{V} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} U_*\mathcal{W}$$

*defines a  $\mathcal{V}$ -enriched adjunction between the  $\mathcal{V}$ -categories  $\mathcal{V}$  and  $U_*\mathcal{W}$ ; i.e., there exists a  $\mathcal{V}$ -natural isomorphism  $UW(Fv, w) \cong \mathcal{V}(v, Uw)$ .*

PROOF. The internal action  $U_{a,b}: U\mathcal{W}(a,b) \rightarrow \mathcal{V}(Ua, Ub)$  of the  $\mathcal{V}$ -functor  $U: U_*\mathcal{W} \rightarrow \mathcal{V}$  is defined by the transpose of the map  $U\text{ev}: U\mathcal{W}(a,b) \times Ua \rightarrow Ub$  defined by applying  $U$  to the counit of the cartesian closure adjunction of  $\mathcal{W}$ . The  $\mathcal{V}$ -functoriality of this map follows from naturality of evaluation in a cartesian closed category.

By the  $\mathcal{V}$ -functoriality of  $U: U_*\mathcal{W} \rightarrow \mathcal{V}$ , the map

$$U\mathcal{W}(Fv, w) \xrightarrow{U_{Fv,w}} \mathcal{V}(UFv, Uw) \xrightarrow{-\circ\eta_v} \mathcal{V}(v, Uw)$$

is  $\mathcal{V}$ -natural in  $w \in U_*\mathcal{W}$  for all  $v \in \mathcal{V}$ . By a general result about enriching adjoints, to construct a compatible  $\mathcal{V}$ -enrichment of  $F$ , we need only demonstrate that this map is an isomorphism in  $\mathcal{V}$ .

We do this by constructing an explicit inverse, namely

$$\mathcal{V}(v, Uw) \xrightarrow{\eta} UF\mathcal{V}(v, Uw) \xrightarrow{U(Fv, Uw)} U\mathcal{W}(Fv, FUw) \xrightarrow{\epsilon_w \circ -} U\mathcal{W}(Fv, w)$$

where the middle map is defined by applying the unenriched functor  $U$  to the action map from the  $\mathcal{W}$ -functor  $F: F_*\mathcal{V} \rightarrow \mathcal{W}$ , which is defined similarly to the  $\mathcal{V}$ -functor  $U: U_*\mathcal{W} \rightarrow \mathcal{V}$ .

The proof that these maps are inverses involves a pair of diagram chases, the first of which demonstrates that the top-right composite reduces to the left-bottom composite, which is the identity:

$$\begin{array}{ccccccc} \mathcal{V}(v, Uw) & \xrightarrow{\eta} & UF\mathcal{V}(v, Uw) & \xrightarrow{U(Fv, Uw)} & U\mathcal{W}(Fv, FUw) & \xrightarrow{\epsilon_w \circ -} & U\mathcal{W}(Fv, w) \\ & & & \searrow U_{Fv, Uw} & \downarrow U_{Fv, FUw} & & \downarrow U_{Fv, w} \\ & & & & \mathcal{V}(UFv, UFUw) & \xrightarrow{U\epsilon_w \circ -} & \mathcal{V}(UFv, Uw) \\ & \searrow \eta_{Uw \circ -} & & & \downarrow -\circ\eta_v & & \downarrow -\circ\eta_v \\ & & & & \mathcal{V}(v, UFUw) & \xrightarrow{U\epsilon_w \circ -} & \mathcal{V}(v, Uw) \end{array}$$

The only subtle point is the commutativity of the trapezoidal region, which expresses the fact that  $\eta: \text{id}_{\mathcal{V}} \Rightarrow UF$  is a *closed natural transformation* between product-preserving functors between cartesian closed categories. This region commutes because the transposed diagram does:

$$\begin{array}{ccccc} \mathcal{V}(v, Uw) \times v & \xlongequal{\quad} & \mathcal{V}(v, Uw) \times v & \xrightarrow{\text{ev}} & Uw \\ \eta \times \eta_v \downarrow & & \eta \downarrow & & \downarrow \eta_{Uw} \\ UF\mathcal{V}(v, Uw) \times UFv & \xrightarrow{\cong} & UF(\mathcal{V}(v, Uw) \times v) & \xrightarrow{UF\text{ev}} & UFUw \end{array}$$

the right-hand square by naturality, and the left-hand square because any natural transformation between product-preserving functors is automatically a monoidal natural transformation. The other diagram chase is similar.  $\square$

PROPOSITION 1.5.9. *Given an adjunction between cartesian closed categories*

$$\mathcal{V} \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{U} \end{array} \mathcal{W}$$

whose left adjoint preserves finite products then if  $\mathcal{C}$  is co/tensored as a  $\mathcal{W}$ -category,  $U_*\mathcal{C}$  is co/tensored as  $\mathcal{V}$ -category with the co/tensor of  $c \in \mathcal{C}$  by  $v \in \mathcal{V}$  defined by

$$v \otimes c := Fv \otimes c \quad \text{and} \quad c^v := c^{Fv}.$$

PROOF. Suppose  $\mathcal{C}$  admits cotensors as a  $\mathcal{W}$ -category. To verify that  $U_*\mathcal{C}$  admits cotensors as a  $\mathcal{V}$ -category we must supply an isomorphism

$$U\mathcal{C}(x, c^{Fv}) \cong (U\mathcal{C}(x, c))^v$$

in  $\mathcal{V}$  that is  $\mathcal{V}$ -natural in  $x$ . By the enriched Yoneda lemma, we can extract this isomorphism from an isomorphism

$$\mathcal{V}(u, U\mathcal{C}(x, c^{Fv})) \cong \mathcal{V}(u, (U\mathcal{C}(x, c))^v)$$

that is  $\mathcal{V}$ -natural in  $u \in \mathcal{V}$ . To that end, by composing the  $\mathcal{V}$ -natural isomorphisms of Lemma 1.5.8, the enriched natural isomorphisms arising from the cartesian closed structure on  $\mathcal{V}$  and on  $U_*\mathcal{W}$ , and the isomorphisms that characterize the cotensor on  $\mathcal{C}$  and express the fact that  $F$  preserves binary products, we have:

$$\begin{aligned} \mathcal{V}(u, U\mathcal{C}(x, c^{Fv})) &\cong U\mathcal{W}(Fu, \mathcal{C}(x, c^{Fv})) \cong U\mathcal{W}(Fu, \mathcal{C}(x, c)^{Fv}) \\ &\cong U\mathcal{W}(Fu \times Fv, \mathcal{C}(x, c)) \cong U\mathcal{W}(F(u \times v), \mathcal{C}(x, c)) \\ &\cong \mathcal{V}(u \times v, U\mathcal{C}(x, c)) \cong \mathcal{V}(u, (U\mathcal{C}(x, c))^v). \end{aligned} \quad \square$$

The general theory of change-of-base will be applied in the following case the next section.

EXAMPLE 1.5.10. Both adjoints of the adjunction

$$sSet \begin{array}{c} \xleftarrow{\quad h \quad} \\ \perp \\ \xrightarrow{\quad} \end{array} Cat$$

of Proposition 1.2.26 preserve finite products. Hence, Proposition 1.5.6 induces a change-of-base adjunction defined by the 2-functors

$$sCat \begin{array}{c} \xleftarrow{\quad h_* \quad} \\ \perp \\ \xrightarrow{\quad} \end{array} 2-Cat$$

that act identically on objects and act by applying the homotopy category functor or nerve functor, respectively, on homs. The right adjoint, which builds a simplicially enriched category from a 2-category, respects the underlying category: the underlying category of objects and 1-cells is identified with the underlying category of objects and 0-arrows. In this case, the functor  $h: sSet \rightarrow Cat$  commutes with the underlying set functors, so in fact both adjoints preserve underlying categories, as is evident from direct computation. In particular, the homotopy 2-category of an  $\infty$ -cosmos has the same underlying 1-category. Since the nerve embedding is fully faithful, 2-categories can be identified as a full subcategory comprised of those simplicial categories whose hom spaces are nerves of categories.

## 1.6. The homotopy 2-category

Small 1-categories define the objects of a strict 2-category  $Cat$  of categories, functors, and natural transformations. Many basic categorical notions — those defined in terms of categories, functors, and natural transformations — can be defined internally to the 2-category  $Cat$ . This suggests a natural avenue for generalization: reinterpreting these same definitions in a generic 2-category using its objects in place of small categories, its 1-cells in place of functors, and its 2-cells in place of natural transformations.

A significant portion of the theory of  $\infty$ -categories in any fixed  $\infty$ -cosmos can be developed by following exactly this outline, working internally to a 2-category that we refer to as the *homotopy 2-category* that we associate to any  $\infty$ -cosmos. The homotopy 2-category of an  $\infty$ -cosmos is a quotient of the full  $\infty$ -cosmos, replacing each quasi-categorical functor space by its homotopy category. Surprisingly, this rather destructive quotienting operation preserves quite a lot of information. This



said, we caution the reader against becoming overly seduced by homotopy 2-categories, which are more of a technical convenience for reducing the complexity of our arguments than a fundamental notion of  $\infty$ -category theory.

Paralleling our discussion of simplicial categories in Definition 1.3.1 and Digression 1.3.5, there are two perspectives on the notion of a 2-category, which can be understood equally as:

- (i) “two-dimensional” categories, with objects; **1-cells**, whose boundary are given by a pair of objects; and **2-cells**, whose boundary are given by a parallel pair of 1-cells between a pair of objects — together with partially defined composition operations governed by this boundary data
- (ii) or as categories enriched over  $\mathcal{C}at$ .

Both notions exist in Mathlib in some form. The notion `i` is called a *strict bicategory* and is defined as a special case of a bicategory, in which the associators and unitors are identities (converted into 2-cells). The general notion of enriched category can be specialized to the case of enriching over the cartesian monoidal category of categories, but the connection between these notions remains to be explored.

**PROPOSITION 1.6.1.** *There is an equivalence between categories enriched in categories and strict bicategories. In particular, each can be converted into the other.*

The homotopy 2-category is most efficiently defined as a category enriched in  $\mathcal{C}at$  by applying the theory of change-of-base developed in §1.5. The homotopy 2-category for the  $\infty$ -cosmos of quasi-categories was first introduced by Joyal in his work on the foundations of quasi-category theory [Joy08].

**DEFINITION 1.6.2** (homotopy 2-category). Let  $\mathcal{K}$  be an  $\infty$ -cosmos. Its **homotopy 2-category** is the 2-category  $\mathfrak{h}\mathcal{K}$  whose

- objects are the objects  $A, B$  of  $\mathcal{K}$ , i.e., the  $\infty$ -categories;
- 1-cells  $f: A \rightarrow B$  are the 0-arrows in the functor space  $\mathrm{Fun}(A, B)$ , i.e., the  $\infty$ -functors;
- and
- 2-cells

$$A \begin{array}{c} \xrightarrow{f} \\ \Downarrow \alpha \\ \xrightarrow{g} \end{array} B$$

are homotopy classes of 1-simplices in  $\mathrm{Fun}(A, B)$ , which we call  **$\infty$ -natural transformations**.

Put another way  $\mathfrak{h}\mathcal{K}$  is the 2-category with the same objects as  $\mathcal{K}$  and with hom-categories defined by

$$\mathfrak{h}\mathrm{Fun}(A, B) := \mathfrak{h}(\mathrm{Fun}(A, B)),$$

that is,  $\mathfrak{h}\mathrm{Fun}(A, B)$  is the homotopy category of the quasi-category  $\mathrm{Fun}(A, B)$ .

**DEFINITION 1.6.3** (underlying category of a 2-category). The **underlying category** of a 2-category is defined by simply forgetting its 2-cells. Note that an  $\infty$ -cosmos  $\mathcal{K}$  and its homotopy 2-category  $\mathfrak{h}\mathcal{K}$  share the same underlying category  $\mathcal{K}_0$  of  $\infty$ -categories and  $\infty$ -functors in  $\mathcal{K}$ .

**LEMMA 1.6.4.** *The underlying category of the homotopy 2-category of an  $\infty$ -cosmos is isomorphic to the underlying category of the  $\infty$ -cosmos.*

We elaborate on the connection between data in the homotopy 2-category and data in the  $\infty$ -cosmos.

LEMMA 1.6.5.

(i) *Every 2-cell*

$$A \begin{array}{c} \xrightarrow{f} \\ \Downarrow \alpha \\ \xrightarrow{g} \end{array} B$$

in the homotopy 2-category of an  $\infty$ -cosmos is represented by a map of quasi-categories as below-left or equivalently by a functor as below-right

$$\begin{array}{ccc} & \mathbb{1} + \mathbb{1} & \\ \swarrow & & \searrow (f,g) \\ 2 & \xrightarrow{\alpha} & \mathbf{Fun}(A, B) \end{array} \quad \Leftrightarrow \quad \begin{array}{ccc} A & \xrightarrow{\ulcorner \alpha \urcorner} & B^2 \\ (g,f) \searrow & & \swarrow (p_1, p_0) \\ & B \times B & \end{array}$$

and two such maps represent the same 2-cell if and only if they are homotopic as 1-simplices in  $\mathbf{Fun}(A, B)$ .

(ii) *Every invertible 2-cell*

$$A \begin{array}{c} \xrightarrow{f} \\ \Downarrow \cong \alpha \\ \xrightarrow{g} \end{array} B$$

in the homotopy 2-category of an  $\infty$ -cosmos is represented by a map of quasi-categories as below-left or equivalently by a functor as below-right

$$\begin{array}{ccc} & \mathbb{1} + \mathbb{1} & \\ \swarrow & & \searrow (f,g) \\ \mathbb{1} & \xrightarrow{\alpha} & \mathbf{Fun}(A, B) \end{array} \quad \Leftrightarrow \quad \begin{array}{ccc} A & \xrightarrow{\ulcorner \alpha \urcorner} & B^{\mathbb{1}} \\ (g,f) \searrow & & \swarrow (p_1, p_0) \\ & B \times B & \end{array}$$

and two such maps represent the same invertible 2-cell if and only if their common restrictions along  $2 \hookrightarrow \mathbb{1}$  are homotopic as 1-simplices in  $\mathbf{Fun}(A, B)$ .

The notion of homotopic 1-simplices referenced here is defined in Lemma 1.2.20. Since the 2-cells in the homotopy 2-category are referred to as  $\infty$ -natural transformations, we refer to the invertible 2-cells in the homotopy 2-category as  **$\infty$ -natural isomorphisms**.

PROOF. The statement i records the definition of the 2-cells in the homotopy 2-category and the universal property (1.3.7) of the simplicial cotensor. For ii, a 2-cell in the homotopy 2-category is **invertible** if and only if it defines an isomorphism in the appropriate hom-category  $\mathbf{hFun}(A, B)$ . By Proposition 1.2.32 it follows that each invertible 2-cell  $\alpha$  is represented by a homotopy coherent isomorphism  $\alpha: \mathbb{1} \rightarrow \mathbf{Fun}(A, B)$ , which similarly internalizes to define a functor  $\ulcorner \alpha \urcorner: A \rightarrow B^{\mathbb{1}}$ .  $\square$

We now begin to relate the simplicially enriched structures of an  $\infty$ -cosmos to the 2-categorical structures in its homotopy 2-category by proving that homotopy 2-categories inherit products from their  $\infty$ -cosmoi that satisfy a 2-categorical universal property. To illustrate, recall that the terminal  $\infty$ -category  $1 \in \mathcal{K}$  has the universal property  $\mathbf{Fun}(X, 1) \cong \mathbb{1}$  for all  $X \in \mathcal{K}$ . Applying the homotopy category functor we see that  $1 \in \mathbf{h}\mathcal{K}$  has the universal property  $\mathbf{hFun}(X, 1) \cong \mathbb{1}$  for all  $X \in \mathbf{h}\mathcal{K}$ , which is expressed by saying that the  $\infty$ -category 1 defines a **2-terminal object** in the homotopy 2-category. This 2-categorical universal property has both a 1-dimensional and a 2-dimensional aspect. Since  $\mathbf{hFun}(X, 1) \cong \mathbb{1}$  is a category with a single object, there exists a unique morphism  $X \rightarrow 1$  in  $\mathcal{K}$ , and since  $\mathbf{hFun}(X, 1) \cong \mathbb{1}$  has only a single morphism, the only 2-cells in  $\mathbf{h}\mathcal{K}$  with codomain 1 are identities.

PROPOSITION 1.6.6 (cartesian (closure)).

- (i) *The homotopy 2-category of any  $\infty$ -cosmos has 2-categorical products.*
- (ii) *The homotopy 2-category of a cartesian closed  $\infty$ -cosmos is cartesian closed as a 2-category.*

PROOF. While the functor  $h: sSet \rightarrow Cat$  only preserves finite products, the restricted functor  $h: QCat \rightarrow Cat$  preserves *all* products on account of the simplified description of the homotopy category of a quasi-category given in Lemma 1.2.28. Thus for any set  $I$  and family of  $\infty$ -categories  $(A_i)_{i \in I}$  in  $\mathcal{K}$ , the homotopy category functor carries the isomorphism of functor spaces to an isomorphism of hom-categories

$$\mathrm{Fun}(X, \prod_{i \in I} A_i) \xrightarrow{\cong} \prod_{i \in I} \mathrm{Fun}(X, A_i) \xrightarrow{h} \mathrm{hFun}(X, \prod_{i \in I} A_i) \xrightarrow{\cong} \prod_{i \in I} \mathrm{hFun}(X, A_i).$$

This proves that the homotopy 2-category  $h\mathcal{K}$  has products whose universal properties have both a 1- and 2-dimensional component, as described in the empty case for terminal objects above.

If  $\mathcal{K}$  is a cartesian closed  $\infty$ -cosmos, then for any triple of  $\infty$ -categories  $A, B, C \in \mathcal{K}$  there exist exponential objects  $C^A, C^B \in \mathcal{K}$  characterized by natural isomorphisms

$$\mathrm{Fun}(A \times B, C) \cong \mathrm{Fun}(A, C^B) \cong \mathrm{Fun}(B, C^A).$$

Passing to homotopy categories we have natural isomorphisms

$$\mathrm{hFun}(A \times B, C) \cong \mathrm{hFun}(A, C^B) \cong \mathrm{hFun}(B, C^A),$$

which demonstrates that  $h\mathcal{K}$  is cartesian closed as a 2-category: functors  $A \times B \rightarrow C$  transpose to define functors  $A \rightarrow C^B$  and  $B \rightarrow C^A$ , and natural transformations transpose similarly.  $\square$

There is a standard definition of *isomorphism* between two objects in any 1-category, preserved by any functor. Similarly, there is a standard definition of *equivalence* between two objects in any 2-category, preserved by any 2-functor:

DEFINITION 1.6.7 (equivalence). An **equivalence** in a 2-category is given by

- a pair of objects  $A$  and  $B$ ;
- a pair of 1-cells  $f: A \rightarrow B$  and  $g: B \rightarrow A$ ; and
- a pair of invertible 2-cells

$$A \begin{array}{c} \xrightarrow{\quad} \\ \cong \Downarrow \alpha \\ \xrightarrow{\quad} \end{array} A \quad \text{and} \quad B \begin{array}{c} \xrightarrow{\quad} \\ \cong \Downarrow \beta \\ \xrightarrow{\quad} \end{array} B$$

When  $A$  and  $B$  are **equivalent**, we write  $A \simeq B$  and refer to the 1-cells  $f$  and  $g$  as **equivalences**, denoted by “ $\xrightarrow{\sim}$ .”

In the case of the homotopy 2-category of an  $\infty$ -cosmos we have a competing definition of equivalence from 1.4.1: namely a 1-cell  $f: A \xrightarrow{\sim} B$  that induces an equivalence  $f_*: \mathrm{Fun}(X, A) \xrightarrow{\sim} \mathrm{Fun}(X, B)$  on functor spaces — or equivalently, by Lemma 1.4.9, a homotopy equivalence defined relative to the interval  $\mathbb{I}$ . Crucially, all three notions of equivalence coincide:

THEOREM 1.6.8 (equivalences are equivalences). *In any  $\infty$ -cosmos  $\mathcal{K}$ , the following are equivalent and characterize what it means for a functor  $f: A \rightarrow B$  between  $\infty$ -categories to define an **equivalence**.*

- (i) *For all  $X \in \mathcal{K}$ , the post-composition map  $f_*: \mathrm{Fun}(X, A) \xrightarrow{\sim} \mathrm{Fun}(X, B)$  defines an equivalence of quasi-categories.*

- (ii) *There exists a functor  $g: B \rightarrow A$  and natural isomorphisms  $\alpha: \text{id}_A \cong gf$  and  $\beta: fg \cong \text{id}_B$  in the homotopy 2-category.*
- (iii) *There exists a functor  $g: B \rightarrow A$  and maps*

$$\begin{array}{ccc}
 & A & \\
 \nearrow & \uparrow \sim \text{ev}_0 & \\
 A & \xrightarrow{\alpha} A^\natural & \\
 \searrow & \downarrow \sim \text{ev}_1 & \\
 & A &
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 & B & \\
 \nearrow fg & \uparrow \sim \text{ev}_0 & \\
 B & \xrightarrow{\beta} B^\natural & \\
 \searrow & \downarrow \sim \text{ev}_1 & \\
 & B &
 \end{array}$$

*in the  $\infty$ -cosmos  $\mathcal{K}$ .*

As an illustrative comparison of 2-categorical and quasi-categorical techniques, rather than appealing to Lemma 1.4.9 to prove  $\text{i} \Leftrightarrow \text{iii}$ , we re-prove it.

PROOF. For  $\text{i} \Rightarrow \text{ii}$ , if the induced map  $f_*: \text{Fun}(X, A) \xrightarrow{\sim} \text{Fun}(X, B)$  defines an equivalence of quasi-categories then the functor  $f_*: \text{hFun}(X, A) \xrightarrow{\sim} \text{hFun}(X, B)$  defines an equivalence of categories, by Lemma 1.2.36. In particular, the equivalence  $f_*: \text{hFun}(B, A) \xrightarrow{\sim} \text{hFun}(B, B)$  is essentially surjective so there exists  $g \in \text{hFun}(B, A)$  and an isomorphism  $\beta: fg \cong \text{id}_B \in \text{hFun}(B, B)$ . Now since  $f_*: \text{hFun}(A, A) \xrightarrow{\sim} \text{hFun}(A, B)$  is fully faithful, the isomorphism  $\beta f: fgf \cong f \in \text{hFun}(A, B)$  can be lifted to define an isomorphism  $\alpha^{-1}: gf \cong \text{id}_A \in \text{hFun}(A, A)$ . This defines the data of a 2-categorical equivalence in Definition 1.6.7.

To see that  $\text{ii} \Rightarrow \text{iii}$  recall from Lemma 1.6.5 that the natural isomorphisms  $\alpha: \text{id}_A \cong gf$  and  $\beta: fg \cong \text{id}_B$  in  $\text{h}\mathcal{K}$  are represented by maps  $\alpha: A \rightarrow A^\natural$  and  $\beta: B \rightarrow B^\natural$  in  $\mathcal{K}$  as in Lemma 1.4.9.

Finally,  $\text{iii} \Rightarrow \text{i}$  since  $\text{Fun}(X, -)$  carries the data of  $\text{iii}$  to the data of an equivalence of quasi-categories as in Definition 1.2.35.  $\square$

It is hard to overstate the importance of Theorem 1.6.8 for the work that follows. The categorical constructions that we introduce for  $\infty$ -categories,  $\infty$ -functors, and  $\infty$ -natural transformations are invariant under 2-categorical equivalence in the homotopy 2-category and the universal properties we develop similarly characterize 2-categorical equivalence classes of  $\infty$ -categories. Theorem 1.6.8 then asserts that such constructions are “homotopically correct”: both invariant under equivalence in the  $\infty$ -cosmos and precisely identifying equivalence classes of objects.

The equivalence invariance of the functor space in the codomain variable is axiomatic, but equivalence invariance in the domain variable is not.<sup>18</sup> Nor is it evident how this could be proven from either  $\text{i}$  or  $\text{iii}$  of Theorem 1.6.8. But using  $\text{ii}$  and 2-categorical techniques, there is now a short proof.

COROLLARY 1.6.9. *Equivalences of  $\infty$ -categories  $A' \xrightarrow{\sim} A$  and  $B \xrightarrow{\sim} B'$  induce an equivalence of functor spaces  $\text{Fun}(A, B) \xrightarrow{\sim} \text{Fun}(A', B')$ .*

PROOF. The representable simplicial functors  $\text{Fun}(A, -): \mathcal{K} \rightarrow \mathcal{QCat}$  and  $\text{Fun}(-, B): \mathcal{K}^{\text{op}} \rightarrow \mathcal{QCat}$  induce 2-functors  $\text{Fun}(A, -): \text{h}\mathcal{K} \rightarrow \text{h}\mathcal{QCat}$  and  $\text{Fun}(-, B): \text{h}\mathcal{K}^{\text{op}} \rightarrow \text{h}\mathcal{QCat}$ , which preserve the 2-categorical equivalences of Definition 1.6.7. By Theorem 1.6.8 this is what we wanted to show.  $\square$

<sup>18</sup>The functor  $\text{Fun}(A, -)$  is a *cosmological functor*, preserving all of the structure of Definition 1.4.1. Cosmological functors then preserve a large class of cosmological notions, including equivalences. These results, however, do not apply to  $\text{Fun}(-, B)$  since this functor is not cosmological.

There is also a standard 2-categorical notion of an isofibration, defined in the statement of Proposition 1.6.10. We now show that any isofibration in an  $\infty$ -cosmos defines an isofibration in its homotopy 2-category.

PROPOSITION 1.6.10 (isofibrations are isofibrations). *An isofibration  $p: E \rightarrowtail B$  in an  $\infty$ -cosmos  $\mathcal{K}$  also defines an **isofibration** in the homotopy 2-category  $\mathfrak{h}\mathcal{K}$ : given any invertible 2-cell as displayed below-left abutting to  $B$  with a specified lift of one of its boundary 1-cells through  $p$ , there exists an invertible 2-cell abutting to  $E$  with this boundary 1-cell as displayed below-right that whiskers with  $p$  to the original 2-cell.*

$$\begin{array}{ccc} X & \xrightarrow{e} & E \\ & \cong \Downarrow \beta & \downarrow p \\ & & B \end{array} \quad \begin{array}{c} \text{= } \end{array} \quad \begin{array}{ccc} X & \xrightarrow{e} & E \\ & \cong \Downarrow \gamma & \downarrow p \\ & & B \end{array}$$

PROOF. The universal property of the statement says that the functor

$$p_*: \mathfrak{h}\mathrm{Fun}(X, E) \rightarrowtail \mathfrak{h}\mathrm{Fun}(X, B)$$

is an isofibration of categories in the sense defined in Proposition 1.4.7. By axiom 1.4.1ii, since  $p: E \rightarrowtail B$  is an isofibration in  $\mathcal{K}$ , the induced map  $p_*: \mathrm{Fun}(X, E) \rightarrowtail \mathrm{Fun}(X, B)$  is an isofibration of quasi-categories. So it suffices to show that the functor  $\mathfrak{h}: \mathcal{QCat} \rightarrow \mathcal{Cat}$  carries isofibrations of quasi-categories to isofibrations of categories.

So let us now consider an isofibration  $p: E \rightarrowtail B$  between quasi-categories. By Proposition 1.2.32, every isomorphism  $\beta$  in the homotopy category  $\mathfrak{h}B$  of the quasi-category  $B$  is represented by a simplicial map  $\beta: \mathbb{1} \rightarrow B$ . By Definition 1.2.34, the lifting problem

$$\begin{array}{ccc} \mathbb{1} & \xrightarrow{e} & E \\ \downarrow & \nearrow \gamma & \downarrow p \\ \mathbb{1} & \xrightarrow{\beta} & B \end{array}$$

can be solved, and the map  $\gamma: \mathbb{1} \rightarrow E$  so produced represents a lift of the isomorphism from  $\mathfrak{h}B$  to an isomorphism in  $\mathfrak{h}E$  with domain  $e$ .  $\square$

CONVENTION 1.6.11 (on isofibrations in homotopy 2-categories). Since the converse to Proposition 1.6.10 does not hold, there is a potential ambiguity when using the term “isofibration” to refer to a map in the homotopy 2-category of an  $\infty$ -cosmos. We adopt the convention that when we declare a map in  $\mathfrak{h}\mathcal{K}$  to be an isofibration we always mean this is the stronger sense of defining an isofibration in  $\mathcal{K}$ . This stronger condition gives us access to the 2-categorical lifting property of Proposition 1.6.10 and also to homotopical properties axiomatized in Definition 1.4.1, which ensure that the strictly defined limits of 1.4.1i are automatically equivalence invariant constructions (see [RV22, 6.2.8, §C.1]).

We conclude this chapter with a final definition that can be extracted from the homotopy 2-category of an  $\infty$ -cosmos. The 1- and 2-cells in the homotopy 2-category from the terminal  $\infty$ -category  $1 \in \mathcal{K}$  to a generic  $\infty$ -category  $A \in \mathcal{K}$  define the objects and morphisms in the homotopy category of the  $\infty$ -category  $A$ .

DEFINITION 1.6.12 (homotopy category of an  $\infty$ -category). The **homotopy category** of an  $\infty$ -category  $A$  in an  $\infty$ -cosmos  $\mathcal{K}$  is defined to be the homotopy category of its underlying quasi-category, that is:

$$\mathbf{h}A := \mathbf{h}\mathrm{Fun}(1, A) := \mathbf{h}(\mathrm{Fun}(1, A)).$$

As we shall discover, homotopy categories generally inherit “derived” analogues of structures present at the level of  $\infty$ -categories.

## Bibliography

- [Bro73] K. S. Brown. Abstract homotopy theory and generalized sheaf cohomology. *Trans. Amer. Math. Soc.*, 186:419–458, 1973.
- [BV73] J. M. Boardman and R. Vogt. *Homotopy Invariant Algebraic Structures on Topological Spaces*, volume 347 of *Lecture Notes in Mathematics*. Springer-Verlag, 1973.
- [Cru08] G. S. H. Cruttwell. *Normed Spaces and the Change of Base for Enriched Categories*. PhD thesis, Dalhousie University, 2008.
- [EK66] S. Eilenberg and G. M. Kelly. Closed categories. In *Proceedings of the Conference on Categorical Algebra (La Jolla 1965)*, page 421–562. Springer-Verlag, 1966.
- [EML45] S. Eilenberg and S. Mac Lane. General theory of natural equivalences. *Trans. Amer. Math. Soc.*, 58:231–294, 1945.
- [GZ67] P. Gabriel and M. Zisman. *Calculus of Fractions and Homotopy Theory*. Number 35 in *Ergebnisse der Math.* Springer-Verlag, 1967.
- [Joy02] A. Joyal. Quasi-categories and Kan complexes. *J. Pure Appl. Algebra*, 175:207–222, 2002.
- [Joy08] A. Joyal. *The Theory of Quasi-Categories and its Applications*. Quadern 45 vol II. Centre de Recerca Matemàtica Barcelona, 2008.
- [Kel89] G. M. Kelly. Elementary observations on 2-categorical limits. *Bull. Aust. Math. Soc.*, 49:301–317, 1989.
- [Qui67] D. G. Quillen. *Homotopical Algebra*, volume 43 of *Lecture Notes in Mathematics*. Springer-Verlag, 1967.
- [Rie16] E. Riehl. *Category Theory in Context*. Aurora Modern Math Originals. Dover Publications, 2016.
- [RV22] Emily Riehl and Dominic Verity. *Elements of  $\infty$ -Category Theory*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2022.
- [Str74] R. H. Street. Elementary cosmoi I. In *Category Seminar (Proc. Sem., Sydney, 1972/1973)*, volume 420 of *Lecture Notes in Math.*, page 134–180. Springer, 1974.