

CS 6316 Machine Learning

Generative Models

Yangfeng Ji

Department of Computer Science
University of Virginia



ENGINEERING

Basic Definition

Data generation process

An idealized process to illustrate the relations among domain set \mathcal{X} , label set \mathcal{Y} , and the training set S

1. the probability distribution \mathcal{D} over the domain set \mathcal{X}
2. sample an instance $x \in \mathcal{X}$ according to \mathcal{D}
3. annotate it using the labeling function f as $y = f(x)$

[From Lecture 02]

Example

Here is an data generation model

$$p(x) = \underbrace{0.6 \cdot \mathcal{N}(x; \mu_+, \Sigma_+)}_{y=+1} + \underbrace{0.4 \cdot \mathcal{N}(x; \mu_-, \Sigma_-)}_{y=-1} \quad (1)$$

with

- ▶ $\mu_+ = [2, 0]^T$
- ▶ $\Sigma_+ = \begin{bmatrix} 1.0 & 0.8 \\ 0.8 & 2.0 \end{bmatrix}$
- ▶ $\mu_- = [-2, 0]^T$
- ▶ $\Sigma_- = \begin{bmatrix} 2.0 & 0.6 \\ 0.6 & 1.0 \end{bmatrix}$

Example (II)

The data generation model can also be represented with the following components

$$p(y = +1) = 0.6 \quad (2)$$

$$p(y = -1) = 1 - p(y = +1) = 0.4 \quad (3)$$

$$p(\mathbf{x} \mid y = +1) = \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_+, \boldsymbol{\Sigma}_+) \quad (4)$$

$$p(\mathbf{x} \mid y = -1) = \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_-, \boldsymbol{\Sigma}_-) \quad (5)$$

Data Generation

The specific data generation process:
for each data point

1. Randomly select a value of $y \in \{+1, -1\}$ based on

$$p(y = +1) = 0.6 \quad p(y = -1) = 0.4 \quad (6)$$

Data Generation

The specific data generation process:
for each data point

1. Randomly select a value of $y \in \{+1, -1\}$ based on

$$p(y = +1) = 0.6 \quad p(y = -1) = 0.4 \quad (6)$$

2. Sample x from the corresponding component based on the value of y

$$p(x | y) = \begin{cases} \mathcal{N}(x; \mu_+, \Sigma_+) & y = +1 \\ \mathcal{N}(x; \mu_-, \Sigma_-) & y = -1 \end{cases} \quad (7)$$

Data Generation

The specific data generation process:
for each data point

1. Randomly select a value of $y \in \{+1, -1\}$ based on

$$p(y = +1) = 0.6 \quad p(y = -1) = 0.4 \quad (6)$$


2. Sample x from the corresponding component based on the value of y

$$p(x | y) = \begin{cases} \mathcal{N}(x; \mu_+, \Sigma_+) & y = +1 \\ \mathcal{N}(x; \mu_-, \Sigma_-) & y = -1 \end{cases} \quad (7)$$

3. Add (x, y) to S , go to step 1

Illustration

With $N = 1000$ samples, here is the plot



- 588 **positive** samples and 412 **negative** samples

Discriminative Models for Classification

- ▶ Discriminative models directly give predictions on the **target** variable (e.g., y)
- ▶ Example: logistic regression

$$p(y \mid x) = \sigma(y\langle w, x \rangle) = \frac{1}{1 + e^{-y\langle w, x \rangle}} \quad (8)$$

where w is the model parameter

Discriminative Models for Classification

- ▶ Discriminative models directly give predictions on the **target** variable (e.g., y)
- ▶ Example: logistic regression

$$p(y \mid x) = \sigma(y\langle w, x \rangle) = \frac{1}{1 + e^{-y\langle w, x \rangle}} \quad (8)$$

where w is the model parameter

- ▶ Other examples
 - ▶ AdaBoost (lecture 05)
 - ▶ SVMs (lecture 07)
 - ▶ Feed-forward neural network (lecture 08)

Generative Models for Classification

- ▶ Basic idea: Building a classifier by *simulating* the data generation process

Generative Models for Classification

- ▶ Basic idea: Building a classifier by *simulating* the data generation process
- ▶ For the binary classification problem, recall the basic components of the data generation process
 - ▶ $p(y)$ where $y \in \{-1, +1\}$
 - ▶ $p(x \mid y = +1)$ where $x \in \mathbb{R}^d$
 - ▶ $p(x \mid y = -1)$ where $x \in \mathbb{R}^d$

Generative Models for Classification

- ▶ Basic idea: Building a classifier by *simulating* the data generation process
- ▶ For the binary classification problem, recall the basic components of the data generation process
 - ▶ $p(y)$ where $y \in \{-1, +1\}$
 - ▶ $p(x \mid y = +1)$ where $x \in \mathbb{R}^d$
 - ▶ $p(x \mid y = -1)$ where $x \in \mathbb{R}^d$
- ▶ Challenge in machine learning: we do **not** know any of them, instead we have the samples **S** from this distribution
 - ▶ This has always been our assumption in machine learning — we have no idea about the true data distribution

Generative Models for Classification (II)

We use a set of distribution $q(\cdot)$ to approximate the true distribution $p(\cdot)$

| Data Generation Model | Generative Model |
|-----------------------|--------------------|
| $p(y)$ | $q(y)$ |
| $p(x \mid y = +1)$ | $q(x \mid y = +1)$ |
| $p(x \mid y = -1)$ | $q(x \mid y = -1)$ |

Learning with Generative Models

1. Define distributions for all components
2. Estimate the parameters for each component distribution

Defining Distributions

A typical way of defining distributions for generative models is based on *our understanding about the data*

Defining Distributions

A typical way of defining distributions for generative models is based on *our understanding about the data*

- Output domain $y \in \{+1, -1\}$: **Bernoulli** distribution

$$p(y) = \text{Bern}(y; \alpha) = \alpha^{\delta(y=+1)}(1 - \alpha)^{\delta(y=-1)} \quad (9)$$

where $\alpha \in (0, 1)$ is the parameter

Defining Distributions

A typical way of defining distributions for generative models is based on *our understanding about the data*

- ▶ Output domain $y \in \{+1, -1\}$: **Bernoulli** distribution

$$p(y) = \text{Bern}(y; \alpha) = \alpha^{\delta(y=+1)}(1 - \alpha)^{\delta(y=-1)} \quad (9)$$

where $\alpha \in (0, 1)$ is the parameter

- ▶ Input domain $x \in \mathbb{R}^d$: **Gaussian** distribution

$$p(x \mid y = +1) = \mathcal{N}(x; \mu_+, \Sigma_+) \quad (10)$$

where μ_+ and Σ_+ are the parameters

Defining Distributions

A typical way of defining distributions for generative models is based on *our understanding about the data*

- ▶ Output domain $y \in \{+1, -1\}$: **Bernoulli** distribution

$$p(y) = \text{Bern}(y; \alpha) = \alpha^{\delta(y=+1)}(1 - \alpha)^{\delta(y=-1)} \quad (9)$$

where $\alpha \in (0, 1)$ is the parameter

- ▶ Input domain $x \in \mathbb{R}^d$: **Gaussian** distribution

$$p(x \mid y = +1) = \mathcal{N}(x; \mu_+, \Sigma_+) \quad (10)$$

where μ_+ and Σ_+ are the parameters

- ▶ Similarly, for $p(x \mid y = -1)$

$$p(x \mid y = -1) = \mathcal{N}(x; \mu_-, \Sigma_-) \quad (11)$$


where μ_- and Σ_- are the parameters

Parameter Estimation

- ▶ The collection of the parameters

$$\theta = \{\alpha, \mu_+, \Sigma_+, \mu_-, \Sigma_-\} \quad (12)$$

- ▶ Training data $S = \{(x_1, y_1), \dots, (x_m, y_m)\}$




Parameter Estimation

- ▶ The collection of the parameters

$$\theta = \{\alpha, \mu_+, \Sigma_+, \mu_-, \Sigma_-\} \quad (12)$$

- ▶ Training data $S = \{(x_1, y_1), \dots, (x_m, y_m)\}$



- ▶ Learning algorithm: Maximum Likelihood Estimation (MLE)

Maximum Likelihood Estimation (MLE)

MLE defined on the whole distribution $q(x, y)$

$$\theta \leftarrow \operatorname{argmax}_{\theta'} \sum_{i=1}^m \log q(x_i, y_i; \theta') \quad (13)$$

Maximum Likelihood Estimation (MLE)

MLE defined on the whole distribution $q(x, y)$

$$\theta \leftarrow \operatorname{argmax}_{\theta'} \sum_{i=1}^m \log q(x_i, y_i; \theta') \quad (13)$$

Based on the chain rule of probability

$$q(x, y; \theta) = q(y; \alpha)q(x \mid y; \mu_y, \Sigma_y), \quad (14)$$

Maximum Likelihood Estimation (MLE)

MLE defined on the whole distribution $q(x, y)$

$$\theta \leftarrow \operatorname{argmax}_{\theta'} \sum_{i=1}^m \log q(x_i, y_i; \theta') \quad (13)$$

Based on the chain rule of probability

$$q(x, y; \theta) = q(y; \alpha)q(x | y; \mu_y, \Sigma_y), \quad (14)$$

Therefore

$$\hat{\theta} \leftarrow \operatorname{argmax}_{\theta} \left\{ \sum_{i=1}^m \log q(y_i; \alpha) + \sum_{i=1}^m \log q(x_i | y_i; \mu_y, \Sigma_y) \right\}$$

the last item has two components, depending on the value of y

MLE: Bernoulli Distribution

Recall the definition of Bernoulli distribution, we have

$$\sum_{i=1}^m \log q(y_i; \alpha) = \sum_{i=1}^m \{\delta(y_i = +1) \log \alpha + \delta(y_i = -1) \log(1-\alpha)\} \quad (15)$$

MLE: Bernoulli Distribution

Recall the definition of Bernoulli distribution, we have

$$\sum_{i=1}^m \log q(y_i; \alpha) = \sum_{i=1}^m \{ \delta(y_i = +1) \log \alpha + \delta(y_i = -1) \log(1-\alpha) \} \quad (15)$$

Then, the value of α can be estimated from

$$\frac{d \sum_{i=1}^m \log q(y_i; \alpha)}{d\alpha} = \frac{\sum_{i=1}^m \delta(y_i = +1)}{\alpha} - \frac{\sum_{i=1}^m \delta(y_i = -1)}{1-\alpha} = 0 \quad (16)$$

MLE: Bernoulli Distribution

Recall the definition of Bernoulli distribution, we have

$$\sum_{i=1}^m \log q(y_i; \alpha) = \sum_{i=1}^m \{\delta(y_i = +1) \log \alpha + \delta(y_i = -1) \log(1-\alpha)\} \quad (15)$$

Then, the value of α can be estimated from

$$\frac{d \sum_{i=1}^m \log q(y_i; \alpha)}{d\alpha} = \frac{\sum_{i=1}^m \delta(y_i = +1)}{\alpha} - \frac{\sum_{i=1}^m \delta(y_i = -1)}{1-\alpha} = 0 \quad (16)$$

therefore,

$$\alpha = \frac{\sum_{i=1}^m \delta(y_i = +1)}{m} \quad (17)$$

MLE: Gaussian Distribution

The definition of multi-variate Gaussian distribution

$$q(x \mid y; \mu, \Sigma) = \frac{1}{(2\pi)^{d/2} |\Sigma|} \exp \left((x - \mu)^\top \Sigma^{-1} (x - \mu) \right) \quad (18)$$

- For $y = +1$, MLE on μ_+ and Σ_+ will only consider the samples x with $y = +1$ (assume it's S_+)

MLE: Gaussian Distribution

The definition of multi-variate Gaussian distribution

$$q(x \mid y; \mu, \Sigma) = \frac{1}{(2\pi)^{d/2} |\Sigma|} \exp((x - \mu)^\top \Sigma^{-1} (x - \mu)) \quad (18)$$

- ▶ For $y = +1$, MLE on μ_+ and Σ_+ will only consider the samples x with $y = +1$ (assume it's S_+)
- ▶ MLE on μ_+

$$\mu = \frac{1}{|S_+|} \sum_{x_i \in S_+} x_i \quad (19)$$

MLE: Gaussian Distribution

The definition of multi-variate Gaussian distribution

$$q(x \mid y; \mu, \Sigma) = \frac{1}{(2\pi)^{d/2} |\Sigma|} \exp((x - \mu)^\top \Sigma^{-1} (x - \mu)) \quad (18)$$

- ▶ For $y = +1$, MLE on μ_+ and Σ_+ will only consider the samples x with $y = +1$ (assume it's S_+)
- ▶ MLE on μ_+

$$\mu = \frac{1}{|S_+|} \sum_{x_i \in S_+} x_i \quad (19)$$

- ▶ MLE on Σ_+

$$\Sigma_+ = \sum_{x_i \in S_+} (x_i - \mu)(x_i - \mu)^\top \quad (20)$$

MLE: Gaussian Distribution

The definition of multi-variate Gaussian distribution

$$q(x \mid y; \mu, \Sigma) = \frac{1}{(2\pi)^{d/2} |\Sigma|} \exp((x - \mu)^\top \Sigma^{-1} (x - \mu)) \quad (18)$$

- ▶ For $y = +1$, MLE on μ_+ and Σ_+ will only consider the samples x with $y = +1$ (assume it's S_+)
- ▶ MLE on μ_+

$$\mu = \frac{1}{|S_+|} \sum_{x_i \in S_+} x_i \quad (19)$$

- ▶ MLE on Σ_+

$$\Sigma_+ = \sum_{x_i \in S_+} (x_i - \mu)(x_i - \mu)^\top \quad (20)$$

- ▶ *Exercise:* prove equations 19 and 20 with $d = 1$

Example: Parameter Estimation

Given $N = 1000$ samples, here are the parameters

| Parameter | $p(\cdot)$ | $q(\cdot)$ |
|------------|--|--|
| μ_+ | $[2, 0]^\top$ | $[1.95, -0.11]^\top$ |
| Σ_+ | $\begin{bmatrix} 1.0 & 0.8 \\ 0.8 & 2.0 \end{bmatrix}$ | $\begin{bmatrix} 0.88 & 0.74 \\ 0.74 & 1.97 \end{bmatrix}$ |
| μ_- | $[-2, 0]^\top$ | $[-2.08, 0.08]^\top$ |
| Σ_- | $\begin{bmatrix} 2.0 & 0.6 \\ 0.6 & 1.0 \end{bmatrix}$ | $\begin{bmatrix} 1.88 & 0.55 \\ 0.55 & 1.07 \end{bmatrix}$ |

Prediction

- ▶ For a new data point x' , the prediction is given as

$$q(y' | x') = \frac{q(y')q(x | y')}{q(x')} \propto q(y')q(x' | y') \quad (21)$$

No need to compute $q(x')$

Prediction

- ▶ For a new data point x' , the prediction is given as

$$q(y' | x') = \frac{q(y')q(x | y')}{q(x')} \propto q(y')q(x' | y') \quad (21)$$

No need to compute $q(x')$

- ▶ Prediction rule

$$y' = \begin{cases} +1 & q(y' = +1 | x') > q(y' = -1 | x') \\ -1 & q(y' = +1 | x') < q(y' = +1 | x') \end{cases} \quad (22)$$

Prediction

- ▶ For a new data point \mathbf{x}' , the prediction is given as

$$q(y' | \mathbf{x}') = \frac{q(y')q(\mathbf{x} | y')}{q(\mathbf{x}')} \propto q(y')q(\mathbf{x}' | y') \quad (21)$$

No need to compute $q(\mathbf{x}')$

- ▶ Prediction rule

$$y' = \begin{cases} +1 & q(y' = +1 | \mathbf{x}') > q(y' = -1 | \mathbf{x}') \\ -1 & q(y' = +1 | \mathbf{x}') < q(y' = +1 | \mathbf{x}') \end{cases} \quad (22)$$

- ▶ Although equation 22 looks like the one used in the Bayes optimal predictor, the prediction power is limited by

$$q(y' | \mathbf{x}') \approx p(y | \mathbf{x}) \quad (23)$$

Again, we don't know $p(\cdot)$

Naive Bayes Classifiers

Number of Parameters

Assume $\mathbf{x} = (x_{\cdot,1}, \dots, x_{\cdot,d}) \in \mathbb{R}^d$, then the number of parameters in $q(\mathbf{x}, y)$

- ▶ $q(y)$: 1 (α)
- ▶ $q(\mathbf{x} \mid y = +1)$:
 - ▶ $\mu_+ \in \mathbb{R}^d$: d parameters
 - ▶ $\Sigma_+ \in \mathbb{R}^{d \times d}$: d^2 parameters
- ▶ $q(\mathbf{x} \mid y = -1)$: $d^2 + d$ parameters

In total, we have $2d^2 + 2d + 1$ parameters

Challenge of Parameter Estimation

- ▶ When $d = 100$, we have $2d^2 + 2d + 1 = 20201$ parameters
- ▶ A close look about the covariance matrix Σ in a multivariate Gaussian distribution

$$\Sigma = \begin{bmatrix} \sigma_{1,1}^2 & \cdots & \sigma_{1,d}^2 \\ \vdots & \ddots & \vdots \\ \sigma_{d,1}^2 & \cdots & \sigma_{d,d}^2 \end{bmatrix} \quad (24)$$

Challenge of Parameter Estimation

- ▶ When $d = 100$, we have $2d^2 + 2d + 1 = 20201$ parameters
- ▶ A close look about the covariance matrix Σ in a multivariate Gaussian distribution

$$\Sigma = \begin{bmatrix} \sigma_{1,1}^2 & \cdots & \sigma_{1,d}^2 \\ \vdots & \ddots & \vdots \\ \sigma_{d,1}^2 & \cdots & \sigma_{d,d}^2 \end{bmatrix} \quad (24)$$

- ▶ To reduce the number of parameters, we assume

$$\sigma_{i,j} = 0 \quad \text{if } i \neq j \quad (25)$$

Diagonal Covariance Matrix

With the diagonal covariance matrix

$$\Sigma = \begin{bmatrix} \sigma_{1,1}^2 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \sigma_{d,d}^2 \end{bmatrix} \quad (26)$$

Now, the multivariate Gaussian distribution can be rewritten with

- ▶ $|\Sigma| = \prod_{j=1}^d \sigma_{j,j}^2$
- ▶ assume $\mu = 0$ for simplicity

$$(x - \mu)^\top \Sigma^{-1} (x - \mu) = \sum_{j=1}^d \frac{(x_{\cdot,j} - \mu_j)^2}{\sigma_{j,j}^2} \quad (27)$$

Diagonal Covariance Matrix (II)

In other words

$$q(\mathbf{x} \mid y, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{j=1}^d q(x_{\cdot,j} \mid y; \mu_j, \sigma_{j,j}^2) \quad (28)$$

Diagonal Covariance Matrix (II)

In other words

$$q(\mathbf{x} \mid y, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{j=1}^d q(x_{\cdot,j} \mid y; \mu_j, \sigma_{j,j}^2) \quad (28)$$

- **Conditional Independence:** Equation 28 means, given y , each component x_j is independent of other components

Diagonal Covariance Matrix (II)

In other words

$$q(\mathbf{x} \mid y, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{j=1}^d q(x_{\cdot,j} \mid y; \mu_j, \sigma_{j,j}^2) \quad (28)$$

- ▶ **Conditional Independence:** Equation 28 means, given y , each component x_j is independent of other components
- ▶ This is a strong and **naive** assumption about $q(\mathbf{x} \mid \cdot)$

Diagonal Covariance Matrix (II)

In other words

$$q(\mathbf{x} \mid y, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{j=1}^d q(x_{\cdot,j} \mid y; \mu_j, \sigma_{j,j}^2) \quad (28)$$

- ▶ **Conditional Independence:** Equation 28 means, given y , each component x_j is independent of other components
- ▶ This is a strong and **naive** assumption about $q(\mathbf{x} \mid \cdot)$
- ▶ Together with $q(y)$, this generative model is called the **Naive Bayes** classifier

Diagonal Covariance Matrix (II)

In other words

$$q(\mathbf{x} \mid y, \boldsymbol{\mu}, \boldsymbol{\Sigma}) = \prod_{j=1}^d q(x_{\cdot,j} \mid y; \mu_j, \sigma_{j,j}^2) \quad (28)$$

- ▶ **Conditional Independence:** Equation 28 means, given y , each component x_j is independent of other components
- ▶ This is a strong and **naive** assumption about $q(\mathbf{x} \mid \cdot)$
- ▶ Together with $q(y)$, this generative model is called the **Naive Bayes** classifier
- ▶ Parameter estimation can be done **per dimension**

Example: Parameter Estimation

Given $N = 1000$ samples, here are the parameters

| Parameter | $p(\cdot)$ | $q(\cdot)$ | Naive Bayes |
|------------|--|--|--|
| μ_+ | $[2, 0]^\top$ | $[1.95, -0.11]^\top$ | $[1.95, -0.11]^\top$ |
| Σ_+ | $\begin{bmatrix} 1.0 & 0.8 \\ 0.8 & 2.0 \end{bmatrix}$ | $\begin{bmatrix} 0.88 & 0.74 \\ 0.74 & 1.97 \end{bmatrix}$ | $\begin{bmatrix} 0.88 & 0 \\ 0 & 1.97 \end{bmatrix}$ |
| μ_- | $[-2, 0]^\top$ | $[-2.08, 0.08]^\top$ | $[-2.08, 0.08]^\top$ |
| Σ_- | $\begin{bmatrix} 2.0 & 0.6 \\ 0.6 & 1.0 \end{bmatrix}$ | $\begin{bmatrix} 1.88 & 0.55 \\ 0.55 & 1.07 \end{bmatrix}$ | $\begin{bmatrix} 1.88 & 0 \\ 0 & 1.07 \end{bmatrix}$ |

Latent Variable Models

EM Algorithm

Reference



Jurafsky, D. and Martin, J. (2019).
Speech and language processing.