


A Combinatorial Journey to the Challenger Deep of Mathematics


Jan Kurkofka













graph



graph




Social networks




Infrastructure




Maths!



Dodecahedron







Farey graph



Petersen graph

Which graphs can be drawn in the plane so that no two edges cross?
are planar



graph

graph



planar?

graph



planar?



yes!

graph



planar?



yes!



draw it

graph




planar?


yes!



Easy to verify

draw it






Euler's formula (1752)

For every planar drawing of a connected graph with n vertices and m edges:

$$n - m + \ell = 2$$

where ℓ is the number of faces of the drawing.

faces: connected regions of the plane minus the drawing



$$n = 4$$

$$m = 6$$

$$\ell = 4$$



Euler's formula (1752)

For every planar drawing of a connected graph with n vertices and m edges:

$$n - m + \ell = 2$$

where ℓ is the number of faces of the drawing.

faces: connected regions of the plane minus the drawing



Corollary A. Every triangulation of the plane with n vertices has $3n - 6$ edges.

Euler's formula (1752)

For every planar drawing of a connected graph with n vertices and m edges:

$$n - m + \ell = 2$$

where ℓ is the number of faces of the drawing.


faces: connected regions of the plane minus the drawing



Corollary A. Every triangulation of the plane with n vertices has $3n - 6$ edges.


Corollary B. K_5 is not planar.

A graph H is a *subgraph* of a graph G if H can be obtained from G by successively deleting edges or isolated vertices.



Fact. Subgraphs of planar graphs are planar.


A graph H is a *subgraph* of a graph G if H can be obtained from G by successively deleting edges or isolated vertices.



Fact. Subgraphs of planar graphs are planar.

Conjecture. Every nonplanar graph contains K_5 as a subgraph.


A graph H is a **minor** of a graph G if H can be obtained from G by successively deleting edges or isolated vertices or contracting edges.




Fact. **Minors** of planar graphs are planar.

Conjecture. Every nonplanar graph contains K_5 as a **minor**.



Life as a Mathematician





Life as a Mathematician



Life as a Mathematician




Life as a Mathematician




Kuratowski's theorem (1930)

For every graph G , the following assertions are equivalent:

- G is planar;
- G contains neither K_5 nor $K_{3,3}$ as a minor.




K_5




$K_{3,3}$







Is there a Kuratowski-type theorem for the torus?




Is there a planar drawing of K_5 on the torus?



Is there a planar drawing of K_5 on the torus?



Are there Kuratowski-type theorems for other surfaces?



Conjecture (allegedly Wagner, 1960s)

For **every** graph-property \mathcal{P} that is closed under taking minors

(e.g. being planar or admitting a drawing on



Conjecture (allegedly Wagner, 1960s)

For **every** graph-property \mathcal{P} that is closed under taking minors

(e.g. being planar or admitting a drawing on



there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.

minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar

forest



minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless


minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless




minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

???

minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

??? ≥ 157


minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

??? ≥ 157

torus



minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

??? ≥ 157



???


minor-closed graph-property \mathcal{P}

excluded minors X_1, \dots, X_k

planar



forest



linkless



planar after deleting ≤ 1 vertex

??? ≥ 157



??? $\geq 17,523$

Conjecture (allegedly Wagner, 1960s)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



Neil Robertson



Paul Seymour

Conjecture (allegedly Wagner, 1960s)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



1983–2004

Neil Robertson



Paul Seymour

Conjecture (allegedly Wagner, 1960s)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



Neil Robertson

1983–2004
20 papers



Paul Seymour

Conjecture (allegedly Wagner, 1960s)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



Neil Robertson

1983–2004
20 papers
> 500 pages



Paul Seymour

Graph-Minor Theorem (Robertson and Seymour)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



Neil Robertson

1983–2004
20 papers
> 500 pages



Paul Seymour

Graph-Minor Theorem (Robertson and Seymour)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.



Neil Robertson

1983–2004
20 papers
> 500 pages



Paul Seymour

Corollary. For every minor-closed graph-property there exists an efficient (cubic time) algorithm for testing whether a given graph exhibits the property.



Graph-Minor Theorem (Robertson and Seymour)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.

Active research

- Graph-Minor Theorem for matroids (write-up phase)
- Graph-Minor Theorem for directed graphs
- Given an explicit \mathcal{P} , find X_1, \dots, X_k explicitly
- Algorithms to compute X_1, \dots, X_k given \mathcal{P}





Graph-Minor Theorem (Robertson and Seymour)

For **every** minor-closed graph-property \mathcal{P} there exist **finitely many** graphs X_1, \dots, X_k such that the following assertions are equivalent:

- G exhibits the property \mathcal{P} ;
- G contains none of the graphs X_1, \dots, X_k as a minor.

Active research

- Graph-Minor Theorem for matroids (write-up phase)
- Graph-Minor Theorem for directed graphs
- Given an explicit \mathcal{P} , find X_1, \dots, X_k explicitly
- Algorithms to compute X_1, \dots, X_k given \mathcal{P}



Thank you!

All feedback welcome :)

