





Yuanpeng Zhang
Neutron Scattering Division
Dec-14-2022

U.S. DEPARTMENT OF


ORNL is managed by UT-Battelle, LLC for the US Department of Energy








- ❖ Unit cell expanded to 10x10x10 supercell and symmetry constraint removed.
- Modeling were driven purely by experimental data.
- Conventional metropolis Monte Carlo routine used.
- Bragg data not included due to the complexity with average structure. Thus, we only focus on the local scale correlation in real space.
- ❖ Both the reciprocal and real space data were included for the model refinement and the weight for each dataset was adjusted on-the-fly in an automatic manner.
- For doped sample, Ti atoms were replaced with Sn in the expanded supercell in a random manner according to the stoichiometry.
- ❖ On average, ~100 moves/atom were generated, among which ~20 moves/atom were accepted.


## Total scattering data fitting















Overall, the fitting quality is reasonable in both spaces.


## **Partial PDFs**























## **ADPs**




No obvious anisotropy was observed in all samples

## Triplets bond angle distribution







- ❖ As the doping level increases, the distortion of the local TiO6 octahedrons increases.
- Meanwhile, the relative orientation in between those corner sharing TiO6 octahedrons is distorted as well.
- ❖ The Yb pyrochlore sublattice is well kept.
- ❖ Distortion of the local tetrahedral environment formed by Ti(Sn) and Yb is enhanced.
- The Sn-only involved angles do not show that obvious change as compared to those Ti-only involved angles.

- Consistence with other results?
- ❖ Multiple RMC runs to check self-consistenc