

India District-Level Soil Dataset

Soil Fertility, Moisture, pH, Nitrogen & Phosphorus by State and District

This document is formatted as one self-contained descriptive paragraph per district, so it can be directly chunked and indexed by a Retrieval-Augmented Generation (RAG) pipeline. Each paragraph repeats the state and district name along with all soil parameters, so a single retrieved chunk is fully interpretable without needing neighbouring context.

1. What This Dataset Contains

This dataset covers 640 districts across 35 Indian states and union territories (administrative boundaries as per Census of India 2011). For each district it records: dominant Soil Type, Soil Fertility (%), Soil Moisture (%), Soil pH, available Nitrogen (kg/ha), available Phosphorus (kg/ha), and a derived Fertility Class (Low/Medium/High).

2. Data Sources and Method

District and state names are taken from the public Census of India 2011 district list. Note that some states have since been reorganised (for example Telangana was carved out of Andhra Pradesh in 2014, and several new districts have been notified between 2011 and 2026); this dataset reflects 2011-era administrative boundaries.

Soil Type for each district is assigned from its state's dominant soil order as classified by ICAR and the National Bureau of Soil Survey & Land Use Planning (NBSS-LUP) - e.g. Alluvial soils across the Indo-Gangetic plains, Black (Regur) soils across the Deccan trap region, Red and Laterite soils across the peninsular plateau and Western Ghats, Arid soils across Rajasthan, and Mountain/Forest soils across the Himalayan states. This classification is real, published soil science and is not fabricated.

Soil Fertility %, Soil Moisture %, Soil pH, Nitrogen (kg/ha) and Phosphorus (kg/ha) are **indicative regional estimates**. They were generated within realistic value ranges published for each soil type, informed by NBSS-LUP soil survey reports and by publicly reported Soil Health Card (SHC) scheme cycle-wise averages (for example, published SHC case studies report soil pH in the 6.5-7.9 band, available nitrogen in roughly the 100-400 kg/ha band, and available phosphorus in roughly the 5-25 kg/ha band, depending on soil type and region). **These figures are not individual laboratory measurements for each specific district** - genuine, per-district lab data is generated per farm holding under India's Soil Health Card scheme and is not published anywhere as a single clean, bulk, district-summary file that can be reliably scraped in bulk. Presenting invented precision as if it were scraped lab data would be misleading, so every value here is explicitly labelled as an indicative estimate.

For genuine lab-grade, per-farm soil test data, consult: the Soil Health Card portal (soilhealth.dac.gov.in); ICAR-NBSS&LUP district soil survey reports; or your State Agriculture Department / local Krishi Vigyan Kendra.

3. How to Use This With a RAG Pipeline

Section 5 below contains one paragraph per district, grouped by state under clear headings. Each paragraph is self-contained (repeats state, district, and all metrics), making it suitable as a single retrieval chunk (roughly 60-90 tokens each) without further splitting. The accompanying India_Soil_Dataset.xlsx file contains the same data in structured tabular form for filtering, joins, or use as a lookup table alongside this

PDF.

4. Soil Type Reference Table

Soil Type	Typical Region	Fertility %	Moisture %	pH	N (kg/ha)	P (kg/ha)
Alluvial Soil	Indo-Gangetic plains	55-80	18-32	6.5-7.8	180-320	12-28
Black Soil (Regur)	Deccan trap region	50-75	20-35	7.0-8.3	150-280	8-20
Red & Black Soil	AP, Karnataka, TN	45-68	14-26	6.2-7.5	140-250	9-22
Red & Yellow Soil	Jharkhand, Chhattisgarh	38-60	12-22	5.5-6.8	110-220	6-16
Red & Laterite Soil	Odisha	35-58	15-28	5.0-6.5	100-210	5-14
Red & Lateritic Soil	NE hill states	35-58	18-32	4.8-6.2	100-200	5-14
Laterite Soil	Kerala, Goa	32-55	20-35	4.5-5.8	90-190	4-12
Arid & Desert Soil	Rajasthan	18-38	3-10	7.5-8.8	40-110	3-10
Mountain & Forest Soil	Himalayan states	40-65	22-38	5.0-6.8	120-240	6-16
Coastal Alluvial Soil	Puducherry, A&N Islands	42-66	24-40	6.8-8.0	130-240	7-18
Coastal Sandy Soil	Lakshadweep	25-42	15-26	6.5-7.8	60-140	4-10

5. District-Wise Soil Profiles

Each entry below is a standalone paragraph: State, District, Soil Type, Fertility %, Moisture %, pH, Nitrogen, Phosphorus, and Fertility Class. Values are indicative regional estimates as described in Section 2.

Andaman & Nicobar Islands

Nicobars district, Andaman & Nicobar Islands. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 60.6% (High fertility class). Soil moisture content is estimated at 31.0%. Soil pH is estimated at 7.83 (on a scale of 0-14). Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 8.0 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

North And Middle Andaman district, Andaman & Nicobar Islands. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 65.4% (High fertility class). Soil moisture content is estimated at 36.2%. Soil pH is estimated at 7.74 (on a scale of 0-14). Available nitrogen is estimated at 144 kg/ha, and available phosphorus is estimated at 12.0 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

South Andaman district, Andaman & Nicobar Islands. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 50.9% (Medium fertility class). Soil moisture content is estimated at 38.8%. Soil pH is estimated at 7.57 (on a scale of 0-14). Available nitrogen is estimated at 220 kg/ha, and available phosphorus is estimated at 11.9 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Andhra Pradesh

Adilabad district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 50.2% (Medium fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 6.28 (on a scale of 0-14). Available nitrogen is estimated at 231 kg/ha, and available phosphorus is estimated at 17.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Anantapur district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 62.4% (High fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 7.46 (on a scale of 0-14). Available nitrogen is estimated at 238 kg/ha, and available phosphorus is estimated at 19.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chittoor district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.5% (Medium fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 6.26 (on a scale of 0-14). Available nitrogen is estimated at 156 kg/ha, and available phosphorus is estimated at 17.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

East Godavari district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 62.1% (High fertility class). Soil moisture content is estimated at 25.6%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 15.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Guntur district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.4% (Medium fertility class). Soil moisture content is estimated at 15.6%. Soil pH is estimated at 6.82 (on a scale of 0-14). Available nitrogen is estimated at 164 kg/ha, and available phosphorus is estimated at 17.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Hyderabad district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 55.1% (High fertility class). Soil moisture content is estimated at 24.0%. Soil pH is estimated at 7.11 (on a scale of 0-14). Available nitrogen is estimated at 174 kg/ha, and available phosphorus is estimated at 19.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Karimnagar district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 63.5% (High fertility class). Soil moisture content is estimated at 18.6%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 10.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Khammam district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.6% (Medium fertility class). Soil moisture content is estimated at 14.1%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 18.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Krishna district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 63.0% (High fertility class). Soil moisture content is estimated at 19.5%. Soil pH is estimated at 6.94 (on a scale of 0-14). Available nitrogen is estimated at 155 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kurnool district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 60.4% (High fertility class). Soil moisture content is estimated at 19.7%. Soil pH is estimated at 6.93 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 17.3 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Mahbubnagar district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 57.7% (High fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 6.6 (on a scale of 0-14). Available nitrogen is estimated at 143 kg/ha, and available phosphorus is estimated at 14.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Medak district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.9% (Medium fertility class). Soil moisture content is estimated at 18.9%. Soil pH is estimated at 7.31 (on a scale of 0-14). Available nitrogen is estimated at 165 kg/ha, and available phosphorus is estimated at 9.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Nalgonda district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 51.5% (Medium fertility class). Soil moisture content is estimated at 17.5%. Soil pH is estimated at 7.06 (on a scale of 0-14). Available nitrogen is estimated at 201 kg/ha, and available phosphorus is estimated at 19.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Nizamabad district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 60.3% (High fertility class). Soil moisture content is estimated at 18.9%. Soil pH is estimated at 7.26 (on a scale of 0-14). Available nitrogen is estimated at 158 kg/ha, and available phosphorus is estimated at 9.3 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Prakasam district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 47.1% (Medium fertility class). Soil moisture content is estimated at 22.7%. Soil pH is estimated at 6.8 (on a scale of 0-14). Available nitrogen is estimated at 157 kg/ha, and available phosphorus is estimated at 15.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Rangareddy district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 48.5% (Medium fertility class). Soil moisture content is estimated at 22.4%. Soil pH is estimated at 6.78 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 12.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Sri Potti Sriramulu Nellore district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 59.5% (High fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 6.31 (on a scale of 0-14). Available nitrogen is estimated at 152 kg/ha, and available phosphorus is estimated at 21.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Srikakulam district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.9% (High fertility class). Soil moisture content is estimated at 22.4%. Soil pH is estimated at 6.55 (on a scale of 0-14). Available nitrogen is estimated at 246 kg/ha, and available phosphorus is estimated at 19.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Visakhapatnam district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 61.5% (High fertility class). Soil moisture content is estimated at 19.4%. Soil pH is estimated at 6.55 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 20.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Vizianagaram district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 55.5% (High fertility class). Soil moisture content is estimated at 16.4%. Soil pH is estimated at 6.6 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Warangal district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.7% (High fertility class). Soil moisture content is estimated at 23.1%. Soil pH is estimated at 7.14 (on a scale of 0-14). Available nitrogen is estimated at 187 kg/ha, and available phosphorus is estimated at 17.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

West Godavari district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 58.4% (High fertility class). Soil moisture content is estimated at 21.8%. Soil pH is estimated at 6.31 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Y.S.R. district, Andhra Pradesh. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 18.0%. Soil pH is estimated at 6.39 (on a scale of 0-14). Available nitrogen is estimated at 151 kg/ha, and available phosphorus is estimated at 16.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Arunachal Pradesh

Anjaw district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 44.3% (Medium fertility class). Soil moisture content is estimated at 36.8%. Soil pH is estimated at 6.05 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 11.9 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Changlang district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 40.6% (Medium fertility class). Soil moisture content is estimated at 37.3%. Soil pH is estimated at 5.87 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 6.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Dibang Valley district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 52.2% (Medium fertility class). Soil moisture content is estimated at 29.9%. Soil pH is

estimated at 6.69 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 10.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

East Kameng district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.7% (Medium fertility class). Soil moisture content is estimated at 27.3%. Soil pH is estimated at 5.94 (on a scale of 0-14). Available nitrogen is estimated at 172 kg/ha, and available phosphorus is estimated at 6.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

East Siang district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 60.7% (High fertility class). Soil moisture content is estimated at 36.3%. Soil pH is estimated at 5.25 (on a scale of 0-14). Available nitrogen is estimated at 186 kg/ha, and available phosphorus is estimated at 7.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kurung Kumey district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.8% (High fertility class). Soil moisture content is estimated at 26.5%. Soil pH is estimated at 6.19 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Lohit district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 42.7% (Medium fertility class). Soil moisture content is estimated at 36.7%. Soil pH is estimated at 5.41 (on a scale of 0-14). Available nitrogen is estimated at 124 kg/ha, and available phosphorus is estimated at 11.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Lower Dibang Valley district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 49.3% (Medium fertility class). Soil moisture content is estimated at 35.3%. Soil pH is estimated at 6.45 (on a scale of 0-14). Available nitrogen is estimated at 158 kg/ha, and available phosphorus is estimated at 15.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Lower Subansiri district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 47.3% (Medium fertility class). Soil moisture content is estimated at 30.2%. Soil pH is estimated at 5.46 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 7.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Papum Pare district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 41.1% (Medium fertility class). Soil moisture content is estimated at 29.0%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 227 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Tawang district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 62.3% (High fertility class). Soil moisture content is estimated at 36.3%. Soil pH is estimated at 5.93 (on a scale of 0-14). Available nitrogen is estimated at 157 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Tirap district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.5% (High fertility class). Soil moisture content is estimated at 28.0%. Soil pH is estimated at 5.17 (on a scale of 0-14). Available nitrogen is estimated at 209 kg/ha, and available phosphorus is estimated at 8.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Upper Siang district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 63.4% (High fertility class). Soil moisture content is estimated at 25.9%. Soil pH is estimated at 5.22 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 7.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Upper Subansiri district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 44.5% (Medium fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 143 kg/ha, and available phosphorus is estimated at 9.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

West Kameng district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 59.4% (High fertility class). Soil moisture content is estimated at 37.5%. Soil pH is estimated at 5.9 (on a scale of 0-14). Available nitrogen is estimated at 137 kg/ha, and available phosphorus is estimated at 6.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

West Siang district, Arunachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 45.7% (Medium fertility class). Soil moisture content is estimated at 24.1%. Soil pH is estimated at 6.22 (on a scale of 0-14). Available nitrogen is estimated at 134 kg/ha, and available phosphorus is estimated at 11.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Assam

Baksa district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.4% (High fertility class). Soil moisture content is estimated at 26.1%. Soil pH is estimated at 6.76 (on a scale of 0-14). Available nitrogen is estimated at 292 kg/ha, and available phosphorus is estimated at 23.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Barpeta district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 19.8%. Soil pH is estimated at 6.66 (on a scale of 0-14). Available nitrogen is estimated at 309 kg/ha, and available phosphorus is estimated at 18.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bongaigaon district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.5% (High fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 7.36 (on a scale of 0-14). Available nitrogen is estimated at 313 kg/ha, and available phosphorus is estimated at 16.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Cachar district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.1% (High fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 268 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Chirang district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.5% (High fertility class). Soil moisture content is estimated at 23.9%. Soil pH is estimated at 7.76 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 26.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Darrang district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.1% (High fertility class). Soil moisture content is estimated at 24.5%. Soil pH is estimated at 7.52 (on a scale of 0-14). Available nitrogen is estimated at 182 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Dhemaji district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.7% (High fertility class). Soil moisture content is estimated at 29.2%. Soil pH is estimated at 6.8 (on a scale of 0-14). Available nitrogen is estimated at 254 kg/ha, and available phosphorus is estimated at 21.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Dhubri district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 76.7% (High fertility class). Soil moisture content is estimated at 26.4%. Soil pH is estimated at 7.04 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 18.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Dibrugarh district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 30.1%. Soil pH is estimated at 7.09 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Dima Hasao district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.7% (High fertility class). Soil moisture content is estimated at 29.4%. Soil pH is estimated at 6.64 (on a scale of 0-14). Available nitrogen is estimated at 189 kg/ha, and available phosphorus is estimated at 21.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Goalpara district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.7% (High fertility class). Soil moisture content is estimated at 29.5%. Soil pH is estimated at 6.9 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 26.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Golaghat district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.1% (High fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 6.7 (on a scale of 0-14). Available nitrogen is estimated at 196 kg/ha, and available phosphorus is estimated at 12.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hailakandi district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 262 kg/ha, and available phosphorus is estimated at 22.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jorhat district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.8% (High fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 7.16 (on a scale of 0-14). Available nitrogen is estimated at 302 kg/ha, and available phosphorus is estimated at 25.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kamrup district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 20.5%. Soil pH is estimated at 6.81 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 21.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kamrup Metropolitan district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.4% (High fertility class). Soil moisture content is estimated at 18.7%. Soil pH is estimated at 6.99 (on a scale of 0-14). Available nitrogen is estimated at 253 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Karbi Anglong district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.8% (High fertility class). Soil moisture content is estimated at 18.7%. Soil pH is estimated at 7.7 (on a scale of 0-14). Available nitrogen is estimated at 193 kg/ha, and available phosphorus is estimated at 25.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Karimganj district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.6% (High fertility class). Soil moisture content is estimated at 31.7%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 289 kg/ha, and available phosphorus is estimated at 22.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kokrajhar district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.5% (High fertility class). Soil moisture content is estimated at 19.9%. Soil pH is estimated at 7.2 (on a scale of 0-14). Available nitrogen is estimated at 251 kg/ha, and available phosphorus is estimated at 25.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Lakhimpur district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.6% (High fertility class). Soil moisture content is estimated at 23.4%. Soil pH is estimated at 7.33 (on a scale of 0-14). Available nitrogen is estimated at 217 kg/ha, and available phosphorus is estimated at 14.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Morigaon district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.9% (High fertility class). Soil moisture content is estimated at 23.8%. Soil pH is estimated at 6.8 (on a scale of 0-14). Available nitrogen is estimated at 231 kg/ha, and available phosphorus is estimated at 17.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Nagaon district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.2% (High fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 7.39 (on a scale of 0-14). Available nitrogen is estimated at 221 kg/ha, and available phosphorus is estimated at 27.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Nalbari district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.9% (High fertility class). Soil moisture content is estimated at 24.7%. Soil pH is estimated at 6.93 (on a scale of 0-14). Available nitrogen is estimated at 254 kg/ha, and available phosphorus is estimated at 25.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sivasagar district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 7.19 (on a scale of 0-14). Available nitrogen is estimated at 268 kg/ha, and available phosphorus is estimated at 16.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sonitpur district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.4% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 7.4 (on a scale of 0-14). Available nitrogen is estimated at 300 kg/ha, and available phosphorus is estimated at 14.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Tinsukia district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.4% (High fertility class). Soil moisture content is estimated at 19.3%. Soil pH is estimated at 7.44 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 27.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Udalguri district, Assam. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.4% (High fertility class). Soil moisture content is estimated at 31.4%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 24.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bihar

Araria district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.9% (High fertility class). Soil moisture content is estimated at 31.2%. Soil pH is estimated at 6.83 (on a scale of 0-14). Available nitrogen is estimated at 262 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Arwal district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.4% (High fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 7.23 (on a scale of 0-14). Available nitrogen is estimated at 260 kg/ha, and available phosphorus is estimated at 19.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Aurangabad district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.1% (High fertility class). Soil moisture content is estimated at 28.7%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 248 kg/ha, and available phosphorus is estimated at 21.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Banka district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.3% (High fertility class). Soil moisture content is estimated at 19.7%. Soil pH is estimated at 6.65 (on a scale of 0-14). Available nitrogen is estimated at 192 kg/ha, and available phosphorus is estimated at 22.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Begusarai district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.5% (High fertility class). Soil moisture content is estimated at 28.8%. Soil pH is estimated at 7.37 (on a scale of 0-14). Available nitrogen is estimated at 226 kg/ha, and available phosphorus is estimated at 26.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bhagalpur district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.1% (High fertility class). Soil moisture content is estimated at 21.8%. Soil pH is estimated at 6.97 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 14.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bhojpur district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.7% (High fertility class). Soil moisture content is estimated at 31.1%. Soil pH is estimated at 7.07 (on a scale of 0-14). Available nitrogen is estimated at 233 kg/ha, and available phosphorus is estimated at 23.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Buxar district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.8% (High fertility class). Soil moisture content is estimated at 31.1%. Soil pH is estimated at 7.51 (on a scale of 0-14). Available nitrogen is estimated at 247 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Darbhanga district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.7% (High fertility class). Soil moisture content is estimated at 28.0%. Soil pH is estimated at 7.74 (on a scale of 0-14). Available nitrogen is estimated at 196 kg/ha, and available phosphorus is estimated at 25.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gaya district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.9% (High fertility class). Soil moisture content is estimated at 19.7%. Soil pH is estimated at 7.26 (on a scale of 0-14). Available nitrogen is estimated at 276 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gopalganj district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.4% (High fertility class). Soil moisture content is estimated at 29.6%. Soil pH is estimated at 6.88 (on a scale of 0-14). Available nitrogen is estimated at 244 kg/ha, and available phosphorus is estimated at 19.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jamui district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.5% (High fertility class). Soil moisture content is estimated at 30.9%. Soil pH is estimated at 7.52 (on a scale of 0-14). Available nitrogen is estimated at 195 kg/ha, and available phosphorus is estimated at 28.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jehanabad district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.0% (High fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 7.59 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 28.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kaimur (Bhabua) district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.6% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 305 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Katihar district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.0% (High fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 7.51 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 26.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Khagaria district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.4% (High fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 6.51 (on a scale of 0-14). Available nitrogen is estimated at 187 kg/ha, and available phosphorus is estimated at 21.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kishanganj district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.0% (High fertility class). Soil moisture content is estimated at 21.3%. Soil pH is estimated at 7.6 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 24.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Lakhisarai district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.2% (High fertility class). Soil moisture content is estimated at 28.8%. Soil pH is estimated at 7.41 (on a scale of 0-14). Available nitrogen is estimated at 297 kg/ha, and available phosphorus is estimated at 12.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Madhepura district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.0% (High fertility class). Soil moisture content is estimated at 19.7%. Soil pH is estimated at 7.16 (on a scale of 0-14). Available nitrogen is estimated at 284 kg/ha, and available phosphorus is estimated at 22.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Madhubani district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 76.3% (High fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 7.46 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 16.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Munger district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.7% (High fertility class). Soil moisture content is estimated at 31.7%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 187 kg/ha, and available phosphorus is estimated at 21.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Muzaffarpur district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 30.4%. Soil pH is estimated at 7.42 (on a scale of 0-14). Available nitrogen is estimated at 204 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Nalanda district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.6% (High fertility class). Soil moisture content is estimated at 31.7%. Soil pH is estimated at 7.1 (on a scale of 0-14). Available nitrogen is estimated at 289 kg/ha, and available phosphorus is estimated at 22.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Nawada district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.3% (High fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 7.73 (on a scale of 0-14). Available nitrogen is estimated at 222 kg/ha, and available phosphorus is estimated at 21.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Pashchim Champaran district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.5% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 20.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Patna district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.1% (High fertility class). Soil moisture content is estimated at 24.6%. Soil pH is estimated at 6.65 (on a scale of 0-14). Available nitrogen is estimated at 198 kg/ha, and available phosphorus is estimated at 16.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Purba Champaran district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.6% (High fertility class). Soil moisture content is estimated at 24.0%. Soil pH is estimated at 7.29 (on a scale of 0-14). Available nitrogen is estimated at 268 kg/ha, and available phosphorus is estimated at 18.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Purnia district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.2% (High fertility class). Soil moisture content is estimated at 21.0%. Soil pH is estimated at 7.26 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 12.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rohtas district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.5% (High fertility class). Soil moisture content is estimated at 24.6%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 260 kg/ha, and available phosphorus is estimated at 21.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Saharsa district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.6% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 7.35 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 24.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Samastipur district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.7% (High fertility class). Soil moisture content is estimated at 24.0%. Soil pH is estimated at 7.31 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 20.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Saran district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.4% (High fertility class). Soil moisture content is estimated at 30.4%. Soil pH is estimated at 7.7 (on a scale of 0-14). Available nitrogen is estimated at 250 kg/ha, and available phosphorus is estimated at 20.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sheikhpura district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.0% (High fertility class). Soil moisture content is estimated at 22.4%. Soil pH is estimated at 7.59 (on a scale of 0-14). Available nitrogen is estimated at 249 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sheohar district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.8% (High fertility class). Soil moisture content is estimated at 29.8%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 17.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sitamarhi district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.3% (High fertility class). Soil moisture content is estimated at 22.4%. Soil pH is estimated at 7.47 (on a scale of 0-14). Available nitrogen is estimated at 182 kg/ha, and available phosphorus is estimated at 25.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Siwan district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 76.4% (High fertility class). Soil moisture content is estimated at 23.2%. Soil pH is estimated at 6.7 (on a scale of 0-14). Available nitrogen is estimated at 264 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Supaul district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.1% (High fertility class). Soil moisture content is estimated at 31.4%. Soil pH is estimated at 7.79 (on a scale of 0-14). Available nitrogen is estimated at 288 kg/ha, and available phosphorus is estimated at 17.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Vaishali district, Bihar. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.2% (High fertility class). Soil moisture content is estimated at 27.9%. Soil pH is estimated at 7.0 (on a scale of 0-14). Available nitrogen is estimated at 269 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Chandigarh

Chandigarh district, Chandigarh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.2% (High fertility class). Soil moisture content is estimated at 25.4%. Soil pH is estimated at 6.71 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 25.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Chhattisgarh

Bastar district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 59.8% (High fertility class). Soil moisture content is estimated at 17.6%. Soil pH is estimated at 6.59 (on a scale of 0-14). Available nitrogen is estimated at 218 kg/ha, and available phosphorus is estimated at 7.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Bijapur district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 47.9% (Medium fertility class). Soil moisture content is estimated at 15.9%. Soil pH is estimated at 5.6 (on a scale of 0-14). Available nitrogen is estimated at 193 kg/ha, and available phosphorus is estimated at 10.3

kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Bilaspur district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 48.3% (Medium fertility class). Soil moisture content is estimated at 13.5%. Soil pH is estimated at 5.74 (on a scale of 0-14). Available nitrogen is estimated at 209 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Dakshin Bastar Dantewada district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 43.1% (Medium fertility class). Soil moisture content is estimated at 14.9%. Soil pH is estimated at 6.14 (on a scale of 0-14). Available nitrogen is estimated at 174 kg/ha, and available phosphorus is estimated at 10.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Dhamtari district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 39.9% (Medium fertility class). Soil moisture content is estimated at 14.4%. Soil pH is estimated at 6.6 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Durg district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 52.7% (Medium fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 5.58 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 11.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Janjgir - Champa district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 45.4% (Medium fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 6.13 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 14.5 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Jashpur district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 49.1% (Medium fertility class). Soil moisture content is estimated at 21.1%. Soil pH is estimated at 6.26 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Kabeerdham district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 49.0% (Medium fertility class). Soil moisture content is estimated at 17.3%. Soil pH is estimated at 5.64 (on a scale of 0-14). Available nitrogen is estimated at 153 kg/ha, and available phosphorus is estimated at 15.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Korba district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 51.9% (Medium fertility class). Soil moisture content is estimated at 13.8%. Soil pH is estimated at 5.94 (on a scale of 0-14). Available nitrogen is estimated at 131 kg/ha, and available phosphorus is estimated at 6.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Koriya district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 58.4% (High fertility class). Soil moisture content is estimated at 16.5%. Soil pH is estimated at 5.9 (on a scale of 0-14). Available nitrogen is estimated at 175 kg/ha, and available phosphorus is estimated at 6.1 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Mahasamund district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 44.1% (Medium fertility class). Soil moisture content is estimated at 19.0%. Soil pH is estimated at 6.32 (on a scale of 0-14). Available nitrogen is estimated at 217 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Narayanpur district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 48.5% (Medium fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 6.67 (on a scale of 0-14). Available nitrogen is estimated at 189 kg/ha, and available phosphorus is estimated at 15.6 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Raigarh district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 55.2% (High fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 5.65 (on a scale of 0-14). Available nitrogen is estimated at 190 kg/ha, and available phosphorus is estimated at 10.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Raipur district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 50.2% (Medium fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 6.76 (on a scale of 0-14). Available nitrogen is estimated at 218 kg/ha, and available phosphorus is estimated at 8.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Rajnandgaon district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 54.1% (Medium fertility class). Soil moisture content is estimated at 19.5%. Soil pH is estimated at 5.95 (on a scale of 0-14). Available nitrogen is estimated at 123 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Surguja district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 55.1% (High fertility class). Soil moisture content is estimated at 16.9%. Soil pH is estimated at 6.78 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Uttar Bastar Kanker district, Chhattisgarh. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 47.1% (Medium fertility class). Soil moisture content is estimated at 19.9%. Soil pH is estimated at 5.61 (on a scale of 0-14). Available nitrogen is estimated at 171 kg/ha, and available phosphorus is estimated at 14.0 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Dadra & Nagar Haveli

Dadra And Nagar Haveli district, Dadra & Nagar Haveli. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 56.3% (High fertility class). Soil moisture content is estimated at 29.5%. Soil pH is estimated at 4.85 (on a scale of 0-14). Available nitrogen is estimated at 137 kg/ha, and available phosphorus is estimated at 5.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Daman & Diu

Daman district, Daman & Diu. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 44.6% (Medium fertility class). Soil moisture content is estimated at 34.8%. Soil pH is estimated at 7.66 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Diu district, Daman & Diu. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 59.7% (High fertility class). Soil moisture content is estimated at 36.8%. Soil pH is estimated at 6.86 (on a scale of 0-14). Available nitrogen is estimated at 155 kg/ha, and available phosphorus is estimated at 13.8 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Delhi

Central district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 76.5% (High fertility class). Soil moisture content is estimated at 18.1%. Soil pH is estimated at 7.17 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

East district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.0% (High fertility class). Soil moisture content is estimated at 30.5%. Soil pH is estimated at 7.37 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 25.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

New Delhi district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.7% (High fertility class). Soil moisture content is estimated at 29.9%. Soil pH is estimated at 6.89 (on a scale of 0-14). Available nitrogen is estimated at 262 kg/ha, and available phosphorus is estimated at 18.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

North district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.9% (High fertility class). Soil moisture content is estimated at 30.4%. Soil pH is estimated at 6.74 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 17.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

North East district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.9% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 7.8 (on a scale of 0-14). Available nitrogen is estimated at 221 kg/ha, and available phosphorus is estimated at 18.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

North West district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.4% (High fertility class). Soil moisture content is estimated at 26.0%. Soil pH is estimated at 7.8 (on a scale of 0-14). Available nitrogen is estimated at 278 kg/ha, and available phosphorus is estimated at 21.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

South district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.8% (High fertility class). Soil moisture content is estimated at 30.8%. Soil pH is estimated at 7.15 (on a scale of 0-14). Available nitrogen is estimated at 198 kg/ha, and available phosphorus is estimated at 17.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

South West district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.7% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 6.52 (on a scale of 0-14). Available nitrogen is estimated at 243 kg/ha, and available phosphorus is estimated at 22.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

West district, Delhi. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.6% (High fertility class). Soil moisture content is estimated at 23.9%. Soil pH is estimated at 7.75 (on a scale of 0-14). Available nitrogen is estimated at 285 kg/ha, and available phosphorus is estimated at 20.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Goa

North Goa district, Goa. Dominant soil type: Laterite Soil. Soil fertility is estimated at 38.5% (Medium fertility class). Soil moisture content is estimated at 33.5%. Soil pH is estimated at 4.81 (on a scale of 0-14). Available nitrogen is estimated at 122 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

South Goa district, Goa. Dominant soil type: Laterite Soil. Soil fertility is estimated at 44.2% (Medium fertility class). Soil moisture content is estimated at 31.1%. Soil pH is estimated at 5.27 (on a scale of 0-14). Available nitrogen is estimated at 155 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Gujarat

Ahmadabad district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.0% (High fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 7.2 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 11.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Amreli district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 64.0% (High fertility class). Soil moisture content is estimated at 31.9%. Soil pH is estimated at 8.02 (on a scale of 0-14). Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Anand district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 74.9% (High fertility class). Soil moisture content is estimated at 30.1%. Soil pH is estimated at 8.06 (on a scale of 0-14). Available nitrogen is estimated at 267 kg/ha, and available phosphorus is estimated at 17.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Banas Kantha district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 54.6% (Medium fertility class). Soil moisture content is estimated at 28.4%. Soil pH is estimated at 7.13 (on a scale of 0-14). Available nitrogen is estimated at 234 kg/ha, and available phosphorus is estimated at 19.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bharuch district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.8% (High fertility class). Soil moisture content is estimated at 26.5%. Soil pH is estimated at 7.05 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 9.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bhavnagar district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.0% (Medium fertility class). Soil moisture content is estimated at 23.7%. Soil pH is estimated at 7.09 (on a scale of 0-14). Available nitrogen is estimated at 209 kg/ha, and available phosphorus is estimated at 14.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Dohad district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.8% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 7.15 (on a scale of 0-14). Available nitrogen is estimated at 199 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Gandhinagar district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 67.5% (High fertility class). Soil moisture content is estimated at 23.1%. Soil pH is estimated at 7.39 (on a scale of 0-14). Available nitrogen is estimated at 201 kg/ha, and available phosphorus is estimated at 13.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Jamnagar district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.0% (Medium fertility class). Soil moisture content is estimated at 21.7%. Soil pH is estimated at 8.12 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 14.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Junagadh district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.2% (High fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 7.56 (on a scale of 0-14). Available nitrogen is estimated at 251 kg/ha, and available phosphorus is estimated at 18.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Kachchh district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.5% (High fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 7.32 (on a scale of 0-14). Available nitrogen is estimated at 212 kg/ha, and available phosphorus is estimated at 16.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Kheda district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.7% (High fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 8.21 (on a scale of 0-14). Available nitrogen is estimated at 156 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Mahesana district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 67.8% (High fertility class). Soil moisture content is estimated at 22.3%. Soil pH is estimated at 7.06 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 19.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Narmada district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.2% (Medium fertility class). Soil moisture content is estimated at 22.8%. Soil pH is estimated at 7.04 (on a scale of 0-14). Available nitrogen is estimated at 164 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Navsari district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.0% (High fertility class). Soil moisture content is estimated at 28.5%. Soil pH is estimated at 7.77 (on a scale of 0-14). Available nitrogen is estimated at 260 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Panch Mahals district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 7.21 (on a scale of 0-14). Available nitrogen is estimated at 250 kg/ha, and available phosphorus is estimated at 18.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Patan district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 33.8%. Soil pH is estimated at 7.59 (on a scale of 0-14). Available nitrogen is estimated at 228 kg/ha, and available phosphorus is estimated at 19.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Porbandar district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 59.0% (High fertility class). Soil moisture content is estimated at 32.2%. Soil pH is estimated at 7.41 (on a scale of 0-14). Available nitrogen is estimated at 253 kg/ha, and available phosphorus is estimated at 15.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Rajkot district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.4% (High fertility class). Soil moisture content is estimated at 26.2%. Soil pH is estimated at 7.41 (on a scale of 0-14). Available nitrogen is estimated at 160 kg/ha, and available phosphorus is estimated at 8.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sabar Kantha district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 58.7% (High fertility class). Soil moisture content is estimated at 20.3%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 244 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Surat district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 68.5% (High fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 8.16 (on a scale of 0-14). Available nitrogen is estimated at 227 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Surendranagar district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 53.6% (Medium fertility class). Soil moisture content is estimated at 30.4%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 19.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Tapi district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.1% (Medium fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 8.29 (on a scale of 0-14). Available nitrogen is estimated at 226 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

The Dangs district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 72.4% (High fertility class). Soil moisture content is estimated at 33.2%. Soil pH is estimated at 7.7 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Vadodara district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.3% (Medium fertility class). Soil moisture content is estimated at 28.9%. Soil pH is estimated at 7.38 (on a scale of 0-14). Available nitrogen is estimated at 236 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Valsad district, Gujarat. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.5% (Medium fertility class). Soil moisture content is estimated at 28.9%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 11.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Haryana

Ambala district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.3% (High fertility class). Soil moisture content is estimated at 25.7%. Soil pH is estimated at 7.02 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 23.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bhiwani district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.2% (High fertility class). Soil moisture content is estimated at 28.6%. Soil pH is estimated at 7.78 (on a scale of 0-14). Available nitrogen is estimated at 238 kg/ha, and available phosphorus is estimated at 20.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Faridabad district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 55.3% (High fertility class). Soil moisture content is estimated at 29.1%. Soil pH is estimated at 7.18 (on a scale of 0-14). Available nitrogen is estimated at 237 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Fatehabad district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.2% (High fertility class). Soil moisture content is estimated at 23.5%. Soil pH is estimated at 7.39 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 27.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gurgaon district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.5% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 7.62 (on a scale of 0-14). Available nitrogen is estimated at 307 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hisar district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.7% (High fertility class). Soil moisture content is estimated at 28.3%. Soil pH is estimated at 7.64 (on a scale of 0-14). Available nitrogen is estimated at 233 kg/ha, and available phosphorus is estimated at 20.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jhajjar district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 28.2%. Soil pH is estimated at 7.52 (on a scale of 0-14). Available nitrogen is estimated at 259 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jind district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.6% (High fertility class). Soil moisture content is estimated at 30.1%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 193 kg/ha, and available phosphorus is estimated at 15.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kaithal district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.6% (High fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 7.77 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 23.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Karnal district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.2% (High fertility class). Soil moisture content is estimated at 23.5%. Soil pH is estimated at 6.91 (on a scale of 0-14). Available nitrogen is estimated at 265 kg/ha, and available phosphorus is estimated at 21.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kurukshetra district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.2% (High fertility class). Soil moisture content is estimated at 26.4%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 245 kg/ha, and available phosphorus is estimated at 15.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mahendragarh district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.4% (High fertility class). Soil moisture content is estimated at 23.5%. Soil pH is estimated at 6.67 (on a scale of 0-14). Available nitrogen is estimated at 202 kg/ha, and available phosphorus is estimated at 23.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mewat district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.5% (High fertility class). Soil moisture content is estimated at 31.4%. Soil pH is estimated at 6.82 (on a scale of 0-14). Available nitrogen is estimated at 193 kg/ha, and available phosphorus is estimated at 24.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Palwal district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 77.0% (High fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 6.76 (on a scale of 0-14). Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 20.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Panchkula district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.7% (High fertility class). Soil moisture content is estimated at 21.6%. Soil pH is estimated at 6.56 (on a scale of 0-14). Available nitrogen is estimated at 247 kg/ha, and available phosphorus is estimated at 27.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Panipat district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 24.9%. Soil pH is estimated at 6.64 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 16.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rewari district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.1% (High fertility class). Soil moisture content is estimated at 30.3%. Soil pH is estimated at 7.67 (on a scale of 0-14). Available nitrogen is estimated at 317 kg/ha, and available phosphorus is estimated at 27.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rohtak district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.8% (High fertility class). Soil moisture content is estimated at 19.0%. Soil pH is estimated at 6.68 (on a scale of 0-14). Available nitrogen is estimated at 222 kg/ha, and available phosphorus is estimated at 20.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sirsa district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.4% (High fertility class). Soil moisture content is estimated at 29.8%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 255 kg/ha, and available phosphorus is estimated at 14.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sonipat district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.3% (High fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 7.61 (on a scale of 0-14). Available nitrogen is estimated at 261 kg/ha, and available phosphorus is estimated at 23.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Yamunanagar district, Haryana. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.4% (High fertility class). Soil moisture content is estimated at 23.2%. Soil pH is estimated at 7.03 (on a scale of 0-14). Available nitrogen is estimated at 286 kg/ha, and available phosphorus is estimated at 24.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Himachal Pradesh

Bilaspur district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 45.2% (Medium fertility class). Soil moisture content is estimated at 37.1%. Soil pH is estimated at 5.22 (on a scale of 0-14). Available nitrogen is estimated at 227 kg/ha, and available phosphorus is estimated at 7.0 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Chamba district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.6% (Medium fertility class). Soil moisture content is estimated at 35.5%. Soil pH is estimated at 5.32 (on a scale of 0-14). Available nitrogen is estimated at 169 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Hamirpur district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.1% (Medium fertility class). Soil moisture content is estimated at 33.4%. Soil pH is estimated at 6.53 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kangra district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 53.3% (Medium fertility class). Soil moisture content is estimated at 26.0%. Soil pH is estimated at 5.44

(on a scale of 0-14). Available nitrogen is estimated at 139 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kinnaur district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 62.2% (High fertility class). Soil moisture content is estimated at 34.4%. Soil pH is estimated at 5.93 (on a scale of 0-14). Available nitrogen is estimated at 178 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kullu district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 53.4% (Medium fertility class). Soil moisture content is estimated at 29.0%. Soil pH is estimated at 5.24 (on a scale of 0-14). Available nitrogen is estimated at 135 kg/ha, and available phosphorus is estimated at 15.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Lahul And Spiti district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 52.1% (Medium fertility class). Soil moisture content is estimated at 37.3%. Soil pH is estimated at 5.29 (on a scale of 0-14). Available nitrogen is estimated at 186 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Mandi district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.3% (Medium fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 5.21 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 9.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Shimla district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 55.2% (High fertility class). Soil moisture content is estimated at 29.4%. Soil pH is estimated at 5.72 (on a scale of 0-14). Available nitrogen is estimated at 183 kg/ha, and available phosphorus is estimated at 7.9 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Sirmaur district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 64.7% (High fertility class). Soil moisture content is estimated at 35.1%. Soil pH is estimated at 6.33 (on a scale of 0-14). Available nitrogen is estimated at 176 kg/ha, and available phosphorus is estimated at 7.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Solan district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 63.0% (High fertility class). Soil moisture content is estimated at 27.5%. Soil pH is estimated at 5.09 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Una district, Himachal Pradesh. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 34.0%. Soil pH is estimated at 6.43 (on a scale of 0-14). Available nitrogen is estimated at 145 kg/ha, and available phosphorus is estimated at 15.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Jammu And Kashmir

Anantnag district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 50.9% (Medium fertility class). Soil moisture content is estimated at 32.2%. Soil pH is

estimated at 5.0 (on a scale of 0-14). Available nitrogen is estimated at 239 kg/ha, and available phosphorus is estimated at 8.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Badgam district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 41.6% (Medium fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 5.23 (on a scale of 0-14). Available nitrogen is estimated at 138 kg/ha, and available phosphorus is estimated at 12.3 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Bandipore district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 49.8% (Medium fertility class). Soil moisture content is estimated at 36.7%. Soil pH is estimated at 5.57 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Baramula district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.5% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 5.85 (on a scale of 0-14). Available nitrogen is estimated at 234 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Doda district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 44.4% (Medium fertility class). Soil moisture content is estimated at 32.8%. Soil pH is estimated at 6.52 (on a scale of 0-14). Available nitrogen is estimated at 124 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Ganderbal district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 62.3% (High fertility class). Soil moisture content is estimated at 34.0%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 183 kg/ha, and available phosphorus is estimated at 12.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Jammu district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 47.6% (Medium fertility class). Soil moisture content is estimated at 37.2%. Soil pH is estimated at 5.66 (on a scale of 0-14). Available nitrogen is estimated at 208 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kargil district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 54.0% (Medium fertility class). Soil moisture content is estimated at 33.5%. Soil pH is estimated at 5.92 (on a scale of 0-14). Available nitrogen is estimated at 228 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kathua district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.2% (High fertility class). Soil moisture content is estimated at 27.5%. Soil pH is estimated at 5.73 (on a scale of 0-14). Available nitrogen is estimated at 172 kg/ha, and available phosphorus is estimated at 7.3 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kishtwar district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 42.3% (Medium fertility class). Soil moisture content is estimated at 32.7%. Soil pH is estimated at 6.18 (on a scale of 0-14). Available nitrogen is estimated at 199 kg/ha, and available phosphorus is estimated at 6.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kulgam district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 48.2% (Medium fertility class). Soil moisture content is estimated at 25.2%. Soil pH is estimated at 6.4 (on a scale of 0-14). Available nitrogen is estimated at 223 kg/ha, and available phosphorus is estimated at 8.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Kupwara district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.7% (High fertility class). Soil moisture content is estimated at 24.1%. Soil pH is estimated at 5.5 (on a scale of 0-14). Available nitrogen is estimated at 136 kg/ha, and available phosphorus is estimated at 8.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Leh(Ladakh) district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 56.9% (High fertility class). Soil moisture content is estimated at 36.6%. Soil pH is estimated at 5.36 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 13.0 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Pulwama district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 63.8% (High fertility class). Soil moisture content is estimated at 31.8%. Soil pH is estimated at 5.46 (on a scale of 0-14). Available nitrogen is estimated at 195 kg/ha, and available phosphorus is estimated at 9.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Punch district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 64.4% (High fertility class). Soil moisture content is estimated at 28.4%. Soil pH is estimated at 6.23 (on a scale of 0-14). Available nitrogen is estimated at 122 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Rajouri district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 61.9% (High fertility class). Soil moisture content is estimated at 31.8%. Soil pH is estimated at 5.8 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 14.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Ramban district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 48.1% (Medium fertility class). Soil moisture content is estimated at 30.7%. Soil pH is estimated at 5.88 (on a scale of 0-14). Available nitrogen is estimated at 195 kg/ha, and available phosphorus is estimated at 9.3 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Reasi district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.1% (Medium fertility class). Soil moisture content is estimated at 36.6%. Soil pH is estimated at 6.66 (on a scale of 0-14). Available nitrogen is estimated at 221 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Samba district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 58.6% (High fertility class). Soil moisture content is estimated at 34.2%. Soil pH is estimated at 6.37 (on a scale of 0-14). Available nitrogen is estimated at 177 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Shupiyan district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 46.0% (Medium fertility class). Soil moisture content is estimated at 32.3%. Soil pH is estimated at 5.52 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 11.9 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Srinagar district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 50.2% (Medium fertility class). Soil moisture content is estimated at 25.5%. Soil pH is estimated at 6.47 (on a scale of 0-14). Available nitrogen is estimated at 223 kg/ha, and available phosphorus is estimated at 7.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Udhampur district, Jammu And Kashmir. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 40.2% (Medium fertility class). Soil moisture content is estimated at 27.4%. Soil pH is estimated at 6.44 (on a scale of 0-14). Available nitrogen is estimated at 220 kg/ha, and available phosphorus is estimated at 7.4 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Jharkhand

Bokaro district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 55.0% (High fertility class). Soil moisture content is estimated at 13.2%. Soil pH is estimated at 6.01 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 7.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Chatra district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 41.4% (Medium fertility class). Soil moisture content is estimated at 16.2%. Soil pH is estimated at 6.31 (on a scale of 0-14). Available nitrogen is estimated at 212 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Deoghar district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 39.5% (Medium fertility class). Soil moisture content is estimated at 13.4%. Soil pH is estimated at 6.53 (on a scale of 0-14). Available nitrogen is estimated at 139 kg/ha, and available phosphorus is estimated at 14.8 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Dhanbad district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 52.6% (Medium fertility class). Soil moisture content is estimated at 13.1%. Soil pH is estimated at 6.58 (on a scale of 0-14). Available nitrogen is estimated at 129 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Dumka district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 50.1% (Medium fertility class). Soil moisture content is estimated at 17.0%. Soil pH is estimated at 6.39 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 15.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Garhwa district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 60.0% (High fertility class). Soil moisture content is estimated at 17.2%. Soil pH is estimated at 5.62 (on a scale of 0-14). Available nitrogen is estimated at 190 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Giridih district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 45.0% (Medium fertility class). Soil moisture content is estimated at 16.6%. Soil pH is estimated at 6.05 (on a scale of 0-14). Available nitrogen is estimated at 115 kg/ha, and available phosphorus is estimated at 12.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Godda district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 45.8% (Medium fertility class). Soil moisture content is estimated at 19.9%. Soil pH is estimated at 6.58 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Gumla district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 45.1% (Medium fertility class). Soil moisture content is estimated at 19.5%. Soil pH is estimated at 6.54 (on a scale of 0-14). Available nitrogen is estimated at 164 kg/ha, and available phosphorus is estimated at 14.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Hazaribagh district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 41.2% (Medium fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 5.62 (on a scale of 0-14). Available nitrogen is estimated at 126 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Jamtara district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 39.5% (Medium fertility class). Soil moisture content is estimated at 12.5%. Soil pH is estimated at 5.5 (on a scale of 0-14). Available nitrogen is estimated at 157 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Khunti district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 38.8% (Medium fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 7.5 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Kodarma district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 56.9% (High fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 6.2 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 6.9 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Latehar district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 46.7% (Medium fertility class). Soil moisture content is estimated at 19.4%. Soil pH is estimated at 5.83 (on a scale of 0-14). Available nitrogen is estimated at 192 kg/ha, and available phosphorus is estimated at 11.0 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Lohardaga district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 54.5% (Medium fertility class). Soil moisture content is estimated at 16.6%. Soil pH is estimated at 5.98 (on a scale of 0-14). Available nitrogen is estimated at 134 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Pakur district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 54.5% (Medium fertility class). Soil moisture content is estimated at 13.2%. Soil pH is estimated at 5.61 (on a scale of 0-14). Available nitrogen is estimated at 129 kg/ha, and available phosphorus is estimated at 7.8 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Palamu district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 16.0%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 125 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Pashchimi Singhbhum district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 47.9% (Medium fertility class). Soil moisture content is estimated at 15.9%. Soil pH is estimated at 6.64 (on a scale of 0-14). Available nitrogen is estimated at 217 kg/ha, and available phosphorus is estimated at 14.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Purbi Singhbhum district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 42.2% (Medium fertility class). Soil moisture content is estimated at 14.2%. Soil pH is estimated at 6.35 (on a scale of 0-14). Available nitrogen is estimated at 141 kg/ha, and available phosphorus is estimated at 13.3 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Ramgarh district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 50.5% (Medium fertility class). Soil moisture content is estimated at 17.5%. Soil pH is estimated at 6.58 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 6.3 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Ranchi district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 39.1% (Medium fertility class). Soil moisture content is estimated at 18.0%. Soil pH is estimated at 6.13 (on a scale of 0-14). Available nitrogen is estimated at 138 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Sahibganj district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 54.7% (Medium fertility class). Soil moisture content is estimated at 20.3%. Soil pH is estimated at 6.23 (on a scale of 0-14). Available nitrogen is estimated at 152 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Saraikela-Kharsawan district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 49.5% (Medium fertility class). Soil moisture content is estimated at 15.1%. Soil pH is estimated at 6.23 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 6.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Simdega district, Jharkhand. Dominant soil type: Red & Yellow Soil. Soil fertility is estimated at 38.0% (Medium fertility class). Soil moisture content is estimated at 14.1%. Soil pH is estimated at 6.66 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 7.7 kg/ha. [Basis: indicative regional estimate for Red & Yellow Soil, not an individual lab measurement.]

Karnataka

Bagalkot district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.5% (High fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 6.82 (on a scale of 0-14). Available nitrogen is estimated at 242 kg/ha, and available phosphorus is estimated at 9.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Bangalore district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.9% (Medium fertility class). Soil moisture content is estimated at 21.0%. Soil pH is estimated at 6.45 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 11.8

kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Bangalore Rural district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 47.3% (Medium fertility class). Soil moisture content is estimated at 16.5%. Soil pH is estimated at 6.75 (on a scale of 0-14). Available nitrogen is estimated at 159 kg/ha, and available phosphorus is estimated at 10.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Belgaum district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.8% (High fertility class). Soil moisture content is estimated at 17.8%. Soil pH is estimated at 6.67 (on a scale of 0-14). Available nitrogen is estimated at 146 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Bellary district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 51.1% (Medium fertility class). Soil moisture content is estimated at 21.3%. Soil pH is estimated at 6.47 (on a scale of 0-14). Available nitrogen is estimated at 237 kg/ha, and available phosphorus is estimated at 21.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Bidar district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 47.3% (Medium fertility class). Soil moisture content is estimated at 15.3%. Soil pH is estimated at 6.35 (on a scale of 0-14). Available nitrogen is estimated at 208 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Bijapur district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.6% (High fertility class). Soil moisture content is estimated at 23.1%. Soil pH is estimated at 7.27 (on a scale of 0-14). Available nitrogen is estimated at 197 kg/ha, and available phosphorus is estimated at 22.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chamarajanagar district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 46.0% (Medium fertility class). Soil moisture content is estimated at 18.8%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 244 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chikkaballapura district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 63.5% (High fertility class). Soil moisture content is estimated at 16.0%. Soil pH is estimated at 7.47 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chikmagalur district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 67.3% (High fertility class). Soil moisture content is estimated at 18.6%. Soil pH is estimated at 6.72 (on a scale of 0-14). Available nitrogen is estimated at 204 kg/ha, and available phosphorus is estimated at 10.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chitradurga district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 52.4% (Medium fertility class). Soil moisture content is estimated at 16.3%. Soil pH is estimated at 6.33 (on a scale of 0-14). Available nitrogen is estimated at 235 kg/ha, and available phosphorus is estimated at 16.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Dakshina Kannada district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 53.2% (Medium fertility class). Soil moisture content is estimated at 18.9%. Soil pH is estimated at 6.76 (on a scale of 0-14). Available nitrogen is estimated at 209 kg/ha, and available phosphorus is estimated at 21.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Davanagere district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 66.5% (High fertility class). Soil moisture content is estimated at 18.7%. Soil pH is estimated at 6.63 (on a scale of

0-14). Available nitrogen is estimated at 189 kg/ha, and available phosphorus is estimated at 16.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Dharwad district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 6.92 (on a scale of 0-14). Available nitrogen is estimated at 175 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Gadag district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 53.7% (Medium fertility class). Soil moisture content is estimated at 15.1%. Soil pH is estimated at 6.42 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 14.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Gulbarga district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 49.9% (Medium fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 7.18 (on a scale of 0-14). Available nitrogen is estimated at 168 kg/ha, and available phosphorus is estimated at 19.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Hassan district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 66.3% (High fertility class). Soil moisture content is estimated at 21.2%. Soil pH is estimated at 7.07 (on a scale of 0-14). Available nitrogen is estimated at 145 kg/ha, and available phosphorus is estimated at 21.3 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Haveri district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 54.0% (Medium fertility class). Soil moisture content is estimated at 25.1%. Soil pH is estimated at 6.95 (on a scale of 0-14). Available nitrogen is estimated at 140 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kodagu district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 60.6% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 6.23 (on a scale of 0-14). Available nitrogen is estimated at 221 kg/ha, and available phosphorus is estimated at 20.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kolar district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 6.21 (on a scale of 0-14). Available nitrogen is estimated at 226 kg/ha, and available phosphorus is estimated at 13.3 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Koppal district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.2% (High fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 6.93 (on a scale of 0-14). Available nitrogen is estimated at 169 kg/ha, and available phosphorus is estimated at 11.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Mandya district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 62.8% (High fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 6.9 (on a scale of 0-14). Available nitrogen is estimated at 246 kg/ha, and available phosphorus is estimated at 21.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Mysore district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.7% (High fertility class). Soil moisture content is estimated at 16.3%. Soil pH is estimated at 6.97 (on a scale of 0-14). Available nitrogen is estimated at 236 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Raichur district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 47.3% (Medium fertility class). Soil moisture content is estimated at 23.7%. Soil pH is estimated at 6.83 (on a scale of 0-14). Available nitrogen is estimated at 202 kg/ha, and available phosphorus is estimated at 21.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Ramanagara district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 59.0% (High fertility class). Soil moisture content is estimated at 19.1%. Soil pH is estimated at 6.68 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Shimoga district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 63.3% (High fertility class). Soil moisture content is estimated at 24.1%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 173 kg/ha, and available phosphorus is estimated at 20.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tumkur district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 45.6% (Medium fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 6.8 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 11.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Udupi district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 60.4% (High fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 7.0 (on a scale of 0-14). Available nitrogen is estimated at 144 kg/ha, and available phosphorus is estimated at 15.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Uttara Kannada district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 57.0% (High fertility class). Soil moisture content is estimated at 16.6%. Soil pH is estimated at 6.38 (on a scale of 0-14). Available nitrogen is estimated at 160 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Yadgir district, Karnataka. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 47.2% (Medium fertility class). Soil moisture content is estimated at 25.4%. Soil pH is estimated at 6.73 (on a scale of 0-14). Available nitrogen is estimated at 228 kg/ha, and available phosphorus is estimated at 21.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kerala

Alappuzha district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 33.9% (Low fertility class). Soil moisture content is estimated at 29.8%. Soil pH is estimated at 5.63 (on a scale of 0-14). Available nitrogen is estimated at 153 kg/ha, and available phosphorus is estimated at 7.4 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Ernakulam district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 41.9% (Medium fertility class). Soil moisture content is estimated at 20.5%. Soil pH is estimated at 5.19 (on a scale of 0-14). Available nitrogen is estimated at 189 kg/ha, and available phosphorus is estimated at 11.7 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Idukki district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 32.9% (Low fertility class). Soil moisture content is estimated at 30.6%. Soil pH is estimated at 4.89 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 4.4 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Kannur district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 53.5% (Medium fertility class). Soil moisture content is estimated at 32.6%. Soil pH is estimated at 4.65 (on a scale of 0-14). Available nitrogen is estimated at 95 kg/ha, and available phosphorus is estimated at 10.7 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Kasaragod district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 35.6% (Medium fertility class). Soil moisture content is estimated at 28.0%. Soil pH is estimated at 5.69 (on a scale of 0-14). Available nitrogen is estimated at 146 kg/ha, and available phosphorus is estimated at 7.2 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Kollam district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 42.2% (Medium fertility class). Soil moisture content is estimated at 20.9%. Soil pH is estimated at 5.76 (on a scale of 0-14). Available nitrogen is estimated at 121 kg/ha, and available phosphorus is estimated at 5.5 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Kottayam district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 38.6% (Medium fertility class). Soil moisture content is estimated at 34.5%. Soil pH is estimated at 4.62 (on a scale of 0-14). Available nitrogen is estimated at 176 kg/ha, and available phosphorus is estimated at 9.7 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Kozhikode district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 51.1% (Medium fertility class). Soil moisture content is estimated at 34.6%. Soil pH is estimated at 5.59 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 5.0 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Malappuram district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 50.3% (Medium fertility class). Soil moisture content is estimated at 28.2%. Soil pH is estimated at 5.21 (on a scale of 0-14). Available nitrogen is estimated at 179 kg/ha, and available phosphorus is estimated at 11.8 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Palakkad district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 48.2% (Medium fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 5.09 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 6.9 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Pathanamthitta district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 41.0% (Medium fertility class). Soil moisture content is estimated at 21.3%. Soil pH is estimated at 4.6 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 7.6 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Thiruvananthapuram district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 53.4% (Medium fertility class). Soil moisture content is estimated at 29.2%. Soil pH is estimated at 4.55 (on a scale of 0-14). Available nitrogen is estimated at 105 kg/ha, and available phosphorus is estimated at 5.6 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Thrissur district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 54.4% (Medium fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 5.17 (on a scale of 0-14). Available nitrogen is estimated at 183 kg/ha, and available phosphorus is estimated at 11.8 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Wayanad district, Kerala. Dominant soil type: Laterite Soil. Soil fertility is estimated at 37.4% (Medium fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 4.7 (on a scale of 0-14). Available nitrogen is estimated at 96 kg/ha, and available phosphorus is estimated at 5.6 kg/ha. [Basis: indicative regional estimate for Laterite Soil, not an individual lab measurement.]

Lakshadweep

Lakshadweep district, Lakshadweep. Dominant soil type: Coastal Sandy Soil. Soil fertility is estimated at 34.7% (Low fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 7.2 (on a scale of 0-14). Available nitrogen is estimated at 110 kg/ha, and available phosphorus is estimated at 5.2 kg/ha. [Basis: indicative regional estimate for Coastal Sandy Soil, not an individual lab measurement.]

Madhya Pradesh

Alirajpur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 70.0% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 8.12 (on a scale of 0-14). Available nitrogen is estimated at 273 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Anuppur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 52.3% (Medium fertility class). Soil moisture content is estimated at 23.4%. Soil pH is estimated at 7.57 (on a scale of 0-14). Available nitrogen is estimated at 272 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Ashoknagar district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.3% (High fertility class). Soil moisture content is estimated at 28.1%. Soil pH is estimated at 7.24 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Balaghat district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 66.8% (High fertility class). Soil moisture content is estimated at 26.5%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 243 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Barwani district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 73.4% (High fertility class). Soil moisture content is estimated at 22.8%. Soil pH is estimated at 7.14 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 14.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Betul district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.7% (High fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 8.29 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bhind district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 60.5% (High fertility class). Soil moisture content is estimated at 21.0%. Soil pH is estimated at 7.77 (on a scale of 0-14). Available nitrogen is estimated at 179 kg/ha, and available phosphorus is estimated at 15.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bhopal district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.3% (Medium fertility class). Soil moisture content is estimated at 34.7%. Soil pH is estimated at 7.61 (on a scale of 0-14). Available nitrogen is estimated at 266 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Burhanpur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.5% (Medium fertility class). Soil moisture content is estimated at 32.3%. Soil pH is estimated at 7.84 (on a scale of 0-14). Available nitrogen is estimated at 251 kg/ha, and available phosphorus is estimated at

13.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Chhatarpur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 65.9% (High fertility class). Soil moisture content is estimated at 32.8%. Soil pH is estimated at 7.82 (on a scale of 0-14). Available nitrogen is estimated at 195 kg/ha, and available phosphorus is estimated at 16.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Chhindwara district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 66.8% (High fertility class). Soil moisture content is estimated at 34.4%. Soil pH is estimated at 7.48 (on a scale of 0-14). Available nitrogen is estimated at 205 kg/ha, and available phosphorus is estimated at 17.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Damoh district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.6% (High fertility class). Soil moisture content is estimated at 31.0%. Soil pH is estimated at 7.6 (on a scale of 0-14). Available nitrogen is estimated at 178 kg/ha, and available phosphorus is estimated at 16.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Datia district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 53.3% (Medium fertility class). Soil moisture content is estimated at 23.0%. Soil pH is estimated at 7.81 (on a scale of 0-14). Available nitrogen is estimated at 247 kg/ha, and available phosphorus is estimated at 18.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Dewas district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.8% (High fertility class). Soil moisture content is estimated at 21.7%. Soil pH is estimated at 8.24 (on a scale of 0-14). Available nitrogen is estimated at 170 kg/ha, and available phosphorus is estimated at 10.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Dhar district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.3% (High fertility class). Soil moisture content is estimated at 27.9%. Soil pH is estimated at 8.15 (on a scale of 0-14). Available nitrogen is estimated at 248 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Dindori district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 54.4% (Medium fertility class). Soil moisture content is estimated at 25.4%. Soil pH is estimated at 7.62 (on a scale of 0-14). Available nitrogen is estimated at 169 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Guna district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.7% (High fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 241 kg/ha, and available phosphorus is estimated at 8.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Gwalior district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.0% (High fertility class). Soil moisture content is estimated at 31.7%. Soil pH is estimated at 7.97 (on a scale of 0-14). Available nitrogen is estimated at 255 kg/ha, and available phosphorus is estimated at 13.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Harda district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 70.5% (High fertility class). Soil moisture content is estimated at 23.6%. Soil pH is estimated at 7.47 (on a scale of 0-14). Available nitrogen is estimated at 172 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Hoshangabad district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 59.8% (High fertility class). Soil moisture content is estimated at 23.6%. Soil pH is estimated at 7.98 (on

a scale of 0-14). Available nitrogen is estimated at 190 kg/ha, and available phosphorus is estimated at 11.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Indore district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 53.3% (Medium fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 7.39 (on a scale of 0-14). Available nitrogen is estimated at 198 kg/ha, and available phosphorus is estimated at 10.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Jabalpur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.1% (High fertility class). Soil moisture content is estimated at 34.0%. Soil pH is estimated at 7.65 (on a scale of 0-14). Available nitrogen is estimated at 233 kg/ha, and available phosphorus is estimated at 18.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Jhabua district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 75.0% (High fertility class). Soil moisture content is estimated at 29.2%. Soil pH is estimated at 8.16 (on a scale of 0-14). Available nitrogen is estimated at 268 kg/ha, and available phosphorus is estimated at 17.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Katni district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 61.5% (High fertility class). Soil moisture content is estimated at 21.2%. Soil pH is estimated at 7.02 (on a scale of 0-14). Available nitrogen is estimated at 212 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Khandwa (East Nimar) district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 60.2% (High fertility class). Soil moisture content is estimated at 28.5%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 258 kg/ha, and available phosphorus is estimated at 9.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Khargone (West Nimar) district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 31.9%. Soil pH is estimated at 8.1 (on a scale of 0-14). Available nitrogen is estimated at 279 kg/ha, and available phosphorus is estimated at 18.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Mandla district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 72.2% (High fertility class). Soil moisture content is estimated at 24.7%. Soil pH is estimated at 7.52 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 15.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Mandsaur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 25.2%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Morena district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 54.9% (Medium fertility class). Soil moisture content is estimated at 20.9%. Soil pH is estimated at 8.23 (on a scale of 0-14). Available nitrogen is estimated at 202 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Narsimhapur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 67.0% (High fertility class). Soil moisture content is estimated at 22.9%. Soil pH is estimated at 7.23 (on a scale of 0-14). Available nitrogen is estimated at 255 kg/ha, and available phosphorus is estimated at 11.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Neemuch district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 72.8% (High fertility class). Soil moisture content is estimated at 22.1%. Soil pH is estimated at 7.88 (on a scale of 0-14). Available nitrogen is estimated at 255 kg/ha, and available phosphorus is estimated at 10.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Panna district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.9% (High fertility class). Soil moisture content is estimated at 34.5%. Soil pH is estimated at 7.34 (on a scale of 0-14). Available nitrogen is estimated at 205 kg/ha, and available phosphorus is estimated at 13.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Raisen district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.3% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 8.0 (on a scale of 0-14). Available nitrogen is estimated at 199 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Rajgarh district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.3% (High fertility class). Soil moisture content is estimated at 20.1%. Soil pH is estimated at 7.74 (on a scale of 0-14). Available nitrogen is estimated at 279 kg/ha, and available phosphorus is estimated at 8.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Ratlam district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.2% (High fertility class). Soil moisture content is estimated at 27.8%. Soil pH is estimated at 8.05 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 12.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Rewa district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 68.7% (High fertility class). Soil moisture content is estimated at 25.6%. Soil pH is estimated at 7.25 (on a scale of 0-14). Available nitrogen is estimated at 175 kg/ha, and available phosphorus is estimated at 12.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sagar district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 62.0% (High fertility class). Soil moisture content is estimated at 32.9%. Soil pH is estimated at 7.83 (on a scale of 0-14). Available nitrogen is estimated at 239 kg/ha, and available phosphorus is estimated at 19.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Satna district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 60.3% (High fertility class). Soil moisture content is estimated at 26.1%. Soil pH is estimated at 8.2 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 16.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sehore district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 68.9% (High fertility class). Soil moisture content is estimated at 21.5%. Soil pH is estimated at 7.61 (on a scale of 0-14). Available nitrogen is estimated at 176 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Seoni district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 61.6% (High fertility class). Soil moisture content is estimated at 24.3%. Soil pH is estimated at 8.03 (on a scale of 0-14). Available nitrogen is estimated at 267 kg/ha, and available phosphorus is estimated at 17.6 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Shahdol district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.7% (High fertility class). Soil moisture content is estimated at 24.1%. Soil pH is estimated at 7.33 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 19.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Shajapur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 60.1% (High fertility class). Soil moisture content is estimated at 23.4%. Soil pH is estimated at 8.1 (on a scale of 0-14). Available nitrogen is estimated at 201 kg/ha, and available phosphorus is estimated at 8.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sheopur district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 54.0% (Medium fertility class). Soil moisture content is estimated at 29.6%. Soil pH is estimated at 8.02 (on a scale of 0-14). Available nitrogen is estimated at 177 kg/ha, and available phosphorus is estimated at 19.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Shivpuri district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 32.7%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 156 kg/ha, and available phosphorus is estimated at 18.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sidhi district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.6% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 7.99 (on a scale of 0-14). Available nitrogen is estimated at 166 kg/ha, and available phosphorus is estimated at 17.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Singrauli district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 30.9%. Soil pH is estimated at 7.57 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 14.3 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Tikamgarh district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 68.6% (High fertility class). Soil moisture content is estimated at 32.3%. Soil pH is estimated at 7.97 (on a scale of 0-14). Available nitrogen is estimated at 187 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Ujjain district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.0% (High fertility class). Soil moisture content is estimated at 27.5%. Soil pH is estimated at 7.66 (on a scale of 0-14). Available nitrogen is estimated at 225 kg/ha, and available phosphorus is estimated at 12.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Umari district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.7% (High fertility class). Soil moisture content is estimated at 33.7%. Soil pH is estimated at 7.32 (on a scale of 0-14). Available nitrogen is estimated at 245 kg/ha, and available phosphorus is estimated at 16.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Vidisha district, Madhya Pradesh. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 63.1% (High fertility class). Soil moisture content is estimated at 22.8%. Soil pH is estimated at 8.03 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 16.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Maharashtra

Ahmadnagar district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 72.9% (High fertility class). Soil moisture content is estimated at 34.8%. Soil pH is estimated at 7.46 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Akola district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 74.7% (High fertility class). Soil moisture content is estimated at 21.4%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 11.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Amravati district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.7% (High fertility class). Soil moisture content is estimated at 25.7%. Soil pH is estimated at 8.27 (on a scale of 0-14). Available nitrogen is estimated at 264 kg/ha, and available phosphorus is estimated at 11.6 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Aurangabad district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.3% (High fertility class). Soil moisture content is estimated at 25.1%. Soil pH is estimated at 7.79 (on a scale of 0-14). Available nitrogen is estimated at 235 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bhandara district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.6% (Medium fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 7.93 (on a scale of 0-14). Available nitrogen is estimated at 223 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Bid district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.4% (Medium fertility class). Soil moisture content is estimated at 25.2%. Soil pH is estimated at 8.12 (on a scale of 0-14). Available nitrogen is estimated at 154 kg/ha, and available phosphorus is estimated at 12.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Buldana district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 73.6% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 7.97 (on a scale of 0-14). Available nitrogen is estimated at 183 kg/ha, and available phosphorus is estimated at 18.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Chandrapur district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 68.7% (High fertility class). Soil moisture content is estimated at 33.3%. Soil pH is estimated at 7.28 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 17.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Dhule district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 23.5%. Soil pH is estimated at 7.04 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Gadchiroli district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 65.9% (High fertility class). Soil moisture content is estimated at 31.2%. Soil pH is estimated at 7.69 (on a scale of 0-14). Available nitrogen is estimated at 211 kg/ha, and available phosphorus is estimated at 13.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Gondiya district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 66.0% (High fertility class). Soil moisture content is estimated at 23.0%. Soil pH is estimated at 8.15 (on a scale of 0-14). Available nitrogen is estimated at 255 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Hingoli district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 64.5% (High fertility class). Soil moisture content is estimated at 22.1%. Soil pH is estimated at 8.26 (on a scale of 0-14). Available nitrogen is estimated at 267 kg/ha, and available phosphorus is estimated at 19.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Jalgaon district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 58.3% (High fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 7.28 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Jalna district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.8% (Medium fertility class). Soil moisture content is estimated at 34.8%. Soil pH is estimated at 7.1 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 17.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Kolhapur district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 64.5% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 7.82 (on a scale of 0-14). Available nitrogen is estimated at 264 kg/ha, and available phosphorus is estimated at 15.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Latur district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 63.1% (High fertility class). Soil moisture content is estimated at 30.7%. Soil pH is estimated at 7.58 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Mumbai district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 54.2% (Medium fertility class). Soil moisture content is estimated at 31.9%. Soil pH is estimated at 7.34 (on a scale of 0-14). Available nitrogen is estimated at 169 kg/ha, and available phosphorus is estimated at 11.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Mumbai Suburban district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 71.1% (High fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 7.89 (on a scale of 0-14). Available nitrogen is estimated at 261 kg/ha, and available phosphorus is estimated at 8.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Nagpur district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 69.1% (High fertility class). Soil moisture content is estimated at 24.6%. Soil pH is estimated at 7.28 (on a scale of 0-14). Available nitrogen is estimated at 239 kg/ha, and available phosphorus is estimated at 18.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Nanded district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.4% (High fertility class). Soil moisture content is estimated at 28.5%. Soil pH is estimated at 7.38 (on a scale of 0-14). Available nitrogen is estimated at 222 kg/ha, and available phosphorus is estimated at 15.5 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Nandurbar district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 65.3% (High fertility class). Soil moisture content is estimated at 29.4%. Soil pH is estimated at 7.18 (on a scale of 0-14). Available nitrogen is estimated at 240 kg/ha, and available phosphorus is estimated at 16.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Nashik district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 67.1% (High fertility class). Soil moisture content is estimated at 21.1%. Soil pH is estimated at 7.58 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 9.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Osmanabad district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 7.97 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 18.0

kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Parbhani district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 63.6% (High fertility class). Soil moisture content is estimated at 31.4%. Soil pH is estimated at 7.61 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Pune district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 73.2% (High fertility class). Soil moisture content is estimated at 27.4%. Soil pH is estimated at 7.69 (on a scale of 0-14). Available nitrogen is estimated at 228 kg/ha, and available phosphorus is estimated at 16.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Raigarh district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 51.8% (Medium fertility class). Soil moisture content is estimated at 27.7%. Soil pH is estimated at 7.63 (on a scale of 0-14). Available nitrogen is estimated at 269 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Ratnagiri district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 61.7% (High fertility class). Soil moisture content is estimated at 34.4%. Soil pH is estimated at 7.59 (on a scale of 0-14). Available nitrogen is estimated at 173 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sangli district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 52.4% (Medium fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 7.56 (on a scale of 0-14). Available nitrogen is estimated at 250 kg/ha, and available phosphorus is estimated at 16.8 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Satara district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 50.8% (Medium fertility class). Soil moisture content is estimated at 28.5%. Soil pH is estimated at 7.2 (on a scale of 0-14). Available nitrogen is estimated at 272 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Sindhudurg district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 65.7% (High fertility class). Soil moisture content is estimated at 24.9%. Soil pH is estimated at 7.83 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 17.1 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Solapur district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 55.8% (High fertility class). Soil moisture content is estimated at 33.6%. Soil pH is estimated at 7.65 (on a scale of 0-14). Available nitrogen is estimated at 248 kg/ha, and available phosphorus is estimated at 18.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Thane district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 75.0% (High fertility class). Soil moisture content is estimated at 23.4%. Soil pH is estimated at 7.37 (on a scale of 0-14). Available nitrogen is estimated at 182 kg/ha, and available phosphorus is estimated at 11.2 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Wardha district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 64.6% (High fertility class). Soil moisture content is estimated at 34.9%. Soil pH is estimated at 8.29 (on a scale of 0-14). Available nitrogen is estimated at 218 kg/ha, and available phosphorus is estimated at 15.7 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Washim district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 71.5% (High fertility class). Soil moisture content is estimated at 34.0%. Soil pH is estimated at 7.27 (on a scale of

0-14). Available nitrogen is estimated at 218 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Yavatmal district, Maharashtra. Dominant soil type: Black Soil (Regur). Soil fertility is estimated at 57.6% (High fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 8.28 (on a scale of 0-14). Available nitrogen is estimated at 278 kg/ha, and available phosphorus is estimated at 18.9 kg/ha. [Basis: indicative regional estimate for Black Soil (Regur), not an individual lab measurement.]

Manipur

Bishnupur district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 53.4% (Medium fertility class). Soil moisture content is estimated at 21.8%. Soil pH is estimated at 5.25 (on a scale of 0-14). Available nitrogen is estimated at 154 kg/ha, and available phosphorus is estimated at 10.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Chandel district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 46.6% (Medium fertility class). Soil moisture content is estimated at 26.9%. Soil pH is estimated at 5.37 (on a scale of 0-14). Available nitrogen is estimated at 174 kg/ha, and available phosphorus is estimated at 8.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Churachandpur district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 42.6% (Medium fertility class). Soil moisture content is estimated at 19.8%. Soil pH is estimated at 5.06 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 9.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Imphal East district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 40.2% (Medium fertility class). Soil moisture content is estimated at 28.6%. Soil pH is estimated at 5.58 (on a scale of 0-14). Available nitrogen is estimated at 102 kg/ha, and available phosphorus is estimated at 7.8 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Imphal West district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 39.1% (Medium fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 5.19 (on a scale of 0-14). Available nitrogen is estimated at 106 kg/ha, and available phosphorus is estimated at 5.5 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Senapati district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 48.4% (Medium fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 5.99 (on a scale of 0-14). Available nitrogen is estimated at 115 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Tamenglong district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 45.2% (Medium fertility class). Soil moisture content is estimated at 24.2%. Soil pH is estimated at 4.9 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 13.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Thoubal district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 35.3% (Medium fertility class). Soil moisture content is estimated at 20.5%. Soil pH is estimated at 6.07 (on a scale of 0-14). Available nitrogen is estimated at 154 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Ukhrul district, Manipur. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 39.7% (Medium fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 5.81 (on a scale of 0-14). Available nitrogen is estimated at 159 kg/ha, and available phosphorus is estimated at 13.6

kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Meghalaya

East Garo Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 46.4% (Medium fertility class). Soil moisture content is estimated at 22.2%. Soil pH is estimated at 5.47 (on a scale of 0-14). Available nitrogen is estimated at 135 kg/ha, and available phosphorus is estimated at 7.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

East Khasi Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 36.1% (Medium fertility class). Soil moisture content is estimated at 22.9%. Soil pH is estimated at 5.8 (on a scale of 0-14). Available nitrogen is estimated at 101 kg/ha, and available phosphorus is estimated at 9.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Jaintia Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 49.8% (Medium fertility class). Soil moisture content is estimated at 26.5%. Soil pH is estimated at 5.89 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 11.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Ribhoi district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 37.4% (Medium fertility class). Soil moisture content is estimated at 24.7%. Soil pH is estimated at 5.64 (on a scale of 0-14). Available nitrogen is estimated at 124 kg/ha, and available phosphorus is estimated at 5.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

South Garo Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 40.1% (Medium fertility class). Soil moisture content is estimated at 24.4%. Soil pH is estimated at 5.38 (on a scale of 0-14). Available nitrogen is estimated at 117 kg/ha, and available phosphorus is estimated at 12.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

West Garo Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 56.3% (High fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 6.1 (on a scale of 0-14). Available nitrogen is estimated at 101 kg/ha, and available phosphorus is estimated at 13.3 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

West Khasi Hills district, Meghalaya. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 56.9% (High fertility class). Soil moisture content is estimated at 18.2%. Soil pH is estimated at 5.87 (on a scale of 0-14). Available nitrogen is estimated at 197 kg/ha, and available phosphorus is estimated at 13.2 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Mizoram

Aizawl district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 57.3% (High fertility class). Soil moisture content is estimated at 25.7%. Soil pH is estimated at 4.91 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Champhai district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 49.1% (Medium fertility class). Soil moisture content is estimated at 23.8%. Soil pH is estimated at 5.6 (on a scale of 0-14). Available nitrogen is estimated at 115 kg/ha, and available phosphorus is estimated at 11.7 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Kolasib district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 36.7% (Medium fertility class). Soil moisture content is estimated at 28.2%. Soil pH is estimated at 5.99 (on a scale of 0-14). Available nitrogen is estimated at 147 kg/ha, and available phosphorus is estimated at 9.6 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Lawngtlai district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 42.2% (Medium fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 5.83 (on a scale of 0-14). Available nitrogen is estimated at 136 kg/ha, and available phosphorus is estimated at 6.6 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Lunglei district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 40.1% (Medium fertility class). Soil moisture content is estimated at 24.5%. Soil pH is estimated at 6.12 (on a scale of 0-14). Available nitrogen is estimated at 173 kg/ha, and available phosphorus is estimated at 6.9 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Mamit district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 54.8% (Medium fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 5.99 (on a scale of 0-14). Available nitrogen is estimated at 124 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Saiha district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 56.3% (High fertility class). Soil moisture content is estimated at 21.6%. Soil pH is estimated at 5.77 (on a scale of 0-14). Available nitrogen is estimated at 198 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Serchhip district, Mizoram. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 37.0% (Medium fertility class). Soil moisture content is estimated at 27.4%. Soil pH is estimated at 5.28 (on a scale of 0-14). Available nitrogen is estimated at 107 kg/ha, and available phosphorus is estimated at 9.3 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Nagaland

Dimapur district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 52.3% (Medium fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 5.27 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 9.7 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Kiphire district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 54.4% (Medium fertility class). Soil moisture content is estimated at 24.2%. Soil pH is estimated at 6.13 (on a scale of 0-14). Available nitrogen is estimated at 165 kg/ha, and available phosphorus is estimated at 6.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Kohima district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 55.4% (High fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 5.0 (on a scale of 0-14). Available nitrogen is estimated at 115 kg/ha, and available phosphorus is estimated at 11.2 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Longleng district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 36.1% (Medium fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 5.0 (on a scale of 0-14). Available nitrogen is estimated at 144 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Mokokchung district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 47.0% (Medium fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 4.84 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 8.4 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Mon district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 50.0% (Medium fertility class). Soil moisture content is estimated at 19.5%. Soil pH is estimated at 4.82 (on a scale of 0-14). Available nitrogen is estimated at 124 kg/ha, and available phosphorus is estimated at 9.1 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Peren district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 44.1% (Medium fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 5.33 (on a scale of 0-14). Available nitrogen is estimated at 153 kg/ha, and available phosphorus is estimated at 10.3 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Phek district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 35.5% (Medium fertility class). Soil moisture content is estimated at 25.2%. Soil pH is estimated at 5.28 (on a scale of 0-14). Available nitrogen is estimated at 142 kg/ha, and available phosphorus is estimated at 5.9 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Tuensang district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 53.0% (Medium fertility class). Soil moisture content is estimated at 31.2%. Soil pH is estimated at 4.83 (on a scale of 0-14). Available nitrogen is estimated at 160 kg/ha, and available phosphorus is estimated at 11.9 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Wokha district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 41.6% (Medium fertility class). Soil moisture content is estimated at 29.0%. Soil pH is estimated at 5.69 (on a scale of 0-14). Available nitrogen is estimated at 192 kg/ha, and available phosphorus is estimated at 13.8 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Zunheboto district, Nagaland. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 36.0% (Medium fertility class). Soil moisture content is estimated at 24.2%. Soil pH is estimated at 5.57 (on a scale of 0-14). Available nitrogen is estimated at 114 kg/ha, and available phosphorus is estimated at 8.0 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Odisha

Anugul district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 57.7% (High fertility class). Soil moisture content is estimated at 15.2%. Soil pH is estimated at 5.68 (on a scale of 0-14). Available nitrogen is estimated at 192 kg/ha, and available phosphorus is estimated at 7.9 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Balangir district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 46.0% (Medium fertility class). Soil moisture content is estimated at 27.7%. Soil pH is estimated at 5.71 (on a scale of 0-14). Available nitrogen is estimated at 114 kg/ha, and available phosphorus is estimated at 5.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Baleshwar district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 56.7% (High fertility class). Soil moisture content is estimated at 18.5%. Soil pH is estimated at 5.76 (on a scale of 0-14). Available nitrogen is estimated at 109 kg/ha, and available phosphorus is estimated at 8.8 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Bargarh district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 21.5%. Soil pH is estimated at 6.46 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Baudh district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 54.8% (Medium fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 5.42 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 8.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Bhadrak district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 47.2% (Medium fertility class). Soil moisture content is estimated at 27.7%. Soil pH is estimated at 5.28 (on a scale of 0-14). Available nitrogen is estimated at 133 kg/ha, and available phosphorus is estimated at 5.3 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Cuttack district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 44.0% (Medium fertility class). Soil moisture content is estimated at 17.6%. Soil pH is estimated at 6.02 (on a scale of 0-14). Available nitrogen is estimated at 119 kg/ha, and available phosphorus is estimated at 11.1 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Debagarh district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 41.1% (Medium fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 5.52 (on a scale of 0-14). Available nitrogen is estimated at 178 kg/ha, and available phosphorus is estimated at 9.2 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Dhenkanal district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 43.3% (Medium fertility class). Soil moisture content is estimated at 26.1%. Soil pH is estimated at 5.86 (on a scale of 0-14). Available nitrogen is estimated at 123 kg/ha, and available phosphorus is estimated at 13.1 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Gajapati district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 42.9% (Medium fertility class). Soil moisture content is estimated at 15.5%. Soil pH is estimated at 5.74 (on a scale of 0-14). Available nitrogen is estimated at 113 kg/ha, and available phosphorus is estimated at 7.1 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Ganjam district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 52.2% (Medium fertility class). Soil moisture content is estimated at 16.7%. Soil pH is estimated at 6.21 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Jagatsinghapur district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 51.3% (Medium fertility class). Soil moisture content is estimated at 24.3%. Soil pH is estimated at 6.49 (on a scale of 0-14). Available nitrogen is estimated at 137 kg/ha, and available phosphorus is estimated at 6.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Jajapur district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 46.9% (Medium fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 5.27 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 5.1 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Jharsuguda district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 56.3% (High fertility class). Soil moisture content is estimated at 18.4%. Soil pH is estimated at 5.06 (on a scale of 0-14). Available nitrogen is estimated at 111 kg/ha, and available phosphorus is estimated at 7.2 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Kalahandi district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 41.3% (Medium fertility class). Soil moisture content is estimated at 15.4%. Soil pH is estimated at 5.42 (on a scale of 0-14). Available nitrogen is estimated at 121 kg/ha, and available phosphorus is estimated at 8.4 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Kandhamal district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 43.9% (Medium fertility class). Soil moisture content is estimated at 19.4%. Soil pH is estimated at 5.49 (on a scale of 0-14). Available nitrogen is estimated at 109 kg/ha, and available phosphorus is estimated at 5.2 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Kendrapara district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 39.6% (Medium fertility class). Soil moisture content is estimated at 24.8%. Soil pH is estimated at 6.49 (on a scale of 0-14). Available nitrogen is estimated at 202 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Kendujhar district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 47.2% (Medium fertility class). Soil moisture content is estimated at 19.1%. Soil pH is estimated at 6.16 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 12.6 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Khordha district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 20.6%. Soil pH is estimated at 6.15 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 7.8 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Koraput district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 16.2%. Soil pH is estimated at 6.49 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 5.8 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Malkangiri district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 50.1% (Medium fertility class). Soil moisture content is estimated at 15.4%. Soil pH is estimated at 6.16 (on a scale of 0-14). Available nitrogen is estimated at 187 kg/ha, and available phosphorus is estimated at 10.4 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Mayurbhanj district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 54.7% (Medium fertility class). Soil moisture content is estimated at 16.3%. Soil pH is estimated at 5.41 (on a scale of 0-14). Available nitrogen is estimated at 145 kg/ha, and available phosphorus is estimated at 5.4 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Nabarangapur district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 45.3% (Medium fertility class). Soil moisture content is estimated at 18.4%. Soil pH is estimated at 5.59 (on a scale of 0-14). Available nitrogen is estimated at 146 kg/ha, and available phosphorus is estimated at 12.9 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Nayagarh district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 41.2% (Medium fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 5.64 (on a scale of 0-14). Available nitrogen is estimated at 136 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Nuapada district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 44.6% (Medium fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 5.75 (on a scale of 0-14). Available nitrogen is estimated at 116 kg/ha, and available phosphorus is estimated at 8.2 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Puri district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 48.0% (Medium fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 6.24 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 13.7 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Rayagada district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 53.0% (Medium fertility class). Soil moisture content is estimated at 27.6%. Soil pH is estimated at 6.26 (on a scale of 0-14). Available nitrogen is estimated at 115 kg/ha, and available phosphorus is estimated at 9.7 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Sambalpur district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 38.9% (Medium fertility class). Soil moisture content is estimated at 25.6%. Soil pH is estimated at 5.88 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 13.4 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Subarnapur district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 37.6% (Medium fertility class). Soil moisture content is estimated at 22.3%. Soil pH is estimated at 6.08 (on a scale of 0-14). Available nitrogen is estimated at 106 kg/ha, and available phosphorus is estimated at 10.4 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Sundargarh district, Odisha. Dominant soil type: Red & Laterite Soil. Soil fertility is estimated at 53.8% (Medium fertility class). Soil moisture content is estimated at 15.9%. Soil pH is estimated at 5.77 (on a scale of 0-14). Available nitrogen is estimated at 128 kg/ha, and available phosphorus is estimated at 11.0 kg/ha. [Basis: indicative regional estimate for Red & Laterite Soil, not an individual lab measurement.]

Pondicherry

Karaikal district, Pondicherry. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 44.8% (Medium fertility class). Soil moisture content is estimated at 39.0%. Soil pH is estimated at 7.87 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 10.6 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Mahe district, Pondicherry. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 35.5%. Soil pH is estimated at 7.33 (on a scale of 0-14). Available nitrogen is estimated at 147 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Pondicherry district, Pondicherry. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 66.0% (High fertility class). Soil moisture content is estimated at 31.3%. Soil pH is estimated at 7.85 (on a scale of 0-14). Available nitrogen is estimated at 183 kg/ha, and available phosphorus is estimated at 17.1 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Yanam district, Pondicherry. Dominant soil type: Coastal Alluvial Soil. Soil fertility is estimated at 44.1% (Medium fertility class). Soil moisture content is estimated at 25.8%. Soil pH is estimated at 7.92 (on a scale of 0-14). Available nitrogen is estimated at 234 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Coastal Alluvial Soil, not an individual lab measurement.]

Punjab

Amritsar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.9% (High fertility class). Soil moisture content is estimated at 24.3%. Soil pH is estimated at 7.04 (on a scale of 0-14).

Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 26.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Barnala district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.3% (High fertility class). Soil moisture content is estimated at 27.9%. Soil pH is estimated at 6.59 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 22.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bathinda district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.5% (High fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 312 kg/ha, and available phosphorus is estimated at 24.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Faridkot district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 76.9% (High fertility class). Soil moisture content is estimated at 19.9%. Soil pH is estimated at 6.92 (on a scale of 0-14). Available nitrogen is estimated at 231 kg/ha, and available phosphorus is estimated at 20.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Fatehgarh Sahib district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.4% (High fertility class). Soil moisture content is estimated at 18.8%. Soil pH is estimated at 6.98 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 18.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Firozpur district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.9% (High fertility class). Soil moisture content is estimated at 21.7%. Soil pH is estimated at 7.32 (on a scale of 0-14). Available nitrogen is estimated at 234 kg/ha, and available phosphorus is estimated at 17.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gurdaspur district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.3% (High fertility class). Soil moisture content is estimated at 28.8%. Soil pH is estimated at 6.99 (on a scale of 0-14). Available nitrogen is estimated at 262 kg/ha, and available phosphorus is estimated at 27.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hoshiarpur district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.0% (High fertility class). Soil moisture content is estimated at 30.7%. Soil pH is estimated at 7.53 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 18.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jalandhar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.5% (High fertility class). Soil moisture content is estimated at 31.8%. Soil pH is estimated at 7.1 (on a scale of 0-14). Available nitrogen is estimated at 192 kg/ha, and available phosphorus is estimated at 21.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kapurthala district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.2% (High fertility class). Soil moisture content is estimated at 28.5%. Soil pH is estimated at 7.16 (on a scale of 0-14). Available nitrogen is estimated at 281 kg/ha, and available phosphorus is estimated at 25.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Ludhiana district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.7% (High fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 7.48 (on a scale of 0-14). Available nitrogen is estimated at 297 kg/ha, and available phosphorus is estimated at 25.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mansa district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 55.9% (High fertility class). Soil moisture content is estimated at 25.8%. Soil pH is estimated at 6.78 (on a scale of 0-14). Available nitrogen is estimated at 300 kg/ha, and available phosphorus is estimated at 22.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Moga district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.5% (High fertility class). Soil moisture content is estimated at 20.6%. Soil pH is estimated at 6.94 (on a scale of 0-14). Available nitrogen is estimated at 278 kg/ha, and available phosphorus is estimated at 15.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Muktsar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.3% (High fertility class). Soil moisture content is estimated at 30.4%. Soil pH is estimated at 7.49 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 16.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Patiala district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.3% (High fertility class). Soil moisture content is estimated at 27.9%. Soil pH is estimated at 7.29 (on a scale of 0-14). Available nitrogen is estimated at 223 kg/ha, and available phosphorus is estimated at 16.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rupnagar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.0% (High fertility class). Soil moisture content is estimated at 28.2%. Soil pH is estimated at 6.56 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 23.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sahibzada Ajit Singh Nagar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.7% (High fertility class). Soil moisture content is estimated at 28.6%. Soil pH is estimated at 7.11 (on a scale of 0-14). Available nitrogen is estimated at 248 kg/ha, and available phosphorus is estimated at 19.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sangrur district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.1% (High fertility class). Soil moisture content is estimated at 25.3%. Soil pH is estimated at 7.06 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 26.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Shahid Bhagat Singh Nagar district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.9% (High fertility class). Soil moisture content is estimated at 19.1%. Soil pH is estimated at 7.0 (on a scale of 0-14). Available nitrogen is estimated at 205 kg/ha, and available phosphorus is estimated at 17.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Tarn Taran district, Punjab. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.2% (High fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 7.49 (on a scale of 0-14). Available nitrogen is estimated at 249 kg/ha, and available phosphorus is estimated at 26.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rajasthan

Ajmer district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 18.7% (Low fertility class). Soil moisture content is estimated at 3.5%. Soil pH is estimated at 8.02 (on a scale of 0-14). Available nitrogen is estimated at 45 kg/ha, and available phosphorus is estimated at 4.8 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Alwar district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 37.7% (Medium fertility class). Soil moisture content is estimated at 4.5%. Soil pH is estimated at 7.76 (on a scale of 0-14). Available nitrogen is estimated at 80 kg/ha, and available phosphorus is estimated at 6.5 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Banswara district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 29.6% (Low fertility class). Soil moisture content is estimated at 5.6%. Soil pH is estimated at 8.57 (on a scale of 0-14). Available nitrogen is estimated at 49 kg/ha, and available phosphorus is estimated at 6.0 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Baran district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 31.8% (Low fertility class). Soil moisture content is estimated at 9.1%. Soil pH is estimated at 7.98 (on a scale of 0-14). Available nitrogen is estimated at 79 kg/ha, and available phosphorus is estimated at 3.5 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Barmer district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 21.4% (Low fertility class). Soil moisture content is estimated at 3.5%. Soil pH is estimated at 8.45 (on a scale of 0-14). Available nitrogen is estimated at 60 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Bharatpur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 19.3% (Low fertility class). Soil moisture content is estimated at 9.0%. Soil pH is estimated at 8.59 (on a scale of 0-14). Available nitrogen is estimated at 46 kg/ha, and available phosphorus is estimated at 3.2 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Bhilwara district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 19.1% (Low fertility class). Soil moisture content is estimated at 9.0%. Soil pH is estimated at 7.81 (on a scale of 0-14). Available nitrogen is estimated at 78 kg/ha, and available phosphorus is estimated at 6.0 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Bikaner district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 33.9% (Low fertility class). Soil moisture content is estimated at 6.4%. Soil pH is estimated at 8.15 (on a scale of 0-14). Available nitrogen is estimated at 109 kg/ha, and available phosphorus is estimated at 4.1 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Bundi district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 34.6% (Low fertility class). Soil moisture content is estimated at 3.4%. Soil pH is estimated at 8.12 (on a scale of 0-14). Available nitrogen is estimated at 102 kg/ha, and available phosphorus is estimated at 7.7 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Chittaurgarh district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 31.7% (Low fertility class). Soil moisture content is estimated at 4.8%. Soil pH is estimated at 8.04 (on a scale of 0-14). Available nitrogen is estimated at 65 kg/ha, and available phosphorus is estimated at 6.7 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Churu district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 27.2% (Low fertility class). Soil moisture content is estimated at 5.3%. Soil pH is estimated at 7.9 (on a scale of 0-14). Available nitrogen is estimated at 57 kg/ha, and available phosphorus is estimated at 5.6 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Dausa district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 21.0% (Low fertility class). Soil moisture content is estimated at 9.1%. Soil pH is estimated at 8.28 (on a scale of 0-14). Available nitrogen is estimated at 49 kg/ha, and available phosphorus is estimated at 7.9 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Dhaulpur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 35.0% (Medium fertility class). Soil moisture content is estimated at 4.7%. Soil pH is estimated at 7.7 (on a scale of 0-14). Available nitrogen is estimated at 40 kg/ha, and available phosphorus is estimated at 3.3 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Dungarpur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 33.2% (Low fertility class). Soil moisture content is estimated at 9.9%. Soil pH is estimated at 7.56 (on a scale of 0-14). Available nitrogen is estimated at 92 kg/ha, and available phosphorus is estimated at 3.5 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Ganganagar district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 29.8% (Low fertility class). Soil moisture content is estimated at 8.3%. Soil pH is estimated at 7.97 (on a scale of 0-14). Available nitrogen is estimated at 40 kg/ha, and available phosphorus is estimated at 7.2 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Hanumangarh district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 37.8% (Medium fertility class). Soil moisture content is estimated at 4.8%. Soil pH is estimated at 7.75 (on a scale of 0-14). Available nitrogen is estimated at 48 kg/ha, and available phosphorus is estimated at 6.9 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jaipur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 29.9% (Low fertility class). Soil moisture content is estimated at 8.5%. Soil pH is estimated at 8.48 (on a scale of 0-14). Available nitrogen is estimated at 102 kg/ha, and available phosphorus is estimated at 8.6 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jaisalmer district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 22.9% (Low fertility class). Soil moisture content is estimated at 6.9%. Soil pH is estimated at 7.89 (on a scale of 0-14). Available nitrogen is estimated at 63 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jalor district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 31.6% (Low fertility class). Soil moisture content is estimated at 9.8%. Soil pH is estimated at 8.04 (on a scale of 0-14). Available nitrogen is estimated at 77 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jhalawar district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 20.1% (Low fertility class). Soil moisture content is estimated at 4.7%. Soil pH is estimated at 8.19 (on a scale of 0-14). Available nitrogen is estimated at 44 kg/ha, and available phosphorus is estimated at 9.5 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jhunjhunun district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 20.5% (Low fertility class). Soil moisture content is estimated at 4.4%. Soil pH is estimated at 8.49 (on a scale of 0-14). Available nitrogen is estimated at 46 kg/ha, and available phosphorus is estimated at 9.3 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Jodhpur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 32.5% (Low fertility class). Soil moisture content is estimated at 3.5%. Soil pH is estimated at 8.4 (on a scale of 0-14). Available nitrogen is estimated at 84 kg/ha, and available phosphorus is estimated at 8.3 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Karauli district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 34.6% (Low fertility class). Soil moisture content is estimated at 8.0%. Soil pH is estimated at 8.26 (on a scale of 0-14). Available nitrogen is estimated at 62 kg/ha, and available phosphorus is estimated at 7.9 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Kota district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 37.1% (Medium fertility class). Soil moisture content is estimated at 7.1%. Soil pH is estimated at 8.36 (on a scale of 0-14). Available nitrogen is estimated at 70 kg/ha, and available phosphorus is estimated at 8.3 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Nagaur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 38.0% (Medium fertility class). Soil moisture content is estimated at 4.1%. Soil pH is estimated at 7.76 (on a scale of 0-14). Available nitrogen is estimated at 77 kg/ha, and available phosphorus is estimated at 6.4 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Pali district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 29.4% (Low fertility class). Soil moisture content is estimated at 3.3%. Soil pH is estimated at 8.03 (on a scale of 0-14). Available nitrogen is estimated at 80 kg/ha, and available phosphorus is estimated at 7.1 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Pratapgarh district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 31.1% (Low fertility class). Soil moisture content is estimated at 8.0%. Soil pH is estimated at 8.33 (on a scale of 0-14). Available nitrogen is estimated at 95 kg/ha, and available phosphorus is estimated at 7.4 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Rajsamand district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 28.8% (Low fertility class). Soil moisture content is estimated at 7.2%. Soil pH is estimated at 7.63 (on a scale of 0-14). Available nitrogen is estimated at 59 kg/ha, and available phosphorus is estimated at 6.7 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Sawai Madhopur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 35.5% (Medium fertility class). Soil moisture content is estimated at 7.8%. Soil pH is estimated at 7.62 (on a scale of 0-14). Available nitrogen is estimated at 81 kg/ha, and available phosphorus is estimated at 4.6 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Sikar district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 32.7% (Low fertility class). Soil moisture content is estimated at 8.8%. Soil pH is estimated at 8.27 (on a scale of 0-14). Available nitrogen is estimated at 81 kg/ha, and available phosphorus is estimated at 5.4 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Sirohi district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 26.9% (Low fertility class). Soil moisture content is estimated at 9.0%. Soil pH is estimated at 8.44 (on a scale of 0-14). Available nitrogen is estimated at 76 kg/ha, and available phosphorus is estimated at 5.6 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Tonk district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 19.0% (Low fertility class). Soil moisture content is estimated at 4.0%. Soil pH is estimated at 7.75 (on a scale of 0-14). Available nitrogen is estimated at 66 kg/ha, and available phosphorus is estimated at 9.6 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Udaipur district, Rajasthan. Dominant soil type: Arid & Desert Soil. Soil fertility is estimated at 28.1% (Low fertility class). Soil moisture content is estimated at 4.6%. Soil pH is estimated at 7.64 (on a scale of 0-14). Available nitrogen is estimated at 63 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Arid & Desert Soil, not an individual lab measurement.]

Sikkim

East District district, Sikkim. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 49.6% (Medium fertility class). Soil moisture content is estimated at 34.4%. Soil pH is estimated at 6.44 (on a scale of 0-14). Available nitrogen is estimated at 133 kg/ha, and available phosphorus is estimated at 12.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

North District district, Sikkim. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 64.0% (High fertility class). Soil moisture content is estimated at 28.4%. Soil pH is estimated at 5.09 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 14.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

South District district, Sikkim. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 53.1% (Medium fertility class). Soil moisture content is estimated at 30.8%. Soil pH is estimated at 5.38 (on a scale of 0-14). Available nitrogen is estimated at 218 kg/ha, and available phosphorus is estimated at 11.0 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

West District district, Sikkim. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 40.8% (Medium fertility class). Soil moisture content is estimated at 32.9%. Soil pH is estimated at 6.71 (on a scale of 0-14). Available nitrogen is estimated at 219 kg/ha, and available phosphorus is estimated at 8.8 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Tamil Nadu

Ariyalur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 46.2% (Medium fertility class). Soil moisture content is estimated at 25.3%. Soil pH is estimated at 7.42 (on a scale of 0-14). Available nitrogen is estimated at 198 kg/ha, and available phosphorus is estimated at 17.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Chennai district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 55.6% (High fertility class). Soil moisture content is estimated at 19.0%. Soil pH is estimated at 6.36 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 12.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Coimbatore district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.2% (High fertility class). Soil moisture content is estimated at 15.7%. Soil pH is estimated at 6.49 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 11.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Cuddalore district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 48.8% (Medium fertility class). Soil moisture content is estimated at 20.8%. Soil pH is estimated at 6.68 (on a scale of 0-14). Available nitrogen is estimated at 208 kg/ha, and available phosphorus is estimated at 20.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Dharmapuri district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 48.5% (Medium fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 7.37 (on a scale of 0-14). Available nitrogen is estimated at 231 kg/ha, and available phosphorus is estimated at 17.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Dindigul district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 59.3% (High fertility class). Soil moisture content is estimated at 21.7%. Soil pH is estimated at 7.1 (on a scale of

0-14). Available nitrogen is estimated at 141 kg/ha, and available phosphorus is estimated at 21.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Erode district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 58.5% (High fertility class). Soil moisture content is estimated at 23.4%. Soil pH is estimated at 6.69 (on a scale of 0-14). Available nitrogen is estimated at 141 kg/ha, and available phosphorus is estimated at 15.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kancheepuram district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 56.7% (High fertility class). Soil moisture content is estimated at 19.2%. Soil pH is estimated at 6.64 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 15.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Kanniyakumari district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 53.6% (Medium fertility class). Soil moisture content is estimated at 25.6%. Soil pH is estimated at 6.42 (on a scale of 0-14). Available nitrogen is estimated at 175 kg/ha, and available phosphorus is estimated at 12.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Karur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 57.5% (High fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 7.04 (on a scale of 0-14). Available nitrogen is estimated at 244 kg/ha, and available phosphorus is estimated at 12.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Krishnagiri district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 60.9% (High fertility class). Soil moisture content is estimated at 14.3%. Soil pH is estimated at 6.54 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Madurai district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 59.2% (High fertility class). Soil moisture content is estimated at 22.1%. Soil pH is estimated at 7.17 (on a scale of 0-14). Available nitrogen is estimated at 196 kg/ha, and available phosphorus is estimated at 16.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Nagapattinam district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 54.1% (Medium fertility class). Soil moisture content is estimated at 14.5%. Soil pH is estimated at 6.33 (on a scale of 0-14). Available nitrogen is estimated at 212 kg/ha, and available phosphorus is estimated at 21.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Namakkal district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 45.4% (Medium fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 7.18 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 16.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Perambalur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 59.3% (High fertility class). Soil moisture content is estimated at 22.4%. Soil pH is estimated at 7.32 (on a scale of 0-14). Available nitrogen is estimated at 153 kg/ha, and available phosphorus is estimated at 21.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Pudukkottai district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 45.6% (Medium fertility class). Soil moisture content is estimated at 24.0%. Soil pH is estimated at 6.3 (on a scale of 0-14). Available nitrogen is estimated at 161 kg/ha, and available phosphorus is estimated at 9.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Ramanathapuram district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 55.2% (High fertility class). Soil moisture content is estimated at 18.0%. Soil pH is estimated at 6.42 (on a scale of 0-14). Available nitrogen is estimated at 189 kg/ha, and available phosphorus is estimated at 14.2 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Salem district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 62.7% (High fertility class). Soil moisture content is estimated at 24.7%. Soil pH is estimated at 7.05 (on a scale of 0-14). Available nitrogen is estimated at 153 kg/ha, and available phosphorus is estimated at 12.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Sivaganga district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 66.0% (High fertility class). Soil moisture content is estimated at 15.1%. Soil pH is estimated at 6.73 (on a scale of 0-14). Available nitrogen is estimated at 243 kg/ha, and available phosphorus is estimated at 21.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Thanjavur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 54.5% (Medium fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 222 kg/ha, and available phosphorus is estimated at 9.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

The Nilgiris district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 48.3% (Medium fertility class). Soil moisture content is estimated at 21.4%. Soil pH is estimated at 6.5 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Theni district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.4% (High fertility class). Soil moisture content is estimated at 23.8%. Soil pH is estimated at 6.84 (on a scale of 0-14). Available nitrogen is estimated at 225 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Thiruvallur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 56.4% (High fertility class). Soil moisture content is estimated at 16.4%. Soil pH is estimated at 7.32 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 20.7 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Thiruvarur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.2% (High fertility class). Soil moisture content is estimated at 25.5%. Soil pH is estimated at 6.5 (on a scale of 0-14). Available nitrogen is estimated at 242 kg/ha, and available phosphorus is estimated at 16.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Thoothukkudi district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 54.7% (Medium fertility class). Soil moisture content is estimated at 20.7%. Soil pH is estimated at 6.65 (on a scale of 0-14). Available nitrogen is estimated at 171 kg/ha, and available phosphorus is estimated at 12.1 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tiruchirappalli district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 46.9% (Medium fertility class). Soil moisture content is estimated at 20.0%. Soil pH is estimated at 7.35 (on a scale of 0-14). Available nitrogen is estimated at 209 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tirunelveli district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 54.9% (Medium fertility class). Soil moisture content is estimated at 19.6%. Soil pH is estimated at 6.47 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 11.6 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tiruppur district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 61.5% (High fertility class). Soil moisture content is estimated at 23.1%. Soil pH is estimated at 6.95 (on a scale of 0-14). Available nitrogen is estimated at 234 kg/ha, and available phosphorus is estimated at 12.8 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tiruvannamalai district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 65.3% (High fertility class). Soil moisture content is estimated at 22.8%. Soil pH is estimated at 7.24 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 20.0 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Vellore district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 67.8% (High fertility class). Soil moisture content is estimated at 19.0%. Soil pH is estimated at 7.26 (on a scale of 0-14). Available nitrogen is estimated at 221 kg/ha, and available phosphorus is estimated at 14.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Viluppuram district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 66.3% (High fertility class). Soil moisture content is estimated at 18.7%. Soil pH is estimated at 6.94 (on a scale of 0-14). Available nitrogen is estimated at 151 kg/ha, and available phosphorus is estimated at 11.9 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Virudhunagar district, Tamil Nadu. Dominant soil type: Red & Black Soil. Soil fertility is estimated at 64.3% (High fertility class). Soil moisture content is estimated at 18.8%. Soil pH is estimated at 6.71 (on a scale of 0-14). Available nitrogen is estimated at 167 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Red & Black Soil, not an individual lab measurement.]

Tripura

Dhalai district, Tripura. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 53.5% (Medium fertility class). Soil moisture content is estimated at 30.5%. Soil pH is estimated at 4.9 (on a scale of 0-14). Available nitrogen is estimated at 157 kg/ha, and available phosphorus is estimated at 9.2 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

North Tripura district, Tripura. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 49.1% (Medium fertility class). Soil moisture content is estimated at 24.3%. Soil pH is estimated at 5.36 (on a scale of 0-14). Available nitrogen is estimated at 147 kg/ha, and available phosphorus is estimated at 10.2 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

South Tripura district, Tripura. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 46.5% (Medium fertility class). Soil moisture content is estimated at 24.6%. Soil pH is estimated at 5.94 (on a scale of 0-14). Available nitrogen is estimated at 174 kg/ha, and available phosphorus is estimated at 8.1 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

West Tripura district, Tripura. Dominant soil type: Red & Lateritic Soil. Soil fertility is estimated at 53.4% (Medium fertility class). Soil moisture content is estimated at 30.0%. Soil pH is estimated at 5.01 (on a scale of 0-14). Available nitrogen is estimated at 158 kg/ha, and available phosphorus is estimated at 7.6 kg/ha. [Basis: indicative regional estimate for Red & Lateritic Soil, not an individual lab measurement.]

Uttar Pradesh

Agra district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 21.3%. Soil pH is estimated at 7.37 (on a scale of 0-14). Available nitrogen is estimated at 215 kg/ha, and available phosphorus is estimated at 20.2 kg/ha.

[Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Aligarh district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.7% (High fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 7.64 (on a scale of 0-14). Available nitrogen is estimated at 318 kg/ha, and available phosphorus is estimated at 26.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Allahabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.1% (High fertility class). Soil moisture content is estimated at 29.5%. Soil pH is estimated at 6.54 (on a scale of 0-14). Available nitrogen is estimated at 235 kg/ha, and available phosphorus is estimated at 25.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Ambedkar Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.2% (High fertility class). Soil moisture content is estimated at 25.5%. Soil pH is estimated at 7.0 (on a scale of 0-14). Available nitrogen is estimated at 293 kg/ha, and available phosphorus is estimated at 23.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Auraiya district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.7% (High fertility class). Soil moisture content is estimated at 30.5%. Soil pH is estimated at 6.8 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 22.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Azamgarh district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.3% (High fertility class). Soil moisture content is estimated at 21.5%. Soil pH is estimated at 6.82 (on a scale of 0-14). Available nitrogen is estimated at 199 kg/ha, and available phosphorus is estimated at 25.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Baghpat district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.7% (High fertility class). Soil moisture content is estimated at 25.0%. Soil pH is estimated at 7.79 (on a scale of 0-14). Available nitrogen is estimated at 181 kg/ha, and available phosphorus is estimated at 20.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bahraich district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 67.8% (High fertility class). Soil moisture content is estimated at 29.3%. Soil pH is estimated at 6.96 (on a scale of 0-14). Available nitrogen is estimated at 237 kg/ha, and available phosphorus is estimated at 14.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Ballia district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 29.0%. Soil pH is estimated at 6.68 (on a scale of 0-14). Available nitrogen is estimated at 235 kg/ha, and available phosphorus is estimated at 26.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Balrampur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 55.1% (High fertility class). Soil moisture content is estimated at 22.2%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 226 kg/ha, and available phosphorus is estimated at 16.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Banda district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.5% (High fertility class). Soil moisture content is estimated at 20.5%. Soil pH is estimated at 7.19 (on a scale of 0-14). Available nitrogen is estimated at 278 kg/ha, and available phosphorus is estimated at 21.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bara Banki district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.3% (High fertility class). Soil moisture content is estimated at 29.8%. Soil pH is estimated at 6.73 (on a scale of

0-14). Available nitrogen is estimated at 287 kg/ha, and available phosphorus is estimated at 17.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bareilly district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.6% (High fertility class). Soil moisture content is estimated at 20.9%. Soil pH is estimated at 6.74 (on a scale of 0-14). Available nitrogen is estimated at 249 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Basti district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.6% (High fertility class). Soil moisture content is estimated at 28.1%. Soil pH is estimated at 7.59 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 26.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bijnor district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.9% (High fertility class). Soil moisture content is estimated at 30.2%. Soil pH is estimated at 7.41 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 20.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Budaun district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.3% (High fertility class). Soil moisture content is estimated at 30.9%. Soil pH is estimated at 6.63 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 13.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bulandshahr district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.6% (High fertility class). Soil moisture content is estimated at 24.2%. Soil pH is estimated at 7.38 (on a scale of 0-14). Available nitrogen is estimated at 297 kg/ha, and available phosphorus is estimated at 27.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Chandauli district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.0% (High fertility class). Soil moisture content is estimated at 29.2%. Soil pH is estimated at 7.65 (on a scale of 0-14). Available nitrogen is estimated at 186 kg/ha, and available phosphorus is estimated at 24.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Chitrakoot district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.9% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 7.17 (on a scale of 0-14). Available nitrogen is estimated at 216 kg/ha, and available phosphorus is estimated at 14.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Deoria district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.9% (High fertility class). Soil moisture content is estimated at 19.4%. Soil pH is estimated at 7.33 (on a scale of 0-14). Available nitrogen is estimated at 229 kg/ha, and available phosphorus is estimated at 15.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Etah district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.3% (High fertility class). Soil moisture content is estimated at 27.6%. Soil pH is estimated at 7.2 (on a scale of 0-14). Available nitrogen is estimated at 310 kg/ha, and available phosphorus is estimated at 18.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Etawah district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.9% (High fertility class). Soil moisture content is estimated at 19.1%. Soil pH is estimated at 7.46 (on a scale of 0-14). Available nitrogen is estimated at 318 kg/ha, and available phosphorus is estimated at 25.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Faizabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.5% (High fertility class). Soil moisture content is estimated at 26.8%. Soil pH is estimated at 6.96 (on a scale of 0-14). Available nitrogen is estimated at 275 kg/ha, and available phosphorus is estimated at 24.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Farrukhabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 80.0% (High fertility class). Soil moisture content is estimated at 24.9%. Soil pH is estimated at 6.58 (on a scale of 0-14). Available nitrogen is estimated at 263 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Fatehpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.1% (High fertility class). Soil moisture content is estimated at 22.6%. Soil pH is estimated at 7.45 (on a scale of 0-14). Available nitrogen is estimated at 205 kg/ha, and available phosphorus is estimated at 14.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Firozabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.1% (High fertility class). Soil moisture content is estimated at 30.2%. Soil pH is estimated at 7.4 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 12.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gautam Buddha Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.0% (High fertility class). Soil moisture content is estimated at 22.0%. Soil pH is estimated at 7.28 (on a scale of 0-14). Available nitrogen is estimated at 240 kg/ha, and available phosphorus is estimated at 26.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Ghaziabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.2% (High fertility class). Soil moisture content is estimated at 27.5%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 313 kg/ha, and available phosphorus is estimated at 27.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Ghazipur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 73.5% (High fertility class). Soil moisture content is estimated at 24.5%. Soil pH is estimated at 6.79 (on a scale of 0-14). Available nitrogen is estimated at 293 kg/ha, and available phosphorus is estimated at 23.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gonda district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.5% (High fertility class). Soil moisture content is estimated at 24.3%. Soil pH is estimated at 7.1 (on a scale of 0-14). Available nitrogen is estimated at 295 kg/ha, and available phosphorus is estimated at 22.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Gorakhpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.1% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 6.85 (on a scale of 0-14). Available nitrogen is estimated at 232 kg/ha, and available phosphorus is estimated at 12.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hamirpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.7% (High fertility class). Soil moisture content is estimated at 24.1%. Soil pH is estimated at 7.07 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 15.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hardoi district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.7% (High fertility class). Soil moisture content is estimated at 21.3%. Soil pH is estimated at 6.58 (on a scale of 0-14). Available nitrogen is estimated at 287 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jalaun district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.5% (High fertility class). Soil moisture content is estimated at 24.0%. Soil pH is estimated at 7.56 (on a scale of 0-14). Available nitrogen is estimated at 224 kg/ha, and available phosphorus is estimated at 24.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jaunpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.5% (High fertility class). Soil moisture content is estimated at 27.3%. Soil pH is estimated at 7.06 (on a scale of 0-14). Available nitrogen is estimated at 284 kg/ha, and available phosphorus is estimated at 19.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jhansi district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.4% (High fertility class). Soil moisture content is estimated at 25.4%. Soil pH is estimated at 6.91 (on a scale of 0-14). Available nitrogen is estimated at 194 kg/ha, and available phosphorus is estimated at 18.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jyotiba Phule Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.1% (High fertility class). Soil moisture content is estimated at 23.7%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 310 kg/ha, and available phosphorus is estimated at 12.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kannauj district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.5% (High fertility class). Soil moisture content is estimated at 24.4%. Soil pH is estimated at 6.53 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 13.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kanpur Dehat district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.4% (High fertility class). Soil moisture content is estimated at 21.8%. Soil pH is estimated at 7.02 (on a scale of 0-14). Available nitrogen is estimated at 254 kg/ha, and available phosphorus is estimated at 12.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kanpur Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.0% (High fertility class). Soil moisture content is estimated at 19.3%. Soil pH is estimated at 7.21 (on a scale of 0-14). Available nitrogen is estimated at 226 kg/ha, and available phosphorus is estimated at 20.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kanshiram Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 60.5% (High fertility class). Soil moisture content is estimated at 18.3%. Soil pH is estimated at 7.72 (on a scale of 0-14). Available nitrogen is estimated at 195 kg/ha, and available phosphorus is estimated at 14.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kaushambi district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.7% (High fertility class). Soil moisture content is estimated at 31.7%. Soil pH is estimated at 7.3 (on a scale of 0-14). Available nitrogen is estimated at 265 kg/ha, and available phosphorus is estimated at 27.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kheri district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.2% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 6.77 (on a scale of 0-14). Available nitrogen is estimated at 267 kg/ha, and available phosphorus is estimated at 18.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kushinagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.6% (High fertility class). Soil moisture content is estimated at 31.0%. Soil pH is estimated at 6.69 (on a scale of 0-14). Available nitrogen is estimated at 319 kg/ha, and available phosphorus is estimated at 18.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Lalitpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.2% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 309 kg/ha, and available phosphorus is estimated at 21.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Lucknow district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.6% (High fertility class). Soil moisture content is estimated at 20.1%. Soil pH is estimated at 7.19 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 19.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mahamaya Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.9% (High fertility class). Soil moisture content is estimated at 21.4%. Soil pH is estimated at 7.61 (on a scale of 0-14). Available nitrogen is estimated at 184 kg/ha, and available phosphorus is estimated at 23.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mahoba district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 80.0% (High fertility class). Soil moisture content is estimated at 18.1%. Soil pH is estimated at 6.71 (on a scale of 0-14). Available nitrogen is estimated at 274 kg/ha, and available phosphorus is estimated at 13.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mahrajganj district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.7% (High fertility class). Soil moisture content is estimated at 27.6%. Soil pH is estimated at 7.26 (on a scale of 0-14). Available nitrogen is estimated at 231 kg/ha, and available phosphorus is estimated at 23.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mainpuri district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 69.9% (High fertility class). Soil moisture content is estimated at 20.9%. Soil pH is estimated at 7.23 (on a scale of 0-14). Available nitrogen is estimated at 303 kg/ha, and available phosphorus is estimated at 22.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mathura district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.8% (High fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 6.61 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 12.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mau district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.7% (High fertility class). Soil moisture content is estimated at 21.1%. Soil pH is estimated at 7.56 (on a scale of 0-14). Available nitrogen is estimated at 203 kg/ha, and available phosphorus is estimated at 16.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Meerut district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.5% (High fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 7.54 (on a scale of 0-14). Available nitrogen is estimated at 314 kg/ha, and available phosphorus is estimated at 26.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Mirzapur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.8% (High fertility class). Soil moisture content is estimated at 28.9%. Soil pH is estimated at 6.69 (on a scale of 0-14). Available nitrogen is estimated at 307 kg/ha, and available phosphorus is estimated at 20.3 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Moradabad district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.4% (High fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 6.52 (on a scale of 0-14). Available nitrogen is estimated at 214 kg/ha, and available phosphorus is estimated at 26.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Muzaffarnagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.9% (High fertility class). Soil moisture content is estimated at 31.6%. Soil pH is estimated at 6.63 (on a scale of 0-14). Available nitrogen is estimated at 262 kg/ha, and available phosphorus is estimated at 16.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Pilibhit district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 68.6% (High fertility class). Soil moisture content is estimated at 22.5%. Soil pH is estimated at 6.87 (on a scale of 0-14). Available nitrogen is estimated at 260 kg/ha, and available phosphorus is estimated at 14.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Pratapgarh district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.7% (High fertility class). Soil moisture content is estimated at 27.1%. Soil pH is estimated at 6.67 (on a scale of 0-14). Available nitrogen is estimated at 310 kg/ha, and available phosphorus is estimated at 20.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rae Bareli district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.5% (High fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 7.42 (on a scale of 0-14). Available nitrogen is estimated at 280 kg/ha, and available phosphorus is estimated at 26.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Rampur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.3% (High fertility class). Soil moisture content is estimated at 21.9%. Soil pH is estimated at 6.63 (on a scale of 0-14). Available nitrogen is estimated at 264 kg/ha, and available phosphorus is estimated at 13.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Saharanpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 61.0% (High fertility class). Soil moisture content is estimated at 18.8%. Soil pH is estimated at 7.21 (on a scale of 0-14). Available nitrogen is estimated at 202 kg/ha, and available phosphorus is estimated at 21.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sant Kabir Nagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.8% (High fertility class). Soil moisture content is estimated at 29.5%. Soil pH is estimated at 6.9 (on a scale of 0-14). Available nitrogen is estimated at 256 kg/ha, and available phosphorus is estimated at 21.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sant Ravidas Nagar (Bhadohi) district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 66.8% (High fertility class). Soil moisture content is estimated at 27.2%. Soil pH is estimated at 6.77 (on a scale of 0-14). Available nitrogen is estimated at 317 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Shahjahanpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 75.7% (High fertility class). Soil moisture content is estimated at 26.3%. Soil pH is estimated at 6.82 (on a scale of 0-14). Available nitrogen is estimated at 284 kg/ha, and available phosphorus is estimated at 13.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Shrawasti district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.4% (High fertility class). Soil moisture content is estimated at 26.3%. Soil pH is estimated at 6.64 (on a scale of 0-14). Available nitrogen is estimated at 191 kg/ha, and available phosphorus is estimated at 24.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Siddharthnagar district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.3% (High fertility class). Soil moisture content is estimated at 30.5%. Soil pH is estimated at 7.49 (on a scale of 0-14). Available nitrogen is estimated at 277 kg/ha, and available phosphorus is estimated at 16.5

kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sitapur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.7% (High fertility class). Soil moisture content is estimated at 31.3%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 225 kg/ha, and available phosphorus is estimated at 15.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sonbhadra district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.8% (High fertility class). Soil moisture content is estimated at 28.4%. Soil pH is estimated at 7.12 (on a scale of 0-14). Available nitrogen is estimated at 259 kg/ha, and available phosphorus is estimated at 17.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Sultanpur district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 58.5% (High fertility class). Soil moisture content is estimated at 21.7%. Soil pH is estimated at 7.62 (on a scale of 0-14). Available nitrogen is estimated at 264 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Unnao district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.7% (High fertility class). Soil moisture content is estimated at 18.9%. Soil pH is estimated at 6.83 (on a scale of 0-14). Available nitrogen is estimated at 212 kg/ha, and available phosphorus is estimated at 27.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Varanasi district, Uttar Pradesh. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.7% (High fertility class). Soil moisture content is estimated at 26.6%. Soil pH is estimated at 6.51 (on a scale of 0-14). Available nitrogen is estimated at 205 kg/ha, and available phosphorus is estimated at 22.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Uttarakhand

Almora district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 49.5% (Medium fertility class). Soil moisture content is estimated at 36.8%. Soil pH is estimated at 6.38 (on a scale of 0-14). Available nitrogen is estimated at 173 kg/ha, and available phosphorus is estimated at 13.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Bageshwar district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 64.4% (High fertility class). Soil moisture content is estimated at 29.4%. Soil pH is estimated at 6.16 (on a scale of 0-14). Available nitrogen is estimated at 230 kg/ha, and available phosphorus is estimated at 8.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Chamoli district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 62.2% (High fertility class). Soil moisture content is estimated at 23.6%. Soil pH is estimated at 6.17 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 12.1 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Champawat district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 52.4% (Medium fertility class). Soil moisture content is estimated at 31.9%. Soil pH is estimated at 5.27 (on a scale of 0-14). Available nitrogen is estimated at 145 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Dehradun district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 54.9% (Medium fertility class). Soil moisture content is estimated at 29.1%. Soil pH is estimated at 5.44

(on a scale of 0-14). Available nitrogen is estimated at 225 kg/ha, and available phosphorus is estimated at 9.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Garhwal district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 51.2% (Medium fertility class). Soil moisture content is estimated at 28.6%. Soil pH is estimated at 5.33 (on a scale of 0-14). Available nitrogen is estimated at 207 kg/ha, and available phosphorus is estimated at 10.5 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Hardwar district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 40.4% (Medium fertility class). Soil moisture content is estimated at 37.4%. Soil pH is estimated at 6.47 (on a scale of 0-14). Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 14.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Nainital district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 62.7% (High fertility class). Soil moisture content is estimated at 27.4%. Soil pH is estimated at 6.1 (on a scale of 0-14). Available nitrogen is estimated at 193 kg/ha, and available phosphorus is estimated at 8.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Pithoragarh district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 45.5% (Medium fertility class). Soil moisture content is estimated at 23.8%. Soil pH is estimated at 6.19 (on a scale of 0-14). Available nitrogen is estimated at 213 kg/ha, and available phosphorus is estimated at 13.6 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Rudraprayag district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 47.9% (Medium fertility class). Soil moisture content is estimated at 25.3%. Soil pH is estimated at 5.82 (on a scale of 0-14). Available nitrogen is estimated at 185 kg/ha, and available phosphorus is estimated at 15.2 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Tehri Garhwal district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 61.9% (High fertility class). Soil moisture content is estimated at 36.7%. Soil pH is estimated at 6.53 (on a scale of 0-14). Available nitrogen is estimated at 235 kg/ha, and available phosphorus is estimated at 10.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Udham Singh Nagar district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 41.2% (Medium fertility class). Soil moisture content is estimated at 23.3%. Soil pH is estimated at 5.7 (on a scale of 0-14). Available nitrogen is estimated at 150 kg/ha, and available phosphorus is estimated at 7.7 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

Uttarkashi district, Uttarakhand. Dominant soil type: Mountain & Forest Soil. Soil fertility is estimated at 63.8% (High fertility class). Soil moisture content is estimated at 25.1%. Soil pH is estimated at 6.73 (on a scale of 0-14). Available nitrogen is estimated at 211 kg/ha, and available phosphorus is estimated at 12.4 kg/ha. [Basis: indicative regional estimate for Mountain & Forest Soil, not an individual lab measurement.]

West Bengal

Bankura district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.9% (High fertility class). Soil moisture content is estimated at 22.6%. Soil pH is estimated at 7.34 (on a scale of 0-14). Available nitrogen is estimated at 206 kg/ha, and available phosphorus is estimated at 27.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Bardhaman district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.5% (High fertility class). Soil moisture content is estimated at 21.6%. Soil pH is estimated at 7.42 (on a scale of 0-14). Available nitrogen is estimated at 271 kg/ha, and available phosphorus is estimated at 19.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Birbhum district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 70.8% (High fertility class). Soil moisture content is estimated at 20.2%. Soil pH is estimated at 7.38 (on a scale of 0-14). Available nitrogen is estimated at 217 kg/ha, and available phosphorus is estimated at 16.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Dakshin Dinajpur district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.1% (High fertility class). Soil moisture content is estimated at 22.7%. Soil pH is estimated at 6.93 (on a scale of 0-14). Available nitrogen is estimated at 243 kg/ha, and available phosphorus is estimated at 15.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Darjiling district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.8% (High fertility class). Soil moisture content is estimated at 25.6%. Soil pH is estimated at 6.67 (on a scale of 0-14). Available nitrogen is estimated at 276 kg/ha, and available phosphorus is estimated at 27.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Haora district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 65.7% (High fertility class). Soil moisture content is estimated at 30.6%. Soil pH is estimated at 6.63 (on a scale of 0-14). Available nitrogen is estimated at 186 kg/ha, and available phosphorus is estimated at 23.5 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Hugli district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 74.0% (High fertility class). Soil moisture content is estimated at 29.8%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 288 kg/ha, and available phosphorus is estimated at 18.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Jalpaiguri district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 62.6% (High fertility class). Soil moisture content is estimated at 20.4%. Soil pH is estimated at 7.79 (on a scale of 0-14). Available nitrogen is estimated at 188 kg/ha, and available phosphorus is estimated at 25.2 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Koch Bihar district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 71.0% (High fertility class). Soil moisture content is estimated at 28.3%. Soil pH is estimated at 7.43 (on a scale of 0-14). Available nitrogen is estimated at 300 kg/ha, and available phosphorus is estimated at 14.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Kolkata district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 55.0% (High fertility class). Soil moisture content is estimated at 26.4%. Soil pH is estimated at 6.96 (on a scale of 0-14). Available nitrogen is estimated at 197 kg/ha, and available phosphorus is estimated at 15.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Maldah district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 63.3% (High fertility class). Soil moisture content is estimated at 19.0%. Soil pH is estimated at 6.57 (on a scale of 0-14). Available nitrogen is estimated at 308 kg/ha, and available phosphorus is estimated at 17.0 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Murshidabad district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.6% (High fertility class). Soil moisture content is estimated at 31.5%. Soil pH is estimated at 6.69 (on a scale of 0-14). Available nitrogen is estimated at 280 kg/ha, and available phosphorus is estimated at 21.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Nadia district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 72.3% (High fertility class). Soil moisture content is estimated at 19.3%. Soil pH is estimated at 6.75 (on a scale of 0-14). Available nitrogen is estimated at 190 kg/ha, and available phosphorus is estimated at 21.7 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

North Twenty Four Parganas district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 79.7% (High fertility class). Soil moisture content is estimated at 27.4%. Soil pH is estimated at 7.42 (on a scale of 0-14). Available nitrogen is estimated at 317 kg/ha, and available phosphorus is estimated at 22.8 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Paschim Medinipur district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 56.0% (High fertility class). Soil moisture content is estimated at 30.1%. Soil pH is estimated at 6.97 (on a scale of 0-14). Available nitrogen is estimated at 180 kg/ha, and available phosphorus is estimated at 20.1 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Purba Medinipur district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 78.4% (High fertility class). Soil moisture content is estimated at 21.6%. Soil pH is estimated at 7.14 (on a scale of 0-14). Available nitrogen is estimated at 204 kg/ha, and available phosphorus is estimated at 12.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Puruliya district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 57.0% (High fertility class). Soil moisture content is estimated at 28.2%. Soil pH is estimated at 7.62 (on a scale of 0-14). Available nitrogen is estimated at 200 kg/ha, and available phosphorus is estimated at 16.6 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

South Twenty Four Parganas district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 59.9% (High fertility class). Soil moisture content is estimated at 18.4%. Soil pH is estimated at 7.22 (on a scale of 0-14). Available nitrogen is estimated at 277 kg/ha, and available phosphorus is estimated at 22.4 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]

Uttar Dinajpur district, West Bengal. Dominant soil type: Alluvial Soil. Soil fertility is estimated at 64.7% (High fertility class). Soil moisture content is estimated at 25.8%. Soil pH is estimated at 6.62 (on a scale of 0-14). Available nitrogen is estimated at 201 kg/ha, and available phosphorus is estimated at 26.9 kg/ha. [Basis: indicative regional estimate for Alluvial Soil, not an individual lab measurement.]