

Fig. 3. Fungal species identified from comparisons of PCR-derived sequences of the ITS1-5.8S-ITS2 region of ribosomal deoxyribonucleic acid. Species recovered from air, soil, and inflorescences from an indoor cannabis organic facility and from inflorescences from an outdoor facility are indicated. Similar colored symbols indicate that particular species was present at each sampling site.

Fungal Species	Sample Source			
	Indoor Air	Indoor Soil	Indoor Flower	Outdoor Flower
<i>Acremonium alternatum</i>	▲	▲		
<i>Alternaria alternata</i>	▲		▲	▲
<i>Alternaria tenuissima</i>	▲		▲	▲
<i>Aspergillus niger</i>	▲	▲	▲	▲
<i>Aspergillus ochraceus</i>	▲	▲	▲	▲
<i>Beauveria bassiana</i>	▲	▲	▲	
<i>Bjerkandera adusta</i>	▲			
<i>Botrytis cinerea</i>	▲		▲	▲
<i>Cercospora canescens</i>	▲			
<i>Chaetomium elatum</i>			▲	▲
<i>Chaetomium brasiliensis</i>	▲		▲	
<i>Chaetomium globosum</i>	▲		▲	▲
<i>Conidiobolus coronatus</i>	▲	▲	▲	
<i>Cladosporium cladosporioides</i>	▲		▲	▲
<i>Cladosporium floccosum</i>	▲			
<i>Diaporthe eres</i>	▲		▲	▲
<i>Epicoccum nigrum</i>	▲		▲	▲
<i>Fusarium avenaceum</i>	▲		▲	
<i>Fusarium oxysporum</i>	▲	▲	▲	▲
<i>Fusarium proliferatum</i>	▲	▲	▲	
<i>Fusarium sporotrichioides</i>	▲		▲	▲
<i>Humicola brevis</i>	▲			
<i>Lasiodiplodia theobromae</i>	▲		▲	
<i>Hydnopolyporus fimbriatus</i>	▲			
<i>Lecanacillium aphanocladii</i>	▲			
<i>Mortierella hyalina</i>	▲	▲		
<i>Mucor circinelloides</i>	▲	▲	▲	▲
<i>Mucor racemosus</i>	▲	▲	▲	▲
<i>Nigrospora oryzae</i>	▲	▲	▲	
<i>Paraphaeosphaeria michotii</i>	▲			
<i>Penicillium brevicompactum</i>	▲	▲	▲	
<i>Penicillium citrinum</i>	▲		▲	▲
<i>Penicillium expansum</i>	▲		▲	▲
<i>Penicillium olsonii</i>	▲	▲	▲	
<i>Penicillium polonicum</i>	▲		▲	
<i>Penicillium spathulatum</i>	▲		▲	
<i>Rhizopus stolonifer</i>	▲	▲	▲	▲
<i>Scedosporium aurantiacum</i>	▲			
<i>Stemphylium versicarium</i>	▲			
<i>Trichoderma harzianum</i>	▲	▲	▲	

images from the present study showed that spores stuck to trichome glands resembled those of *Botrytis*, *Aspergillus*, *Cladosporium*, as well as other fungi that may be present in inflorescences while adhered to the sticky surfaces of stigmas and trichome glands. The swab method of sampling likely recovered a portion of these fungi that were present as viable epiphytes. There is no previously published information to demonstrate if the resinous content of cannabis trichome glands is inhibitory to fungal spore germination but the re-

covery of viable colonies suggests it may not be. The effects of resinous trichome secretions on fungal spore germination is worthy of further study.

Sampling of inflorescences revealed that multiple species of fungi were present at any given time, many of which are previously unreported from cannabis inflorescences. In particular, *Conidiobolus coronatus*, *Nigrospora oryzae*, *Epicoccum nigrum*, *Mucor circinelloides*, and *Mucor racemosus* were recovered in this study. None of these species were previously observed