




Figure 22. Comparative efficacy of reduced-risk products at managing powdery mildew development on cannabis genotype ‘MP’. (a–d) Disease was rated according to the scale shown, from 0 (a) to 3 (d). (e) Products were applied as preventative treatments at days 0, 7, and 14 of the flowering period. (f) Products were applied as a curative treatment once at day 42 of the flowering period after the onset of disease development. The trials were conducted during the spring growing season.

A summary of the IDM approaches that can be used against four important pathogens of cannabis is provided in Table 1.

Table 1. Summary of IDM strategies for four important pathogens affecting cannabis plants.

	HLVd Stunting Disease	<i>Fusarium/Pythium</i> Root and Crown Rot	<i>Botrytis cinerea</i> Bud Rot	Powdery Mildew
Prevention	Test propagative materials and stock plants; utilize pathogen-free planting materials.	Test propagative materials and stock plants; utilize pathogen-free planting materials.	Reduce canopy humidity by adjusting planting density and enhancing air circulation.	Maintain an even climate above 21 °C and vaporize sulfur nightly.
Sanitation	Clean equipment and bench surfaces; destroy diseased plants.	Clean equipment and bench surfaces; actively remove dead or diseased tissues.	Fog growing environment with reduced-risk products prior to planting.	Fog growing environment with reduced-risk products prior to planting.
Protection	Isolate propagative materials and stock plants in controlled access areas.	Apply <i>Trichoderma harzianum</i> and <i>Gliocladium cantenulatum</i> as a drench to rooted cuttings and plants.	Apply Rootshield HC® on developing inflorescences from day 14 to day 28 of flowering.	Preventatively spray reduced-risk products, such as Suffoil-X and Regalia Maxx, on susceptible genotypes.
Monitoring	Scout regularly for symptoms; routinely sample water and suspect plants.	Scout regularly for symptoms; routinely sample water and suspect plants.	Conduct daily scouting for bud rot from the sixth week of flowering onwards.	Conduct weekly scouting at all plant development stages.
Eradication	Immediately remove and safely dispose of diseased plants at all stages of growth.	Immediately remove and safely dispose of diseased plants at all stages of growth.	Remove and dispose of infected inflorescences; perform post-drying bud rot severity checks.	Remove infected leaves and perform targeted sprays with reduced-risk products.
Genotype Selection	Avoid highly susceptible genotypes; evaluate tolerant genotypes.	Avoid highly susceptible genotypes; evaluate tolerant genotypes.	Avoid planting highly susceptible genotypes during <i>B. cinerea</i> -prone periods; evaluate tolerant genotypes.	Avoid highly susceptible genotypes; evaluate tolerant genotypes.