

sessile-capitate and were present at a high density (Fig. 4a, b). There was a mixture of sessile-capitate trichomes with bulbous trichomes and nonglandular hairs on other bract tissues (Fig. 4b). A range of stalk lengths could be seen on capitate trichomes (Fig. 4f). At 6 weeks,

stalked-capitate trichomes that formed along the edge of a bract had very long stalks, sometimes reaching up to 1,000 μm (Fig. 4g, h). The heads of capitate-stalked trichomes were fully formed at this time and surrounded by a cuticle (Fig. 4i). In Fig. 4c-e, presecretory (sessile)




Fig. 4 Capitate trichome development on cannabis strain SQ, lower bract surface, at 3 and 6 weeks of inflorescence development. **a** At 3 weeks, most capitate trichomes at the tip of the bract are sessile. **b** Close-up view showing a mixture of sessile-capitate trichomes with bulbous trichomes and nonglandular hairs. There are no stalked-capitate trichomes. **c–e** Presecretory (sessile) trichomes showing the outlines of presecretory disc cells. **f** A range of stalk lengths have developed on capitate trichomes, and sessile trichomes can be seen in the background. **g** At 6 weeks, stalked-capitate trichomes are densely packed together on the underside of the bracts and have formed long stalks. **h** Some stalked-capitate trichomes that form at the edge of a bract have extremely long stalks, up to 1000 μm (arrow). **i** Magnified view of capitate trichomes showing fully formed heads surrounded by a cuticle