

(Fig. 2d, e). The sessile-capitate trichomes, together with early development stalked-capitate trichomes, could be observed adjacent to one another (Fig. 2f, g). On

6-week-old bract tissues, fully developed stalked-capitate trichomes with enlarged heads surrounded by a cuticle were present (Fig. 2h, i).




Fig. 2 The different trichome types on *Cannabis* bract tissues observed in this study. All images are obtained from strain MD, lower bract surface, at 3 and 6 weeks of inflorescence development. **a** A stalked capitate trichome in the foreground (arrow) compared with small bulbous trichomes in the background. **b, c** Bulbous trichomes formed on short pedicels. **d** Nonglandular trichomes (ha) formed next to a bulbous (bu) and a sessile-capitate trichome (sc). In the background, abundant stomata (st) can be seen on raised epidermal cells. **e** Nonglandular hairs (ha) interspersed with sessile-capitate trichomes on a young bract. **f** Magnified view of a sessile-capitate trichome with fully formed head surrounded by a cuticle. **g** Presecretory sessile-capitate (arrow) and mature stalked-capitate trichomes with fully formed heads adjacent to one another. **h** Mature stalked-capitate trichomes formed in large numbers. Note the presence of small bulbous trichomes (arrow) and nonglandular trichomes in the background. **i** Magnified view of the head of a stalked-capitate trichome with fully formed cuticle surrounding the head