



Fig. 6 (Colour online) Disease symptoms 3 weeks after inoculation of rooted cuttings of two cannabis strains with isolates selected to represent four *Pythium* species. In the left panel (a, c, e, g) is strain 'Hash Plant'; in the right panel (b, d, f, h) is strain 'White Rhino'. The four species inoculated were *P. myriotylum* (a, b), *P. ultimum* (c, d), *P. dissotocum* (e, f) and *P. catenulatum* (g, h). In each photo, noninoculated plants are shown on the left. The isolates used are shown in Table 1.

and subjected to intermittent periods of daily flooding for two weeks (Fig. 1c) before being transferred to the flowering room. Under these conditions, water-saturated and damaged roots emerging from the bottom of the coco block (Fig. 1d) are likely to be predisposed to infection by *Pythium* spp. In saturated soils and temperatures of 26–30°C, *P. myriotylum* destroyed ginger rhizomes within 7 days (Stirling et al. 2009).

The symptoms of infection caused by *Pythium* spp. observed on cannabis plants in commercial facilities are consistent with those on other hydroponically grown greenhouse crops infected by *Pythium* spp., which may include an initial darkening (greening) of leaves on affected cucumber and pepper plants (Punja and Yip 2003; Sutton et al. 2006) followed by wilting and plant collapse. Total plant collapse due to *Pythium* infection was not uncommon during the hot summer