

1 Frame 12 – Functions of Complex Variables

1.1 Functions

If S is a set of complex numbers, then a **function** f is a rule that assigns a complex number w to each z in S . The number w is called the **value** of f at z . We denote it as

$$w = f(z)$$

The set S is called the **domain of definition** of f . Note that we need both a rule (f) and a domain (S) for a function to be well defined.

Suppose that $w = u + iv$ and $z = x + iy$. Then,

$$u + iv = f(x + iy)$$

Then, we can express $f(z)$ as a pair of real functions of x and y :

$$f(z) = u(x, y) + iv(x, y)$$

Alternatively, we could use polar coordinates to write

$$u + iv = f(re^{i\theta})$$

so

$$f(z) = u(r, \theta) + iv(r, \theta)$$

Example: the function $f(z) = z^2$ can be written as

$$\begin{aligned} f(x + iy) &= (x + iy)^2 \\ &= (x^2 - y^2) + i2xy \end{aligned}$$

so

$$\begin{aligned} u(x, y) &= x^2 - y^2 \\ v(x, y) &= 2xy \end{aligned}$$

In polar coordinates,

$$\begin{aligned} f(x + iy) &= (re^{i\theta})^2 \\ &= r^2 e^{i2\theta} \\ &= r^2 \cos 2\theta + ir^2 \sin 2\theta \end{aligned}$$

so

$$\begin{aligned} u(r, \theta) &= r^2 \cos 2\theta \\ v(r, \theta) &= r^2 \sin 2\theta \end{aligned}$$

1.2 Real-Valued Functions

We say that f is a **real-valued function** if v is zero everywhere.

Example: one real-valued function is

$$f(z) = |z|^2 = x^2 + y^2 + i0$$

1.3 Polynomials

If n is a non-negative integer and $a_0, a_1, a_2, \dots, a_n$ are complex numbers with $a_n \neq 0$, then the function

$$P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$$

is a **polynomial** of degree n . Note that this sum has a finite number of terms and that the domain of definition is the entire z plane.

As in real numbers, a **rational function** is a quotient of two polynomials:

$$R(z) = \frac{P(z)}{Q(z)}$$

A rational function is defined everywhere that $Q(z) \neq 0$.

1.4 Multi-Valued Functions

A generalization of a function is a rule that assigns more than one value to a point z . These **multiple-valued functions** are usually studied by taking one of the possible values at each point and constructing a single-valued function.

Example: we know that we can write

$$z^{1/2} = \pm \sqrt{r}e^{i\theta/2}$$

where we denoted $-\pi < \theta \leq \pi$ as the **principal value** of $\arg z$. To turn this into a single valued function, we can choose the positive value of r and write

$$f(z) = \sqrt{r}e^{i\theta/2}$$

Then, f is well-defined on the entire plane.