

1 Frame 37 – Derivatives with Real Variables

1.1 Definition

In the previous chapter, we looked at derivatives of complex functions of a complex variable z . Now, we look at the derivatives of a complex-valued function of a real variable t . If we write our function as

$$w(t) = u(t) + iv(t)$$

where u and v are real-valued, then we can define the derivative of w at a point t as

$$w'(t) = \frac{d}{dt}w(t) = u'(t) + iv'(t)$$

provided that u' and v' exist at t .

1.2 Properties

If $z_0 = x_0 + iy_0$ is a complex constant, then we can show that

$$\begin{aligned}\frac{d}{dt}[z_0 w(t)] &= [(x_0 + iy_0)(u(t) + iv(t))]' \\ &= [x_0 u(t) - y_0 v(t)]' + i[y_0 u(t) + x_0 v(t)]' \\ &= [x_0 u'(t) - y_0 v'(t)] + i[y_0 u'(t) + x_0 v'(t)] \\ &= z_0 w'(t)\end{aligned}$$

as we expect.

Next, if z_0 is still a complex constant, the derivative of $e^{z_0 t}$ is

$$\begin{aligned}\frac{d}{dt}e^{z_0 t} &= \frac{d}{dt}e^{x_0 t}(\cos y_0 t + i \sin y_0 t) \\ &= \frac{d}{dt}e^{x_0 t} \cos y_0 t + i \frac{d}{dt}e^{x_0 t} \sin y_0 t \\ &= (x_0 + iy_0)(e^{x_0 t} \cos y_0 t + i e^{x_0 t} \sin y_0 t) \\ &= z_0 e^{z_0 t}\end{aligned}$$

Many other rules carry over from standard calculus. However, some rules no longer apply. For instance, in calculus, the mean value theorem for derivatives states that

$$w'(c) = \frac{w(b) - w(a)}{b - a}$$

for some c in the interval $a \leq c \leq b$ as long as w is continuous. However, this is easily disproved by the function

$$w(t) = e^{it}$$

If $a = 0$ and $b = 2\pi$, then $w(a) = w(b) = 1$ and we expect to find a point c in $[0, 2\pi]$ such that $w'(c) = 0$. However, no such points exist – the magnitude of the derivative is always 1.

2 Frame 38 – Definite Integrals of Complex Functions

2.1 Definitions

If $w(t)$ is a complex-valued function of a real variable t , as in the previous section

$$w(t) = u(t) + iv(t)$$

then we define the **definite integral** of $w(t)$ over the interval $a \leq t \leq b$ as

$$\int_a^b w(t)dt = \int_a^b u(t)dt + i \int_a^b v(t)dt$$

provided the two right-side integrals exist. Then,

$$\begin{aligned}\Re \left[\int_a^b w(t)dt \right] &= \int_a^b \Re[w(t)]dt \\ \Im \left[\int_a^b w(t)dt \right] &= \int_a^b \Im[w(t)]dt\end{aligned}$$

Improper integrals over unbounded intervals are defined similarly.

The two real integrals will exist as long as u and v are **piecewise continuous** on the interval $[a, b]$ – that is, continuous everywhere in the interval except possibly for a finite number of points where it has one-sided limits. When u and v are piecewise continuous, we say that w is also piecewise continuous.

2.2 Properties

The most common rules of integrals from calculus apply here as well:

- $\int z_0 w(t)dt = z_0 \int w(t)$
- $\int w_1(t) + w_2(t)dt = \int w_1(t)dt + \int w_2(t)dt$
- $\int_a^b w(t)dt = - \int_b^a w(t)dt$
- $\int_a^b w(t)dt = \int_a^c w(t)dt + \int_c^b w(t)dt$

We can also extend the fundamental theorem of calculus to complex integrals. Suppose that two functions

$$\begin{aligned}w(t) &= u(t) + iv(t) \\ W(t) &= U(t) + iV(t)\end{aligned}$$

are continuous on the interval $[a, b]$ and $W'(t) = w(t)$ when $a \leq t \leq b$. Then, we can write

$$\int_a^b w(t)dt = W(b) - W(a) = W(t)\Big|_a^b$$

Example: noting that the derivative of $\frac{1}{i}e^{it}$ is

$$\frac{d}{dt} \left(\frac{1}{i}e^{it} \right) = \frac{1}{i}ie^{it} = e^{it}$$

we can evaluate $\int e^{it}dt$ as

$$\begin{aligned} \int_0^{\pi/4} e^{it}dt &= \frac{e^{it}}{i} \Big|_0^{\pi/4} \\ &= \frac{1}{i} \left[e^{\pi/4} - 1 \right] \\ &= \frac{1}{i} \left[\frac{1}{\sqrt{2}} - 1 + \frac{i}{\sqrt{2}} \right] \\ &= \frac{1}{\sqrt{2}} + i \left(1 - \frac{1}{\sqrt{2}} \right) \end{aligned}$$

As in the previous section, the mean value theorem for integrals does not apply. We can show this by finding the integral $\int_0^{2\pi} e^{it}dt = 0$, even though the function is never zero on this interval.

3 Frame 39 – Contours

3.1 Definitions

In calculus, integrals are defined on intervals of the real line. In complex analysis, we instead use curves in the complex plane.

An **arc** is a set of points $z = (x, y)$ in the complex plane such that the functions

$$x = x(t), \quad y = y(t); \quad z = z(t) = x(t) + iy(t)$$

are continuous functions of the parameter t , where $a \leq t \leq b$. This definition is a continuous mapping of the interval $a \leq t \leq b$ into the z plane.

We say that an arc is **simple** if it does not cross itself; ie:

$$z(t_1) \neq z(t_2) \quad \text{for all } t_1 \neq t_2$$

If a simple arc starts and ends at the same point ($z(a) = z(b)$), it is called a **simple closed curve**. These curves are **positively oriented** when they are oriented in the counterclockwise direction.

Example: the unit circle

$$z = e^{i\theta}$$

where $0 \leq \theta \leq 2\pi$ is a positively oriented simple closed curve centered at the origin with a radius of 1. A more general circle is

$$z = z_0 + Re^{i\theta}$$

which is centered at z_0 and has a radius of R .

3.2 Uniqueness

Note that the parametric representation for any arc is not unique. If we know a function ϕ such that

$$t = \phi(\tau)$$

maps the interval $\alpha \leq \tau \leq \beta$ onto the interval $a \leq t \leq b$. Then, the two equations

$$z(t) \quad (a \leq t \leq b)$$

and

$$z(\phi(t)) \quad (\alpha \leq t \leq \beta)$$

represent the same arc.

3.3 Smoothness

Suppose that the real and imaginary components of z are differentiable, and their derivatives are continuous. Then, the arc $z(t)$ is a **differentiable arc**, and

$$|z'(t)| = |x'(t) + iy'(t)| = \sqrt{[x'(t)]^2 + [y'(t)]^2}$$

is integrable. This allows us to find the length of an arc as

$$L = \int_a^b |z'(t)| dt$$

If an arc is differentiable and $z'(t)$ is never zero (except maybe at $t = a$ or $t = b$), then we call the arc a **smooth arc**. We can write the unit tangent vector

$$\mathbf{T} = \frac{z'(t)}{|z'(t)|}$$

which has an angle of inclination of $\arg z'(t)$.

A **contour** is an arc which consists of a finite number of smooth arcs joined together. Specifically, if $z(t)$ represents a contour, then $z(t)$ is continuous and $z'(t)$ is piecewise continuous. If a contour is also a simple closed arc, we call it a **simple closed contour**.

The points on a simple closed arc are the boundary points of two different domains:

- The interior of the arc, which is bounded;
- The exterior of the arc, which is unbounded.

4 Frame 40 – Contour Integrals

4.1 Definitions and conditions

We can now integrate a complex function f along a contour C , which starts and ends at points z_1 and z_2 , respectively. This is effectively a line integral. These integrals can be written as

$$\int_C f(z)dz$$

or, if the integral does not depend on the path taken,

$$\int_{z_1}^{z_2} f(z)dz$$

This integral (along a complex path) represents an integral with respect to a real parameter t . If the contour C is written as $z(t)$ on the interval $a \leq t \leq b$, then the integral represented is

$$\int_C f(z)dz = \int_a^b f[z(t)]z'(t)dt$$

Since $z'(t)$ must be piecewise continuous, this integral exists as long as $f[z(t)]$ is also piecewise continuous on this interval.

4.2 Basic properties

From the definition and the properties of integrals, we can write

$$\int_C z_0 f(z)dz = z_0 \int_C f(z)dz$$

and

$$\int_C [f(z) + g(z)]dz = \int_C f(z)dz + \int_C g(z)dz$$

We can also create a new contour $-C$ that consists of the points in C in reversed order – this contour extends from z_2 to z_1 . Integrating along this reversed contour, we find that

$$\begin{aligned} \int_{-C} f(z)dz &= \int_{-b}^{-a} f[z(-t)] \frac{d}{dt} z(-t)dt \\ &= - \int_{-b}^{-a} f[z(-t)]z'(-t)dt \\ &= - \int_a^b f[z(t)]z'(t)dt \\ &= - \int_C f(z)dz \end{aligned}$$

We can also split up a contour C into multiple legs C_1, C_2, \dots . If we can write a contour this way, then we say that $C = C_1 + C_2$. The contour integral along C can then be written as

$$\int_C f(z)dz = \int_{C_1} f(z)dz + \int_{C_2} f(z)dz$$

5 Frame 41 – Examples of Contour Integrals

This section will show several specific examples of contour integrals.

5.1 Example 1

Suppose that the contour C is the right hand half of the circle $|z| = 2$:

$$z = 2e^{i\theta}, \quad \left(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}\right)$$

Then,

$$\begin{aligned} \int_C \bar{z} dz &= \int_{-\pi/2}^{\pi/2} \overline{2e^{i\theta}} (2e^{i\theta})' d\theta \\ &= 4i \int_{-\pi/2}^{\pi/2} e^{-i\theta} e^{i\theta} d\theta \\ &= 4i \int_{-\pi/2}^{\pi/2} d\theta \\ &= 4\pi i \end{aligned}$$

Also, note that all of the points on this semicircle satisfy

$$z\bar{z} = |z|^2 = 4$$

so we can see from this result that

$$\int_C \frac{1}{z} dz = \pi i$$

5.2 Example 2

Suppose that the points O , A , and B are 0 , i , and $1 + i$, respectively. Then, if C_1 is the polyline OAB and

$$f(z) = y - x - i3x^2$$

then the contour integral of f along C_1 is

$$\begin{aligned}
\int_{C_1} f(z)dz &= \int_{OA} f(z)dz + \int_{AB} f(z)dz \\
&= \int_0^1 yidy + \int_0^1 (1-x-i3x^2)dx \\
&= \frac{i}{2} + \int_0^1 (1-x)dx - 3i \int_0^1 x^2 dx \\
&= \frac{i}{2} + \frac{1}{2} - i \\
&= \frac{1-i}{2}
\end{aligned}$$

Next, if C_2 is the line OB , the contour integral along this curve is

$$\begin{aligned}
\int_{C_2} f(z)dz &= \int_0^1 -i3x^2(1+i)dx \\
&= 3(1-i) \int_0^1 x^2 \\
&= 1-i
\end{aligned}$$

Finally, the integral of f over the simple closed contour $OABO$ is $C_1 - C_2$, which is

$$\int_{OABO} f(z)dz = \frac{-1+i}{2}$$

5.3 Example 3

Suppose that C is any arbitrary smooth arc from a fixed point z_1 to another point z_2 :

$$z = z(t) \quad (a \leq t \leq b)$$

The contour integral of $f(z) = z$ along this curve is

$$\begin{aligned}
\int_C z dz &= \int_a^b z(t)z'(t)dt \\
&= \int_a^b \frac{d}{dt} \frac{[z(t)]^2}{2} dt \\
&= \left. \frac{[z(t)]^2}{2} \right|_a^b \\
&= \frac{z_2^2 - z_1^2}{2}
\end{aligned}$$

Note that this integral only depends on the endpoints of C and not the path. This lets us write

$$\int_{z_1}^{z_2} z dz = \frac{z_2^2 - z_1^2}{2}$$

This holds when C is not a smooth contour. Since all contours are sums of finite numbers of smooth arcs, this expression holds for each arc in C , leading to the same final expression.

Also, note that the integral of $f(z) = z$ around any closed contour in the plane is zero.

6 Frame 42 – Examples with Branch Cuts

A contour integral's path can include a point on a branch cut. The following two examples show this.

6.1 Example 1

Suppose we want to integrate the function

$$f(z) = z^{1/2} = e^{\frac{1}{2} \log z} \quad (0 < \arg z < 2\pi)$$

on the semicircle

$$z = 3e^{i\theta} \quad (0 \leq \theta \leq \pi)$$

Although the function is not defined at $\theta = 0$, we can still write

$$f[z(\theta)] = e^{\frac{1}{2}(\ln 3 + i\theta)} = \sqrt{3}e^{i\theta/2}$$

and the right hand limit of this function exists at $\theta = 0$. Thus, the integrand exists as long as we define the missing point as

$$f[z(0)]z'(0) = i3\sqrt{3}$$

Then,

$$\begin{aligned} \int_C f(z)dz &= 3\sqrt{3} \int_0^\pi e^{i3\theta/2} \\ &= 3\sqrt{3} \frac{2}{3i} e^{i3\theta/2} \Big|_0^\pi \\ &= -\frac{2}{3i}(1+i) \\ &= -2\sqrt{3}(1+i) \end{aligned}$$

6.2 Example 2

Suppose that we want to integrate the function

$$f(z) = z^{a-1} = e^{(a-1) \operatorname{Log} z} \quad (-\pi < \operatorname{Arg} z < \pi)$$

on the positively oriented circle

$$z = Re^{i\theta} \quad (-\pi \leq \theta \leq \pi)$$

The contour integral is

$$\begin{aligned}
 \int_C z^{a-1} dz &= \int_{-\pi}^{\pi} iR^a e^{ia\theta} d\theta \\
 &= iR^a \int_{-\pi}^{\pi} e^{ia\theta} d\theta \\
 &= iR^a \left(\frac{e^{ia\theta}}{ia} \right)_{-\pi}^{\pi} \\
 &= i \frac{2R^a}{a} \frac{e^{ia\pi} - e^{-ia\pi}}{2i} \\
 &= i \frac{2R^a}{a} \sin a\pi
 \end{aligned}$$

Note that if a is a non-zero integer, this integral is zero; if $a = 0$, this integral reduces to

$$\int_C \frac{dz}{z} = 2\pi i$$

7 Frame 43 – Upper Bounds for Contour Integrals

We can put a bound on the modulus of a contour integral by observing that

$$\left| \int_a^b w(t) dt \right| \leq \int_a^b |w(t)| dt$$

7.1 Theorem

Suppose that C is a contour with a length of L and that $f(z)$ is a function that is piecewise continuous on C . If $M \geq 0$ is a real constant such that

$$|f(z)| \leq M$$

for all points on C , then

$$\left| \int_C f(z) dz \right| \leq ML$$

Note that such a number M will always exist because f is continuous on C .

7.2 Examples

Example: Suppose that

$$f(z) = \frac{z+4}{z^3-1}$$

and the contour C is a quarter circle with a radius of 2 in the first quadrant (running from $z = 2$ to $z = 2i$). Since $|z| = 2$ at all points on this contour, we can write that

$$|z+4| \leq |z| + 4 = 6$$

and

$$|z^3-1| \geq |z|^3 - 1 = 7$$

Since the length of the contour is $L = \pi$, we can write the upper bound

$$\left| \int_C \frac{z+4}{z^3-1} dz \right| \leq \frac{6\pi}{7}$$

Example: suppose that C_R is the semicircular contour

$$z = Re^{i\theta} \quad (0 \leq \theta \leq \pi)$$

and f is the function

$$f(z) = \frac{z^{1/2}}{z^2+1}$$

where $z^{1/2}$ denotes the branch $-\pi/2 < \theta < 3\pi/2$. Anywhere on this semicircle,

$$|z^{1/2}| = \sqrt{R}$$

and

$$|z^2 + 1| \geq ||z^2| - 1| = R^2 - 1$$

Since the contour has a length of πR , the contour integral of f along C can be limited by

$$\begin{aligned} \int_C f(z) dz &\leq \frac{\sqrt{R}}{R^2 - 1} \cdot \pi R \\ &= \frac{\pi R^{3/2}}{R^2 - 1} \\ &= \frac{\pi/\sqrt{R}}{1 - (1/R^2)} \end{aligned}$$

As R approaches infinity, this bound approaches zero, so

$$\lim_{R \rightarrow \infty} \int_C f(z) dz = 0$$