

Problem Sequence - Solutions

This document will be filled up with the solutions from the problem sequence.

Solution 1. (Greg)

From Axiom 1, there are three cases to consider:

1. $p = 0$

p is equal to the only element of M . From the definition of limit point, every segment containing p must contain a point of M different from p . However, there are no points in M different from p , so p must not be a limit point of M .

2. $p > 0$

From Axiom 3, there exists a point a such that $0 < a < p$. There also exists a point b such that $b > p$. Since $p > 0$ and $b > p$, Axiom 2 tells us that $b > 0$. We can then form the segment $S = (a, b)$. Since $a > 0$ and $b > 0$, 0 is not between a and b , so S does not contain 0. However, $a < p < b$, so S contains p . S is a segment containing p that does not contain any element of M , so p is not a limit point of M .

3. $p < 0$

(symmetric to the $p > 0$ case)

Therefore, regardless of our choice of p , we can construct a segment that contradicts the requirements in the limit point definition, so p is not a limit point of M .

Solution 2. (Jeff)

According to Definition 4, we can prove that p is not a limit point of M if we can construct a segment containing p but not a different point of M . Construct this segment as follows:

1. If all points of M are on the opposite side of p , choose any value as an endpoint.
2. If any point of M is on the same side of p , choose a point between the nearest point of M and p . (Axiom 3 confirms that there will be such a point.)

This segment contains p but will not contain any points of M (with the exception of p , if p is 0 or 1). Therefore, we have found a segment that does not fulfill the requirements of Definition 4, so p is not a limit point of M .

Solution 3. (Erin, solved after Problem 4)

Let $a = 0$ and $b = 1$. Then, according to Problem 4, b is a limit point of (a, b) . Therefore, 1 is a limit point of $(0, 1)$.

Solution 4. (Erin)

We will prove that b is a limit point of (a, b) ; The proof for a is similar.

Construct a segment (p, q) containing b ($p < b < q$). According to Axiom 1, there are three cases, and we will deal with two of them simultaneously:

1. $p < a$

In this case, $p < a < b < q$, so $p < (a, b) < q$. Since (p, q) contains every point of (a, b) , we have found points that satisfy Definition 4.

2. $p \geq a$

In this case, $a \leq p < b < q$. According to Axiom 3, there is a point d between p and b . The inequality is then $a \leq p < d < b < q$. Then,

- $a < d < b$, so d is in (a, b) .
- $p < d < q$, so d is in (p, q) .

This means that (p, q) contains d , which is a point of (a, b) .

Therefore, every (p, q) containing b also contains a point of (a, b) , so b is a limit point of (a, b) .

Solution 5. (Greg)

Choose any point p from S . Construct a segment (x, y) that contains p (ie: $x < p < y$). Put no other condition on y . According to Axiom 1, one of these three cases is true:

1. $x > a$

Choose a point q between x and p ($x < q < p$; this exists by Axiom 3). Apply Axiom 2 three times:

- $a < x$ and $x < q$, so $a < q$
- $q < p$ and $p < b$, so $q < b$
- $q < p$ and $p < y$, so $q < y$

so $a < q < b$ and $x < q < y$. Therefore, q is an element of S inside (a, b) that is different from p .

2. $x = a$

Repeat the proof for $x > a$ with one change:

- $a = x$ and $x < q$, so $a < q$

3. $x < a$

Choose q between a and p . Change:

- $x < a$ and $a < q$, so $x < q$

(same conclusion as $x > a$)

In each of these three cases, every (x, y) contains a point q from (a, b) . Therefore, we have satisfied Definition 4, so p is a limit point of (a, b) .

Solution 6.

Solution 7. (Amber)

Solution 8. (Greg)

Consider a point p . According to Axiom 4, there exists a largest integer M_x and a smallest integer M_y such that $M_x < p < M_y$. Then, choose points x and y from Axiom 3 such that $M_x < x < p$ and $p < y < M_y$, and consider the segment $S = (x, y)$.

We will try to find an integer different from p inside S . Axiom 1 gives us three cases:

1. $n = p$ (note: this is only possible if p is an integer)

Here, n is not different from p , so we have not found an integer different from p .

2. $n < p$

Note that n must satisfy $n \leq M_x$; if $n > M_x$, we have contradicted Axiom 4. Then, from Axiom 2, $n \leq M_x$ and $M_x < x$, so $n < x$. This shows that n is not between x and y , so S does not contain n .

3. $n > p$

This case is symmetric to the $n < p$ case.

Therefore, S is a segment containing p that does not contain any integers different from p . Since every element of M is an integer, we have proven that there exists a segment S for every point p without any other points of M , so M has no limit points.

Solution 9.

Solution 10.

Solution 11. (Greg)

Construct a segment $S = (x, y)$ that contains 0 ($x < 0 < y$). Since the reciprocal of a positive number is positive, every element M_i of M is positive, so $x < M_i$ from Axiom 2.

Then, we will attempt to find an element of M that is less than y . Since every element of M is of the form $\frac{1}{n}$ for some integer n , we are looking for $\frac{1}{n} < y$. From the properties of reciprocals, this is equivalent to $\frac{1}{y} < n$. According to Axiom 4, the point $\frac{1}{y}$ has an integer greater than it, so we can find an n that satisfies this inequality.

Therefore, every S containing 0 also contains a point of M , so 0 is a limit point of M .

Solution 12.**Solution 13.** (Erin)

First, note that the set $H \cap K$ is a subset of H because every point of $H \cap K$ must be a point of H . Similarly, $H \cap K \subseteq K$.

Then, from Problem 7, since p is a limit point of $H \cap K$, p is also a limit point of H ; identically, p is a limit point of K .

Solution 14.**Solution 15.****Solution 16.** (Fernando)

Let $S_1 = (a_1, b_1)$ and $S_2 = (a_2, b_2)$. Since p is in both segments, $a_1 < p < b_1$ and $a_2 < p < b_2$. From Axiom 2, we know that:

- $a_1 < b_1$
- $a_1 < b_2$
- $a_2 < b_1$
- $a_2 < b_2$

Then, there are only 4 segments that could be the result of the intersection $S_1 \cap S_2$:

- (a_1, b_1)
- (a_1, b_2)
- (a_2, b_1)
- (a_2, b_2)

Since p is in all of these segments, we have shown that the resulting segment will always contain p .

Solution 17. (Greg)

p is not a limit point of H , so there is a segment $S_H = (H_a, H_b)$ that contains p but no point of H different from p . Likewise, there is a segment $S_K(K_a, K_b)$ that contains p but no different point of K . From these two segments, construct $S = S_H \cap S_K$.

Since p was in both of the original segments, Problem 16 tells us that S contains p . However, since there were no elements of H or K in these segments (except possibly p), S contains no elements of H or K . Finally, because every element of $H \cup K$ is an element of H or K , we know that S contains no elements of $H \cup K$.

Therefore, S is a segment that contains p but no other elements of $H \cup K$. We have found a counterexample to the definition of a limit point, so p is not a limit point of $H \cup K$.

Solution 18.