$v \in \text{ext}(L_{k,\alpha})$ and every n, there is a word x such that $|x| > |u_n|$ and $xv \in \text{ext}(L_{k,\alpha})$; since \mathbf{u} contains the factor xv, there is an occurrence of v in \mathbf{u} outside the prefix u_n . Hence v occurs in \mathbf{u} infinitely many times. Thus \mathbf{u} is recurrent and we proved this implication.

Now turn to the backward implication. For arbitrary words $u, v \in \text{ext}(L_{k,\alpha})$ each of them occurs in **u** infinitely often, so we can find a factor of the form uwv. This factor also occurs in **u** infinitely often, allowing us to find arbitrarily long words x, y such that xuwvy is a factor of **u**. Then $uwv \in \text{ext}(L_{k,\alpha})$. Hence we proved the Restivo-Salemi property for $L_{k,\alpha}$.

Remark 39. It is worth mentioning that for small binary languages Theorem 37 works in an extremal form. Since $2^+ \le \alpha \le 7/3$ implies $\mathsf{ext}(L_{2,\alpha}) = \mathsf{Fac}(\mathbf{t})$, the language $L_{2,\alpha}$ trivially has the Restivo-Salemi property; as we know from Corollary 3, all α -power-free infinite binary words contain all words of $\mathsf{ext}(L_{2,\alpha})$ as factors.

The following conjecture is based on extensive numerical studies.

Conjecture 40. [30, Conjecture 1] All power-free languages satisfy the Restivo-Salemi property.

As an approach to Conjecture 40, we suggest the following.

Open Question 41. Prove the converse of Corollary 38.

References

- [1] J.-P. Allouche and J. O. Shallit. The ubiquitous Prouhet-Thue-Morse sequence. In C. Ding, T. Helleseth, and H. Niederreiter, editors, *Sequences and Their Applications*, *Proceedings of SETA '98*, pages 1–16. Springer-Verlag, 1999.
- [2] J.-P. Allouche and J. Shallit. Automatic Sequences: Theory, Applications, Generalizations. Cambridge University Press, 2003.
- [3] S. E. Arshon. Proof of the existence of asymmetric infinite sequences. *Mat. Sbornik* **2** (1937), 769–779. In Russian, with French abstract.
- [4] S. V. Avgustinovich. The number of different subwords of given length in the Morse-Hedlund sequence. Sibirsk. Zh. Issled. Oper. 1 (1994), 3–7,103. In Russian. English translation in A. D. Korshunov, ed., Discrete Analysis and Operations Research, Kluwer, 1996, pp. 1-5.
- [5] S. Brlek. Enumeration of factors in the Thue-Morse word. *Disc. Appl. Math.* **24** (1989), 83–96.
- [6] A. Carpi and A. de Luca. Special factors, periodicity, and an application to Sturmian words. Acta Inf. 36(12) (2000), 983–1006.