

Formaldehyde S1 · frozen X11 common-slice surfaces + X07 full electronic dipole derivatives
Zero-temperature, one-local-well double-harmonic/linear-HT model · no component normalization
HT activation: total is nonnegative
Total, σ = 0.02 eV
HT, σ = 0.02 eV
FC-only, raw
Total, σ = 0.01 eV
Oscillator-strength density / eV⁻¹
Photon energy / eV
×10⁻³⁰ FC-only numerical symmetry residual
Σ f_FC = 2.847e-31
Raw FC oscillator-strength density / eV⁻¹
Photon energy / eV
×10⁻¹⁷ Signed interference term, not absorption
Positive
Negative
Interference contribution / eV⁻¹
Photon energy / eV
×10⁻³ Initial-mode contributions to HT closure
Σₓ |∂μₓ/∂Qₖ|² / (2ωg,k)
Includes zero-point coordinate variance
Integrated HT dipole strength / (e bohr)²
Ground normal mode index
Closure and raw-tail convergence
Relative dipole-strength remainder
Change in integrated f
Profile change, larger of two σ
Closure gate
Relative remainder or change
Maximum total vibrational quanta
×10⁻⁵ Largest activated lines: full Cartesian sum
(2,0,0,0,0,0)
(2,0,0,1,0,0)
(3,0,0,1,0,0)
Total oscillator strength f
Energy above model 0–0 / eV