

MAHC 10-FALL RISK ASSESSMENT TOOL

Member Name:	
Member ID:	
Date of Assessment	
UAS Assessment Type	

Instructions: The UAS RN must complete the Fall Assessment each time they complete an UAS-NY; and scoring should be based on assessment finding within the context of the question protocols listed with each question.

Required Core Elements Assess one point for each core element “yes”	Points
Age 65+	
Diagnosis (3 or more co-existing) Includes only documented medical diagnosis	
Prior history of falls within 3 months An unintentional change in position resulting in coming to rest on the ground or at a lower level	
Incontinence Inability to make it to the bathroom or commode in timely manner Includes frequency,urgency, and/or nocturia.	
Visual impairment Includes but not limited to, macular degeneration, diabetic retinopathies, visual field loss,age related changes, decline in visual acuity, accommodation, glare tolerance, depth perception, and night vision or not wearing prescribed glasses or having the correct prescription.	
Impaired functional mobility May include patients who need help with IADLS or ADLS or have gait or transfer problems,arthritis, pain, fear of falling, foot problems, impaired sensation, impaired coordination or improper use of assistive devices.	
Environmental hazards May include but not limited to, poor illumination, equipment tubing, inappropriate footwear, pets, hard to reach items, floor surfaces that are uneven or cluttered, or outdoor entry and exits.	
Poly Pharmacy (4 or more prescriptions – any type) All PRESCRIPTIONS including prescriptions for OTC meds. Drugs highly associated with fallrisk include but not limited to, sedatives, anti-depressants, tranquilizers, narcotics, antihypertensives, cardiac meds, corticosteroids, anti-anxiety drugs, anticholinergic drugs, and hypoglycemic drugs.	
Pain affecting level of function Pain often affects an individual’s desire or ability to move or pain can be a factor in depression or compliance with safety recommendations	
Cognitive impairment Could include patients with dementia, Alzheimer’s or stroke patients or patients who are confused, use poor judgment, have decreased comprehension, impulsivity, memory deficits. Consider patients ability to adhere to the plan of care.	
Add up the points and enter a Total	
A score of 4 or more is considered at risk for falling	
UAS Assessor Name:	

MAHC 10-FALL RISK ASSESSMENT TOOL

Required Interventions for All Members:	Optional Interventions for Members with Risk Score of ≥ 4 , and/or to address any specific risk identified on Mach 10:
☐ Home Safety Assessed, Recommendation Reviewed with Member/Caregiver	☐ Recommend CM consults with PCP to modify/change Medication Regimen as member is on high-risk medications
☐ Medication Reconciliation	☐ Recommend CM notifies MD of Fall Risk and if applicable, schedule an appointment with the PCP for member
☐ Education on high-risk meds	☐ Recommend CM discuss Vision Evaluation with member and coordinated Eye Exam appointment as needed
☐ Member/Caregiver Education Universal Fall Prevention	☐ Referral for Skilled Services : ☐ SN ☐ PT ☐ OT ☐ MSW
☐ PERS Reviewed with member/caregiver, if applicable	☐ Referral to Integra Social Worker
Optional Interventions for All Members Select all applicable recommendations below:	☐ Recommend CM consults with PCP regarding Evaluation, Diagnosis and Management of member's incontinence
☐ Developed/Reviewed/ Revised as Necessary Back - Up Care Emergency Plan	☐ Recommend Development of an individualized Toileting Plans for Urinary Incontinence
☐ Recommend PERS	☐ Recommend Development of Continance Care and BowelManagement Program
☐ CM to Request UAS Assessment, if applicable	☐ Recommend Heavy Duty Housekeeping
☐ N/A	☐ Recommend Home Modification
	☐ N/A

Universal Falls Prevention Strategies

- **Ensure** mobility devices are kept within reach. If you use a cane, walker or a mobility device to help you walk, **use it!**
- If you wear glasses or hearing aids, **use them.**
- **Wear** appropriate footwear. Wear well-fitting shoes with good support, both inside and outside the house. **Avoid** going barefoot, wearing shoes or slippers that do not have backs on them.
- **Remove clutter** -Keep the floor and stairs clear of objects such as books, boxes, blankets, papers, shoes, clothing and other tripping hazards.
- **Arrange** furniture so that it creates plenty of room to walk freely. Move coffee tables, magazine racks, plant stands from high-traffic areas. Get help to move the furniture, so your path is clear.
- **Avoid** low chairs that are difficult to sit in and get out of easily. Use stable chairs with armrests to help you get up
- **Arrange** electrical/phone cords and Oxygen tubing so that they are not a tripping hazard
- **Install** grab bars next to the toilet, the tub or shower to prevent slipping. Place nonslip strips or a rubber mat on the floor of the tub or shower.
- **Be sure** you have adequate lighting throughout your house.
- **Keep** an easy to use lamp close to your bed.
- **Use** nightlights to light dark hallways and the pathway from the bedroom to the bathroom at night.
- **Remove** throw rugs or secure them firmly to the floor.
- **Keep** floor surfaces clean and dry; immediately clean spilled liquids, grease or food.
- **Use** caution when carrying items while walking.
- **Avoid distractions while walking.** Concentrate on the task of walking.
- **Stay active** to maintain overall strength and endurance. Talk to your PCP about starting an exercise program such as Silver Slippers, Healthy Moves.
- **Know** your limitations. If there is a task you cannot complete with ease, do not risk a fall by trying to complete it. **ASK for Help!**