

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY

LOGIC AND COMPUTATION PROJECT

Coq Modules for Relational Calculus

(Ver.0.1)

Hisaharu TANAKA Saga University

Shuichi INOKUCHI Fukuoka Institute of Technology Toshiaki MATSUSHIMA Kyushu University

Yoshihiro MIZOGUCHI Kyushu University

May 3, 2016

 ${\bf Repository:}\ \ {\tt https://github.com/KyushuUniversityMathematics/RelationalCalculus}$

Contents

1	Libi	eary Basic_Notations	4
	1.1	このライブラリについて	4
	1.2	定義	4
	1.3	関数の定義	7
	1.4	関係の公理・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・	7
		1.4.1 Dedekind 圏の公理	8
		1.4.2 排中律	12
			12
			12
			13
			13
		_,	
2	Libi		14
	2.1	定義	14
	2.2	関数の定義	15
	2.3	関係の公理	15
		2.3.1 Dedekind 圏の公理	16
		2.3.2 排中律	24
			25
		2.3.4 選択公理	26
		2.3.5 関係の有理性	27
		2.3.6 直和と直積	30
3	Libi	σ	35
	3.1		35
		3.1.1 和関係, 共通関係	35
		3.1.2 分配法則	45
		**** • · · ·	46
	3.2	Heyting 代数に関する補題	48
	3.3	補関係に関する補題・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・	55
	3 4		50

4	Libr		62
	4.1	関係計算の基本的な性質・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・・	62
	4.2	1 1	73
	4.3	逆関係に関する補題	75
	4.4		80
	4.5	単域と Tarski の定理	85
5	Libr	ary Functions_Mappings	90
	5.1		90
	5.2		01
	5.3	,	09
6	Libr	ary Tactics 1	12
	6.1	· ·	12
	6.2		13
	6.3		14
7	Libr		17
	7.1	Dedekind formula に関する補題	17
	7.2	Dedekind formula と全関係	18
	7.3	Dedekind formula と恒等関係	21
8	Libr	ary Conjugate	25
	8.1	共役性の定義 1	25
	8.2	共役の例 \ldots	26
9	Libr	ary Domain	35
	9.1	定義域の定義	35
	9.2	定義域の性質	36
		9.2.1 基本的な性質 1	36
			38
		9.2.3 その他の性質	42
			48
10	Libr	ary Residual 1	51
	10.1	剰余合成関係の性質 1	51
		10.1.1 基本的な性質 1	51
		10.1.2 単調性と分配法則	54
			57
	10.2		60
			61
			71
			71
			76

11	Libr	rary So	chroder													178
	11.1	Schröd	ler 圏の 性	生質 .				 •	 	•		 •			 	178
12	Libr	ary Su	ım_Pro	duct												186
	12.1	関係の	直和 .						 						 	186
		12.1.1	入射対,	関係直	1和0)定	莨		 						 	 186
		12.1.2	関係直和	1の性質	質 .				 						 	188
		12.1.3	分配法則	钊					 						 	 192
	12.2	関係の	直積 .						 						 	194
		12.2.1	射影対,	関係直	1積の)定	莨		 						 	 194
			関係直和													
			鋭敏性													
			分配法則													
13	Libr	ary Po	$\operatorname{oint}_{-}\mathbf{A}\mathbf{x}$	iom												214
	13.1	I-点 .							 						 	214
		13.1.1	I-点の定	⋮義					 						 	214
		13.1.2	I-点の性	Ł質					 						 	215
	13.2	I-点に	関する諸	公理 .					 						 	 217
			点公理													
		13.2.2	全域性的	公理 .					 						 	 221
		13.2.3	その他の	の公理					 						 	 224
	13.3		の補題													

Chapter 1

Library Basic_Notations

Require Export Ssreflect.ssreflect Ssreflect.eqtype MathComp.bigop.
Require Export Logic.ClassicalFacts.

1.1 このライブラリについて

- このライブラリは河原康雄先生の"関係の理論 Dedekind 圏概説 -" をもとに制作されている.
- 現状サポートしているのは、
 - 1.4 節大半, 1.5 1.6 節全部
 - 2.1 2.3 節全部, 2.4 2.5 節大半, 2.6 節全部, 2.7 節大半, 2.8 節有理性
 - 4.2 4.3 節全部, 4.4 4.5 節大半, 4.6 節命題 4.6.1, 4.7 節大半, 4.8 節命題 4.8.1, 4.9 節全部
 - -5.1 節大半, 5.2 5.3 節一部

といったところである.

● 第 3 章以外でカバーしていない箇所は、基礎的もしくは自明な補題、Coq での定式 化が難しい定義や補題、証明の正当性が示しきれなかった補題、汎用性の低い一部 の記号などである。

1.2 定義

- A, B を eqType として, A から B への関係の型を (Rel A B) と書き, $A \to B \to Prop$ として定義する. 本文中では型 (Rel A B) を $A \to B$ と書く.
- 関係 $\alpha:A\to B$ の逆関係 $\alpha^{\sharp}:B\to A$ は (inverse α) で, Coq では (α #) と記述する.

- 2 つの関係 $\alpha:A\to B,\beta:B\to C$ の合成関係 $\alpha\cdot\beta(\mathrm{or}\ \alpha\beta):A\to C$ は $(\mathrm{composite}\ \alpha\ \beta)$ で、 $(\alpha\ \cdot\ \beta)$ と記述する.
- 剰余合成関係 $\alpha \triangleright \beta : A \rightarrow C$ は (residual $\alpha \beta$) で, $(\alpha \beta)$ と記述する.
- 恒等関係 $id_A: A \rightarrow A$ は (identity A) で, (Id A) と記述する.
- 空関係 $\phi_{AB}: A \rightarrow B$ は (empty AB) で, (AB) と記述する.
- 全関係 $\nabla_{AB}: A \rightarrow B$ は (universal AB) で, (AB) と記述する.
- 2 つの関係 $\alpha:A\to B,\,\beta:A\to B$ の和関係 $\alpha\sqcup\beta:A\to B$ は $(\operatorname{cup}\ \alpha\ \beta)$ で、 $(\alpha\quad \beta)$ と記述する.
- 共通関係 $\alpha \sqcap \beta : A \to B$ は (cap $\alpha \beta$) で, $(\alpha \quad \beta)$ と記述する.
- 相対擬補関係 $\alpha \Rightarrow \beta : A \rightarrow B$ は $(\operatorname{rpc} \alpha \beta)$ で, $(\alpha >> \beta)$ と記述する.
- 関係 $\alpha:A\to B$ の補関係 $\alpha^-:A\to B$ は (complement α) で, Coq では $(\alpha ^\circ)$ と記述する.
- 2 つの関係 $\alpha:A \to B$, $\beta:A \to B$ の差関係 $\alpha-\beta:A \to B$ は (difference α β) で, $(\alpha --\beta)$ と記述する.
- (cupP) と (capP) は添字付の和関係と共通関係であり、述語 P に対し、 $\{f(\alpha) \mid P(\alpha)\}$ の和関係、共通関係を表す.
- また、1点集合 I = {*} は i と表記する.
- ◆ なお,通常の共通関係,和関係も添字付のもので表現することができるため、ここでは それを用いて表記する.
- 後で述べるように、剰余合成 $\alpha \triangleright \beta$ も $(\alpha \cdot \beta^-)^-$ のように表現することは可能だが、"剰余合成が存在すれば、それは $(\alpha \cdot \beta^-)^-$ に等しい" というレベルのものであるため、剰余合成に関する公理はやはり必要となる.

表 1.1 に関係の表記についてまとめる.

Axiom prop_extensionality_ok: prop_extensionality.

Definition $Rel(A B : eqType) := A \rightarrow B \rightarrow Prop.$

Module Type Relation.

Parameter inverse: $(\forall A B : eqType, Rel A B \rightarrow Rel B A)$.

Notation "a #" := $(inverse _ a)$ (at level 20).

Parameter composite: $(\forall A B C : eqType, Rel A B \rightarrow Rel B C \rightarrow Rel A C)$.

Notation "a' \cdot 'b" := $(composite _ _ _ a \ b)$ (at level 50).

Parameter residual: $(\forall A B C : eqType, Rel A B \rightarrow Rel B C \rightarrow Rel A C)$.

Notation "a', b" := $(residual _ _ _ a \ b)$ (at level 50).

	数式	Coq	Notation
逆関係	$lpha^{\sharp}$	(inverse α)	(α #)
合成関係	$\alpha \cdot \beta$	(composite $\alpha \beta$)	$(\alpha \cdot \beta)$
剰余合成関係	$\alpha \rhd \beta$	$(exttt{residual} \ lpha \ eta)$	$(\alpha \qquad \beta)$
恒等関係	id_A	(identity A)	$(\operatorname{Id}\ A)$
空関係	ϕ_{AB}	$(\mathtt{empty}\ A\ B)$	(AB)
全関係	∇_{AB}	$(\mathtt{universal}\ A\ B)$	(AB)
和関係	$\alpha \sqcup \beta$	$(\operatorname{cup} \ \alpha \ \beta)$	$(\alpha \qquad \beta)$
共通関係	$\alpha \sqcap \beta$	$(extsf{cap} \ lpha \ eta)$	$(\alpha \qquad \beta)$
相対擬補関係	$\alpha \Rightarrow \beta$	$(\operatorname{\mathtt{rpc}}\ \alpha\ \beta)$	$(\alpha >> \beta)$
補関係	α^{-}	$(\mathtt{complement}\ \alpha)$	(α ^)
差関係	$\alpha - \beta$	(difference $\alpha \beta$)	$(\alpha \beta)$
添字付和関係	$\sqcup_{P(\alpha)} f(\alpha)$	$(\operatorname{cupP}\ P\ f)$	$(-\{P\} f)$
添字付共通関係	$\sqcap_{P(\alpha)} f(\alpha)$	$(\mathtt{capP}\ P\ f)$	$(-\{P\} f)$

Table 1.1: 関係の表記について

```
Parameter identity: (\forall A : eqType, Rel A A).
Notation "'Id'" := identity.
Parameter empty : (\forall A B : eqType, Rel A B).
Notation "' := empty.
Parameter universal : (\forall A B : eqType, Rel A B).
Notation "' := universal.
Parameter include: (\forall A B : eqType, Rel A B \rightarrow Rel A B \rightarrow Prop).
Notation "a', b" := (include \_ \_ a \ b) (at level 50).
Parameter cupP: (\forall A B C D : eqType, (Rel C D \rightarrow Prop) \rightarrow (Rel C D \rightarrow Rel A B) \rightarrow
Rel\ A\ B).
                \{ p' \}' f'' := (cupP_{-} - p f) \text{ (at level 50)}.
Parameter capP: (\forall A \ B \ C \ D: eqType, (Rel \ C \ D \rightarrow Prop) \rightarrow (Rel \ C \ D \rightarrow Rel \ A \ B) \rightarrow
Rel\ A\ B).
                \{', p'\}'  f" := (capP_{---} p f) (at level 50).
Notation "'
Definition cup \{A B : eqType\} (alpha beta : Rel A B)
       _{\text{loc}} = \{\text{fun } gamma : Rel } A B \Rightarrow gamma = alpha \lor gamma = beta \} id.
Notation "a', b" := (cup \ a \ b) (at level 50).
Definition cap {A B : eqType} (alpha beta : Rel A B)
       \{\text{fun } gamma : Rel \ A \ B \Rightarrow gamma = alpha \lor gamma = \text{beta}\} \ id.
Notation "a', b" := (cap \ a \ b) (at level 50).
Parameter rpc: (\forall A B : eqType, Rel A B \rightarrow Rel A B \rightarrow Rel A B).
Notation "a'»' b" := (rpc - a b) (at level 50).
Definition complement \{A \ B : eqType\} (alpha : Rel \ A \ B) := alpha \gg
                                                                                     A B.
```

```
Notation "a '^'" := (complement\ a) (at level 20).

Definition difference\ \{A\ B: eqType\}\ (alpha\ beta: Rel\ A\ B) := alpha\ beta^*.

Notation "a - b" := (difference\ a\ b) (at level 50).

Notation "i'" := unit\_eqType.
```

1.3 関数の定義

 $\alpha:A\to B$ に対し、全域性 total_r、一価性 univalent_r、関数 function_r、全射 surjective_r、単射 injective_r、全単射 bijection_r を以下のように定義する.

```
• total_r : id_A \sqsubseteq \alpha \cdot \alpha^{\sharp}
```

- univalent_r : $\alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- function_r : $id_A \sqsubseteq \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- surjection_r : $id_A \sqsubseteq \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha = id_B$
- injection_r : $id_A = \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- bijection_r : $id_A = \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha = id_B$

```
Definition total_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B) := (Id \ A) \quad (alpha \bullet alpha \#).
Definition univalent_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B) := (alpha \# \bullet alpha) \quad (Id \ B).
Definition function_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B)
:= (total_r \ alpha) \land (univalent_r \ alpha).
Definition surjection_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B)
:= (function_r \ alpha) \land (total_r \ (alpha \#)).
Definition injection_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B)
:= (function_r \ alpha) \land (univalent_r \ (alpha \#)).
Definition bijection_r \ \{A \ B : eqType\} \ (alpha : Rel \ A \ B)
:= (function_r \ alpha) \land (total_r \ (alpha \#)) \land (univalent_r \ (alpha \#)).
```

1.4 関係の公理

今後の諸定理の証明は、原則以下の公理群、およびそれらから導かれる補題のみを用いて行っていくことにする.

1.4.1 Dedekind 圏の公理

Axiom 1 (comp_id_l, comp_id_r) Let $\alpha : A \rightarrow B$. Then,

$$id_A \cdot \alpha = \alpha \cdot id_B = \alpha.$$

Definition $axiom1a := \forall (A \ B : eqType)(alpha : Rel \ A \ B), Id \ A \bullet alpha = alpha.$

Axiom $comp_{-}id_{-}l$: axiom1a.

Definition $axiom1b := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha \cdot Id \ B = alpha.$

Axiom $comp_id_r$: axiom1b.

Axiom 2 (comp_assoc) Let $\alpha : A \rightarrow B$, $\beta : B \rightarrow C$, and $\gamma : C \rightarrow D$. Then,

$$(\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).$$

Definition axiom2 :=

 $\forall (A \ B \ C \ D : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ C \ D),$

 $(alpha \cdot beta) \cdot gamma = alpha \cdot (beta \cdot gamma).$

Axiom $comp_assoc: axiom2$.

Axiom 3 (inc_refl) Let $\alpha : A \rightarrow B$. Then,

 $\alpha \sqsubset \alpha$.

Definition $axiom3 := \forall (A B : eqType)(alpha : Rel A B), alpha alpha.$

Axiom $inc_refl: axiom3$.

Axiom 4 (inc_trans) Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$\alpha \sqsubseteq \beta \land \beta \sqsubseteq \gamma \Rightarrow \alpha \sqsubseteq \gamma.$$

Definition $axiom4 := \forall (A B : eqType)(alpha beta gamma : Rel A B),$

alpha beta \rightarrow beta $qamma \rightarrow alpha$ qamma.

Axiom $inc_trans: axiom4$.

Axiom 5 (inc_antisym) Let $\alpha, \beta : A \rightarrow B$. Then,

$$\alpha \sqsubseteq \beta \land \beta \sqsubseteq \alpha \Rightarrow \alpha = \beta.$$

Definition $axiom5 := \forall (A B : eqType)(alpha beta : Rel A B),$

alpha beta \rightarrow beta $alpha \rightarrow alpha =$ beta.

Axiom $inc_antisym : axiom5$.

Axiom 6 (inc_empty_alpha) Let $\alpha : A \rightarrow B$. Then,

 $\phi_{AB} \sqsubseteq \alpha$.

Definition $axiom6 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), A \ B \ alpha.$ Axiom $inc_empty_alpha : axiom6.$

Axiom 7 (inc_alpha_universal) Let $\alpha : A \rightarrow B$. Then,

 $\alpha \sqsubseteq \nabla_{AB}$.

Definition $axiom 7 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha$ Axiom $inc_alpha_universal : axiom 7$.

Axiom 8 (inc_capP, inc_cap)

- 1. $\mathbf{inc_capP}$: Let $\alpha: A \to B$, $f: (C \to D) \to (A \to B)$ and P: predicate. Then, $\alpha \sqsubseteq (\sqcap_{P(\beta)} f(\beta)) \Leftrightarrow \forall \beta: C \to D, P(\beta) \Rightarrow \alpha \sqsubseteq f(\beta).$
- 2. inc_cap: Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$\alpha \sqsubseteq (\beta \sqcap \gamma) \Leftrightarrow \alpha \sqsubseteq \beta \land \alpha \sqsubseteq \gamma.$$

```
Definition axiom8a :=
 \forall (A B C D : eqType)
 (alpha: Rel \ A \ B)(f: Rel \ C \ D \rightarrow Rel \ A \ B)(P: Rel \ C \ D \rightarrow Prop),
            (-\{P\} f) \leftrightarrow \forall \text{ beta} : Rel \ C \ D, P \text{ beta} \rightarrow alpha
Axiom inc\_capP : axiom8a.
Definition axiom8b := \forall (A B : eqType)(alpha beta qamma : Rel A B),
                        gamma) \leftrightarrow (alpha \quad beta) \land (alpha)
Lemma inc\_cap: axiom8b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
rewrite inc\_capP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

Axiom 9 (inc_cupP, inc_cup)

1. $\mathbf{inc_cupP}$: Let $\alpha: A \to B$, $f: (C \to D) \to (A \to B)$ and P: predicate. Then, $(\sqcup_{P(\beta)} f(\beta)) \sqsubseteq \alpha \Leftrightarrow \forall \beta: C \to D, P(\beta) \Rightarrow f(\beta) \sqsubseteq \alpha$.

2. inc_cup: Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

 $(\beta \sqcup \gamma) \sqsubseteq \alpha \Leftrightarrow \beta \sqsubseteq \alpha \land \gamma \sqsubseteq \alpha.$

```
Definition axiom9a :=
 \forall (A \ B \ C \ D : eqType)
 (alpha: Rel\ A\ B)(f: Rel\ C\ D \rightarrow Rel\ A\ B)(P: Rel\ C\ D \rightarrow Prop),
                   alpha \leftrightarrow \forall beta : Rel\ C\ D, P beta \rightarrow f beta
Axiom inc\_cupP: axiom9a.
Definition axiom9b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             gamma)
                           alpha \leftrightarrow (\texttt{beta} \quad alpha) \land (gamma)
Lemma inc\_cup: axiom9b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
rewrite inc\_cupP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

```
Axiom 10 (inc_rpc) Let \alpha, \beta, \gamma : A \to B. Then,

\alpha \sqsubset (\beta \Rightarrow \gamma) \Leftrightarrow (\alpha \sqcap \beta) \sqsubset \gamma.
```

Definition $axiom10 := \forall (A \ B : eqType)(alpha \ beta \ gamma : Rel \ A \ B),$ $alpha \ (beta \gg gamma) \leftrightarrow (alpha \ beta) \ gamma.$ Axiom $inc_rpc : axiom10$.

Axiom 11 (inv_invol) Let $\alpha : A \to B$. Then, $(\alpha^{\sharp})^{\sharp} = \alpha.$

Definition $axiom11 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), (alpha \#) \# = alpha.$ Axiom $inv_invol : axiom11$.

Axiom 12 (comp_inv) Let $\alpha : A \rightarrow B$ and $\beta : B \rightarrow C$. Then,

$$(\alpha \cdot \beta)^{\sharp} = \beta^{\sharp} \cdot \alpha^{\sharp}.$$

Definition $axiom12 := \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C),$ $(alpha \cdot beta) \# = (beta \# \cdot alpha \#).$

Axiom $comp_inv : axiom12$.

Axiom 13 (inc_inv) Let $\alpha, \beta : A \rightarrow B$. Then,

$$\alpha \sqsubseteq \beta \Rightarrow \alpha^{\sharp} \sqsubseteq \beta^{\sharp}.$$

Definition axiom13 :=

 $\forall (A \ B : eqType)(alpha \ \mathsf{beta} : Rel \ A \ B), \ alpha \ \ \mathsf{beta} \to alpha \ \# \ \ \mathsf{beta} \ \#.$ Axiom $inc_inv : axiom13$.

Axiom 14 (dedekind) Let $\alpha: A \rightarrow B$, $\beta: B \rightarrow C$, and $\gamma: A \rightarrow C$. Then,

$$(\alpha \cdot \beta) \sqcap \gamma \sqsubseteq (\alpha \sqcap (\gamma \cdot \beta^{\sharp})) \cdot (\beta \sqcap (\alpha^{\sharp} \cdot \gamma)).$$

Definition axiom14 :=

 $\forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),$ $((alpha \cdot beta) \quad gamma)$

 $((alpha \quad (gamma \cdot beta \#)) \cdot (beta \quad (alpha \# \cdot gamma))).$

Axiom dedekind : axiom 14.

Axiom 15 (inc_residual) Let $\alpha: A \to B$, $\beta: B \to C$, and $\gamma: A \to C$. Then,

$$\gamma \sqsubseteq (\alpha \rhd \beta) \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.$$

Definition axiom15 :=

 $\forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),$

 $qamma \quad (alpha \quad beta) \leftrightarrow (alpha \# \cdot qamma) \quad beta.$

Axiom $inc_residual : axiom15$.

1.4.2 排中律

Dedekind 圏の公理のほかに、以下の"排中律"を仮定すれば、与えられる圏は Schröder 圏となり、Bool 代数の性質も満たされる. ちなみに剰余合成は補関係から定義可能なので、本来 Schröder 圏には剰余合成に関する公理は存在しない.

Axiom 16 (complement_classic) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcup \alpha^- = \nabla_{AB}$$

Definition $axiom16 := \forall (A B : eqType)(alpha : Rel A B),$

alpha alpha $^{\circ} = AB$.

Axiom complement_classic : axiom16.

1.4.3 単域

1 点集合 *I* が単域となるための条件は

$$\phi_{II} \neq id_I \wedge id_I = \nabla_{II} \wedge \forall X, \nabla_{XI} \cdot \nabla_{IX} = \nabla_{XX}$$

だが、Rel の定義から左 2 つは証明できるため、右の式だけ仮定する.

Axiom 17 (unit_universal)

$$\nabla_{AI} \cdot \nabla_{IA} = \nabla_{AA}$$

Definition $axiom17 := \forall (A : eqType), A i \cdot i A = A A.$

Axiom $unit_universal : axiom17$.

1.4.4 選択公理

この "選択公理" を仮定すれば、排中律と単域の存在 (厳密には全域性公理) を利用して点公理を導出できる.

Axiom 18 (axiom_of_choice) Let $\alpha : A \rightarrow B$ be a total relation. Then,

$$\exists \beta : A \to B, \beta \sqsubseteq \alpha.$$

Definition $axiom18 := \forall (A B : eqType)(alpha : Rel A B),$

 $total_r \ alpha \rightarrow \exists \ \mathtt{beta} : \ Rel \ A \ B, \ function_r \ \mathtt{beta} \wedge \ \mathtt{beta} \quad \ \ alpha.$

Axiom $axiom_of_choice : axiom18$.

1.4.5 関係の有理性

集合論では色々インポートしながら頑張って証明したので、できればそちらもご覧ください.

Axiom 19 (rationality) Let $\alpha : A \rightarrow B$. Then,

$$\exists R, \exists f: R \to A, \exists g: R \to B, \alpha = f^{\sharp} \cdot g \land f \cdot f^{\sharp} \sqcap g \cdot g^{\sharp} = id_R.$$

Definition $axiom19 := \forall (A \ B : eqType)(alpha : Rel \ A \ B),$ $\exists (R : eqType)(f : Rel \ R \ A)(g : Rel \ R \ B),$ $function_r \ f \land function_r \ g \land alpha = f \ \# \ \bullet \ g \land ((f \ \bullet f \ \#) \ (g \ \bullet g \ \#)) = Id \ R.$ Axiom rationality : axiom19.

1.4.6 直和と直積

任意の直和に対して、入射対が存在することを仮定する.

Axiom 20 (pair_of_inclusions) $\exists j : A \to A + B, \exists k : B \to A + B,$

$$j \cdot j^{\sharp} = id_A \wedge k \cdot k^{\sharp} = id_B \wedge j \cdot k^{\sharp} = \phi_{AB} \wedge j^{\sharp} \cdot j \sqcup k^{\sharp} \cdot k = id_{A+B}.$$

Definition axiom20 :=

 \forall $(A \ B : eqType), \exists (j : Rel \ A \ (sum_eqType \ A \ B))(k : Rel \ B \ (sum_eqType \ A \ B)),$ $<math>j \cdot j \# = Id \ A \wedge k \cdot k \# = Id \ B \wedge j \cdot k \# = A \ B \wedge$ $(j \# \cdot j) \quad (k \# \cdot k) = Id \ (sum_eqType \ A \ B).$ Axiom $pair_of_inclusions : axiom20$.

任意の直積に対して、射影対が存在することを仮定する.

実は有理性公理 $(Axiom\ 19)$ があれば直積の公理は必要ないのだが、 $Axiom\ 19$ の準用では直積が "存在する" ことまでしか示してくれないので、"直積として $prod_eqType\ A\ B$ を用いてよい" ことを公理の中に含めたものを用意しておく.

Axiom 21 (pair_of_projections) $\exists p : A \times B \rightarrow A, \exists q : A \times B \rightarrow B,$

$$p^{\sharp} \cdot q = \nabla_{AB} \wedge p \cdot p^{\sharp} \cap q \cdot q^{\sharp} = id_{A \times B}.$$

Definition axiom21 :=

 \forall (A B : eqType), \exists (p : Rel (prod_eqType A B) A)(q : Rel (prod_eqType A B) B), p # • q = A B ∧ (p • p #) (q • q #) = Id (prod_eqType A B) ∧ univalent_r p ∧ univalent_r q.

Axiom $pair_of_projections: axiom21$.

End Relation.

Chapter 2

Library Basic_Notations_Set

```
Require Export Basic_Notations.

Require Import Logic.FunctionalExtensionality.

Require Import Logic.Classical_Prop.

Require Import Logic.IndefiniteDescription.

Require Import Logic.ProofIrrelevance.
```

2.1 定義

この章では、関係を集合論的に定義した場合の定義、およびその定義で諸公理が成立することを示す、公理名や記号などは Basic_Notations と同じものを使用する.

```
Module Rel\_Set <: Relation.
Definition inverse {A B : eqType} (alpha : Rel A B) : Rel B A
 := (\mathbf{fun} \ (b : B)(a : A) \Rightarrow alpha \ a \ b).
Notation "a \#" := (inverse a) (at level 20).
Definition composite {A B C : eqType} (alpha : Rel A B) (beta : Rel B C) : Rel A C
 := (fun (a : A)(c : C) \Rightarrow \exists b : B, alpha \ a \ b \land beta \ b \ c).
Notation "a' \cdot b" := (composite\ a\ b) (at level 50).
Definition residual {A B C : eqType} (alpha : Rel A B) (beta : Rel B C) : Rel A C
 := (fun (a : A)(c : C) \Rightarrow \forall b : B, alpha a b \rightarrow beta b c).
Notation "a', b" := (residual \ a \ b) (at level 50).
Definition identity (A : eqType) : Rel A A := (fun \ a \ a\theta : A \Rightarrow a = a\theta).
Notation "'Id'" := identity.
Definition empty (A B : eqType) : Rel A B := (fun (a : A)(b : B) \Rightarrow False).
Notation "' := empty.
Definition universal (A B : eqType) : Rel A B := (fun (a : A)(b : B) \Rightarrow True).
Notation "' := universal.
Definition include \{A B : eqType\} (alpha beta : Rel A B) : Prop
 := (\forall (a : A)(b : B), alpha \ a \ b \rightarrow beta \ a \ b).
```

```
Notation "a', b" := (include \ a \ b) (at level 50).
Definition cupP \{A \ B \ C \ D : eqType\}\ (P : Rel \ C \ D \rightarrow Prop)\ (f : Rel \ C \ D \rightarrow Rel \ A \ B)
: Rel \ A \ B
:= (\mathbf{fun} \ (a : A)(b : B) \Rightarrow \exists \ alpha : Rel \ C \ D, P \ alpha \wedge (f \ alpha) \ a \ b).
Notation "' _{\{'p'\}'} f" := (cupP \ p \ f) (at level 50).
Definition cap P \{A \ B \ C \ D : eqType\} \ (P : Rel \ C \ D \rightarrow Prop) \ (f : Rel \ C \ D \rightarrow Rel \ A \ B) :
Rel A B
 := (\mathbf{fun} \ (a : A)(b : B) \Rightarrow \forall \ alpha : Rel \ C \ D, P \ alpha \rightarrow (f \ alpha) \ a \ b).
Notation "' _{-}{' p '}' f" := (capP \ p \ f) (at level 50).
Definition cup \{A B : eqType\} (alpha beta : Rel A B)
       _{\text{-}}\{\text{fun } qamma : Rel \ A \ B \Rightarrow qamma = alpha \lor qamma = \text{beta}\} \ id.
Notation "a', b" := (cup \ a \ b) (at level 50).
Definition cap \{A B : eqType\} (alpha beta : Rel A B)
        \{\text{fun } qamma : Rel \ A \ B \Rightarrow qamma = alpha \lor qamma = \text{beta}\} \ id.
Notation "a', b" := (cap \ a \ b) (at level 50).
Definition rpc \{A B : eqType\} (alpha beta : Rel A B) : Rel A B
 := (fun (a : A)(b : B) \Rightarrow alpha \ a \ b \rightarrow beta \ a \ b).
Notation "a'»' b" := (rpc \ a \ b) (at level 50).
Definition complement \{A \ B : eqType\} (alpha : Rel \ A \ B) := alpha \gg
                                                                                         A B.
Notation "a '^' := (complement \ a) (at level 20).
Definition difference \{A \ B : eqType\}\ (alpha \ beta : Rel \ A \ B) := alpha
                                                                                          beta ^.
Notation "a - b" := (difference a b) (at level 50).
Notation "'i'" := unit\_eqType.
```

2.2 関数の定義

```
Definition total\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (Id \ A) (alpha \cdot alpha \#).

Definition univalent\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (alpha \# \cdot alpha) (Id \ B).

Definition function\_r {A \ B : eqType} (alpha : Rel \ A \ B)

:= (function\_r \ alpha) \land (funivalent\_r \ alpha).

Definition function\_r \ alpha) \land (funivalent\_r \ alpha).
```

2.3 関係の公理

今後の諸定理の証明は、原則以下の公理群、およびそれらから導かれる補題のみを用いて行っていくことにする.

Lemma 1 (comp_id_l, comp_id_r) Let $\alpha : A \rightarrow B$. Then,

2.3.1 Dedekind 圏の公理

```
id_A \cdot \alpha = \alpha \cdot id_B = \alpha.
Definition axiom1a := \forall (A B : eqType)(alpha : Rel A B), Id A • alpha = alpha.
Lemma comp\_id\_l: axiom1a.
Proof.
move \Rightarrow A \ B \ alpha.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_{-}extensionality_{-}ok.
split.
elim \Rightarrow a\theta.
elim \Rightarrow H H0.
rewrite H.
apply H0.
move \Rightarrow H.
\exists a.
split.
by [].
apply H.
Qed.
Definition axiom1b := \forall (A B : eqType)(alpha : Rel A B), alpha • Id B = alpha.
Lemma comp_{-}id_{-}r : axiom1b.
Proof.
move \Rightarrow A \ B \ alpha.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split.
elim \Rightarrow b\theta.
elim \Rightarrow H H0.
rewrite -H0.
apply H.
move \Rightarrow H.
\exists b.
```

split. apply H.

Qed.

```
by [].
Qed.
  Lemma 2 (comp_assoc) Let \alpha: A \to B, \beta: B \to C, and \gamma: C \to D. Then,
                                          (\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).
Definition axiom2 :=
 \forall (A \ B \ C \ D : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ C \ D),
 (alpha \cdot beta) \cdot gamma = alpha \cdot (beta \cdot gamma).
Lemma comp\_assoc: axiom2.
Proof.
move \Rightarrow A B C D alpha beta gamma.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow d.
apply prop_extensionality_ok.
split.
elim \Rightarrow c.
elim \Rightarrow H H0.
elim H \Rightarrow b \ H1.
\exists b.
split.
apply H1.
\exists c.
split.
apply H1.
apply H0.
elim \Rightarrow b.
elim \Rightarrow H.
elim \Rightarrow c H0.
\exists c.
split.
\exists b.
split.
apply H.
apply H0.
apply H\theta.
```

```
Lemma 3 (inc_refl) Let \alpha : A \rightarrow B. Then,
                                                       \alpha \sqsubset \alpha.
Definition axiom3 := \forall (A B : eqType)(alpha : Rel A B), alpha
                                                                                           alpha.
Lemma inc\_refl: axiom3.
Proof.
by [rewrite /axiom3/include].
Qed.
  Lemma 4 (inc_trans) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                           \alpha \sqsubseteq \beta \land \beta \sqsubseteq \gamma \Rightarrow \alpha \sqsubseteq \gamma.
Definition axiom4 := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             \mathtt{beta} \to \mathtt{beta}
                                   gamma \rightarrow alpha
                                                              qamma.
Lemma inc\_trans : axiom4.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma \ H \ H0 \ a \ b \ H1.
apply (H0 - (H - H1)).
Qed.
  Lemma 5 (inc_antisym) Let \alpha, \beta : A \rightarrow B. Then,
                                           \alpha \sqsubseteq \beta \land \beta \sqsubseteq \alpha \Rightarrow \alpha = \beta.
Definition axiom5 := \forall (A B : eqType)(alpha beta : Rel A B),
 alpha
             beta \rightarrow beta
                                    alpha \rightarrow alpha = beta.
Lemma inc\_antisym : axiom5.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ H \ H0.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split.
apply H.
apply H0.
```

Qed.

Lemma 6 (inc_empty_alpha) Let $\alpha : A \rightarrow B$. Then,

 $\phi_{AB} \sqsubseteq \alpha$.

Definition $axiom6 := \forall (A B : eqType)(alpha : Rel A B), A B alpha.$

Lemma inc_empty_alpha : axiom6.

Proof.

move $\Rightarrow A \ B \ alpha \ a \ b$.

apply False_ind.

Qed.

Lemma 7 (inc_alpha_universal) Let $\alpha : A \rightarrow B$. Then,

 $\alpha \sqsubseteq \nabla_{AB}$.

Definition $axiom 7 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha A \ B.$

Lemma $inc_alpha_universal: axiom 7$.

Proof.

move $\Rightarrow A \ B \ alpha \ a \ b \ H$.

apply I.

Qed.

Lemma 8 (inc_capP, inc_cap)

1. $inc_capP : Let \ \alpha : A \rightarrow B, \ f : (C \rightarrow D) \rightarrow (A \rightarrow B) \ and \ P : predicate. Then,$

$$\alpha \sqsubseteq (\sqcap_{P(\beta)} f(\beta)) \Leftrightarrow \forall \beta : C \to D, P(\beta) \Rightarrow \alpha \sqsubseteq f(\beta).$$

2. inc_cap: Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$\alpha \sqsubset (\beta \sqcap \gamma) \Leftrightarrow \alpha \sqsubset \beta \land \alpha \sqsubset \gamma.$$

Definition axiom8a :=

 $\forall (A \ B \ C \ D : eqType)$

 $(alpha: Rel\ A\ B)(f: Rel\ C\ D \rightarrow Rel\ A\ B)(P: Rel\ C\ D \rightarrow Prop),$

alpha ($_{-}\{P\} f$) $\leftrightarrow \forall$ beta: $Rel\ C\ D$, P beta $\to alpha$ f beta.

Lemma inc_capP : axiom8a.

Proof.

move $\Rightarrow A B C D alpha f P$.

 $split; move \Rightarrow H.$

move \Rightarrow beta $H0 \ a \ b \ H1$.

apply (H - H1 - H0).

move $\Rightarrow a \ b \ H0$ beta H1.

```
\overline{\text{apply}} (H - H1 - H0).
Definition axiom8b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
                       qamma) \leftrightarrow (alpha
                                               beta) \land (alpha)
           (beta
                                                                      qamma).
Lemma inc\_cap: axiom8b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ qamma.
rewrite inc\_capP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

Lemma 9 (inc_cupP, inc_cup)

```
1. \operatorname{inc\_cupP} : \operatorname{Let} \alpha : A \to B, \ f : (C \to D) \to (A \to B) \ and \ P : \ predicate. \ Then,
(\sqcup_{P(\beta)} f(\beta)) \sqsubseteq \alpha \Leftrightarrow \forall \beta : C \to D, P(\beta) \Rightarrow f(\beta) \sqsubseteq \alpha.
```

2. inc_cup: Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$(\beta \sqcup \gamma) \sqsubseteq \alpha \Leftrightarrow \beta \sqsubseteq \alpha \land \gamma \sqsubseteq \alpha.$$

```
Definition axiom9a :=
 \forall (A B C D : eqType)
 (alpha: Rel \ A \ B)(f: Rel \ C \ D \rightarrow Rel \ A \ B)(P: Rel \ C \ D \rightarrow Prop),
                    alpha \leftrightarrow \forall beta : Rel\ C\ D,\ P\ beta \rightarrow f\ beta
 ( _{P} f)
                                                                                   alpha.
Lemma inc\_cupP: axiom9a.
Proof.
move \Rightarrow A B C D alpha f P.
split.
move \Rightarrow H beta H0 \ a \ b H1.
apply H.
∃ beta.
split.
apply H0.
apply H1.
move \Rightarrow H \ a \ b.
elim \Rightarrow beta.
elim \Rightarrow H0 \ H1.
```

```
apply (H \text{ beta } H0 - H1).
Definition axiom9b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             qamma)
                            alpha \leftrightarrow (\texttt{beta})
                                                    alpha) \wedge (qamma)
                                                                                alpha).
Lemma inc\_cup: axiom9b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ qamma.
rewrite inc\_cupP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
  Lemma 10 (inc_rpc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                        \alpha \sqsubseteq (\beta \Rightarrow \gamma) \Leftrightarrow (\alpha \sqcap \beta) \sqsubseteq \gamma.
Definition axiom 10 := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             (beta \gg gamma) \leftrightarrow (alpha)
                                                   beta)
                                                                qamma.
Lemma inc\_rpc: axiom10.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
split; move \Rightarrow H.
move \Rightarrow a \ b \ H0.
apply H.
apply H0.
by [left].
apply H0.
by [right].
\mathtt{move} \Rightarrow a \ b \ H0 \ H1.
apply H.
move \Rightarrow delta.
case \Rightarrow H2; rewrite H2.
apply H0.
apply H1.
Qed.
```

```
Lemma 11 (inv_invol) Let \alpha : A \rightarrow B. Then,
                                                       (\alpha^{\sharp})^{\sharp} = \alpha.
Definition axiom11 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), (alpha \#) \# = alpha.
Lemma inv\_invol: axiom11.
Proof.
by [move \Rightarrow A \ B \ alpha].
Qed.
  Lemma 12 (comp_inv) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                                 (\alpha \cdot \beta)^{\sharp} = \beta^{\sharp} \cdot \alpha^{\sharp}.
Definition axiom12 := \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C),
 (alpha \cdot beta) \# = (beta \# \cdot alpha \#).
Lemma comp_inv : axiom12.
Proof.
move \Rightarrow A B C alpha beta.
apply functional_extensionality.
move \Rightarrow c.
apply functional_extensionality.
move \Rightarrow a.
apply prop_extensionality_ok.
split; elim \Rightarrow b.
elim \Rightarrow H H0.
\exists b.
split.
apply H0.
apply H.
elim \Rightarrow H H0.
\exists b.
split.
apply H0.
apply H.
Qed.
  Lemma 13 (inc_inv) Let \alpha, \beta : A \rightarrow B. Then,
                                                 \alpha \sqsubseteq \beta \Rightarrow \alpha^{\sharp} \sqsubseteq \beta^{\sharp}.
```

```
Definition axiom 13 :=  \forall (A B : eqType)(alpha beta : Rel A B), alpha beta <math>\rightarrow alpha \# beta \#.
```

```
Lemma inc_inv : axiom13.
move \Rightarrow A \ B \ alpha \ beta \ H \ b \ a \ H\theta.
apply (H - H0).
Qed.
  Lemma 14 (dedekind) Let \alpha: A \rightarrow B, \beta: B \rightarrow C, and \gamma: A \rightarrow C. Then,
                                 (\alpha \cdot \beta) \sqcap \gamma \sqsubseteq (\alpha \sqcap (\gamma \cdot \beta^{\sharp})) \cdot (\beta \sqcap (\alpha^{\sharp} \cdot \gamma)).
Definition axiom14 :=
 \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),
 ((alpha • beta)
                           gamma)
                    (gamma \cdot beta \#)) \cdot (beta
                                                                (alpha \# \bullet gamma))).
     ((alpha
Lemma dedekind: axiom 14.
Proof.
move \Rightarrow A B C alpha beta gamma a c H.
assert (\exists b : B, alpha \ a \ b \land beta \ b \ c).
apply H.
by [left].
elim H\theta \Rightarrow b.
\mathtt{elim} \Rightarrow \mathit{H1}\ \mathit{H2}.
\exists b.
repeat split.
move \Rightarrow delta H3.
case H3 \Rightarrow H4; rewrite H4.
apply H1.
unfold id.
\exists c.
split.
apply H.
by [right].
apply H2.
move \Rightarrow delta H3.
case H3 \Rightarrow H4; rewrite H4.
apply H2.
unfold id.
\exists a.
split.
apply H1.
apply H.
by [right].
Qed.
```

```
Lemma 15 (inc_residual) Let \alpha: A \to B, \beta: B \to C, and \gamma: A \to C. Then, \gamma \sqsubseteq (\alpha \rhd \beta) \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.
```

```
Definition axiom15 :=
 \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),
                          \texttt{beta}) \leftrightarrow (alpha \# \bullet gamma)
Lemma inc_residual: axiom15.
Proof.
move \Rightarrow A B C alpha beta gamma.
split; move \Rightarrow H.
move \Rightarrow b c.
elim \Rightarrow a H0.
apply (H \ a).
apply H0.
apply H0.
move \Rightarrow a \ c \ H0 \ b \ H1.
apply H.
\exists a.
split.
apply H1.
apply H0.
Qed.
```

2.3.2 排中律

Dedekind 圏の公理のほかに、以下の"排中律"を仮定すれば、与えられる圏は Schröder 圏となり、Bool 代数の性質も満たされる. ちなみに剰余合成は補関係から定義可能なので、本来 Schröder 圏には剰余合成に関する公理は存在しない.

Lemma 16 (complement_classic) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcup \alpha^- = \nabla_{AB}$$

```
Definition axiom16 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha \quad alpha \ \hat{\ } = A \ B.
Lemma complement\_classic : axiom16.
Proof.
move \Rightarrow A \ B \ alpha.
apply functional\_extensionality.
move \Rightarrow a.
apply functional\_extensionality.
```

```
move \Rightarrow b.

apply prop\_extensionality\_ok.

split; move \Rightarrow H.

apply I.

case (classic\ (alpha\ a\ b)) \Rightarrow H0.

\exists\ alpha.

split.

by [left].

apply H0.

\exists\ (\mathbf{fun}\ (a0:A)\ (b0:B) \Rightarrow alpha\ a0\ b0 \rightarrow False).

split.

by [right].

apply H0.

Qed.
```

2.3.3 単域

1点集合 / が単域となるための条件は

$$\phi_{II} \neq id_I \wedge id_I = \nabla_{II} \wedge \forall X, \nabla_{XI} \cdot \nabla_{IX} = \nabla_{XX}$$

だが、Rel の定義から左2 つは証明できるため、右の式だけ仮定する.

Lemma 17 (unit_universal)

$$\nabla_{AI} \cdot \nabla_{IA} = \nabla_{AA}$$

```
Definition axiom 17 := \forall (A : eqType),
                                                                           A A.
                                                   A i \cdot i A =
Lemma unit\_universal : axiom17.
Proof.
move \Rightarrow A.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow a\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
apply I.
\exists tt.
by [].
Qed.
```

2.3.4 選択公理

この "選択公理" を仮定すれば、排中律と単域の存在 (厳密には全域性公理) を利用して点公理を導出できる. 証明には集合論の選択公理を用いる.

Lemma 18 (axiom_of_choice) Let $\alpha : A \rightarrow B$ be a total relation. Then,

$$\exists \beta : A \to B, \beta \sqsubseteq \alpha.$$

```
Definition axiom18 := \forall (A B : eqType)(alpha : Rel A B),
 total\_r \ alpha \rightarrow \exists \ \mathtt{beta} : \ Rel \ A \ B, \ function\_r \ \mathtt{beta} \wedge \ \mathtt{beta}
                                                                                      alpha.
Lemma axiom_of_choice: axiom18.
Proof.
move \Rightarrow A \ B \ alpha.
\verb"rewrite" / function\_r/total\_r/univalent\_r/identity/include/composite/inverse.
move \Rightarrow H.
assert (\forall a : A, \{b : B \mid alpha \ a \ b\}).
move \Rightarrow a.
apply constructive_indefinite_description.
move: (H \ a \ a \ (Logic.eq\_refl \ a)).
elim \Rightarrow b H0.
\exists b.
apply H0.
\exists (fun (a:A)(b:B) \Rightarrow b = sval(X a)).
repeat split.
move \Rightarrow a \ a\theta \ H\theta.
\exists (sval (X a)).
by [rewrite H\theta].
move \Rightarrow b \ b\theta.
elim \Rightarrow a.
elim \Rightarrow H0 \ H1.
by [rewrite H0 H1].
move \Rightarrow a \ b \ H0.
rewrite H0.
apply proj2_sig.
Qed.
```

2.3.5 関係の有理性

集合の選択公理 (Logic.IndefiniteDescription) や証明の一意性 (Logic.ProofIrrelevance) を仮定すれば, 集合論上ならごり押しで証明できる. 旧ライブラリの頃は無理だと諦めて Axiom を追加していたが, Standard Library のインポートだけで解けた. 正直びっくり.

Lemma 19 (rationality) Let $\alpha : A \rightarrow B$. Then,

$$\exists R, \exists f: R \to A, \exists g: R \to B, \alpha = f^{\sharp} \cdot g \land f \cdot f^{\sharp} \sqcap g \cdot g^{\sharp} = id_R.$$

この付近は、ごり押しのための補題. 命題の真偽を選択公理で bool 値に変換したり、部分集合の元から上位集合の元を生成する sval (proj1_sig) の単射性を示したりしている.

```
Lemma is\_true\_inv0: \forall P: Prop, \exists b: bool, P \leftrightarrow is\_true b.
Proof.
move \Rightarrow P.
case (classic P); move \Rightarrow H.
\exists true.
split; move \Rightarrow H0.
by [].
apply H.
\exists false.
split; move \Rightarrow H0.
apply False_ind.
apply (H H\theta).
discriminate H0.
Definition is\_true\_inv : Prop \rightarrow bool.
move \Rightarrow P.
move: (is\_true\_inv0 \ P) \Rightarrow H.
apply constructive\_indefinite\_description in H.
apply H.
Defined.
Lemma is\_true\_id : \forall P : Prop, is\_true (is\_true\_inv P) \leftrightarrow P.
Proof.
move \Rightarrow P.
unfold is\_true\_inv.
move: (constructive\_indefinite\_description (fun b : bool \Rightarrow P \leftrightarrow is\_true b) (is\_true\_inv0)
(P)) \Rightarrow x\theta.
apply (@sig\_ind\ bool\ (fun\ b \Rightarrow (P \leftrightarrow is\_true\ b))\ (fun\ y \Rightarrow is\_true\ (let\ (x,\_) := y\ in\ x)
\leftrightarrow P)).
```

```
\mathtt{move} \Rightarrow x \ H.
apply iff_sym.
apply H.
Qed.
Lemma sval\_inv : \forall (A : Type)(P : A \rightarrow Prop)(x : sig P)(a : A), a = sval x \rightarrow \exists (H : P a),
x = exist P a H.
Proof.
move \Rightarrow A P x a H0.
rewrite H0.
\exists (proj2\_siq x).
apply (@sig\_ind \ A \ P \ (fun \ y \Rightarrow y = exist \ P \ (sval \ y) \ (proj2\_sig \ y))).
move \Rightarrow a\theta H.
by [simpl].
Qed.
Lemma sval\_injective : \forall (A : Type)(P : A \rightarrow Prop)(x \ y : sig \ P), sval \ x = sval \ y \rightarrow x = y.
Proof.
move \Rightarrow A P x y H.
move: (sval\_inv \ A \ P \ y \ (sval \ x) \ H).
elim \Rightarrow H0 \ H1.
rewrite H1.
assert (H0 = proj2\_siq x).
apply proof_irrelevance.
rewrite H2.
apply (@sig\_ind \ A \ P \ (fun \ y \Rightarrow y = exist \ P \ (sval \ y) \ (proj2\_sig \ y))).
move \Rightarrow a\theta H3.
by [simpl].
Qed.
Definition axiom 19 := \forall (A B : eqType)(alpha : Rel A B),
 \exists (R : eqType)(f : Rel \ R \ A)(g : Rel \ R \ B),
 function\_r\ f \land function\_r\ g \land alpha = f \# \bullet g \land ((f \bullet f \#) \quad (g \bullet g \#)) = Id\ R.
Lemma rationality: axiom19.
Proof.
move \Rightarrow A \ B \ alpha.
rewrite /function_r/total_r/univalent_r/cap/capP/identity/composite/inverse/include.
\exists (sig\_eqType (fun \ x : prod\_eqType \ A \ B \Rightarrow is\_true\_inv (alpha (fst \ x) (snd \ x)))).
\exists (fun x \ a \Rightarrow a = (fst \ (sval \ x))).
\exists (\mathbf{fun} \ x \ b \Rightarrow b = (snd \ (sval \ x))).
simpl.
repeat split.
move \Rightarrow x \ x\theta \ H.
\exists (fst (sval x)).
```

```
repeat split.
by [rewrite H].
move \Rightarrow a \ a\theta.
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
move \Rightarrow x \ x\theta \ H.
\exists (snd (sval x)).
repeat split.
by [rewrite H].
move \Rightarrow b \ b\theta.
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split; move \Rightarrow H.
assert (is\_true (is\_true\_inv (alpha (fst (a,b)) (snd (a,b))))).
simpl.
apply is\_true\_id.
apply H.
\exists (exist (fun \ x \Rightarrow (is\_true \ (is\_true\_inv \ (alpha \ (fst \ x) \ (snd \ x))))) (a,b) \ H0).
by [simpl].
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 H1.
apply is_true_id.
apply (@sig\_ind (A \times B) (fun x \Rightarrow is\_true (is\_true\_inv (alpha (fst x) (snd x)))) (fun x
\Rightarrow is_true (is_true_inv (alpha (fst (sval x)) (snd (sval x))))).
simpl.
by [move \Rightarrow x\theta].
apply functional_extensionality.
move \Rightarrow y.
apply functional_extensionality.
move \Rightarrow y\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
apply sval_injective.
move: (H \text{ (fun } a \ c : \{x : A \times B \mid is\_true \ (is\_true\_inv \ (alpha \ (fst \ x) \ (snd \ x)))\} \Rightarrow \exists \ b :
```

```
A, b = fst (sval \ a) \land b = fst (sval \ c)) (or_introl \ Logic.eq_refl)).
move: (H \text{ (fun } a \text{ } c \text{ : } \{x \text{ : } A \times B \mid is\_true \text{ (} is\_true\_inv \text{ (} alpha \text{ (} fst \text{ } x \text{) (} snd \text{ } x \text{))})\} \Rightarrow \exists b :
B, b = snd (sval \ a) \land b = snd (sval \ c)) (or\_intror \ Logic.eq\_reft)).
unfold id.
clear H.
elim \Rightarrow b.
elim \Rightarrow H H0.
elim \Rightarrow a.
elim \Rightarrow H1 H2.
rewrite (surjective_pairing (sval y0)) -H0 -H2 H H1.
apply surjective_pairing.
rewrite H.
move \Rightarrow beta H\theta.
case H0 \Rightarrow H1; rewrite H1; unfold id.
\exists (fst (sval y0)).
repeat split.
\exists (snd (sval y\theta)).
repeat split.
Qed.
```

2.3.6 直和と直積

```
任意の直和に対して、入射対が存在することを仮定する。
Lemma 20 (pair_of_inclusions) \exists j:A \to A+B, \exists k:B \to A+B, j\cdot j^\sharp = id_A \wedge k \cdot k^\sharp = id_B \wedge j \cdot k^\sharp = \phi_{AB} \wedge j^\sharp \cdot j \sqcup k^\sharp \cdot k = id_{A+B}.
```

```
Definition axiom20 :=  \forall (A \ B : eqType), \exists (j : Rel \ A \ (sum\_eqType \ A \ B))(k : Rel \ B \ (sum\_eqType \ A \ B)),  j \cdot j \# = Id \ A \wedge k \cdot k \# = Id \ B \wedge j \cdot k \# = A \ B \wedge (j \# \cdot j) \quad (k \# \cdot k) = Id \ (sum\_eqType \ A \ B).
Lemma pair\_of\_inclusions : axiom20.
Proof.
move \Rightarrow A \ B.
\exists \ (fun \ (a : A)(x : sum\_eqType \ A \ B) \Rightarrow x = inl \ a).
\exists \ (fun \ (b : B)(x : sum\_eqType \ A \ B) \Rightarrow x = inr \ b).
repeat split.
apply functional\_extensionality.
move \Rightarrow a.
apply functional\_extensionality.
move \Rightarrow a0.
```

```
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
by [injection H1].
\exists (inl a).
repeat split.
by [rewrite H].
apply functional_extensionality.
move \Rightarrow b.
apply functional_extensionality.
move \Rightarrow b\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
by [injection H1].
\exists (inr \ b).
repeat split.
by [rewrite H].
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
discriminate H1.
apply False_ind.
apply H.
apply functional_extensionality.
move \Rightarrow x.
apply functional_extensionality.
move \Rightarrow x\theta.
apply prop_extensionality_ok.
split.
elim \Rightarrow alpha.
elim \Rightarrow H0 \ H1.
```

```
case H0 \Rightarrow H2; rewrite H2 in H1.
elim H1 \Rightarrow a.
elim \Rightarrow H3 H4.
by [rewrite H3 H4].
elim H1 \Rightarrow b.
elim \Rightarrow H3 H4.
by [rewrite H3 H4].
assert ((\exists a : A, x = inl \ a) \lor (\exists b : B, x = inr \ b)).
\mathtt{move}: x.
apply sum_ind.
move \Rightarrow a.
left.
by [\exists a].
move \Rightarrow b.
right.
by [\exists b].
case H.
elim \Rightarrow a H0 H1.
\exists (fun x \ x\theta \Rightarrow \exists \ a\theta : A, (x = inl \ a\theta \land x\theta = inl \ a\theta)).
split.
by [left].
\exists a.
by [rewrite -H1 H0].
elim \Rightarrow b H0 H1.
\exists (\mathbf{fun} \ x \ x\theta \Rightarrow \exists \ b\theta : B, (x = inr \ b\theta \land x\theta = inr \ b\theta)).
split.
by [right].
\exists b.
by [rewrite -H1 H0].
Qed.
```

任意の直積に対して、射影対が存在することを仮定する.

実は有理性公理 ($Axiom\ 19$) があれば直積の公理は必要ないのだが、 $Axiom\ 19$ の準用では直積が "存在する" ことまでしか示してくれないので、"直積として $prod_eqType\ A\ B$ を用いてよい" ことを公理の中に含めたものを用意しておく.

Lemma 21 (pair_of_projections) $\exists p: A \times B \to A, \exists q: A \times B \to B,$

$$p^{\sharp} \cdot q = \nabla_{AB} \wedge p \cdot p^{\sharp} \sqcap q \cdot q^{\sharp} = id_{A \times B}.$$

```
Definition axiom21 :=
```

```
\forall (A \ B : eqType), \exists (p : Rel (prod_eqType \ A \ B) \ A)(q : Rel (prod_eqType \ A \ B) \ B), 
 <math>p \# \cdot q = A \ B \land (p \cdot p \#) \quad (q \cdot q \#) = Id (prod_eqType \ A \ B) \land univalent\_r \ p
```

```
\land univalent_r g.
Lemma pair_of_projections: axiom21.
Proof.
move \Rightarrow A B.
\exists (fun (x : prod\_eqType \ A \ B)(a : A) <math>\Rightarrow a = (fst \ x)).
\exists (\mathbf{fun} \ (x : prod\_eqType \ A \ B)(b : B) \Rightarrow b = (snd \ x)).
split.
apply functional\_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split; move \Rightarrow H.
apply I.
\exists (a,b).
by [simpl].
split.
apply functional_extensionality.
move \Rightarrow x.
apply functional_extensionality.
move \Rightarrow x\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
move: (H (fun a c: prod_eqType A B \Rightarrow \exists b: A, b = fst a \land b = fst c) (or_introl
Logic.eq\_refl).
move: (H \text{ (fun } a \ c : prod_eqType } A \ B \Rightarrow \exists \ b : B, \ b = snd \ a \land b = snd \ c) \text{ (or\_intror)}
Logic.eg\_refl).
unfold id.
clear H.
elim \Rightarrow b.
elim \Rightarrow H H0.
elim \Rightarrow a.
elim \Rightarrow H1 H2.
rewrite (surjective_pairing x0) -H0 -H2 H H1.
apply surjective_pairing.
rewrite H.
move \Rightarrow alpha H0.
case H0 \Rightarrow H1; rewrite H1; unfold id.
\exists (fst \ x\theta).
repeat split.
\exists (snd x0).
repeat split.
```

```
\begin{array}{l} \operatorname{split.} \\ \operatorname{move} \Rightarrow a \ a\theta. \\ \operatorname{elim} \Rightarrow x. \\ \operatorname{elim} \Rightarrow H \ H\theta. \\ \operatorname{by} [\operatorname{rewrite} H \ H\theta]. \\ \operatorname{move} \Rightarrow b \ b\theta. \\ \operatorname{elim} \Rightarrow x. \\ \operatorname{elim} \Rightarrow H \ H\theta. \\ \operatorname{by} [\operatorname{rewrite} H \ H\theta]. \\ \operatorname{Qed.} \\ \\ \operatorname{End} \ Rel\_Set. \\ \end{array}
```

Chapter 3

Library Basic_Lemmas

```
Require Import Basic\_Notations\_Set.
Require Import Logic.Classical\_Prop.
Module main (def : Relation).
Import def.
```

3.1 束論に関する補題

3.1.1 和関係, 共通関係

```
Lemma 22 (cap_l) Let \alpha, \beta : A \rightarrow B. Then,
                                                 \alpha \sqcap \beta \sqsubseteq \alpha.
Lemma cap_l {A B : eqType} {alpha beta : Rel A B}: (alpha
                                                                               beta)
                                                                                           alpha.
Proof.
assert ((alpha
                       beta)
                                (alpha
                                              beta)).
apply inc\_reft.
apply inc_-cap in H.
apply H.
Qed.
  Lemma 23 (cap_r) Let \alpha, \beta : A \rightarrow B. Then,
                                                 \alpha \sqcap \beta \sqsubseteq \beta.
Lemma cap_r \{A \ B : eqType\} \{alpha \ \mathbf{beta} : Rel \ A \ B\}: (alpha
                                                                                           beta.
Proof.
assert ((alpha
                       beta)
                                  (alpha
                                              beta)).
apply inc\_reft.
```

```
apply inc\_cap in H.
apply H.
Qed.
  Lemma 24 (cup_l) Let \alpha, \beta : A \rightarrow B. Then,
                                                \alpha \sqsubseteq \alpha \sqcup \beta.
Lemma cup_l {A B : eqType} {alpha beta : Rel A B}: alpha
                                                                             (alpha
                                                                                         beta).
Proof.
assert ((alpha
                      beta)
                                 (alpha
                                             beta)).
apply inc_refl.
apply inc_-cup in H.
apply H.
Qed.
  Lemma 25 (cup_r) Let \alpha, \beta : A \rightarrow B. Then,
                                                \beta \sqsubseteq \alpha \sqcup \beta.
Lemma cup_r \{A B : eqType\} \{alpha \text{ beta} : Rel A B\}: beta
                                                                            (alpha
                                                                                        beta).
Proof.
assert ((alpha
                      beta)
                                 (alpha
                                             beta)).
apply inc_refl.
apply inc\_cup in H.
apply H.
Qed.
  Lemma 26 (inc_def1) Let \alpha, \beta : A \rightarrow B. Then,
                                          \alpha = \alpha \sqcap \beta \Leftrightarrow \alpha \sqsubseteq \beta.
Lemma inc\_def1 {A B : eqType} {alpha beta : Rel A B}:
 alpha = alpha
                   \mathtt{beta} \leftrightarrow alpha
Proof.
split; move \Rightarrow H.
assert (alpha
                     (alpha
                                 beta)).
rewrite -H.
apply inc\_reft.
apply inc\_cap in H0.
apply H0.
apply inc\_antisym.
apply inc_-cap.
```

```
split.
apply inc\_reft.
apply H.
apply cap_{-}l.
Qed.
  Lemma 27 (inc_def2) Let \alpha, \beta : A \rightarrow B. Then,
                                         \beta = \alpha \sqcup \beta \Leftrightarrow \alpha \sqsubset \beta.
Lemma inc\_def2 {A B : eqType} {alpha beta : Rel A B}:
 beta = alpha
                   \mathtt{beta} \leftrightarrow alpha
Proof.
split; move \Rightarrow H.
assert ((alpha
                     beta)
                              beta).
rewrite -H.
apply inc\_reft.
apply inc_-cup in H0.
apply H0.
apply inc\_antisym.
                     beta) (alpha beta)).
assert ((alpha)
apply inc\_reft.
apply cup_r.
apply inc_-cup.
split.
apply H.
apply inc\_reft.
Qed.
  Lemma 28 (cap_assoc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                      (\alpha \sqcap \beta) \sqcap \gamma = \alpha \sqcap (\beta \sqcap \gamma).
Lemma cap\_assoc {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha
            beta)
                       gamma = alpha
                                              (beta
                                                         gamma).
Proof.
apply inc\_antisym.
rewrite inc\_cap.
split.
apply (inc_trans _ _ _ (alpha
                                     beta)).
apply cap_{-}l.
apply cap_{-}l.
rewrite inc\_cap.
```

```
split.
apply (inc_trans _ _ _ (alpha
                                    beta)).
apply cap_{-}l.
apply cap_r.
apply cap_{-}r.
rewrite inc\_cap.
split.
rewrite inc_-cap.
split.
apply cap_{-}l.
apply (inc_trans _ _ _ (beta
                                   qamma)).
apply cap_{-}r.
apply cap_l.
apply (inc_trans _ _ _ (beta
                                   qamma)).
apply cap_{-}r.
apply cap_{-}r.
Qed.
  Lemma 29 (cup_assoc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                  (\alpha \sqcup \beta) \sqcup \gamma = \alpha \sqcup (\beta \sqcup \gamma).
Lemma cup_assoc {A B : eqType} {alpha beta qamma : Rel A B}:
                     gamma = alpha
 (alpha
            beta)
                                          (beta
                                                     qamma).
Proof.
apply inc\_antisym.
rewrite inc\_cup.
split.
rewrite inc\_cup.
split.
apply cup_{-}l.
apply (inc_trans _ _ _ (beta
                                   gamma)).
apply cup_{-}l.
apply cup_r.
apply (inc_trans _ _ _ (beta
                                   gamma)).
apply cup_r.
apply cup_{-}r.
rewrite inc\_cup.
apply (inc_trans _ _ _ (alpha
                                  beta)).
apply cup_{-}l.
apply cup_{-}l.
rewrite inc_-cup.
```

```
split.
apply (inc_trans _ _ _ (alpha
                                     beta)).
apply cup_{-}r.
apply cup_{-}l.
apply cup_{-}r.
Qed.
  Lemma 30 (cap_comm) Let \alpha, \beta : A \rightarrow B. Then,
                                          \alpha \sqcap \beta = \beta \sqcap \alpha.
Lemma cap\_comm {A B : eqType} {alpha beta : Rel A B}: alpha
                                                                                               alpha.
                                                                            beta = beta
Proof.
apply inc\_antisym.
rewrite inc\_cap.
split.
apply cap_{-}r.
apply cap_{-}l.
rewrite inc\_cap.
split.
apply cap_r.
apply cap_{-}l.
Qed.
  Lemma 31 (cup_comm) Let \alpha, \beta : A \rightarrow B. Then,
                                          \alpha \sqcup \beta = \beta \sqcup \alpha.
Lemma cup\_comm {A B : eqType} {alpha beta : Rel A B}: alpha beta = beta
                                                                                               alpha.
Proof.
apply inc\_antisym.
rewrite inc\_cup.
split.
apply cup_{-}r.
apply cup_{-}l.
rewrite inc\_cup.
split.
apply cup_r.
apply cup_{-}l.
Qed.
```

```
Lemma 32 (cup_cap_abs) Let \alpha, \beta : A \rightarrow B. Then,
                                           \alpha \sqcup (\alpha \sqcap \beta) = \alpha.
Lemma cup\_cap\_abs {A B : eqType} {alpha beta : Rel A B}:
           (alpha
                      beta) = alpha.
Proof.
move: (@cap_l - alpha beta) \Rightarrow H.
apply inc\_def2 in H.
by [rewrite cup\_comm - H].
Qed.
  Lemma 33 (cap_cup_abs) Let \alpha, \beta : A \rightarrow B. Then,
                                           \alpha \sqcap (\alpha \sqcup \beta) = \alpha.
Lemma cap\_cup\_abs {A B : eqType} {alpha beta : Rel A B}:
 alpha
            (alpha
                       beta) = alpha.
Proof.
move: (@cup_l - alpha beta) \Rightarrow H.
apply inc\_def1 in H.
by [rewrite -H].
Qed.
  Lemma 34 (cap_idem) Let \alpha : A \rightarrow B. Then,
                                              \alpha \sqcap \alpha = \alpha.
Lemma cap\_idem \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha = alpha.
apply inc\_antisym.
apply cap_{-}l.
apply inc_-cap.
split; apply inc\_refl.
Qed.
  Lemma 35 (cup_idem) Let \alpha : A \rightarrow B. Then,
                                              \alpha \sqcup \alpha = \alpha.
Lemma cup\_idem \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha = alpha.
Proof.
apply inc\_antisym.
```

```
apply inc_-cup.
split; apply inc\_refl.
apply cup_{-}l.
Qed.
  Lemma 36 (cap_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                                    \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha' \sqcap \beta'.
Lemma cap_inc_compat {A B : eqType} {alpha alpha' beta beta' : Rel A B}:
             alpha' \rightarrow \mathtt{beta} \qquad beta' \rightarrow (alpha)
                                                             beta)
                                                                          (alpha'
                                                                                         beta').
Proof.
move \Rightarrow H H0.
rewrite -inc_{-}def1.
apply inc\_def1 in H.
apply inc\_def1 in H0.
rewrite cap\_assoc -(@cap\_assoc _ _ beta).
rewrite (@cap\_comm\_\_beta).
rewrite cap\_assoc -(@cap\_assoc _ _ alpha).
by [rewrite -H -H\theta].
Qed.
  Lemma 37 (cap_inc_compat_l) Let \alpha, \beta, \beta' : A \rightarrow B. Then,
                                           \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha \sqcap \beta'.
Lemma cap\_inc\_compat\_l {A B : eqType} {alpha beta beta' : Rel A B}:
            beta' \rightarrow (alpha \quad beta) \quad (alpha)
 beta
                                                              beta').
Proof.
move \Rightarrow H.
apply (@cap_inc_compat _ _ _ _ (@inc_reft _ _ alpha) H).
Qed.
  Lemma 38 (cap_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                           \alpha \sqsubseteq \alpha' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha' \sqcap \beta.
Lemma cap\_inc\_compat\_r {A B : eqType} {alpha \ alpha' \ beta : Rel \ A \ B}:
 alpha
             alpha' \rightarrow (alpha \quad beta)
                                                 (alpha')
                                                                  beta).
Proof.
move \Rightarrow H.
apply (@cap\_inc\_compat\_\_\_\_\_H (@inc\_refl\_\_beta)).
Qed.
```

```
Lemma 39 (cup_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                                     \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha' \sqcup \beta'.
Lemma cup\_inc\_compat {A \ B : eqType} {alpha \ alpha' \ beta \ beta' : Rel \ A \ B}:
             alpha' \rightarrow beta \qquad beta' \rightarrow (alpha \qquad beta)
                                                                           (alpha'
                                                                                          beta').
Proof.
move \Rightarrow H H0.
rewrite -inc_{-}def2.
apply inc\_def2 in H.
apply inc\_def2 in H0.
rewrite cup\_assoc -(@cup\_assoc _ _ beta).
rewrite (@cup_comm _ _ beta).
rewrite cup\_assoc -(@cup\_assoc _ _ alpha).
by [rewrite -H - H\theta].
Qed.
  Lemma 40 (cup_inc_compat_l) Let \alpha, \beta, \beta' : A \rightarrow B. Then,
                                           \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha \sqcup \beta'.
Lemma cup\_inc\_compat\_l {A B : eqType} {alpha beta beta' : Rel A B}:
             beta' \rightarrow (alpha \quad beta) \quad (alpha)
                                                               beta').
 beta
Proof.
move \Rightarrow H.
apply (@cup_inc_compat _ _ _ _ (@inc_reft _ _ alpha) H).
Qed.
  Lemma 41 (cup_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                           \alpha \sqsubseteq \alpha' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha' \sqcup \beta.
Lemma cup\_inc\_compat\_r {A \ B : eqType} {alpha \ alpha' \ beta : Rel \ A \ B}:
 alpha
              alpha' \rightarrow (alpha \quad beta)
                                                  (alpha')
                                                                   beta).
Proof.
move \Rightarrow H.
apply (@cup\_inc\_compat\_\_\_\_\_H (@inc\_refl\_\_beta)).
Qed.
  Lemma 42 (cap_empty) Let \alpha : A \rightarrow B. Then,
                                                  \alpha \sqcap \phi_{AB} = \phi_{AB}.
```

```
A B =
Lemma cap\_empty \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha
                                                                               A B.
Proof.
apply inc\_antisym.
apply cap_r.
apply inc\_empty\_alpha.
Qed.
```

Lemma 43 (cup_empty) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcup \phi_{AB} = \alpha$$
.

```
Lemma cup\_empty {A B : eqType} {alpha : Rel A B}: alpha
                                                                  A B = alpha.
apply inc\_antisym.
apply inc_-cup.
split.
apply inc\_reft.
apply inc\_empty\_alpha.
apply cup_{-}l.
Qed.
```

Lemma 44 (cap_universal) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcap \nabla_{AB} = \alpha$$
.

```
Lemma cap\_universal \{A B : eqType\} \{alpha : Rel A B\}: alpha
                                                                      A B = alpha.
Proof.
apply inc\_antisym.
apply cap_-l.
apply inc\_cap.
split.
apply inc_refl.
apply inc\_alpha\_universal.
Qed.
```

Lemma 45 (cup_universal) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcup \nabla_{AB} = \nabla_{AB}$$
.

Lemma $cup_universal\ \{A\ B: eqType\}\ \{alpha: Rel\ A\ B\}: alpha$ A B =A B. apply $inc_antisym$. apply inc_-cup .

```
split.
apply inc\_alpha\_universal.
apply inc\_reft.
apply cup_r.
Qed.
  Lemma 46 (inc_lower) Let \alpha, \beta : A \rightarrow B. Then,
                                   \alpha = \beta \Leftrightarrow (\forall \gamma : A \to B, \gamma \sqsubseteq \alpha \Leftrightarrow \gamma \sqsubseteq \beta).
Lemma inc\_lower {A \ B : eqType} {alpha \ beta : Rel \ A \ B}:
  alpha = beta \leftrightarrow (\forall \ gamma : Rel \ A \ B, \ gamma
                                                                       alpha \leftrightarrow gamma
                                                                                                    beta).
Proof.
split; move \Rightarrow H.
move \Rightarrow gamma.
by [rewrite H].
apply inc\_antisym.
rewrite -H.
apply inc\_reft.
rewrite H.
apply inc\_reft.
Qed.
  Lemma 47 (inc_upper) Let \alpha, \beta : A \rightarrow B. Then,
                                   \alpha = \beta \Leftrightarrow (\forall \gamma : A \to B, \alpha \sqsubseteq \gamma \Leftrightarrow \beta \sqsubseteq \gamma).
Lemma inc\_upper \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 alpha = beta \leftrightarrow (\forall \ gamma : Rel \ A \ B, \ alpha \qquad gamma \leftrightarrow beta
                                                                                                gamma).
Proof.
split; move \Rightarrow H.
move \Rightarrow gamma.
by [rewrite H].
apply inc\_antisym.
rewrite H.
apply inc_refl.
rewrite -H.
apply inc_refl.
Qed.
```

3.1.2 分配法則

```
Lemma 48 (cap_cup_distr_l) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                   \alpha \sqcap (\beta \sqcup \gamma) = (\alpha \sqcap \beta) \sqcup (\alpha \sqcap \gamma).
Lemma cap\_cup\_distr\_l {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
            (beta
                       gamma) = (alpha
                                                 beta)
 alpha
                                                            (alpha
                                                                        gamma).
Proof.
apply inc\_upper.
move \Rightarrow delta.
split; move \Rightarrow H.
rewrite cap\_comm (@cap\_comm _ _ _ gamma).
apply inc\_cup.
rewrite -inc\_rpc -inc\_rpc.
apply inc\_cup.
rewrite inc_rpc cap_comm.
apply H.
rewrite cap\_comm -inc\_rpc.
apply inc_-cup.
rewrite inc\_rpc inc\_rpc.
apply inc_-cup.
rewrite cap_comm (@cap_comm _ _ gamma).
apply H.
Qed.
  Lemma 49 (cap_cup_distr_r) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                   (\alpha \sqcup \beta) \sqcap \gamma = (\alpha \sqcap \gamma) \sqcup (\beta \sqcap \gamma).
Lemma cap\_cup\_distr\_r {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
                        qamma = (alpha
 (alpha
             beta)
                                                qamma)
                                                               (beta
                                                                           qamma).
Proof.
rewrite (@cap\_comm\_\_ (alpha beta)) (@cap\_comm\_\_ alpha) (@cap\_comm\_\_ beta).
apply cap\_cup\_distr\_l.
Qed.
  Lemma 50 (cup_cap_distr_l) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                   \alpha \sqcup (\beta \sqcap \gamma) = (\alpha \sqcup \beta) \sqcap (\alpha \sqcup \gamma).
Lemma cup\_cap\_distr\_l {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
                       gamma) = (alpha
                                                 beta)
                                                            (alpha
 alpha
            (beta
                                                                        gamma).
```

Proof.

rewrite $cap_cup_distr_l$.

rewrite $(@cap_comm___(alpha \ beta))$ cap_cup_abs $(@cap_comm___(alpha \ beta))$.

rewrite $cap_cup_distr_l$.

rewrite $-cup_assoc$ $(@cap_comm___gamma)$ cup_cap_abs .

by [rewrite cap_comm].

Qed.

Lemma 51 (cup_cap_distr_r) Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$(\alpha \sqcap \beta) \sqcup \gamma = (\alpha \sqcup \gamma) \sqcap (\beta \sqcup \gamma).$$

Lemma $cup_cap_distr_r$ { $A \ B : eqType$ } { $alpha \ beta \ gamma : Rel \ A \ B$ }: ($alpha \ beta$) $gamma = (alpha \ gamma)$ (beta gamma). Proof. rewrite ($@cup_comm__(alpha \ beta)$) ($@cup_comm__alpha$) ($@cup_comm__beta$). apply $cup_cap_distr_l$. Qed.

Lemma 52 (cap_cup_unique) Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$\alpha \sqcap \beta = \alpha \sqcap \gamma \land \alpha \sqcup \beta = \alpha \sqcup \gamma \Rightarrow \beta = \gamma.$$

 $gamma \rightarrow beta = gamma.$

Lemma cap_cup_unique {AB: eqType} {alpha beta $gamma: Rel\ A\ B$ }: alpha beta = alpha $gamma \rightarrow alpha$ beta = alpha gamma. Proof.

move $\Rightarrow H\ H0$.

rewrite $-(@cap_cup_abs___$ beta alpha) $cup_comm\ H0$.

rewrite $cap_cup_distr_l$.

rewrite $cap_cup_distr_r$.

rewrite $H0\ cap_comm\ cup_comm$.

apply cap_cup_abs .

Qed.

3.1.3 原子性

空関係でない $\alpha: A \rightarrow B$ が、任意の $\beta: A \rightarrow B$ について

$$\beta \sqsubseteq \alpha \Rightarrow \beta = \phi_{AB} \lor \beta = \alpha$$

を満たすとき, α は原子的 (atomic) であると言われる.

```
Definition atomic \{A \ B : eqType\} (alpha : Rel \ A \ B) :=
               A B \wedge (\forall beta : Rel A B, beta)
 alpha \neq
                                                           alpha \rightarrow \mathtt{beta} =
                                                                                   A B \vee beta = alpha).
  Lemma 53 (atomic_cap_empty) Let \alpha, \beta : A \to B are atomic and \alpha \neq \beta. Then,
                                               \alpha \sqcap \beta = \phi_{AB}.
Lemma atomic\_cap\_empty {A B : eqType} {alpha beta : Rel A B}:
 atomic\ alpha 	o atomic\ beta 	o alpha 
eq beta 	o alpha
                                                                                       A B.
Proof.
move \Rightarrow H H0.
apply or_{-}to_{-}imply.
case (classic (alpha
                                          (A B)); move \Rightarrow H1.
                            beta =
right.
apply H1.
left.
move \Rightarrow H2.
apply H2.
apply inc\_antisym.
apply inc\_def1.
elim H \Rightarrow H3 H4.
case (H4 (alpha
                        beta) (cap_l); move \Rightarrow H5.
apply False_ind.
apply (H1 \ H5).
by [rewrite H5].
apply inc_-def1.
elim H0 \Rightarrow H3 H4.
case (H4 (alpha
                        beta) (cap_r); move \Rightarrow H5.
apply False_ind.
apply (H1 \ H5).
by [rewrite cap\_comm\ H5].
Qed.
  Lemma 54 (atomic_cup) Let \alpha, \beta, \gamma : A \rightarrow B and \alpha is atomic. Then,
                                     \alpha \sqsubseteq \beta \sqcup \gamma \Rightarrow \alpha \sqsubseteq \beta \lor \alpha \sqsubseteq \gamma.
Lemma atomic\_cup \{A \ B : eqType\} \{alpha \ beta \ gamma : Rel \ A \ B\}:
                                           qamma) \rightarrow alpha
 atomic \ alpha \rightarrow alpha
                                (beta
                                                                     beta \vee alpha
                                                                                          gamma.
Proof.
move \Rightarrow H H0.
apply inc\_def1 in H0.
rewrite cap\_cup\_distr\_l in H0.
```

```
elim H \Rightarrow H1 H2.
rewrite H0 in H1.
assert (alpha
                   beta \neq
                              A B \vee alpha
                                                gamma \neq
                                                              A B).
apply not\_and\_or.
elim \Rightarrow H3 H4.
rewrite H3 H4 in H1.
apply H1.
by [rewrite cup_-empty].
case H3; move \Rightarrow H4.
left.
apply inc\_def1.
case (H2 (alpha
                     beta) (cap_{-}l); move \Rightarrow H5.
apply False_ind.
apply (H4 H5).
by |rewrite H5|.
right.
apply inc\_def1.
case (H2 (alpha
                     gamma) (cap_{-}l); move \Rightarrow H5.
apply False_ind.
apply (H4 H5).
by [rewrite H5].
Qed.
```

3.2 Heyting 代数に関する補題

```
Lemma 55 (rpc_universal) Let \alpha:A\to B. Then, (\alpha\Rightarrow\alpha)=\nabla_{AB}. Lemma rpc\_universal \{A\ B: eqType\} \{alpha: Rel\ A\ B\}: (alpha \gg alpha)=A\ B. Proof. apply inc\_lower. move \Rightarrow gamma. split; move \Rightarrow H. apply inc\_alpha\_universal. apply inc\_alpha\_universal. apply inc\_rpc. apply cap\_r. Qed.
```

```
Lemma 56 (rpc_r) Let \alpha, \beta : A \rightarrow B. Then,
                                            (\alpha \Rightarrow \beta) \sqcap \beta = \beta.
Lemma rpc_r \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: (alpha \gg beta)
                                                                                        beta = beta.
Proof.
                    (alpha \gg beta)).
assert (beta
apply inc\_rpc.
apply cap_{-}l.
apply inc\_def1 in H.
by [rewrite cap\_comm - H].
Qed.
  Lemma 57 (inc_def3) Let \alpha, \beta : A \rightarrow B. Then,
                                       (\alpha \Rightarrow \beta) = \nabla_{AB} \Leftrightarrow \alpha \sqsubseteq \beta.
Lemma inc\_def3 {A B : eqType} {alpha beta : Rel A B}:
 (alpha \gg beta) =
                       A B \leftrightarrow alpha
                                               beta.
Proof.
split; move \Rightarrow H.
rewrite -(@rpc_universal _ _ alpha) in H.
assert((alpha * alpha)
                               (alpha \gg beta)).
rewrite H.
apply inc\_reft.
apply inc\_rpc in H0.
rewrite rpc_{-}r in H0.
apply H0.
apply inc\_antisym.
apply inc\_alpha\_universal.
rewrite -(@rpc_universal _ _ alpha).
apply inc_-rpc.
rewrite rpc_{-}r.
apply H.
Qed.
  Lemma 58 (rpc_l) Let \alpha, \beta : A \rightarrow B. Then,
                                         \alpha \sqcap (\alpha \Rightarrow \beta) = \alpha \sqcap \beta.
Lemma rpc_l {A B : eqType} {alpha beta : Rel A B}:
            (alpha \gg beta) = alpha
 alpha
Proof.
```

```
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_cap.
apply inc\_cap in H.
split.
apply H.
elim H \Rightarrow H0 \ H1.
apply inc\_rpc in H1.
rewrite -(@cap\_idem \_ \_ gamma).
apply (inc_trans _ _ _ (gamma
                                          alpha)).
apply cap\_inc\_compat.
apply inc\_reft.
apply H0.
apply H1.
apply inc_-cap.
apply inc\_cap in H.
split.
apply H.
apply inc\_rpc.
apply (inc\_trans \_ \_ \_ gamma).
apply cap_{-}l.
apply H.
Qed.
  Lemma 59 (rpc_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                              \alpha' \sqsubseteq \alpha \land \beta \sqsubseteq \beta' \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha' \Rightarrow \beta').
Lemma rpc\_inc\_compat {A B : eqType} {alpha \ alpha' \ beta \ beta' : Rel \ A \ B}:
             alpha \rightarrow beta beta' \rightarrow (alpha \gg beta) (alpha' \gg beta').
 alpha'
Proof.
move \Rightarrow H H0.
apply inc\_rpc.
apply (@inc_trans _ _ _ ((alpha » beta)
                                                     alpha)).
apply (@cap\_inc\_compat\_l\_\_\_\_\_H).
rewrite cap\_comm\ rpc\_l.
apply @inc\_trans \_ \_ \_ beta.
apply cap_r.
apply H0.
Qed.
```

Lemma 60 (rpc_inc_compat_l) Let $\alpha, \beta, \beta' : A \rightarrow B$. Then,

```
\beta \sqsubseteq \beta' \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha \Rightarrow \beta').
Lemma rpc\_inc\_compat\_l {A B : eqType} {alpha beta beta' : Rel A B}:
             beta' \rightarrow (alpha \gg beta)
                                                 (alpha \gg beta').
Proof.
move \Rightarrow H.
apply (@rpc\_inc\_compat\_\_\_\_\_ (@inc\_refl\_\_alpha) H).
Qed.
  Lemma 61 (rpc_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                         \alpha' \sqsubseteq \alpha \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha' \Rightarrow \beta).
Lemma rpc\_inc\_compat\_r {A B : eqType} {alpha \ alpha' \ beta : Rel \ A \ B}:
               alpha \rightarrow (alpha \gg beta)
                                                    (alpha' \gg beta).
Proof.
move \Rightarrow H.
apply (@rpc_inc_compat _ _ _ _ H (@inc_reft _ _ beta)).
Qed.
```

Lemma 62 (rpc_universal_alpha) Let $\alpha : A \rightarrow B$. Then,

 $\nabla_{AB} \Rightarrow \alpha = \alpha$.

Lemma $rpc_universal_alpha$ { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }: $A \ B \gg alpha = alpha$. Proof. apply inc_lower . move $\Rightarrow gamma$. split; move $\Rightarrow H$.

apply inc_rpc in H.

rewrite $cap_universal$ in H.

apply H.

apply inc_-rpc .

rewrite cap_universal.

apply H.

Qed.

Lemma 63 (rpc_lemma1) Let $\alpha, \beta, \gamma : A \rightarrow B$. Then,

$$(\alpha \Rightarrow \beta) \sqsubseteq ((\alpha \sqcap \gamma) \Rightarrow (\beta \sqcap \gamma)).$$

```
Lemma rpc\_lemma1 {A B : eqType} {alpha beta gamma : Rel A B}:
                                     gamma) » (beta
 (alpha \gg beta)
                        ((alpha
                                                               qamma)).
Proof.
apply inc\_rpc.
rewrite -cap_assoc (@cap_comm _ _ alpha).
rewrite rpc_{-}l.
apply cap\_inc\_compat\_r.
apply cap_{-}r.
Qed.
  Lemma 64 (rpc_lemma2) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                 (\alpha \Rightarrow \beta) \sqcap (\alpha \Rightarrow \gamma) = (\alpha \Rightarrow (\beta \sqcap \gamma)).
Lemma rpc\_lemma2 {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha \gg beta)
                      (alpha \gg gamma) = alpha \gg (beta)
                                                                        qamma).
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
rewrite inc\_rpc.
apply inc\_cap in H.
apply inc\_cap.
rewrite -inc\_rpc -inc\_rpc.
apply H.
apply inc_-cap.
rewrite inc_rpc inc_rpc.
apply inc_{-}cap.
rewrite -inc\_rpc.
apply H.
Qed.
  Lemma 65 (rpc_lemma3) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                              (\alpha \Rightarrow \beta) \sqcap (\beta \Rightarrow \gamma) \sqsubseteq ((\alpha \sqcup \beta) \Rightarrow (\beta \sqcap \gamma)).
Lemma rpc\_lemma3 {A B : eqType} {alpha beta gamma : Rel A B}:
 ((alpha \gg beta)
                         (beta » gamma))
                                                    ((alpha beta) * (beta)
                                                                                        qamma)).
Proof.
apply inc\_rpc.
rewrite cap\_cup\_distr\_l.
rewrite cap_comm -cap_assoc rpc_l.
\texttt{rewrite} \ (@\mathit{cap\_assoc} \ \_ \ \_ \ \_ \ \_ \ \texttt{beta}) \ (@\mathit{cap\_comm} \ \_ \ \_ \ (\texttt{beta} \ \ \ \mathit{gamma})) \ - \mathit{cap\_assoc} \ \mathit{rpc\_r}.
```

```
rewrite cap_assoc rpc_l.
apply inc\_cup.
split.
apply cap_r.
apply inc\_reft.
Qed.
  Lemma 66 (rpc_lemma4) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                   (\alpha \Rightarrow \beta) \sqcap (\beta \Rightarrow \gamma) \sqsubseteq (\alpha \Rightarrow \gamma).
Lemma rpc\_lemma4 {A B : eqType} {alpha beta gamma : Rel A B}:
                     (beta \gg gamma))
 ((alpha \gg beta)
                                                (alpha \gg gamma).
Proof.
apply (@inc_trans _ _ _ ((alpha beta) » (beta
                                                               qamma))).
apply rpc\_lemma3.
apply rpc\_inc\_compat.
apply cup_{-}l.
apply cap_{-}r.
Qed.
  Lemma 67 (rpc_lemma5) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                    \alpha \Rightarrow (\beta \Rightarrow \gamma) = (\alpha \sqcap \beta) \Rightarrow \gamma.
Lemma rpc\_lemma5 {A B : eqType} {alpha beta gamma : Rel A B}:
 alpha » (beta » qamma) = (alpha
                                            beta) » gamma.
Proof.
apply inc_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_rpc.
rewrite -cap\_assoc.
rewrite -inc\_rpc -inc\_rpc.
apply H.
rewrite inc\_rpc inc\_rpc.
rewrite cap\_assoc.
apply inc\_rpc.
apply H.
Qed.
```

```
Lemma 68 (rpc_lemma6) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                  \alpha \Rightarrow (\beta \Rightarrow \gamma) \sqsubset (\alpha \Rightarrow \beta) \Rightarrow (\alpha \Rightarrow \gamma).
Lemma rpc\_lemma6 {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha » (beta » gamma))
                                       ((alpha \gg beta) \gg (alpha \gg gamma)).
Proof.
rewrite inc\_rpc inc\_rpc.
rewrite cap\_assoc (@cap\_comm\_\_\_alpha).
rewrite rpc_{-}l.
rewrite - cap_assoc (@cap_comm _ _ alpha).
rewrite rpc_{-}l.
rewrite cap\_assoc (@cap\_comm\_\_\__beta).
rewrite rpc_{-}l.
rewrite -cap\_assoc.
apply cap_r.
Qed.
  Lemma 69 (rpc_lemma7) Let \alpha, \beta, \gamma, \delta : A \rightarrow B and \beta \sqsubseteq \alpha \sqsubseteq \gamma. Then,
              (\alpha \sqcap \delta = \beta) \land (\alpha \sqcup \delta = \gamma) \Leftrightarrow (\gamma \sqsubset \alpha \sqcup (\alpha \Rightarrow \beta)) \land (\delta = \gamma \sqcap (\alpha \Rightarrow \beta)).
Lemma rpc\_lemma { A B : eqType } { alpha \ beta \ gamma \ delta : Rel \ A \ B }:
beta
          alpha \rightarrow alpha
                                 qamma \rightarrow (alpha)
                                                            delta = beta \land alpha
                                                                                              delta = qamma
                   (alpha
                                (alpha \gg beta)) \land delta = qamma
                                                                               (alpha \gg beta)).
 \leftrightarrow qamma
Proof.
move \Rightarrow H H0.
split; elim; move \Rightarrow H1 H2; split.
rewrite -H2.
apply cup\_inc\_compat\_l.
apply inc\_rpc.
rewrite cap\_comm\ H1.
apply inc\_reft.
rewrite -H2.
rewrite cap\_cup\_distr\_r\ rpc\_l.
assert (delta
                       (alpha \gg beta).
apply inc\_rpc.
rewrite cap\_comm\ H1.
apply inc\_reft.
apply inc\_def1 in H3.
rewrite -H3 -H1.
rewrite - cap_assoc cap_idem.
```

```
by [rewrite cap_comm cup_comm cup_cap_abs].
rewrite H2.
rewrite (@cap_comm _ _ gamma) -cap_assoc rpc_l.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ (beta gamma)).
apply cap\_inc\_compat\_r.
apply cap_r.
apply cap_{-}l.
move: (@inc\_trans \_ \_ \_ \_ H H0) \Rightarrow H3.
apply inc\_def1 in H.
apply inc\_def1 in H3.
rewrite cap\_comm in H.
rewrite -H -H3.
apply inc\_reft.
rewrite H2.
rewrite cup\_cap\_distr\_l.
apply inc\_def2 in H0.
rewrite -H0.
apply inc\_def1 in H1.
by [rewrite -H1].
Qed.
```

3.3 補関係に関する補題

Lemma 70 (complement_universal)

$$\nabla_{AB}^{-} = \phi_{AB}$$
.

Lemma 71 (complement_alpha_universal) Let $\alpha : A \rightarrow B$. Then,

$$\alpha^- = \nabla_{AB} \Leftrightarrow \alpha = \phi_{AB}.$$

Lemma complement_alpha_universal $\{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha ^= A \ B \leftrightarrow alpha = A \ B.$ Proof.
split; move $\Rightarrow H$.

apply $inc_antisym$.

Qed.

```
rewrite -(@cap_universal _ _ alpha) cap_comm.

apply inc_rpc.

rewrite -H.

apply inc_refl.

apply inc_empty_alpha.

apply inc_antisym.

apply inc_alpha_universal.

apply inc_rpc.

rewrite cap_comm cap_universal.

rewrite H.

apply inc_refl.

Qed.
```

Lemma 72 (complement_empty)

$$\phi_{AB}^{-} = \nabla_{AB}$$
.

by [apply complement_alpha_universal]. Qed.

Lemma 73 (complement_invol_inc) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqsubseteq (\alpha^-)^-$$
.

Lemma $complement_invol_inc$ { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }: alpha (alpha ^) ^. Proof.

apply inc_rpc .

rewrite cap_comm .

apply inc_rpc .

apply inc_reft .

Qed.

Lemma 74 (cap_complement_empty) Let $\alpha : A \rightarrow B$. Then,

$$\alpha \sqcap \alpha^- = \phi_{AB}$$
.

Proof.

apply $inc_antisym$.

rewrite cap_comm .

```
apply inc_rpc.
apply inc_refl.
apply inc_empty_alpha.
Qed.
```

Lemma 75 (complement_invol) Let $\alpha : A \rightarrow B$. Then,

$$(\alpha^-)^- = \alpha.$$

```
Lemma complement_invol \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: (alpha ^) ^= alpha. Proof.

rewrite -(@cap_universal _ _ ((alpha ^) ^)).

rewrite -(@complement_classic _ _ alpha).

rewrite cap_cup_distr_l.

rewrite (@cap_comm _ _ _ (alpha ^)) cap_complement_empty.

rewrite cup_empty cap_comm.

apply Logic.eq_sym.

apply inc_def1.

apply complement_invol_inc.

Qed.
```

Lemma 76 (complement_move) Let $\alpha, \beta : A \rightarrow B$. Then,

$$\alpha = \beta^- \Leftrightarrow \alpha^- = \beta$$
.

```
Lemma complement\_move \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: alpha = beta ^ \leftrightarrow alpha ^ = beta.

Proof.

split; move \Rightarrow H.

by [rewrite H \ complement\_invol].

by [rewrite -H \ complement\_invol].

Qed.
```

Lemma 77 (contraposition) Let $\alpha, \beta : A \rightarrow B$. Then,

$$(\alpha \Rightarrow \beta) = (\beta^- \Rightarrow \alpha^-).$$

```
replace (alpha » beta) with ((alpha ^) ^ » (beta ^) ^).
apply inc_rpc.
apply rpc_lemma4.
by [rewrite complement_invol complement_invol].
Qed.
```

```
Lemma 78 (de_morgan1) Let \alpha, \beta : A \to B. Then, (\alpha \sqcup \beta)^- = \alpha^- \sqcap \beta^-.
```

```
Lemma de\_morgan1 {A B : eqType} {alpha beta : Rel A B}:
            beta) \hat{} = alpha \hat{}
 (alpha
                                 beta ^.
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_cap.
rewrite inc\_rpc inc\_rpc.
apply inc_-cup.
rewrite -cap\_cup\_distr\_l.
apply inc\_rpc.
apply H.
apply inc\_rpc.
rewrite cap_-cup_-distr_-l.
apply inc\_cup.
rewrite -inc\_rpc -inc\_rpc.
apply inc_-cap.
apply H.
Qed.
```

```
Lemma 79 (de_morgan2) Let \alpha, \beta : A \rightarrow B. Then,
```

$$(\alpha \sqcap \beta)^- = \alpha^- \sqcup \beta^-.$$

```
Lemma de\_morgan2 {A B : eqType} {alpha beta : Rel A B}: (alpha beta) \hat{} = alpha beta \hat{}. Proof.
```

by [rewrite -complement_move de_morgan1 complement_invol complement_invol]. Qed.

```
Lemma 80 (cup_to_rpc) Let \alpha, \beta : A \rightarrow B. Then,
                                         \alpha^- \sqcup \beta = (\alpha \Rightarrow \beta).
Lemma cup\_to\_rpc {A B : eqType} {alpha beta : Rel A B}:
             beta = alpha \gg beta.
Proof.
apply inc\_antisym.
apply inc_rpc.
rewrite cap\_cup\_distr\_r cap\_comm.
rewrite cap_complement_empty cup_comm cup_empty.
apply cap_{-}l.
rewrite -(@cap_universal _ _ (alpha » beta)) cap_comm.
rewrite -(@complement_classic _ alpha).
rewrite cap\_cup\_distr\_r\ cup\_comm.
apply cup\_inc\_compat.
apply cap_{-}l.
rewrite rpc_-l.
apply cap_{-}r.
Qed.
  Lemma 81 (beta_contradiction) Let \alpha, \beta : A \rightarrow B. Then,
                                    (\alpha \Rightarrow \beta) \sqcap (\alpha \Rightarrow \beta^{-}) = \alpha^{-}.
Lemma beta\_contradiction \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 (alpha \gg beta)
                      (alpha \gg beta \hat{}) = alpha \hat{}.
Proof.
rewrite -cup_to_rpc -cup_to_rpc.
rewrite -cup\_cap\_distr\_l.
by [rewrite cap_complement_empty cup_empty].
Qed.
```

3.4 Bool 代数に関する補題

```
Lemma 82 (bool_lemma1) Let \alpha, \beta : A \to B. Then, \alpha \sqsubseteq \beta \Leftrightarrow \nabla_{AB} = \alpha^- \sqcup \beta.
```

```
Lemma bool\_lemma1 {A B : eqType} {alpha beta : Rel A B}: alpha beta \leftrightarrow A B = alpha beta.
```

```
Proof.
split; move \Rightarrow H.
apply inc\_antisym.
rewrite -(@complement_classic _ _ alpha) cup_comm.
apply cup\_inc\_compat\_l.
apply H.
apply inc\_alpha\_universal.
apply inc\_def3.
rewrite H.
apply (Logic.eq_sym cup_to_rpc).
Qed.
  Lemma 83 (bool_lemma2) Let \alpha, \beta : A \rightarrow B. Then,
                                     \alpha \sqsubseteq \beta \Leftrightarrow \alpha \sqcap \beta^- = \phi_{AB}.
Lemma bool_lemma2 {A B : eqType} {alpha beta : Rel A B}:
           beta \leftrightarrow alpha
                             beta ^ =
 alpha
Proof.
split; move \Rightarrow H.
rewrite -(@cap_universal _ _ (alpha beta ^)).
apply bool\_lemma1 in H.
rewrite H.
rewrite cap\_cup\_distr\_l.
rewrite (@cap_comm _ _ alpha) cap_assoc cap_complement_empty cap_empty.
rewrite cap_comm -cap_assoc cap_complement_empty cap_comm cap_empty.
by [rewrite cup_-empty].
rewrite -(@cap_universal _ _ alpha).
rewrite -(@complement_classic _ _ beta).
rewrite cap\_cup\_distr\_l.
rewrite H cup\_empty.
apply cap_r.
Qed.
  Lemma 84 (bool_lemma3) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                    \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \alpha \sqcap \beta^- \sqsubseteq \gamma.
Lemma bool_lemma3 {A B : eqType} {alpha beta gamma : Rel A B}:
                     gamma) \leftrightarrow (alpha)
 alpha
           (beta
                                            beta ^)
Proof.
split; move \Rightarrow H.
apply (@inc_trans _ _ _ ((beta gamma) beta ^)).
```

```
apply cap\_inc\_compat\_r.
apply H.
rewrite cap_-cup_-distr_-r.
rewrite cap_complement_empty cup_comm cup_empty.
apply cap_{-}l.
                                                    beta ^))).
apply (@inc_trans _ _ _ (beta
                                        (alpha
rewrite cup\_cap\_distr\_l.
rewrite complement_classic cap_universal.
apply cup_r.
apply cup\_inc\_compat\_l.
apply H.
Qed.
  Lemma 85 (bool_lemma4) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                     \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \beta^- \sqsubseteq \alpha^- \sqcup \gamma.
Lemma bool\_lemma4 {A B : eqType} {alpha beta gamma : Rel A B}:
 alpha
            (beta
                      gamma) \leftrightarrow \texttt{beta} (alpha ^
                                                              qamma).
Proof.
rewrite bool_lemma3.
rewrite cap\_comm.
apply iff_sym.
                       alpha) with (beta ^ (alpha ^) ^).
replace (beta ^
apply bool_lemma3.
by [rewrite complement_invol].
Qed.
  Lemma 86 (bool_lemma5) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                 \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \nabla_{AB} = \alpha^- \sqcup \beta \sqcup \gamma.
Lemma bool_lemma5 {A B : eqType} {alpha beta gamma : Rel A B}:
                      qamma) \leftrightarrow AB = (alpha \hat{beta})
 alpha
                                                                          qamma.
Proof.
rewrite bool\_lemma1.
by [rewrite cup\_assoc].
Qed.
End main.
```

Chapter 4

Library Relation_Properties

```
Require Import Basic\_Notations\_Set.
Require Import Basic\_Lemmas.
Require Import Logic.FunctionalExtensionality.
Require Import Logic.Classical\_Prop.
Module main~(def:Relation).
Import def.
Module Basic\_Lemmas:=Basic\_Lemmas.main~def.
Import Basic\_Lemmas.
```

4.1 関係計算の基本的な性質

```
Lemma 87 (RelAB_unique) \phi_{AB} = \nabla_{AB} \Leftrightarrow \forall \alpha, \beta : A \to B, \alpha = \beta. Lemma RelAB_unique \ \{A \ B : eqType\}: A \ B = A \ B \Leftrightarrow (\forall \ alpha \ \mathsf{beta} : Rel \ A \ B, \ alpha = \mathsf{beta}).
```

```
A \ B = A \ B \leftrightarrow (\forall \ alpha \ \mathsf{beta} : Rel \ A \ B, \ alp
Proof.

split; move \Rightarrow H.

move \Rightarrow alpha \ \mathsf{beta}.

replace \mathsf{beta} \ \mathsf{with} \ (A \ B).

apply inc\_antisym.

rewrite H.

apply inc\_alpha\_universal.

apply inc\_antisym.

apply inc\_antisym.

apply inc\_antisym.

apply inc\_antisym.

apply inc\_empty\_alpha.

rewrite H.
```

```
apply inc\_alpha\_universal. apply H. Qed.
```

```
Lemma 88 (either_empty)
```

```
\phi_{AB} = \nabla_{AB} \Leftrightarrow A = \emptyset \lor B = \emptyset.
Lemma either\_empty \{A \ B : eqType\}:
                                                    A B =
                                                                  A \ B \leftrightarrow (A \rightarrow False) \lor (B \rightarrow False).
Proof.
rewrite RelAB\_unique.
split; move \Rightarrow H.
case (classic (\exists \_: A, True)).
elim \Rightarrow a H0.
right.
move \Rightarrow b.
remember (fun (\_: A) (\_: B) \Rightarrow True) as T.
remember (fun (\_: A) (\_: B) \Rightarrow False) as F.
move: (H \ T \ F) \Rightarrow H1.
assert (T \ a \ b = F \ a \ b).
by [rewrite H1].
rewrite HeqT HeqF in H2.
rewrite -H2.
apply I.
move \Rightarrow H0.
left.
move \Rightarrow a.
apply H0.
\exists a.
apply I.
move \Rightarrow alpha beta.
assert (A \rightarrow B \rightarrow False).
move \Rightarrow a \ b.
case H; move \Rightarrow H\theta.
apply (H0 \ a).
apply (H0\ b).
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply False_ind.
apply (H0 \ a \ b).
Qed.
```

```
Lemma 89 (unit_empty_not_universal)
```

```
\phi_{II} \neq \nabla_{II}.
```

```
Lemma unit\_empty\_not\_universal: i i \neq i i.

Proof.

move \Rightarrow H.

apply either\_empty in H.

case H; move \Rightarrow H0.

apply (H0\ tt).

apply (H0\ tt).

Qed.

Lemma 90 (unit\_empty\_or\_universal) Let\ \alpha: I \to I. Then,

\alpha = \phi_{II} \lor \alpha = \nabla_{II}.

Lemma unit\_empty\_or\_universal\ \{alpha: Rel\ i\ i\}: alpha = i\ i\ \lor
```

```
Lemma unit\_empty\_or\_universal \{alpha : Rel \ i \ i\}: alpha = i \ i \lor alpha =
                                                                                           i i.
assert (\forall beta : Rel\ i\ i, beta = (fun (_ _ : i) \Rightarrow True) \lor beta = (fun (_ _ : i) \Rightarrow False)).
move \Rightarrow beta.
case (classic (beta tt tt)); move \Rightarrow H.
left.
apply functional_extensionality.
induction x.
apply functional_extensionality.
induction x.
apply prop_extensionality_ok.
split; move \Rightarrow H0.
apply I.
apply H.
right.
apply functional_extensionality.
induction x.
apply functional_extensionality.
induction x.
apply prop_extensionality_ok.
split.
apply H.
apply False_ind.
assert ((fun \_ : i \Rightarrow True) \neq (fun \_ : i \Rightarrow False)).
move \Rightarrow H0.
remember (fun \_ : i \Rightarrow True) as T.
```

```
remember (fun \_ : i \Rightarrow False) as F.
assert (T tt tt = F tt tt).
by [rewrite H\theta].
rewrite HeqT HeqF in H1.
rewrite -H1.
apply I.
case (H(i)); move \Rightarrow H1.
case (H(i)); move \Rightarrow H2.
apply False_ind.
apply unit_empty_not_universal.
by [rewrite H1 H2].
case (H \ alpha); move \Rightarrow H3.
left.
by [rewrite H3 H1].
right.
by [rewrite H3 H2].
case (H(i)); move \Rightarrow H2.
case (H \ alpha); move \Rightarrow H3.
right.
by [rewrite H3 H2].
left.
by [rewrite H3 H1].
apply False_ind.
apply unit_empty_not_universal.
by [rewrite H1 H2].
Qed.
```

Lemma 91 (unit_identity_is_universal)

```
id_I = \nabla_{II}.
```

```
Lemma unit\_identity\_is\_universal: Id\ i = i\ i.

Proof.

case (@unit\_empty\_or\_universal\ (Id\ i)); move \Rightarrow H.

apply False\_ind.

assert (Id\ i\ (i\ i\ \#\ i\ i)).

rewrite H.

apply inc\_empty\_alpha.

apply inc\_empty\_alpha.

apply inc\_residual\ in\ H0.

rewrite inv\_invol\ comp\_id\_r\ in\ H0.

apply unit\_empty\_not\_universal.

apply inc\_antisym.

apply inc\_empty\_alpha.
```

```
apply H0. apply H.
```

Qed.

Lemma 92 (unit_identity_not_empty)

 $id_I \neq \phi_{II}$.

Proof.

move $\Rightarrow H$.

apply $unit_empty_not_universal$.

rewrite -H.

apply $unit_identity_is_universal$.

Qed.

Lemma 93 (cupP_False) Let $f:(C \to D) \to (A \to B)$ and $P(\alpha) :=$ "False". Then,

$$\sqcup_{P(\alpha)} f(\alpha) = \phi_{AB}.$$

 $_{-}\{$ fun $_{-}: Rel \ C \ D \Rightarrow False<math>\} \ f = A \ B.$

Proof.

apply $inc_antisym$.

apply inc_cupP .

 $move \Rightarrow beta$.

apply False_ind.

apply inc_empty_alpha .

Qed.

Lemma 94 (capP_False) Let $f:(C \rightarrow D) \rightarrow (A \rightarrow B)$ and $P(\alpha) :=$ "False". Then,

$$\sqcap_{P(\alpha)} f(\alpha) = \nabla_{AB}.$$

Lemma capP-False $\{A \ B \ C \ D : eqType\} \ \{f : Rel \ C \ D \rightarrow Rel \ A \ B\}$:

 $_{\mathbf{I}} = \{ \mathbf{fun} : Rel \ C \ D \Rightarrow False \} \ f = A \ B.$

Proof.

apply $inc_antisym$.

apply $inc_alpha_universal$.

apply inc_capP .

 $move \Rightarrow beta.$

apply False_ind.

Qed.

```
Lemma 95 (cupP_eq) Let f, g: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                   (\forall \alpha: C \to D, P(\alpha) \Rightarrow f(\alpha) = g(\alpha)) \Rightarrow \sqcup_{P(\alpha)} f(\alpha) = \sqcup_{P(\alpha)} g(\alpha).
Lemma cupP_-eq {A B C D : eqType}
 \{f g : Rel \ C \ D \rightarrow Rel \ A \ B\} \ \{P : Rel \ C \ D \rightarrow Prop\}:
 (\forall alpha : Rel \ C \ D, P \ alpha \rightarrow f \ alpha = g \ alpha) \rightarrow -\{P\} \ f = -\{P\} \ g.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc\_cupP.
move \Rightarrow beta H0.
rewrite (H - H0).
move : beta H0.
apply inc\_cupP.
apply inc_refl.
apply inc\_cupP.
move \Rightarrow beta H\theta.
rewrite -(H - H\theta).
move : beta H0.
apply inc\_cupP.
apply inc\_reft.
Qed.
  Lemma 96 (capP_eq) Let f, g: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                   (\forall \alpha : C \to D, P(\alpha) \Rightarrow f(\alpha) = g(\alpha)) \Rightarrow \sqcap_{P(\alpha)} f(\alpha) = \sqcap_{P(\alpha)} g(\alpha).
Lemma capP_{-}eq \{A \ B \ C \ D : eqType\}
 \{f g : Rel \ C \ D \rightarrow Rel \ A \ B\} \ \{P : Rel \ C \ D \rightarrow Prop\}:
 (\forall alpha : Rel \ C \ D, P \ alpha \rightarrow f \ alpha = g \ alpha) \rightarrow \_\{P\} \ f = \_\{P\} \ q.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc\_capP.
move \Rightarrow beta H\theta.
rewrite -(H - H0).
move : beta H0.
apply inc\_capP.
apply inc_refl.
apply inc\_capP.
```

move \Rightarrow beta $H\theta$.

```
rewrite (H - H\theta).
move: beta H0.
apply inc\_capP.
apply inc\_reft.
Qed.
  Lemma 97 (cap_cupP_distr_l) Let \alpha: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                   \alpha \sqcap (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \sqcap f(\beta)).
Lemma cap\_cupP\_distr\_l \{A \ B \ C \ D : eqType\}
 \{alpha: Rel\ A\ B\}\ \{f: Rel\ C\ D \rightarrow Rel\ A\ B\}\ \{P: Rel\ C\ D \rightarrow Prop\}:
 alpha
            ( -\{P\} f) = -\{P\}  (fun beta : Rel \ C \ D \Rightarrow alpha
Proof.
apply inc\_upper.
move \Rightarrow qamma.
split; move \Rightarrow H.
apply inc\_cupP.
move \Rightarrow beta H\theta.
apply (@inc_trans _ _ _ (alpha
                                            _{-}\{P\}\ f)).
apply cap\_inc\_compat\_l.
move: H0.
apply inc\_cupP.
apply inc\_reft.
apply H.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow (alpha f beta)
                                                                            qamma).
apply inc\_cupP.
apply H.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow f beta (alpha \gg gamma)).
move \Rightarrow beta H1.
rewrite inc\_rpc\ cap\_comm.
apply (H0 - H1).
rewrite cap\_comm -inc\_rpc.
apply inc\_cupP.
apply H1.
Qed.
  Lemma 98 (cap_cupP_distr_r) Let \beta: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                   (\sqcup_{P(\alpha)} f(\alpha)) \sqcap \beta = \sqcup_{P(\alpha)} (f(\alpha) \sqcap \beta).
```

Lemma $cap_cupP_distr_r \{A \ B \ C \ D : eqType\}$

```
\{ beta : Rel \ A \ B \} \{ f : Rel \ C \ D \rightarrow Rel \ A \ B \} \{ P : Rel \ C \ D \rightarrow Prop \} :
                              \{P\} (fun alpha : Rel C D \Rightarrow f alpha
 (-\{P\}f)
                beta =
Proof.
rewrite cap\_comm.
replace (fun alpha: Rel\ C\ D \Rightarrow f\ alpha beta) with (fun alpha: Rel\ C\ D \Rightarrow beta
f alpha).
apply cap\_cupP\_distr\_l.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite cap\_comm].
Qed.
  Lemma 99 (cup_capP_distr_l) Let \alpha: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                 \alpha \sqcup (\sqcap_{P(\beta)} f(\beta)) = \sqcap_{P(\beta)} (\alpha \sqcup f(\beta)).
Lemma cup\_capP\_distr\_l \{A \ B \ C \ D : eqType\}
 \{alpha: Rel\ A\ B\}\ \{f: Rel\ C\ D \rightarrow Rel\ A\ B\}\ \{P: Rel\ C\ D \rightarrow Prop\}:
            ( _{P} f) = _{P} (fun beta : Rel C D \Rightarrow alpha f beta).
 alpha
Proof.
apply inc\_lower.
move \Rightarrow qamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply (@inc\_trans \_ \_ \_ (alpha \_ \{P\} f)).
apply H.
apply cup\_inc\_compat\_l.
move: H0.
apply inc\_capP.
apply inc\_reft.
rewrite bool_lemma3.
assert (\forall beta : Rel\ C\ D,\ P\ beta \rightarrow gamma (alpha
                                                                       f beta)).
apply inc\_capP.
apply H.
apply inc\_capP.
move \Rightarrow beta H1.
rewrite -bool_lemma3.
apply (H0 - H1).
Qed.
```

```
predicate. Then,
                                  (\sqcap_{P(\alpha)} f(\alpha)) \sqcup \beta = \sqcap_{P(\alpha)} (f(\alpha) \sqcup \beta).
Lemma cup\_capP\_distr\_r \{A \ B \ C \ D : eqType\}
 \{ beta : Rel \ A \ B \} \{ f : Rel \ C \ D \rightarrow Rel \ A \ B \} \{ P : Rel \ C \ D \rightarrow Prop \} :
                 beta = _{-}\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha
Proof.
rewrite cup\_comm.
replace (fun alpha : Rel \ C \ D \Rightarrow f \ alpha
                                                      beta) with (fun alpha : Rel \ C \ D \Rightarrow beta
f alpha).
apply cup\_capP\_distr\_l.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite cup\_comm].
Qed.
  Lemma 101 (de_morgan3) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                      (\sqcup_{P(\alpha)} f(\alpha))^- = (\sqcap_{P(\alpha)} f(\alpha)^-).
Lemma de\_morgan3
 \{A \ B \ C \ D : eqType\} \{f : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Prop\}:
 ( -\{P\} f)^{\circ} = -\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha^{\circ}).
Proof.
apply inc\_lower.
move \Rightarrow gamma.
rewrite inc\_capP.
split; move \Rightarrow H.
move \Rightarrow beta H0.
rewrite bool_lemma1 -de_morgan2 complement_move complement_universal.
apply bool\_lemma2 in H.
apply inc\_antisym.
apply inc\_empty\_alpha.
rewrite - H complement_invol.
apply cap\_inc\_compat\_l.
move: H0.
apply inc\_cupP.
apply inc_refl.
rewrite bool_lemma2 complement_invol.
rewrite cap\_cupP\_distr\_l.
apply inc\_antisym.
```

Lemma 100 (cup_capP_distr_r) Let $\beta: A \rightarrow B$, $f: (C \rightarrow D) \rightarrow (A \rightarrow B)$ and P:

```
apply inc\_cupP.
move \Rightarrow beta H\theta.
rewrite -inc_-rpc.
apply (H - H\theta).
apply inc\_empty\_alpha.
Qed.
  Lemma 102 (de_morgan4) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                        (\sqcap_{P(\alpha)} f(\alpha))^- = (\sqcup_{P(\alpha)} f(\alpha)^-).
Lemma de\_morgan4
 \{A \ B \ C \ D : eqType\} \{f : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Prop\}:
 ( -\{P\} f)^{=} -\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha^{=}).
Proof.
rewrite -complement_move de_morgan3.
replace (fun alpha : Rel \ C \ D \Rightarrow (f \ alpha \ \hat{}) \ \hat{}) with f.
by ||.
{\tt apply} \ functional\_extensionality.
move \Rightarrow x.
by [rewrite complement_invol].
Qed.
  Lemma 103 (cup_to_cupP) Let f:(C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                        f(\alpha) \sqcup f(\beta) = \sqcup_{\gamma = \alpha \lor \gamma = \beta} f(\gamma).
Lemma cup\_to\_cupP
 \{A \ B \ C \ D : eqType\} \{alpha \ \mathsf{beta} : Rel \ C \ D\} \{f : Rel \ C \ D \to Rel \ A \ B\}:
                f \text{ beta}) = -\{\text{fun } gamma : Rel \ C \ D \Rightarrow gamma = alpha \lor gamma = \text{beta}\}
 (f alpha
f.
Proof.
apply inc\_upper.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_cupP.
apply inc\_cup in H.
move \Rightarrow gamma\ H0.
case H0 \Rightarrow H1.
rewrite H1.
apply H.
rewrite H1.
```

apply H.

```
apply inc\_cup.
assert (\forall gamma : Rel \ C \ D, gamma = alpha \lor gamma = beta \rightarrow f \ gamma
apply inc\_cupP.
apply H.
split.
apply (H0 \ alpha).
by [left].
apply (H0 \text{ beta}).
by [right].
Qed.
  Lemma 104 (cap_to_capP) Let f:(C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                     f(\alpha) \sqcap f(\beta) = \sqcap_{\gamma = \alpha \lor \gamma = \beta} f(\gamma).
Lemma cap\_to\_capP
 \{A \ B \ C \ D : eqType\} \{alpha \ \mathsf{beta} : Rel \ C \ D\} \{f : Rel \ C \ D \to Rel \ A \ B\}:
               f \text{ beta}) = -\{\text{fun } gamma : Rel \ C \ D \Rightarrow gamma = alpha \lor gamma = \text{beta}\}
f.
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_capP.
apply inc\_cap in H.
move \Rightarrow gamma H0.
case H0 \Rightarrow H1.
rewrite H1.
apply H.
rewrite H1.
apply H.
apply inc\_cap.
assert (\forall gamma : Rel \ C \ D, gamma = alpha \lor gamma = beta \rightarrow delta f gamma).
apply inc\_capP.
apply H.
split.
apply (H0 \ alpha).
by [left].
apply (H0 \text{ beta}).
by [right].
Qed.
```

4.2 comp_inc_compat と派生補題

apply inc_-inv .

apply H. Qed.

```
Lemma 105 (comp_inc_compat_ab_ab') Let \alpha : A \to B and \beta, \beta' : B \to C. Then,
                                          \beta \sqsubseteq \beta' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha \cdot \beta'.
Lemma comp\_inc\_compat\_ab\_ab'
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ beta' : Rel \ B \ C\}:
            beta' \rightarrow (alpha \cdot beta') (alpha · beta').
Proof.
move \Rightarrow H.
replace (alpha \cdot beta) with ((alpha \#) \# \cdot beta).
apply inc\_residual.
apply (@inc\_trans \_ \_ \_ beta').
apply H.
apply inc\_residual.
rewrite inv_-invol.
apply inc\_reft.
by [rewrite inv\_invol].
Qed.
  Lemma 106 (comp_inc_compat_ab_a'b) Let \alpha, \alpha' : A \to B and \beta : B \to C. Then,
                                          \alpha \sqsubseteq \alpha' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha' \cdot \beta.
Lemma comp_inc_compat_ab_a'b
 \{A \ B \ C : eqType\} \ \{alpha \ alpha' : Rel \ A \ B\} \ \{beta : Rel \ B \ C\}:
             alpha' \rightarrow (alpha \cdot beta) \quad (alpha' \cdot beta).
 alpha
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ (alpha • beta)).
rewrite -(@inv_invol _ _ (alpha' • beta)).
apply inc_{-}inv.
rewrite comp_{-}inv \ comp_{-}inv.
apply comp\_inc\_compat\_ab\_ab'.
```

```
Lemma 107 (comp_inc_compat) Let \alpha, \alpha' : A \to B and \beta, \beta' : B \to C. Then,
                                     \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha' \cdot \beta'.
Lemma comp\_inc\_compat
 \{A \ B \ C : eqType\} \ \{alpha \ alpha' : Rel \ A \ B\} \ \{beta \ beta' : Rel \ B \ C\}:
                                     beta' \rightarrow (alpha \cdot beta') \quad (alpha' \cdot beta').
 alpha
             alpha' \rightarrow \mathtt{beta}
Proof.
move \Rightarrow H H0.
apply (@inc_trans _ _ _ (alpha' • beta)).
apply (@comp\_inc\_compat\_ab\_a'b\_\_\_\_\_H).
apply (@comp\_inc\_compat\_ab\_ab'\_\_\_\__H0).
Qed.
  Lemma 108 (comp_inc_compat_ab_a) Let \alpha : A \rightarrow B and \beta : B \rightarrow B. Then,
                                            \beta \sqsubseteq id_B \Rightarrow \alpha \cdot \beta \sqsubseteq \beta.
Lemma comp\_inc\_compat\_ab\_a {A \ B : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ B}:
            Id B \rightarrow (alpha \cdot beta)
                                                alpha.
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_ab'\_\_\_alpha\_\_H) \Rightarrow H0.
rewrite comp_{-}id_{-}r in H0.
apply H0.
Qed.
  Lemma 109 (comp_inc_compat_a_ab) Let \alpha : A \rightarrow B and \beta : B \rightarrow B. Then,
                                            id_B \sqsubseteq \beta \Rightarrow \beta \sqsubseteq \alpha \cdot \beta.
Lemma comp_inc_compat_a_ab {A B : eqType} {alpha : Rel A B} {beta : Rel B B}:
 Id\ B
            beta \rightarrow alpha
                                   (alpha • beta).
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_ab'\_\_\_alpha\_\_H) \Rightarrow H0.
rewrite comp_{-}id_{-}r in H0.
apply H0.
Qed.
```

```
Lemma 110 (comp_inc_compat_ab_b) Let \alpha : A \rightarrow A and \beta : A \rightarrow B. Then,
                                             \alpha \sqsubseteq id_A \Rightarrow \alpha \cdot \beta \sqsubseteq \beta.
Lemma comp\_inc\_compat\_ab\_b {A \ B : eqType} {alpha : Rel \ A \ A} {beta : Rel \ A \ B}:
             Id A \rightarrow (alpha \cdot beta)
                                                  beta.
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_a"b\_\_\_\_\_ beta H) \Rightarrow H0.
rewrite comp_{-}id_{-}l in H0.
apply H0.
Qed.
  Lemma 111 (comp_inc_compat_b_ab) Let \alpha : A \rightarrow A and \beta : A \rightarrow B. Then,
                                             id_A \sqsubseteq \alpha \Rightarrow \beta \sqsubseteq \alpha \cdot \beta.
Lemma comp\_inc\_compat\_b\_ab {A \ B : eqType} {alpha : Rel \ A \ A} {beta : Rel \ A \ B}:
             alpha \rightarrow \mathtt{beta}
                                   (alpha • beta).
 Id A
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_a"b\_\_\_\_\_ beta H) \Rightarrow H0.
rewrite comp_{-}id_{-}l in H0.
apply H0.
Qed.
          逆関係に関する補題
4.3
  Lemma 112 (inv_move) Let \alpha : A \rightarrow B and \beta : B \rightarrow A. Then,
                                                \alpha = \beta^{\sharp} \Leftrightarrow \alpha^{\sharp} = \beta.
Lemma inv\_move \{A \ B : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ A\}:
 alpha = \mathtt{beta} \ \# \leftrightarrow alpha \ \# = \mathtt{beta}.
Proof.
split; move \Rightarrow H.
by [rewrite H inv_invol].
```

by [rewrite -H inv_invol].

Qed.

```
Lemma 113 (comp_inv_inv) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                            \alpha \cdot \beta = (\beta^{\sharp} \cdot \alpha^{\sharp})^{\sharp}.
alpha • beta = (beta # • alpha #) #.
Proof.
apply inv_move.
apply comp_{-}inv.
Qed.
  Lemma 114 (inv_inc_move) Let \alpha : A \rightarrow B and \beta : B \rightarrow A. Then,
                                            \alpha \sqsubseteq \beta^{\sharp} \Leftrightarrow \alpha^{\sharp} \sqsubseteq \beta.
Lemma inv\_inc\_move \{A \ B : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ A\}:
            beta \# \leftrightarrow alpha \#
                                        beta.
Proof.
split; move \Rightarrow H.
rewrite -(@inv_invol_{-} beta).
apply inc_{-}inv.
apply H.
rewrite -(@inv_invol _ _ alpha).
apply inc_{-}inv.
apply H.
Qed.
  Lemma 115 (inv_invol2) Let \alpha, \beta : A \rightarrow B. Then,
                                            \alpha^{\sharp} = \beta^{\sharp} \Rightarrow \alpha = \beta.
Lemma inv\_invol2 {A B : eqType} {alpha beta : Rel A B}:
 alpha \# = \mathtt{beta} \# \to alpha = \mathtt{beta}.
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ alpha) -(@inv_invol _ _ beta).
apply f_equal.
apply H.
Qed.
```

Lemma 116 (inv_inc_invol) Let $\alpha, \beta : A \rightarrow B$. Then,

```
\alpha^{\sharp} \sqsubseteq \beta^{\sharp} \Rightarrow \alpha \sqsubseteq \beta.
Lemma inv\_inc\_invol {A B : eqType} {alpha beta : Rel A B}:
               \mathtt{beta} \ \# \to alpha
 alpha \#
                                       beta.
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ alpha) -(@inv_invol _ _ beta).
apply inc_inv.
apply H.
Qed.
  Lemma 117 (inv_cupP_distr, inv_cup_distr) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and
  P: predicate. Then,
                                     (\sqcup_{P(\alpha)} f(\alpha))^{\sharp} = (\sqcup_{P(\alpha)} f(\alpha)^{\sharp}).
Lemma inv\_cupP\_distr {A B C D : eqType} {f : Rel C D \rightarrow Rel A B} {P : Rel C D \rightarrow
Prop}:
 ( _{P} f) # = ( _{P} (fun alpha : Rel C D \Rightarrow f alpha #)).
Proof.
apply inc\_antisym.
rewrite -inv\_inc\_move.
apply inc\_cupP.
assert (\forall beta : Rel\ C\ D,\ P\ beta \to f\ beta # _{{}^{-}}{P}\ (fun\ alpha:\ Rel\ C\ D \Rightarrow f
alpha \#)).
apply inc\_cupP.
apply inc_refl.
move \Rightarrow beta H\theta.
rewrite inv\_inc\_move.
apply (H - H\theta).
apply inc\_cupP.
move \Rightarrow beta H0.
apply inc_{-}inv.
move: H0.
apply inc\_cupP.
apply inc\_reft.
Qed.
Lemma inv\_cup\_distr {A B : eqType} {alpha beta : Rel A B}:
             beta) \# = alpha \# beta \#.
 (alpha
Proof.
by [rewrite cup\_to\_cupP -inv\_cupP\_distr -cup\_to\_cupP].
```

Qed.

```
Lemma 118 (inv_capP_distr, inv_cap_distr) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P
  : predicate. Then,
                                      (\sqcap_{P(\alpha)} f(\alpha))^{\sharp} = (\sqcap_{P(\alpha)} f(\alpha)^{\sharp}).
Lemma inv\_capP\_distr {A B C D : eqType} {f : Rel\ C\ D \rightarrow Rel\ A\ B} {P : Rel\ C\ D \rightarrow Rel\ A\ B}
 ( -\{P\} f) \# = ( -\{P\} (\mathbf{fun} \ alpha : Rel \ C \ D \Rightarrow f \ alpha \#)).
Proof.
apply inc\_antisym.
apply inc\_capP.
move \Rightarrow beta H.
apply inc_{-}inv.
move: H.
apply inc\_capP.
apply inc\_reft.
rewrite inv\_inc\_move.
apply inc\_capP.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow _{{P}} (fun\ alpha : Rel\ C\ D \Rightarrow f\ alpha\ \#)
                                                                                                              f
beta \#).
apply inc\_capP.
apply inc\_reft.
move \Rightarrow beta H\theta.
rewrite -inv\_inc\_move.
apply (H - H\theta).
Qed.
Lemma inv\_cap\_distr {A B : eqType} {alpha beta : Rel A B}:
 (alpha
             beta) \# = alpha \# beta \#.
Proof.
by [rewrite cap_to_capP -inv_capP_distr -cap_to_capP].
Qed.
  Lemma 119 (rpc_inv_distr) Let \alpha, \beta : A \rightarrow B. Then,
                                           (\alpha \Rightarrow \beta)^{\sharp} = \alpha^{\sharp} \Rightarrow \beta^{\sharp}.
Lemma rpc\_inv\_distr {A B : eqType} {alpha beta : Rel A B}:
 (alpha \gg beta) \# = alpha \# \gg beta \#.
Proof.
apply inc_lower.
move \Rightarrow gamma.
```

$$\label{eq:split} \begin{split} & \text{split}; \, \texttt{move} \Rightarrow H. \\ & \text{apply } \, inc_rpc. \\ & \text{rewrite } \, inv_inc_move \, inv_cap_distr \, inv_invol. \\ & \text{rewrite } \, -inc_rpc \, -inv_inc_move. \\ & \text{apply } \, H. \\ & \text{rewrite } \, inv_inc_move \, inc_rpc. \\ & \text{rewrite } \, -(@inv_invol \, _ \, alpha) \, -inv_cap_distr \, -inv_inc_move. \\ & \text{apply } \, inc_rpc. \\ & \text{apply } \, inc_rpc. \\ & \text{apply } \, H. \\ & \text{Qed.} \end{split}$$

Lemma 120 (inv_empty)

$$\phi_{AB}^{\sharp} = \phi_{BA}.$$

Lemma inv_empty { $A\ B: eqType$ }: $A\ B\ \# = B\ A.$ Proof. apply $inc_antisym.$ rewrite $-inv_inc_move.$

apply inc_empty_alpha . apply inc_empty_alpha .

Qed.

Lemma 121 (inv_universal)

$$abla_{AB}^{\sharp} =
abla_{BA}.$$

Lemma $inv_universal\ \{A\ B: eqType\}: A\ B\ \# = B\ A.$

Proof.

apply $inc_antisym$.

apply $inc_alpha_universal$.

rewrite inv_inc_move .

apply $inc_alpha_universal$.

Qed.

Lemma 122 (inv_id)

$$id_A^{\sharp} = id_A.$$

Proof.

replace $(Id \ A \ \#)$ with $((Id \ A \ \#) \ \# \cdot Id \ A \ \#)$.

by [rewrite -comp_inv comp_id_l inv_invol].

by [rewrite inv_invol comp_id_l].

Qed.

Lemma 123 (inv_complement) Let $\alpha : A \rightarrow B$. Then,

$$(\alpha^{-})^{\sharp} = (\alpha^{\sharp})^{-}.$$

Lemma $inv_complement \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: (alpha ^) \# = (alpha \#) ^.$ apply $inc_antisym$. apply inc_rpc . rewrite $-inv_cap_distr$. rewrite cap_comm -inv_inc_move inv_empty. rewrite cap_complement_empty. apply inc_reft . rewrite inv_inc_move . apply inc_rpc . replace $(((alpha \#) \hat{}) \# alpha)$ with $(((alpha \#) \hat{}) \# (alpha \#) \#)$. $\verb"rewrite" - inv_cap_distr".$ rewrite cap_comm -inv_inc_move inv_empty. rewrite cap_complement_empty. apply inc_reft . by [rewrite $inv_{-}invol$]. Qed.

Lemma 124 (inv_difference_distr) Let $\alpha, \beta : A \rightarrow B$. Then,

$$(\alpha - \beta)^{\sharp} = \alpha^{\sharp} - \beta^{\sharp}.$$

Lemma $inv_difference_distr$ { $A \ B : eqType$ } { $alpha \ beta : Rel \ A \ B$ }: (alpha - beta) $\# = alpha \ \# - beta \ \#$.

Proof.

rewrite inv_cap_distr .

by [rewrite $inv_complement$].

Qed.

4.4 合成に関する補題

Lemma 125 (comp_cupP_distr_l, comp_cup_distr_l) Let $\alpha : A \rightarrow B$, $f : (D \rightarrow E) \rightarrow (B \rightarrow C)$ and P : predicate. Then,

$$\alpha \cdot (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \cdot f(\beta)).$$

Lemma $comp_cupP_distr_l$ {A B C D E : eqType}

```
\{alpha: Rel\ A\ B\}\ \{f: Rel\ D\ E \rightarrow Rel\ B\ C\}\ \{P: Rel\ D\ E \rightarrow Prop\}:
 alpha \cdot ( _{P} f) = _{P} (fun beta : Rel D E \Rightarrow (alpha \cdot f beta)).
Proof.
apply inc\_upper.
move \Rightarrow gamma.
split; move \Rightarrow H.
rewrite -(@inv\_invol\_\_alpha) in H.
apply inc\_residual in H.
apply inc\_cupP.
assert (\forall beta : Rel\ D\ E, P beta \rightarrow f beta (alpha\ \#\ qamma)).
apply inc\_cupP.
apply H.
move \Rightarrow beta H1.
rewrite -(@inv_invol _ _ alpha).
apply inc\_residual.
apply (H0 - H1).
rewrite -(@inv_invol _ _ alpha).
apply inc\_residual.
apply inc\_cupP.
assert (\forall beta : Rel\ D\ E, P beta \rightarrow (alpha • f beta)
                                                                       qamma).
apply inc\_cupP.
apply H.
move \Rightarrow beta H1.
apply inc_residual.
rewrite inv\_invol.
apply (H0 - H1).
Qed.
Lemma comp\_cup\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
 alpha • (beta
                      gamma) = (alpha \cdot beta)
                                                          (alpha \cdot gamma).
by [rewrite cup\_to\_cupP -comp\_cupP\_distr\_l -cup\_to\_cupP].
Qed.
  Lemma 126 (comp_cupP_distr_r, comp_cup_distr_r) Let f:(D \rightarrow E) \rightarrow (A \rightarrow E)
  B), \beta: B \rightarrow C and P: predicate. Then,
                                  (\sqcup_{P(\alpha)} f(\alpha)) \cdot \beta = \sqcup_{P(\alpha)} (f(\alpha) \cdot \beta).
Lemma comp\_cupP\_distr\_r {A B C D E : eqType}
 \{ beta : Rel \ B \ C \} \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \{ P : Rel \ D \ E \rightarrow Prop \} :
 ( \{P\} f) \cdot beta = \{P\} (fun \ alpha : Rel \ D \ E \Rightarrow (f \ alpha \cdot beta)).
```

```
Proof.
replace (fun alpha: Rel D E \Rightarrow f alpha • beta) with (fun alpha: Rel D E \Rightarrow (beta #
• f alpha #) #).
rewrite -inv\_cupP\_distr.
rewrite -comp\_cupP\_distr\_l.
rewrite -inv\_cupP\_distr.
rewrite comp_{-}inv.
by [rewrite inv_invol inv_invol].
apply functional_extensionality.
move \Rightarrow x.
rewrite comp_{-}inv.
by [rewrite inv_invol inv_invol].
Qed.
Lemma comp\_cup\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
            beta) • gamma = (alpha • gamma)
 (alpha
                                                          (beta • gamma).
Proof.
by [rewrite (@cup\_to\_cupP\_\_\_\_\_\_id) comp\_cupP\_distr\_r -cup\_to\_cupP].
Qed.
  Lemma 127 (comp_capP_distr) Let \alpha: A \to B, \gamma: C \to D, f: (E \to F) \to (B \to F)
  C) and P: predicate. Then,
                             \alpha \cdot (\sqcap_{P(\beta)} f(\beta)) \cdot \gamma \sqsubseteq \sqcap_{P(\beta)} (\alpha \cdot f(\beta) \cdot \gamma).
Lemma comp\_capP\_distr {A B C D E F : eqType}
 {alpha : Rel \ A \ B} \{gamma : Rel \ C \ D\}
 \{f: Rel\ E\ F \to Rel\ B\ C\}\ \{P: Rel\ E\ F \to \operatorname{Prop}\}:
 (alpha \cdot ( _{\{P\}} f)) \cdot gamma
       \{P\} (fun beta : Rel\ E\ F \Rightarrow ((alpha \cdot f beta) \cdot gamma)).
Proof.
apply inc\_capP.
move \Rightarrow beta H.
apply comp\_inc\_compat\_ab\_a'b.
apply comp_inc_compat_ab_ab'.
move: H.
apply inc\_capP.
apply inc\_reft.
Qed.
```

```
Lemma 128 (comp_capP_distr_l, comp_cap_distr_l) Let \alpha : A \rightarrow B, f : (D \rightarrow B)
  E) \rightarrow (B \rightarrow C) and P: predicate. Then,
                                    \alpha \cdot (\sqcap_{P(\beta)} f(\beta)) \sqsubseteq \sqcap_{P(\beta)} (\alpha \cdot f(\beta)).
Lemma comp\_capP\_distr\_l \{A \ B \ C \ D \ E : eqType\}
 \{alpha : Rel \ A \ B\} \{f : Rel \ D \ E \rightarrow Rel \ B \ C\} \{P : Rel \ D \ E \rightarrow Prop\}:
                               \{P\} (fun beta : Rel\ D\ E \Rightarrow (alpha \cdot f \text{ beta})).
 (alpha \cdot (-\{P\}f))
Proof.
move: (@comp\_capP\_distr\_\_\_\_ alpha (Id C) f P) \Rightarrow H.
rewrite comp_{-}id_{-}r in H.
replace (fun beta: Rel\ D\ E \Rightarrow (alpha \cdot f \text{ beta}) \cdot Id\ C) with (fun beta: Rel\ D\ E \Rightarrow
(alpha \cdot f beta)) in H.
apply H.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite comp_{-}id_{-}r].
Qed.
Lemma comp\_cap\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
 (alpha • (beta
                       qamma))
                                        ((alpha \cdot beta) (alpha \cdot gamma)).
Proof.
rewrite cap\_to\_capP (@cap\_to\_capP\_\_\_\_\_id).
apply comp\_capP\_distr\_l.
Qed.
  Lemma 129 (comp_capP_distr_r, comp_cap_distr_r) Let \beta: B \rightarrow C, f: (D \rightarrow C)
  (E) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                    (\sqcap_{P(\alpha)} f(\alpha)) \cdot \beta \sqsubseteq \sqcap_{P(\alpha)} (f(\alpha) \cdot \beta).
{\tt Lemma}\ comp\_capP\_distr\_r
 \{A \ B \ C \ D \ E : eqType\} \ \{ beta : Rel \ B \ C \} \ \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \ \{ P : Rel \ D \ E \rightarrow Rel \ A \ B \} \} 
Prop}:
 (( _{P} f) \cdot beta) _{P} (fun alpha : Rel D E \Rightarrow (f alpha · beta)).
Proof.
move: (@comp\_capP\_distr\_\_\_\_\_(Id\ A)\ beta\ f\ P) \Rightarrow H.
rewrite comp_{-}id_{-}l in H.
replace (fun alpha: Rel D E \Rightarrow (Id A • f alpha) • beta) with (fun alpha: Rel D E
\Rightarrow f \ alpha \cdot beta) \ in \ H.
apply H.
apply functional_extensionality.
```

```
move \Rightarrow x.
by [rewrite comp_{-}id_{-}l].
Qed.
Lemma comp\_cap\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
             beta) • qamma) ((alpha • qamma) (beta • qamma)).
Proof.
rewrite (@cap\_to\_capP\_\_\_\_\_id) (@cap\_to\_capP\_\_\_\_\_\_(fun x \Rightarrow x \cdot gamma)).
apply comp\_capP\_distr\_r.
Qed.
  Lemma 130 (comp_empty_l, comp_empty_r) Let \alpha : A \rightarrow B, \beta : B \rightarrow C. Then,
                                    \alpha \cdot \phi_{BC} = \phi_{AB} \cdot \beta = \phi_{AC}.
Lemma comp\_empty\_r \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\}: alpha  • B \ C =
apply inc\_antisym.
rewrite -(@inv_invol _ _ alpha).
apply inc\_residual.
apply inc\_empty\_alpha.
apply inc\_empty\_alpha.
Qed.
Lemma comp\_empty\_l \{A \ B \ C : eqType\} \{ beta : Rel \ B \ C \}: A \ B \cdot beta = A \ C.
Proof.
rewrite -(@inv_invol_{-} ( AB \cdot beta)).
rewrite -inv_move comp_inv inv_empty inv_empty.
apply comp\_empty\_r.
Qed.
  Lemma 131 (comp_either_empty) Let \alpha : A \rightarrow B, \beta : B \rightarrow C. Then,
                               \alpha = \phi_{AB} \vee \beta = \phi_{BC} \Rightarrow \alpha \cdot \beta = \phi_{AC}.
Lemma comp_either_empty {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
              A \ B \lor \mathtt{beta} = B \ C \to alpha \bullet \mathtt{beta} = A \ C.
 alpha =
Proof.
case; move \Rightarrow H.
rewrite H.
apply comp\_empty\_l.
rewrite H.
apply comp\_empty\_r.
```

Qed.

```
\alpha \cdot \beta \neq \phi_{AC} \Rightarrow \alpha \neq \phi_{AB} \wedge \beta \neq \phi_{BC}. Lemma comp\_neither\_empty \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}: alpha \cdot beta \neq A \ C \rightarrow alpha \neq A \ B \wedge beta \neq B \ C. Proof. move \Rightarrow H. split; move \Rightarrow H0. apply H. rewrite H0. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_r. Qed.
```

Lemma 132 (comp_neither_empty) Let $\alpha : A \to B$, $\beta : B \to C$. Then,

4.5 単域と Tarski の定理

```
\nabla_{IA} \cdot \alpha \cdot \nabla_{BI} = id_I.
```

Lemma 133 (lemma_for_tarski1) Let $\alpha : A \to B$ and $\alpha \neq \phi_{AB}$. Then,

```
Lemma lemma\_for\_tarski1 \{A B : eqType\} \{alpha : Rel A B\}:
             A B \rightarrow ((i A \cdot alpha) \cdot B i) = Id i.
 alpha \neq
Proof.
move \Rightarrow H.
              i A \cdot alpha \cdot B i \neq i i.
assert (((
move \Rightarrow H0.
apply H.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((Ai \cdot ((iA \cdot alpha) \cdot Bi)) \cdot iB)).
rewrite comp_assoc comp_assoc unit_universal.
rewrite -comp_assoc -comp_assoc unit_universal.
apply (@inc\_trans \_ \_ \_ ((Id A \cdot alpha) \cdot Id B)).
rewrite comp\_id\_l comp\_id\_r.
apply inc\_reft.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
```

```
apply inc\_alpha\_universal.

rewrite H0\ comp\_empty\_r\ comp\_empty\_l.

apply inc\_refl.

apply inc\_empty\_alpha.

case (@unit\_empty\_or\_universal (( i A · alpha) · B i)); move \Rightarrow H1.

apply False\_ind.

apply (H0\ H1).

rewrite unit\_identity\_is\_universal.

apply H1.

Qed.
```

Lemma 134 (lemma_for_tarski2)

$$\nabla_{AI} \cdot \nabla_{IB} = \nabla_{AB}$$
.

Lemma $lemma_for_tarski2$ {A B : eqType}: A $i \cdot i B = i$ A B. Proof. apply $inc_antisym$. apply $inc_alpha_universal$. apply $@inc_trans___$ ($AA \cdot$ A B)). apply $(@inc_trans _ _ _ (Id A \cdot A B))$. rewrite $comp_{-}id_{-}l$. apply $inc_refl.$ apply $comp_inc_compat_ab_a'b$. apply $inc_alpha_universal$. rewrite -(@unit_universal A) comp_assoc. apply $comp_inc_compat_ab_ab$ '. apply $inc_alpha_universal$. Qed.

Lemma 135 (tarski) Let $\alpha : A \rightarrow B$ and $\alpha \neq \phi_{AB}$. Then,

$$\nabla_{AA} \cdot \alpha \cdot \nabla_{BB} = \nabla_{AB}.$$

```
Lemma tarski \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha \neq A \ B \rightarrow ((A \ A \cdot alpha) \cdot B \ B) = A \ B.

Proof.

move \Rightarrow H.

rewrite -(@unit\_universal \ A) - (@unit\_universal \ B).

move : (@lemma\_for\_tarski1 \_ alpha \ H) \Rightarrow H0.

rewrite -comp\_assoc \ (@comp\_assoc \_ \_ (A \ i)) \ (@comp\_assoc \_ (A \ i)).

rewrite H0 \ comp\_id\_r.

apply lemma\_for\_tarski2.
```

Qed.

```
Lemma 136 (comp_universal1) Let B \neq \emptyset. Then,
```

$$\nabla_{AB} \cdot \nabla_{BC} = \nabla_{AC}$$
.

```
Lemma comp\_universal\ \{A\ B\ C: eqType\}: B \rightarrow A\ B \cdot B\ C =
                                                                          A C.
Proof.
move \Rightarrow b.
replace (AB) with (AB \cdot BB).
rewrite -(@lemma_for_tarski2 A B) -(@lemma_for_tarski2 B C).
rewrite (@comp_assoc _ _ _ ( A i)) (@comp_assoc _ _ _ ( A i)) -(@comp_assoc _
---(Bi).
rewrite lemma\_for\_tarski1.
rewrite comp_{-}id_{-}l.
apply lemma_for_tarski2.
apply not\_eq\_sym.
move \Rightarrow H.
apply either\_empty in H.
case H; move \Rightarrow H0.
apply (H0\ b).
apply (H0\ b).
apply inc\_antisym.
apply inc_alpha_universal.
apply (@inc\_trans \_ \_ \_ ( AB \cdot Id B)).
rewrite comp_{-}id_{-}r.
apply inc\_reft.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
```

Lemma 137 (comp_universal2)

$$\nabla_{IA}^{\sharp} \cdot \nabla_{IB} = \nabla_{AB}.$$

Lemma $comp_universal2$ { $A \ B : eqType$ }: $i \ A \ \# \bullet$ $i \ B = A \ B$. Proof. rewrite $inv_universal$. apply $lemma_for_tarski2$. Qed.

Lemma 138 (empty_equivalence1, empty_equivalence2, empty_equivalence3)

```
A = \emptyset \Leftrightarrow \nabla_{IA} = \phi_{IA} \Leftrightarrow \nabla_{AA} = \phi_{AA} \Leftrightarrow id_A = \phi_{AA}.
```

```
Lemma empty_equivalence1 \{A : eqType\}: (A \rightarrow False) \leftrightarrow i A =
Proof.
move: (@either\_empty \ i \ A) \Rightarrow H.
split; move \Rightarrow H0.
apply Logic.eq_sym.
apply H.
right.
apply H0.
apply Logic.eq\_sym in H0.
apply H in H0.
case H0.
move \Rightarrow H1 H2.
apply H1.
apply tt.
by [].
Qed.
Lemma empty\_equivalence2 \{A : eqType\}: (A \rightarrow False) \leftrightarrow
                                                                  A A =
                                                                               A A.
Proof.
move: (@either\_empty\ A\ A) \Rightarrow H.
split; move \Rightarrow H0.
apply Logic.eq_sym.
apply H.
left.
apply H0.
apply Logic.eq_sym in H0.
apply H in H\theta.
case H0.
by [].
by [].
Lemma empty_equivalence3 \{A: eqType\}: (A \rightarrow False) \leftrightarrow Id A = AA.
split; move \Rightarrow H.
assert ( AA =
                        A A).
apply empty_equivalence2.
apply H.
apply RelAB\_unique.
apply Logic.eq_sym.
```

```
apply H0.
assert ( AA = AA).
by [rewrite -(@comp\_id\_r\_\_ ( AA)) H comp\_empty\_r].
apply either\_empty in H0.
case H0.
by [].
by [].
Qed.
End main.
```

Chapter 5

Library Functions_Mappings

```
Require Import Basic\_Notations\_Set.
Require Import Basic\_Lemmas.
Require Import Relation\_Properties.
Require Import Logic.FunctionalExtensionality.

Module main (def:Relation).
Import def.
Module Basic\_Lemmas := Basic\_Lemmas.main \ def.
Module Relation\_Properties := Relation\_Properties.main \ def.
Import Basic\_Lemmas \ Relation\_Properties.
```

5.1 全域性,一価性,写像に関する補題

```
Lemma 139 (id_function) id_A: A \rightarrow A is a function.

Lemma id\_function \{A: eqType\}: function\_r \ (Id\ A).

Proof.

rewrite /function\_r/total\_r/univalent\_r.

rewrite inv\_id\ comp\_id\_l.

split.

apply inc\_refl.

apply inc\_refl.

Qed.
```

```
Lemma unit\_function \{A : eqType\}: function\_r (A i). Proof.
```

Lemma 140 (unit_function) $\nabla_{AI}: A \rightarrow I$ is a function.

rewrite $/function_r/total_r/univalent_r$.

```
rewrite inv_universal lemma_for_tarski2 unit_identity_is_universal.
split.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
Qed.
  Lemma 141 (total_comp) Let \alpha: A \rightarrow B and \beta: B \rightarrow C be total relations, then
  \alpha \cdot \beta is also a total relation.
Lemma total\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 total_r \ alpha \rightarrow total_r \ beta \rightarrow total_r \ (alpha \cdot beta).
Proof.
rewrite /total_r.
move \Rightarrow H H0.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply @inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H0.
Qed.
  Lemma 142 (univalent_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be univalent relations,
  then \alpha \cdot \beta is also a univalent relation.
Lemma univalent\_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (alpha \cdot beta).
Proof.
rewrite /univalent_r.
move \Rightarrow H H0.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ (alpha #)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_ab\_b.
apply H.
Qed.
  Lemma 143 (function_comp) Let \alpha: A \to B and \beta: B \to C be functions, then \alpha \cdot \beta
  is also a function.
Lemma function\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 function\_r \ alpha \rightarrow function\_r \ beta \rightarrow function\_r \ (alpha \ \cdot \ beta).
Proof.
elim \Rightarrow H H0.
```

```
elim \Rightarrow H1 H2.
split.
apply (total_comp H H1).
apply (univalent_comp H0 H2).
Qed.
  Lemma 144 (total_comp2) Let \alpha: A \to B, \beta: B \to C and \alpha \cdot \beta be a total relation,
  then \alpha is also a total relation.
Lemma total\_comp2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total_r (alpha \cdot beta) \rightarrow total_r alpha.
Proof.
move \Rightarrow H.
apply inc\_def1 in H.
rewrite comp_inv cap_comm comp_assoc in H.
rewrite /total_r.
rewrite H.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
apply comp\_inc\_compat.
apply cap_{-}l.
rewrite comp_{-}id_{-}r.
apply cap_r.
Qed.
  Lemma 145 (univalent_comp2) Let \alpha: A \rightarrow B, \beta: B \rightarrow C, \alpha \cdot \beta be a univalent
  relation and \alpha^{\sharp} be a total relation, then \beta is a univalent relation.
Lemma univalent\_comp2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r (alpha \cdot beta) \rightarrow total_r (alpha \#) \rightarrow univalent_r beta.
Proof.
move \Rightarrow H H0.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
rewrite /total_{-}r in H0.
rewrite inv_invol in H0.
apply (comp\_inc\_compat\_b\_ab\ H0).
Qed.
  Lemma 146 (total_inc) Let \alpha : A \rightarrow B be a total relation and \alpha \sqsubseteq \beta, then \beta is also
  a total relation.
```

```
total\_r \ alpha \rightarrow alpha
                               beta \rightarrow total_r beta.
Proof.
move \Rightarrow H H0.
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat.
apply H0.
apply (@inc_inv_i - H\theta).
Qed.
  Lemma 147 (univalent_inc) Let \alpha : A \to B be a univalent relation and \beta \sqsubseteq \alpha, then
  \beta is also a univalent relation.
Lemma univalent_inc {A B : eqType} {alpha beta : Rel A B}:
 univalent_r \ alpha \rightarrow beta \qquad alpha \rightarrow univalent_r \ beta.
Proof.
move \Rightarrow H H0.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat.
apply (@inc_inv_{-} - H0).
apply H0.
Qed.
  Lemma 148 (function_inc) Let \alpha, \beta: A \to B be functions and \alpha \sqsubseteq \beta. Then,
                                                \alpha = \beta.
Lemma function\_inc \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 function\_r \ alpha \rightarrow function\_r \ beta \rightarrow alpha
                                                          beta \rightarrow alpha = beta.
Proof.
move \Rightarrow H H0 H1.
apply inc\_antisym.
apply H1.
apply @inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot beta)).
apply comp\_inc\_compat\_b\_ab.
apply H.
move: (@inc_inv_{-1} - H1) \Rightarrow H2.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot beta \#) \cdot beta)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply H2.
rewrite comp_-assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
```

Qed.

```
Lemma 149 (total_universal) If \nabla_{IB} be a total relation, then
```

$$\nabla_{AB} \cdot \nabla_{BC} = \nabla_{AC}.$$

```
Lemma total\_universal \{A \ B \ C : eqType\}:
 total\_r \ (i \ B) \rightarrow A B \cdot
                                   B C =
Proof.
move \Rightarrow H.
rewrite -(@lemma_for_tarski2 A B) -(@lemma_for_tarski2 B C).
rewrite comp\_assoc -(@comp\_assoc _ _ _ ( i B)).
replace ( i B •
                       B i) with (Id i).
rewrite comp_{-}id_{-}l.
apply lemma_for_tarski2.
apply inc\_antisym.
rewrite /total_{-}r in H.
rewrite inv\_universal in H.
apply H.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
```

Lemma 150 (function_rel_inv_rel) Let $\alpha : A \to B$ be function. Then,

$$\alpha \cdot \alpha^{\sharp} \cdot \alpha = \alpha.$$

```
Lemma function_rel_inv_rel {A B : eqType} {alpha : Rel A B}: function_r alpha \rightarrow (alpha • alpha #) • alpha = alpha.

Proof.

move \Rightarrow H.

apply inc\_antisym.

rewrite comp\_assoc.

apply comp\_inc\_compat\_ab\_a.

apply H.

apply comp\_inc\_compat\_b\_ab.

apply H.

Qed.
```

 $(E \to F) \to (B \to C)$ and P: predicate. Then,

```
f \cdot (\sqcap_{P(\theta)} \theta(\alpha)) \cdot q^{\sharp} = \sqcap_{P(\alpha)} (f \cdot \theta(\alpha) \cdot q^{\sharp}).
Lemma function\_capP\_distr {A \ B \ C \ D \ E \ F : eqType}
 \{f: Rel\ A\ B\}\ \{g: Rel\ D\ C\}\ \{theta: Rel\ E\ F \to Rel\ B\ C\}\ \{P: Rel\ E\ F \to Prop\}:
 function\_r \ f \rightarrow function\_r \ g \rightarrow
 (f \cdot ( _{\{P\}} theta)) \cdot g \# =
    _{-}\{P\} (fun alpha : Rel E F \Rightarrow (f • theta alpha) • g \#).
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
apply inc\_antisym.
apply comp\_capP\_distr.
apply (@inc\_trans \_ \_ \_ (((f \cdot f \#) \cdot \_ \{P\} (fun \ alpha : Rel \ E \ F \Rightarrow (f \cdot theta \ alpha)))
• g \# )) • (g • g \# ))).
apply (@inc\_trans \_ \_ \_ ((f \cdot f \#) \cdot ( \_\{P\} (fun \ alpha : Rel \ E \ F \Rightarrow (f \cdot theta \ alpha)))
· g \#)))).
apply (comp\_inc\_compat\_b\_ab\ H).
apply (comp\_inc\_compat\_a\_ab\ H1).
\texttt{rewrite} \ (@ \textit{comp\_assoc} \ \_\_\_\_\_ \ (f \ \#)) \ \textit{comp\_assoc} \ -(@ \textit{comp\_assoc} \ \_\_\_\_\_ \ g) \ -\textit{comp\_assoc}.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc_trans _ _ _ ( _{P} (fun alpha : Rel E F \Rightarrow (f # • ((f • theta alpha) • g
\#)) \cdot g))).
apply comp\_capP\_distr.
replace (fun alpha: Rel E F \Rightarrow (f # \cdot ((f \cdot theta alpha) \cdot g #)) \cdot g) with (fun alpha
: Rel\ E\ F \Rightarrow ((f \# \bullet f) \bullet theta\ alpha) \bullet (g \# \bullet g)).
apply inc\_capP.
move \Rightarrow beta H3.
apply (@inc\_trans\_\_\_((f \# \bullet f) \bullet theta beta)).
apply (@inc\_trans \_ \_ \_ (((f \# \cdot f) \cdot theta beta) \cdot (g \# \cdot g))).
move: beta H3.
apply inc\_capP.
apply inc_refl.
apply (comp\_inc\_compat\_ab\_a\ H2).
apply (comp\_inc\_compat\_ab\_b\ H0).
apply functional\_extensionality.
move \Rightarrow x.
by [rewrite comp_assoc comp_assoc comp_assoc comp_assoc].
Qed.
```

Lemma 151 (function_capP_distr) Let $f: A \to B, g: D \to C$ be functions, $\theta:$

Let $f: A \to B, g: D \to C$ be functions and $\alpha, \beta: B \to C$. Then,

```
f \cdot (\alpha \sqcap \beta) \cdot a^{\sharp} = (f \cdot \alpha \cdot a^{\sharp}) \sqcap (f \cdot \beta \cdot a^{\sharp}).
Lemma function_cap_distr
 \{A \ B \ C \ D : eqType\} \{f : Rel \ A \ B\} \{alpha \ beta : Rel \ B \ C\} \{g : Rel \ D \ C\}:
 function\_r f \rightarrow function\_r g \rightarrow
 (f \cdot (alpha \quad beta)) \cdot g \# = ((f \cdot alpha) \cdot g \#) \quad ((f \cdot beta) \cdot g \#).
Proof.
rewrite (@cap\_to\_capP\_\_\_\_\_id) (@cap\_to\_capP\_\_\_\_\_\_(fun x \Rightarrow (f \cdot x) \cdot g)
apply function\_capP\_distr.
Qed.
Lemma function\_cap\_distr\_l
 \{A \ B \ C : eqType\} \ \{f : Rel \ A \ B\} \ \{alpha \ beta : Rel \ B \ C\}:
 function_r f \rightarrow
 f \cdot (alpha \quad beta) = (f \cdot alpha) \quad (f \cdot beta).
Proof.
move: (@id\_function \ C) \Rightarrow H.
move \Rightarrow H0.
apply (@function\_cap\_distr\_\_\_\_f \ alpha \ beta) in H.
rewrite inv_id comp_id_r comp_id_r comp_id_r in H.
apply H.
apply H0.
Qed.
Lemma function_cap_distr_r
 \{B\ C\ D: eqType\}\ \{alpha\ \mathsf{beta}: Rel\ B\ C\}\ \{g: Rel\ D\ C\}:
 function_r q \rightarrow
             beta) • g \# = (alpha • g \#) (beta • g \#).
 (alpha
Proof.
move: (@id\_function B) \Rightarrow H.
move \Rightarrow H0.
apply (@function\_cap\_distr\_\_\_\_ alpha beta g) in H.
rewrite comp\_id\_l comp\_id\_l comp\_id\_l in H.
apply H.
apply H0.
Qed.
```

Lemma 152 (function_cap_distr, function_cap_distr_l, function_cap_distr_r)

```
Lemma 153 (function_move1) Let \alpha : A \rightarrow B be a function, \beta : B \rightarrow C and
  \gamma: A \rightarrow C. Then,
                                         \gamma \sqsubseteq \alpha \cdot \beta \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.
Lemma function_move1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C} {gamma :
Rel\ A\ C:
 function\_r \ alpha \rightarrow (gamma \ (alpha \cdot beta) \leftrightarrow (alpha \# \cdot gamma)
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply @inc\_trans \_ \_ \_ ((alpha \# \cdot alpha) \cdot beta)).
rewrite comp_assoc.
apply (comp\_inc\_compat\_ab\_ab' H0).
apply comp\_inc\_compat\_ab\_b.
apply H.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot gamma)).
apply comp\_inc\_compat\_b\_ab.
apply H.
rewrite comp_{-}assoc.
apply (comp\_inc\_compat\_ab\_ab' H0).
Qed.
  Lemma 154 (function_move2) Let \beta: B \rightarrow C be a function, \alpha: A \rightarrow B and
  \gamma: A \to C. Then,
                                         \alpha \cdot \beta \sqsubseteq \gamma \Leftrightarrow \alpha \sqsubseteq \gamma \cdot \beta^{\sharp}.
Lemma function_move2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C} {qamma :
Rel\ A\ C:
 function\_r \ \mathsf{beta} \to ((alpha \cdot \mathsf{beta}) \ gamma \leftrightarrow alpha
                                                                         (gamma \cdot beta \#)).
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot beta) \cdot beta \#)).
rewrite comp\_assoc.
apply comp\_inc\_compat\_a\_ab.
apply H.
apply (comp\_inc\_compat\_ab\_a'b\ H0).
apply (@inc\_trans \_ \_ \_ ((gamma \cdot beta \#) \cdot beta)).
apply (comp\_inc\_compat\_ab\_a'b H0).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply H.
```

Qed.

 $\alpha, \beta: B \rightarrow C$. Then,

```
f \cdot (\alpha \Rightarrow \beta) \cdot q^{\sharp} = (f \cdot \alpha \cdot q^{\sharp}) \Rightarrow (f \cdot \beta \cdot q^{\sharp}).
Lemma function\_rpc\_distr
 \{A \ B \ C \ D : eqType\} \{f : Rel \ A \ B\} \{alpha \ beta : Rel \ B \ C\} \{g : Rel \ D \ C\}:
 function\_r \ f \rightarrow function\_r \ g \rightarrow
 (f \cdot (alpha \otimes beta)) \cdot q \# = ((f \cdot alpha) \cdot q \#) \otimes ((f \cdot beta) \cdot q \#).
Proof.
move \Rightarrow H H0.
apply inc_lower.
move \Rightarrow qamma.
split; move \Rightarrow H1.
apply inc_rpc.
apply (function_move2 H0).
apply (function\_move1 \ H).
apply (@inc\_trans\_\_\_(((f \# \cdot gamma) \cdot g) \ ((f \# \cdot ((f \cdot alpha) \cdot g \#)) \cdot g))).
rewrite -comp\_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (@comp\_cap\_distr\_r \_ \_ \_ \_ \_)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_cap\_distr\_l.
apply (function\_move2 \ H0) in H1.
apply (function\_move1 \ H) in H1.
rewrite -inc\_rpc\ comp\_assoc.
apply @inc\_trans \_ \_ \_ \_ H1).
apply rpc\_inc\_compat\_r.
rewrite comp_assoc comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ (alpha \cdot (g \# \cdot g))).
apply comp\_inc\_compat\_ab\_b.
apply H.
apply comp\_inc\_compat\_ab\_a.
apply H0.
apply (function_move2 H0).
apply (function\_move1 \ H).
apply inc\_rpc.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
apply (@inc\_trans \_ \_ \_ (f \# \cdot ((gamma \cdot g) ((f \#) \# \cdot alpha)))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_l.
rewrite inv_invol.
```

Lemma 155 (function_rpc_distr) Let $f: A \rightarrow B, q: D \rightarrow C$ be functions and

```
apply @inc\_trans\_\_\_((f \# \cdot (gamma ((f \cdot alpha) \cdot g \#))) \cdot g)).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
apply (function_move2 H0).
apply (function\_move1 \ H).
rewrite -inc_rpc -comp_assoc.
apply H1.
Qed.
  Lemma 156 (function_inv_rel1, function_inv_rel2) Let f: A \to B be a function.
  Then,
                         f^{\sharp} \cdot f = id_B \sqcap f^{\sharp} \cdot \nabla_{AA} \cdot f = id_B \sqcap \nabla_{BA} \cdot f.
Lemma function\_inv\_rel1 \{A B : eqType\} \{f : Rel A B\}:
 function\_r f \to f \# \cdot f = Id B \quad ((f \# \cdot A A) \cdot f).
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc_-cap.
split.
apply H.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_a\_ab.
apply inc\_alpha\_universal.
                                    (BA \cdot f)).
apply (@inc_trans _ _ _ (Id B
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite cap\_comm.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite comp_id_l comp_id_r cap_comm inv_universal.
rewrite cap_universal cap_universal.
apply inc\_reft.
Qed.
Lemma function\_inv\_rel2 {A B : eqType} {f : Rel A B}:
function\_r f \rightarrow f \# \cdot f = Id B \quad (BA \cdot f).
Proof.
move \Rightarrow H.
apply inc\_antisym.
```

```
rewrite (@function_inv_rel1 _ _ _ H).
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite cap\_comm.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite comp_id_l comp_id_r cap_comm inv_universal.
rewrite cap_universal cap_universal.
apply inc\_reft.
Qed.
  Lemma 157 (function_dedekind1, function_dedekind2) Let f: A \rightarrow B be a
  function, \mu: C \to A and \rho: C \to B. Then,
                       (\mu \sqcap \rho \cdot f^{\sharp}) \cdot f = \mu \cdot f \sqcap \rho \wedge \rho \cdot f^{\sharp} \cdot f = \nabla_{CA} \cdot f \sqcap \rho.
Lemma function_dedekind1
 \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{mu : Rel \ C \ A\} \{rho : Rel \ C \ B\}:
 function_r f \rightarrow (mu \quad (rho \cdot f \#)) \cdot f = (mu \cdot f)
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
Qed.
Lemma function_dedekind2 \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{rho : Rel \ C \ B\}:
 function\_r f \rightarrow (rho \cdot f \#) \cdot f = (CA \cdot f) rho.
Proof.
move \Rightarrow H.
move: (@function\_dedekind1 \_ \_ \_ f ( CA) rho H) \Rightarrow H0.
rewrite cap_comm cap_universal in H0.
apply H0.
Qed.
```

Lemma 158 (square_diagram) In below figure, $f \cdot x = g \cdot y \Leftrightarrow f^{\sharp} \cdot g \sqsubseteq x \cdot y^{\sharp}.$ $X \xrightarrow{f} A$ $\downarrow x$ $B \xrightarrow{y} D$

```
Lemma square\_diagram \{X \ A \ B \ D : eqType\}
 \{f : Rel \ X \ A\} \{g : Rel \ X \ B\} \{x : Rel \ A \ D\} \{y : Rel \ B \ D\}:
 function\_r \ f \rightarrow function\_r \ g \rightarrow function\_r \ x \rightarrow function\_r \ y \rightarrow
 (f \cdot x = g \cdot y \leftrightarrow (f \# \cdot g) \quad (x \cdot y \#)).
Proof.
move \Rightarrow H H0 H1 H2.
split; move \Rightarrow H3.
rewrite -(function_move1 H) -comp_assoc -(function_move2 H2) H3.
apply inc\_reft.
apply Logic.eq_sym.
apply function_inc.
apply (function_comp H0 H2).
apply (function_comp H H1).
rewrite (function_move2 H2) comp_assoc (function_move1 H).
apply H3.
Qed.
```

5.2 全射, 単射に関する補題

Lemma 159 (surjection_comp) Let $\alpha : A \to B$ and $\beta : B \to C$ be surjections, then $\alpha \cdot \beta$ is also a surjection.

```
Lemma surjection\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}: surjection\_r \ alpha \rightarrow surjection\_r \ beta \rightarrow surjection\_r \ (alpha \cdot beta).

Proof.

rewrite /surjection\_r.

elim \Rightarrow H \ H0.

elim \Rightarrow H1 \ H2.

split.

apply (function\_comp \ H \ H1).

rewrite comp\_inv.

apply (total\_comp \ H2 \ H0).
```

Qed.

```
Lemma 160 (injection_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be injections, then
  \alpha \cdot \beta is also an injection.
Lemma injection\_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 injection\_r \ alpha \rightarrow injection\_r \ beta \rightarrow injection\_r \ (alpha \cdot beta).
Proof.
rewrite /injection_r.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (function_comp H H1).
rewrite comp_{-}inv.
apply (univalent_comp H2 H0).
Qed.
  Lemma 161 (bijection_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be bijections, then
  \alpha \cdot \beta is also a bijection.
Lemma bijection\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 bijection\_r \ alpha \rightarrow bijection\_r \ beta \rightarrow bijection\_r \ (alpha \cdot beta).
Proof.
rewrite /bijection_r.
elim \Rightarrow H.
elim \Rightarrow H0 \ H1.
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
split.
apply (function\_comp\ H\ H2).
rewrite comp_{-}inv.
split.
apply (total\_comp\ H3\ H0).
apply (univalent_comp H4 H1).
Qed.
  Lemma 162 (surjection_unique1) Let e: A \rightarrow B be a surjection, f: A \rightarrow C be a
  function and e \cdot e^{\sharp} \sqsubseteq f \cdot f^{\sharp}, then there exists a unique function g : B \to C s.t. f = eg.
Lemma surjection\_unique1 \{A \ B \ C : eqType\} \{e : Rel \ A \ B\} \{f : Rel \ A \ C\}:
 surjection\_r \ e \rightarrow function\_r \ f \rightarrow (e \ \cdot \ e \ \#)
                                                            (f \cdot f \#) \rightarrow
 (\exists ! \ g : Rel \ B \ C, function\_r \ g \land f = e \cdot g).
Proof.
```

```
rewrite /surjection_r/function_r/total_r/univalent_r.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 \ H3 \ H4.
\exists (e \# \cdot f).
repeat split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ f).
apply @inc\_trans \_ \_ \_ \_ H1).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H2.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
apply (@inc\_trans\_\_\_(f \# \cdot ((f \cdot f \#) \cdot f))).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_a'b H_4).
rewrite comp\_assoc -comp\_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H3).
apply function_inc.
split.
apply H2.
apply H3.
split.
rewrite /total_r.
rewrite comp_inv comp_inv inv_invol.
rewrite -(@comp_assoc _ _ _ e) (@comp_assoc _ _ _ e) (@comp_assoc _ _ _ f)
-(@comp\_assoc\_\_\_f).
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_a\_ab.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply (comp\_inc\_compat\_a\_ab\ H).
rewrite /univalent_r.
rewrite comp_inv comp_inv inv_invol.
 \textbf{rewrite} \ (@comp\_assoc \_\_\_\_ e) \ - (@comp\_assoc \_\_\_\_ e) \ comp\_assoc \ - (@comp\_assoc - e) \ comp\_assoc \ - (@comp\_
- - - - f).
apply (@inc\_trans \_ \_ \_ (f \# \cdot (((f \cdot f \#) \cdot (f \cdot f \#)) \cdot f))).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat.
apply H_4.
apply H_4.
rewrite comp\_assoc (@comp\_assoc _ _ _ _ f) -(@comp\_assoc _ _ _ _ (f \#)) -(@comp\_assoc
```

```
---(f \#)) (@comp\_assoc\_--(f \#)) - (@comp\_assoc\_--(f \#)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H3).
rewrite -comp_-assoc.
apply (comp\_inc\_compat\_b\_ab\ H).
move \Rightarrow q.
elim.
elim \Rightarrow H5 H6 H7.
replace g with (e \# \cdot (e \cdot g)).
apply f_equal.
apply H\gamma.
rewrite - comp_assoc.
apply inc\_antisym.
apply (comp\_inc\_compat\_ab\_b\ H0).
rewrite inv_{-}invol in H1.
apply (comp\_inc\_compat\_b\_ab\ H1).
Qed.
  Lemma 163 (surjection_unique2) Let e: A \rightarrow B be a surjection, f: A \rightarrow C be a
  function and e \cdot e^{\sharp} = f \cdot f^{\sharp}, then function e^{\sharp} f is an injection.
Lemma surjection\_unique2 \{A \ B \ C : eqType\} \{e : Rel \ A \ B\} \{f : Rel \ A \ C\}:
 surjection\_r \ e \rightarrow function\_r \ f \rightarrow (e \cdot e \#) = (f \cdot f \#) \rightarrow injection\_r \ (e \# \cdot f).
Proof.
rewrite /surjection_r/injection_r/function_r/total_r/univalent_r.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 \ H3 \ H4.
repeat split.
rewrite comp\_inv \ comp\_assoc \ -(@comp\_assoc \ \_ \ \_ \ \_ \ f).
apply (@inc\_trans \_ \_ \_ \_ H1).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H2.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
rewrite H_4.
rewrite comp_assoc -comp_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H3).
rewrite inv_invol comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
rewrite -H4.
```

```
rewrite comp_assoc -comp_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp\_inc\_compat\_ab\_a.
apply H0.
Qed.
  Lemma 164 (injection_unique1) Let m: B \rightarrow A be an injection, f: C \rightarrow A be a
  function and f^{\sharp} \cdot f \sqsubseteq m^{\sharp} \cdot m, then there exists a unique function g: C \to B s.t. f = gm.
Lemma injection_unique1 \{A \ B \ C : eqType\} \{m : Rel \ B \ A\} \{f : Rel \ C \ A\}:
 injection\_r \ m \rightarrow function\_r \ f \rightarrow (f \ \# \ \cdot \ f)
                                                     (m \# \cdot m) \rightarrow
 (\exists ! \ g : Rel \ C \ B, function\_r \ g \land f = g \cdot m).
Proof.
rewrite /injection_r/function_r/total_r/univalent_r.
elim.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 H3 H4.
\exists (f \cdot m \#).
repeat split.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ m).
apply (@inc\_trans\_\_\_(f \cdot ((f \# \cdot f) \cdot f \#))).
rewrite comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply (comp\_inc\_compat\_a\_ab H2).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_a'b\ H4).
rewrite comp\_inv \ comp\_assoc \ -(@comp\_assoc \ \_ \ \_ \ \_ \ f).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
rewrite comp\_assoc.
apply Logic.eq_sym.
apply function_inc.
split.
rewrite /total_{-}r.
rewrite comp_inv comp_inv inv_invol.
apply @inc\_trans \_ \_ \_ \_ H2).
apply comp\_inc\_compat.
apply (@inc\_trans\_\_\_(f \cdot (f \# \cdot f))).
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_b\_ab\ H2).
apply (comp_inc_compat_ab_ab' H4).
apply (@inc\_trans \_ \_ \_ ((f \# \cdot f) \cdot f \#)).
```

```
rewrite comp_assoc.
apply (comp\_inc\_compat\_a\_ab\ H2).
apply (comp\_inc\_compat\_ab\_a'b H_4).
rewrite /univalent_r.
rewrite comp_inv comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H0).
split.
apply H2.
apply H3.
apply (comp\_inc\_compat\_ab\_a\ H0).
move \Rightarrow q.
elim.
elim \Rightarrow H5 \ H6 \ H7.
rewrite H7 comp\_assoc.
apply inc\_antisym.
rewrite inv_{-}invol in H1.
apply (comp\_inc\_compat\_ab\_a\ H1).
apply (comp\_inc\_compat\_a\_ab\ H).
Qed.
  Lemma 165 (injection_unique2) Let m: B \rightarrow A be an injection, f: C \rightarrow A be a
  function and f^{\sharp} \cdot f = m^{\sharp} \cdot m, then function f \cdot m^{\sharp} is a surjection.
Lemma injection\_unique2 \{A \ B \ C : eqType\} \{m : Rel \ B \ A\} \{f : Rel \ C \ A\}:
 injection\_r \ m \to function\_r \ f \to (f \# \bullet f) = (m \# \bullet m) \to surjection\_r \ (f \bullet m \#).
Proof.
rewrite /surjection_r/injection_r/function_r/total_r/univalent_r.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 \ H3 \ H4.
repeat split.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ m).
apply (@inc\_trans\_\_\_(f \cdot ((f \# \cdot f) \cdot f \#))).
rewrite comp\_assoc -comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply (comp\_inc\_compat\_a\_ab H2).
apply comp\_inc\_compat\_ab\_ab'.
rewrite H_4.
apply inc\_reft.
rewrite comp\_inv \ comp\_assoc \ -(@comp\_assoc \ \_ \ \_ \ \_ \ f).
```

```
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
rewrite inv_invol comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_ab\_ab'.
rewrite H4 comp_assoc.
apply (comp\_inc\_compat\_a\_ab\ H).
Qed.
  Lemma 166 (bijection_inv) Let \alpha: A \to B, \beta: B \to A, \alpha \cdot \beta = id_A and \beta \cdot \alpha = id_B,
  then \alpha and \beta are bijections and \beta = \alpha^{\sharp}.
Lemma bijection_inv {A B : eqType} {alpha : Rel A B} {beta : Rel B A}:
 alpha • beta = Id\ A \rightarrow beta • alpha = Id\ B \rightarrow bijection\_r\ alpha \land bijection\_r\ beta \land
beta = alpha \#.
Proof.
move \Rightarrow H H0.
move: (@id\_function A) \Rightarrow H1.
move: (@id_function B) \Rightarrow H2.
assert (bijection\_r \ alpha \land bijection\_r \ beta).
assert (total_r \ alpha \wedge total_r \ (alpha \#) \wedge total_r \ beta \wedge total_r \ (beta \#)).
repeat split.
apply (@total\_comp2 \_ \_ \_ \_ beta).
rewrite H.
apply H1.
apply (@total\_comp2\_\_\_\_(beta \#)).
rewrite -comp_inv H0 inv_id.
apply H2.
apply (@total\_comp2\_\_\_\_alpha).
rewrite H0.
apply H2.
apply (@total\_comp2\_\_\_\_ (alpha \#)).
rewrite -comp_inv H inv_id.
apply H1.
repeat split.
apply H3.
apply (@univalent_comp2 _ _ _ beta).
rewrite H0.
apply H2.
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ (beta \#)).
```

CHAPTER 5. LIBRARY FUNCTIONS_MAPPINGS

```
rewrite -comp_inv H inv_id.
apply H1.
rewrite inv_-invol.
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ alpha).
rewrite H.
apply H1.
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ (alpha \#)).
rewrite -comp_inv H0 inv_id.
apply H2.
rewrite inv_invol.
apply H3.
split.
apply H3.
split.
apply H3.
rewrite -(@comp\_id\_r\_\_\_beta) -(@comp\_id\_l\_\_\_(alpha \#)).
rewrite -H0 comp_assoc.
apply f_equal.
apply inc\_antisym.
apply H3.
rewrite comp_inv_inv -inv_inc_move inv_id.
apply H3.
Qed.
  Lemma 167 (bijection_inv_corollary) Let \alpha : A \to B be a bijection, then \alpha^{\sharp} is also
  a bijection.
Lemma bijection_inv_corollary \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 bijection\_r \ alpha \rightarrow bijection\_r \ (alpha \#).
Proof.
move: (@bijection\_inv \_ \_ alpha (alpha \#)) \Rightarrow H.
move \Rightarrow H0.
rewrite /bijection\_r/function\_r/total\_r/univalent\_r in H0.
rewrite inv_{-}invol in H0.
apply H.
apply inc\_antisym.
apply H0.
apply H0.
```

```
apply inc\_antisym. apply H0. apply H0. Qed.
```

5.3 有理性から導かれる系

```
Lemma 168 (rationality_corollary1) Let u: A \to A and u \sqsubseteq id_A. Then, \exists R, \exists j: R \rightarrowtail A, u = j^{\sharp} \cdot j.
```

```
Lemma rationality_corollary1 \{A : eqType\} \{u : Rel A A\}:
       Id A \to \exists (R : eqType)(j : Rel R A), injection_r j \land u = j \# \bullet j.
Proof.
move: (rationality \_ \_ u).
elim \Rightarrow R.
elim \Rightarrow f.
elim \Rightarrow q.
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2 H3.
\exists R.
\exists f.
assert (g = f).
apply (function\_inc\ H0\ H).
apply (@inc\_trans \_ \_ \_ ((f \cdot f \#) \cdot g)).
apply comp\_inc\_compat\_b\_ab.
apply H.
rewrite comp_assoc -H1.
apply (comp\_inc\_compat\_ab\_a\ H3).
rewrite H4 in H1.
rewrite H_4 cap_idem in H_2.
split.
split.
apply H.
rewrite /univalent_r.
rewrite inv_invol H2.
apply inc\_reft.
apply H1.
Qed.
```

CHAPTER 5. LIBRARY FUNCTIONS_MAPPINGS

Lemma 169 (rationality_corollary2) Let $f: A \to B$ be a function. Then,

```
\exists e: A \rightarrow R, \exists m: R \rightarrow B, f = e \cdot m.
Lemma rationality\_corollary2 {A B : eqType} {f : Rel A B}:
 function_r f \to \exists (R : eqType)(e : Rel A R)(m : Rel R B), surjection_r e \land injection_r
m.
Proof.
elim \Rightarrow H H0.
move: (@rationality\_corollary1\_(f \# \bullet f) H0).
elim \Rightarrow R.
elim \Rightarrow m.
elim \Rightarrow H1 H2.
\exists R.
\exists (f \cdot m \#).
\exists m.
split.
apply (injection_unique2 H1 (conj H H0) H2).
apply H1.
Qed.
  Lemma 170 (axiom_of_subobjects) Let u: A \rightarrow A and u \subseteq id_A. Then,
                                \exists R, \exists j : R \to A, j^{\sharp} \cdot j = u \land j \cdot j^{\sharp} = id_R.
Lemma axiom\_of\_subobjects \{A : eqType\} \{u : Rel A A\}:
        Id A \rightarrow \exists (R : eqType)(j : Rel R A), j \# \bullet j = u \land j \bullet j \# = Id R.
 u
Proof.
move \Rightarrow H.
elim(rationality\_corollary1\ H) \Rightarrow R.
elim \Rightarrow j H0.
\exists R.
\exists j.
split.
apply Logic.eq_sym.
apply H0.
apply inc\_antisym.
replace (j \cdot j \#) with ((j \#) \# \cdot j \#).
apply H0.
by [rewrite inv\_invol].
apply H0.
```

Qed.

CHAPTER 5. LIBRARY FUNCTIONS_MAPPINGS

End main.

Chapter 6

Library Tactics

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Module main (def:Relation).

Import def.

Module Basic\_Lemmas := Basic\_Lemmas.main \ def.

Module Relation\_Properties := Relation\_Properties.main \ def.

Import Basic\_Lemmas \ Relation\_Properties.
```

6.1 Tactic 用の補題

```
\alpha = \beta の形では自動計算がしづらいので、事前に \alpha \sqsubseteq \beta \land \beta \sqsubseteq \alpha の形に変換しておく.
```

```
Lemma inc\_antisym\_eq {A \ B : eqType} {alpha \ beta : Rel \ A \ B}: alpha = beta \leftrightarrow alpha \ beta \land beta \ alpha.

Proof. split; move \Rightarrow H. rewrite H. split; apply inc\_refl. apply inc\_antisym; apply H. Qed.
```

6.2 Tactic

```
ここでは以下の 5 tactics を実装している.
```

- Rel_simpl_rewrite ... 関数などの定義の書き換え
- Rel_simpl_intro ... 命題間の関係の整理, inc_antisym_eq の書き換え
- Rel_simpl_comp_inc ... comp_inc_compat 関連の補題の適用
- Rel_simpl ... 証明のための各種動作, 上記 3 tactics を全て含む
- Rel_trans ... Rel_simpl に inc_trans を組み込んだもの, 引数が必要

```
Ltac Rel_simpl_rewrite :=
 rewrite /bijection_r/surjection_r/injection_r;
 rewrite /function_r/total_r/univalent_r.
Ltac Rel_simpl_intro :=
 Rel\_simpl\_rewrite;
 repeat match goal with
                |[\_:\_\vdash(\_\land\_)\rightarrow\_]\Rightarrow elim
                | [\_: \_ \vdash \_ \rightarrow \_] \Rightarrow intro
                |[\_:\_\vdash\_\land\_]\Rightarrow split
                |[\_:\_\vdash\_\leftrightarrow\_]\Rightarrow split
                | [\_: \_ \vdash \_ = \_] \Rightarrow rewrite inc\_antisym\_eq
                | [H : \_ = \_ \vdash \_] \Rightarrow \text{rewrite } inc\_antisym\_eq \text{ in } H
             end.
Ltac Rel_simpl_comp_inc :=
 repeat match goal with
                 \begin{array}{ll} \mid \left[ \begin{array}{ccc} H:?g & Id \_ \vdash (?f & \cdot ?g) & ?f \end{array} \right] \Rightarrow \operatorname{apply} \left( comp\_inc\_compat\_ab\_a \ H \right) \\ \mid \left[ \begin{array}{ccc} H:?g & Id \_ \vdash (?g & \cdot ?f) & ?f \end{array} \right] \Rightarrow \operatorname{apply} \left( comp\_inc\_compat\_ab\_b \ H \right) \\ \end{array} 
                |[H:Id\_]?g\vdash?f (?f\cdot?g)]\Rightarrow \operatorname{apply}(comp\_inc\_compat\_a\_ab\ H)
                | [H: Id \_ ?g \vdash ?f (?g \cdot ?f)] \Rightarrow apply (comp\_inc\_compat\_b\_ab H)
                | [\_: \_ \vdash \_] \Rightarrow \text{repeat rewrite -} comp\_assoc; (apply comp\_inc\_compat\_ab\_a'b
|| apply comp_inc_compat_b_ab || apply comp_inc_compat_ab_b)
                | [ \_ : \_ \vdash \_ ] \Rightarrow \text{repeat rewrite } comp\_assoc; (apply <math>comp\_inc\_compat\_ab\_ab'
|| apply comp_inc_compat_a_ab || apply comp_inc_compat_ab_a)
                | [ \_ : \_ \vdash (\_ \cdot \_) \quad (\_ \cdot \_) | \Rightarrow apply comp\_inc\_compat
             end.
Ltac Rel\_simpl :=
 Rel\_simpl\_intro;
 repeat match goal with
                | [f: Rel\_, H:\_ \vdash\_] \Rightarrow \text{rewrite} (@inv\_invol\_\_f) \text{ in } H
             end;
```

```
repeat match goal with
                                ?f \mid \Rightarrow apply inc\_reft
             | [\_: \_ \vdash ?f]
             \mid [H:?P \vdash ?P] \Rightarrow \text{apply } H
             | [H:?f] ?q, H0:?q
                                             ?h \vdash ?f
                                                             ?h \Rightarrow apply (@inc\_trans \_ \_ \_ \_ H)
H0)
             [ [ : ] \vdash ( \_ \# ) \quad ( \_ \# ) ] \Rightarrow apply inc\_inv
             | [A:eqType, B:eqType, C:eqType \vdash \_] \Rightarrow rewrite (@comp\_inv A B C)
             | [f : Rel \_ \_ \vdash \_] \Rightarrow rewrite (@inv\_invol \_ \_ f)
             | [H:(Id\_) \_ \vdash (Id\_) \_| \Rightarrow apply (@inc\_trans\_\_\_\_H)
             | [H:\_ (Id\_) \vdash \_ (Id\_) | \Rightarrow apply (fun H' \Rightarrow (@inc\_trans\_\_\_\_H') |
H))
             [ [ : \_ \vdash \_ ] \Rightarrow Rel\_simpl\_comp\_inc
           end.
Ltac Rel_trans f :=
 apply @inc\_trans \_ \_ _ f); Rel\_simpl.
```

6.3 実験

Functions_Mappings.v の補題には、単一の tactic のみで解けるものも多い.

```
Lemma total\_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total_r \ alpha \rightarrow total_r \ beta \rightarrow total_r \ (alpha \cdot beta).
Proof.
Rel\_simpl.
Qed.
Lemma univalent_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (alpha \cdot beta).
Proof.
Rel\_simpl.
Qed.
Lemma function\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
function\_r \ alpha \rightarrow function\_r \ beta \rightarrow function\_r \ (alpha \cdot beta).
Proof.
Rel\_simpl.
Qed.
Lemma univalent\_comp2 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 univalent_r (alpha \cdot beta) \rightarrow total_r (alpha \#) \rightarrow univalent_r beta.
Proof.
Rel\_simpl.
```

CHAPTER 6. LIBRARY TACTICS

```
Qed.
Lemma total\_inc {A B : eqType} {alpha beta : Rel A B}:
 total\_r \ alpha \rightarrow alpha \quad beta \rightarrow total\_r \ beta.
Proof.
Rel\_simpl.
Qed.
Lemma univalent\_inc \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 univalent_r \ alpha \rightarrow beta \qquad alpha \rightarrow univalent_r \ beta.
Proof.
Rel\_simpl.
Qed.
Lemma function\_inc \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
function\_r \ alpha \rightarrow function\_r \ \mathtt{beta} \rightarrow alpha
                                                            beta \rightarrow alpha = beta.
Proof.
Rel\_simpl.
Rel\_trans\ ((alpha \cdot alpha \#) \cdot beta).
Qed.
Lemma function_move1 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Beta : Rel \ B \ C\} \}
Rel\ A\ C:
                                      (alpha \cdot beta) \leftrightarrow (alpha \# \cdot gamma)
 function_r \ alpha \rightarrow (gamma)
Proof.
Rel\_simpl.
Rel\_trans ((alpha \# \bullet alpha) \bullet beta).
Rel\_trans\ ((alpha \cdot alpha \#) \cdot gamma).
Qed.
Lemma function_move2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C} {qamma :
Rel\ A\ C:
 function\_r \text{ beta} \rightarrow ((alpha \cdot \text{beta})
                                                 gamma \leftrightarrow alpha
                                                                          (gamma \cdot beta \#)).
Proof.
Rel\_simpl.
Rel\_trans ((alpha \cdot beta) · beta #).
Rel\_trans ((qamma \cdot beta \#) · beta).
Qed.
Lemma surjection\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 surjection\_r \ alpha \rightarrow surjection\_r \ beta \rightarrow surjection\_r \ (alpha \cdot beta).
Proof.
Rel\_simpl.
Qed.
Lemma injection_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 injection\_r \ alpha \rightarrow injection\_r \ beta \rightarrow injection\_r \ (alpha \cdot beta).
Proof.
```

CHAPTER 6. LIBRARY TACTICS

```
 \begin{array}{l} \textit{Rel\_simpl.} \\ \textit{Qed.} \\ \\ \textit{Lemma bijection\_comp} \; \{A \; B \; C : eqType\} \; \{alpha : Rel \; A \; B\} \; \{\texttt{beta} : Rel \; B \; C\} : \\ & \textit{bijection\_r alpha} \; \rightarrow \; \textit{bijection\_r beta} \; \rightarrow \; \textit{bijection\_r} \; (alpha \; \cdot \; \texttt{beta}). \\ \\ \textit{Proof.} \\ & \textit{Rel\_simpl.} \\ \\ \textit{Qed.} \\ \\ \textit{Lemma bijection\_inv\_corollary} \; \{A \; B : eqType\} \; \{alpha : Rel \; A \; B\} : \\ & \textit{bijection\_r alpha} \; \rightarrow \; \textit{bijection\_r} \; (alpha \; \#). \\ \\ \textit{Proof.} \\ & \textit{Rel\_simpl.} \\ \\ \textit{Qed.} \\ \\ \textit{End } \; \textit{main.} \\ \end{array}
```

Chapter 7

Qed.

Library Dedekind

```
Require Import Basic\_Notations\_Set.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Module main\ (def:Relation).

Import def.

Module Basic\_Lemmas := Basic\_Lemmas.main\ def.

Module Relation\_Properties := Relation\_Properties.main\ def.

Module Functions\_Mappings := Functions\_Mappings.main\ def.

Import Basic\_Lemmas\ Relation\_Properties\ Functions\_Mappings.
```

7.1 Dedekind formula に関する補題

```
Lemma dedekind1 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C} {gamma : Rel \ A \ C}: ((alpha \cdot beta) gamma) (alpha \cdot (beta \ (alpha \# \cdot gamma))).

Proof.

apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_)).

apply comp\_inc\_compat\_ab\_a'b.

apply cap\_l.
```

Lemma 171 (dedekind1) Let $\alpha: A \to B$, $\beta: B \to C$ and $\gamma: A \to C$. Then

```
Lemma 172 (dedekind2) Let \alpha: A \to B, \beta: B \to C and \gamma: A \to C. Then \alpha \cdot \beta \sqcap \gamma \sqsubseteq (\alpha \sqcap \gamma \cdot \beta^{\sharp}) \cdot \beta.
```

```
Lemma dedekind2 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C} {gamma : Rel \ A \ C}: ((alpha \cdot beta) gamma) ((alpha \quad (gamma \cdot beta \#)) beta). Proof. apply (@inc\_trans\_\_\_\_ (@dedekind\_\_\_\_\_)). apply comp\_inc\_compat\_ab\_ab'. apply cap\_l. Qed.
```

Lemma 173 (relation_rel_inv_rel) Let $\alpha : A \rightarrow B$. Then

$$\alpha \sqsubseteq \alpha \cdot \alpha^{\sharp} \cdot \alpha$$
.

```
Lemma relation\_rel\_inv\_rel {A \ B : eqType} {alpha : Rel \ A \ B}: alpha ((alpha \cdot alpha \#) • alpha).

Proof.

move : (@dedekind1 \_ \_ alpha \ (Id \ B) \ alpha) \Rightarrow H.

rewrite comp\_id\_r \ cap\_idem \ in \ H.

apply (@inc\_trans \_ \_ \_ \_ H).

rewrite comp\_assoc.

apply comp\_inc\_compat\_ab\_ab'.

apply cap\_r.

Qed.
```

7.2 Dedekind formula と全関係

```
Lemma 174 (dedekind_universal1) Let \alpha : B \rightarrow C. Then
```

$$\nabla_{AC} \cdot \alpha^{\sharp} \cdot \alpha = \nabla_{AB} \cdot \alpha.$$

```
Lemma dedekind\_universal1 {A \ B \ C : eqType} {alpha : Rel \ B \ C}: ( A \ C \cdot alpha \ \#) • alpha = A \ B \cdot alpha.

Proof.

apply inc\_antisym.

apply comp\_inc\_compat\_ab\_a'b.

apply inc\_alpha\_universal.

apply (@inc\_trans\_\_\_ ( A \ B \cdot ((alpha \cdot alpha \ \#) \cdot alpha))).
```

```
apply comp_inc_compat_ab_ab'.
apply relation_rel_inv_rel.
rewrite -comp\_assoc -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
  Lemma 175 (dedekind_universal2a, dedekind_universal2b,
  dedekind_universal2c) Let \alpha : A \rightarrow B and \beta : C \rightarrow B. Then
                   \nabla_{IC} \cdot \beta \sqsubseteq \nabla_{IA} \cdot \alpha \Leftrightarrow \nabla_{CC} \cdot \beta \sqsubseteq \nabla_{CA} \cdot \alpha \Leftrightarrow \beta \sqsubseteq \beta \cdot \alpha^{\sharp} \cdot \alpha.
Lemma dedekind\_universal2a \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ C \ B\}:
 (i \ C \cdot beta) \ (i \ A \cdot alpha) \rightarrow (C \ C \cdot beta) \ (C \ A \cdot alpha).
Proof.
move \Rightarrow H.
rewrite -unit_universal -(@lemma_for_tarski2 C A).
rewrite comp_assoc comp_assoc.
apply (comp\_inc\_compat\_ab\_ab', H).
Qed.
Lemma dedekind_universal2b {A B C : eqType} {alpha : Rel A B} {beta : Rel C B}:
 (CC \cdot beta) \quad (CA \cdot alpha) \rightarrow beta \quad ((beta \cdot alpha \#) \cdot alpha).
Proof.
move \Rightarrow H.
apply (@inc\_trans\_\_\_ (beta)).
apply inc_{-}cap.
split.
apply inc\_reft.
apply comp\_inc\_compat\_b\_ab.
apply inc\_alpha\_universal.
                                       (CA \cdot alpha)).
apply (@inc_trans _ _ _ (beta
apply (cap\_inc\_compat\_l\ H).
rewrite cap\_comm.
apply (@inc\_trans\_\_\_\_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_r.
Qed.
Lemma dedekind\_universal2c {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ C \ B}:
           ((beta \cdot alpha \#) \cdot alpha) \rightarrow (i \ C \cdot beta) \ (i \ A \cdot alpha).
 beta
Proof.
move \Rightarrow H.
```

```
apply (@inc\_trans \_ \_ \_ ( i C \cdot ((beta \cdot alpha \#) \cdot alpha))).
apply (comp\_inc\_compat\_ab\_ab' H).
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
  Lemma 176 (dedekind_universal3a, dedekind_universal3b) Let \alpha:A \rightarrow B and
  \beta: A \rightarrow C. Then
                   \beta \cdot \nabla_{CI} \sqsubseteq \alpha \cdot \nabla_{BI} \Leftrightarrow \beta \cdot \nabla_{CC} \sqsubseteq \alpha \cdot \nabla_{BC} \Leftrightarrow \beta \sqsubseteq \alpha \cdot \alpha^{\sharp} \cdot \beta.
Lemma dedekind\_universal3a {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ A \ C}:
                        (alpha \cdot B i) \leftrightarrow (beta \cdot C C) \quad (alpha \cdot C C)
               C(i)
Proof.
split; move \Rightarrow H.
apply inv_inc_invol.
rewrite comp_inv comp_inv inv_universal inv_universal.
apply dedekind_universal2a.
apply inv_inc_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_universal inv_universal.
apply dedekind_universal2c.
apply dedekind_universal2b.
apply inv_inc_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
Qed.
Lemma dedekind\_universal3b {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
               C(i)
                      (alpha \cdot B i) \leftrightarrow beta \quad ((alpha \cdot alpha \#) \cdot beta).
 (beta •
Proof.
split; move \Rightarrow H.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv -comp_assoc.
apply dedekind_universal2b.
apply dedekind_universal2a.
apply inv_inc_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
apply inv\_inc\_invol.
```

```
rewrite comp\_inv comp\_inv inv\_universal inv\_universal. apply dedekind\_universal2c. rewrite -comp\_inv -comp\_inv -comp\_assoc. apply inc\_inv. apply H. Qed.
```

```
Lemma 177 (universal_total) Let \alpha : A \rightarrow B. Then
```

```
\alpha \cdot \nabla_{BI} = \nabla_{AI} \Leftrightarrow "\alpha \text{ is total"}.
```

```
Lemma universal_total \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha \cdot B \ i = A \ i \leftrightarrow total\_r \ alpha.

Proof.

move : (@dedekind\_universal3b\_\_\_ alpha \ (Id \ A)) \Rightarrow H.

rewrite comp\_id\_l \ comp\_id\_r \ in \ H.

rewrite /total\_r.

rewrite -H.

split; move \Rightarrow H0.

rewrite H0.

apply inc\_refl.

apply inc\_antisym.

apply inc\_alpha\_universal.

apply H0.

Qed.
```

7.3 Dedekind formula と恒等関係

```
Lemma 178 (dedekind_id1) Let \alpha:A\to A. Then \alpha\sqsubseteq id_A\Rightarrow \alpha^\sharp=\alpha.
```

 $Id A \rightarrow alpha \# = alpha.$

```
Lemma dedekind\_id1 {A: eqType} {alpha: Rel\ A\ A}: alpha Proof.

move \Rightarrow H.

assert (alpha \# alpha).

move : (@dedekind1\_\_\_(alpha \#) (Id\ A) (Id\ A)) \Rightarrow H0.

rewrite comp\_id\_r\ comp\_id\_r\ inv\_invol\ in\ H0.

replace (alpha \# Id\ A) with (alpha \#) in H0.

replace (Id\ A\ alpha) with alpha in H0.

apply (@inc\_trans\_\_\_(alpha \# alpha)).

apply H0.
```

apply $cap_{-}r$.

Qed.

```
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_inc_move.
rewrite inv_{-}id.
apply H.
rewrite cap\_comm.
apply inc_-def1.
apply H.
apply inc_-def1.
rewrite -inv\_inc\_move.
rewrite inv_{-}id.
apply H.
apply inc\_antisym.
apply H0.
apply inv\_inc\_move.
apply H\theta.
Qed.
  Lemma 179 (dedekind_id2) Let \alpha : A \rightarrow A. Then
                                       \alpha \sqsubseteq id_A \Rightarrow \alpha \cdot \alpha = \alpha.
Lemma dedekind\_id2 \{A : eqType\} \{alpha : Rel A A\}:
           Id A \rightarrow alpha \cdot alpha = alpha.
 alpha
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply (comp\_inc\_compat\_ab\_a\ H).
move: (dedekind_id1 \ H) \Rightarrow H0.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot Id A) Id A)).
rewrite comp_{-}id_{-}r.
apply inc\_cap.
split.
apply inc\_reft.
apply H.
apply (@inc_trans _ _ _ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite H0 \ comp_{-}id_{-}r.
```

Lemma 180 (dedekind_id3) Let $\alpha, \beta : A \rightarrow A$. Then

```
\alpha \sqsubseteq id_A \wedge \beta \sqsubseteq id_A \Rightarrow \alpha \cdot \beta = \alpha \sqcap \beta.
Lemma dedekind\_id3 {A: eqType} {alpha beta : Rel\ A\ A}:
             Id A \rightarrow beta \qquad Id A \rightarrow alpha \cdot beta = alpha
 alpha
                                                                             beta.
Proof.
move \Rightarrow H H0.
apply inc\_antisym.
apply inc\_cap.
split.
apply (comp\_inc\_compat\_ab\_a\ H0).
apply (comp\_inc\_compat\_ab\_b\ H).
replace (alpha
                       beta) with ((alpha)
                                                   beta) • (alpha
apply comp\_inc\_compat.
apply cap_{-}l.
apply cap_r.
apply dedekind_id2.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply cap_{-}l.
Qed.
  Lemma 181 (dedekind_id4) Let \alpha, \beta : A \rightarrow A. Then
                       \alpha \sqsubseteq id_A \land \beta \sqsubseteq id_A \Rightarrow (\alpha \rhd \beta) \sqcap id_A = (\alpha \Rightarrow \beta) \sqcap id_A.
Lemma dedekind\_id4 {A : eqType} {alpha beta : Rel A A}:
                                                       beta) Id A = (alpha \gg beta)
 alpha
            Id A \rightarrow beta \qquad Id A \rightarrow (alpha)
                                                                                                     Id\ A.
Proof.
move \Rightarrow H H0.
apply inc\_lower.
move \Rightarrow gamma.
rewrite inc\_cap inc\_cap.
split; elim \Rightarrow H1 H2.
split.
rewrite inc\_rpc\ cap\_comm.
rewrite -(@dedekind\_id3 \_ \_ \_ H H2).
rewrite -(@dedekind_id1 - H).
apply inc\_residual.
apply H1.
apply H2.
split.
```

```
rewrite inc\_residual (@dedekind\_id1 \_ \_ H) (@dedekind\_id3 \_ \_ \_ H H2).

rewrite cap\_comm - inc\_rpc.

apply H1.

apply H2.

Qed.

End main.
```

Chapter 8

Library Conjugate

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Require Import Dedekind.

Module main (def:Relation).

Import def.

Module Basic\_Lemmas:=Basic\_Lemmas.main def.

Module Relation\_Properties:=Relation\_Properties.main def.

Module Functions\_Mappings:=Functions\_Mappings.main def.

Module Dedekind:=Dedekind.main def.

Import Basic\_Lemmas Relation\_Properties Functions\_Mappings Dedekind.
```

8.1 共役性の定義

条件 P を満たす関係 $\alpha:A\to B$ と条件 Q を満たす関係 $\beta:A'\to B'$ が変換 $\alpha=\phi(\beta),\beta=\psi(\alpha)$ によって、1 対 1 (全射的) に対応することを、図式

$$\frac{\alpha: A \to B \{P\}}{\beta: A' \to B' \{Q\}} \frac{\alpha = \phi(\beta)}{\beta = \psi(\alpha)}$$

によって表す。また、Coq では以下のように表すことにする.

```
Definition conjugate
(A \ B \ C \ D : eqType) \ (P : Rel \ A \ B \to Prop) \ (Q : Rel \ C \ D \to Prop)
(phi : Rel \ C \ D \to Rel \ A \ B) \ (psi : Rel \ A \ B \to Rel \ C \ D) :=
(\forall \ alpha : Rel \ A \ B, \ P \ alpha \to Q \ (psi \ alpha) \land phi \ (psi \ alpha) = alpha)
\land (\forall \ beta : Rel \ C \ D, \ Q \ beta \to P \ (phi \ beta) \land psi \ (phi \ beta) = beta).
```

さらに、上の図式において条件 P または Q が不要な場合には、以下の $True_r$ を代入する.

Definition $True_r \{A \ B : eqType\} := fun_ : Rel \ A \ B \Rightarrow True.$

8.2 共役の例

Lemma 182 (inv_conjugate) Inverse relation (*) makes conjugate. That is,

$$\frac{\alpha: A \to B}{\beta: B \to A} \frac{\alpha = \beta^{\sharp}}{\beta = \alpha^{\sharp}}.$$

Lemma 183 (injection_conjugate) Let $j: C \rightarrow B$ be an injection. Then,

$$\frac{f:A\to B\ \{f^{\sharp}\cdot f\sqsubseteq j^{\sharp}\cdot j\}}{h:A\to C}\ \frac{f=h\cdot j}{h=f\cdot j^{\sharp}}$$

```
Lemma injection_conjugate \{A \ B \ C : eqType\} \ \{j : Rel \ C \ B\}: injection_r \ j \rightarrow conjugate \ A \ B \ A \ C \ (\mathbf{fun} \ f : Rel \ A \ B \Rightarrow ((f \# \bullet f) \ (j \# \bullet j)) \land function_r \ f) \ (\mathbf{fun} \ h : Rel \ A \ C \Rightarrow function_r \ h) \ (\mathbf{fun} \ h : Rel \ A \ C \Rightarrow h \bullet j) \ (\mathbf{fun} \ f : Rel \ A \ B \Rightarrow f \bullet j \#).

Proof.
elim.
elim \Rightarrow H \ H0 \ H1.
split.
move \Rightarrow alpha.
```

```
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
assert (function_r (alpha \cdot j \#)).
split.
apply (@inc\_trans \_ \_ \_ \_ H3).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ j).
apply (@inc\_trans \_ \_ \_ (alpha \cdot ((alpha \# \cdot alpha) \cdot alpha \#))).
rewrite comp_assoc -comp_assoc.
apply (comp\_inc\_compat\_a\_ab\ H3).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_a'b H2).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_b.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply H0.
split.
apply H5.
apply function_inc.
apply function_comp.
apply H5.
split.
apply H.
apply H0.
split.
apply H3.
apply H_4.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
move \Rightarrow beta.
elim \Rightarrow H2 H3.
assert (function_r (beta \cdot j)).
split.
apply (@inc\_trans \_ \_ \_ \_ H2).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ j).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
```

```
apply (comp_inc_compat_ab_b H3).
split.
split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
apply H_4.
rewrite comp_{-}assoc.
replace (j \cdot j \#) with (Id \ C).
apply comp_{-}id_{-}r.
apply inc\_antisym.
apply H.
rewrite /univalent_r in H1.
rewrite inv_invol in H1.
apply H1.
Qed.
  Lemma 184 (injection_conjugate_corollary1, injection_conjugate_corollary2)
  Let j: C \rightarrow B be an injection and f: A \rightarrow B be a function. Then,
              f^{\sharp} \cdot f \sqsubseteq j^{\sharp} \cdot j \Leftrightarrow (\exists! h : A \to C, f = h \cdot j) \Leftrightarrow (\exists h' : A \to C, f \sqsubseteq h' \cdot j).
Lemma injection_conjugate_corollary1 \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{j : Rel \ C \ B\}:
 injection\_r \ j \rightarrow function\_r \ f \rightarrow
 ((f \# \bullet f) \quad (j \# \bullet j) \leftrightarrow \exists ! \ h : Rel \ A \ C, function\_r \ h \land f = h \bullet j).
Proof.
move \Rightarrow H H0.
move: (@injection\_conjugate\ A\_\_\_\ H).
elim \Rightarrow H1 H2.
split; move \Rightarrow H3.
\exists (f \cdot j \#).
split.
move: (H1 f (conj H3 H0)).
elim \Rightarrow H4 H5.
split.
apply H4.
by [rewrite H5].
move \Rightarrow h.
elim \Rightarrow H4 H5.
rewrite H5 comp\_assoc.
replace (j \cdot j \#) with (Id \ C).
apply comp_{-}id_{-}r.
rewrite /injection\_r/function\_r/univalent\_r in H.
```

```
rewrite inv\_invol in H.
apply inc\_antisym.
apply H.
apply H.
elim H3 \Rightarrow h.
elim.
elim \Rightarrow H4 \ H5 \ H6.
rewrite H5 comp_inv comp_assoc -(@comp_assoc _ _ _ _ h).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_b.
apply H_4.
Qed.
Lemma injection\_conjugate\_corollary2 \{A B C : eqType\} \{f : Rel A B\} \{j : Rel C B\}:
 injection\_r \ j \rightarrow function\_r \ f \rightarrow
 ((f \# \cdot f) \quad (j \# \cdot j) \leftrightarrow \exists h' : Rel \ A \ C, f \quad (h' \cdot j)).
Proof.
move \Rightarrow H H0.
split; move \Rightarrow H1.
apply (injection_conjugate_corollary1 H H0) in H1.
elim H1 \Rightarrow h.
elim.
elim \Rightarrow H2 \ H3 \ H4.
\exists h.
rewrite H3.
apply inc\_reft.
elim H1 \Rightarrow h' H2.
replace (f \# \cdot f) with (f \# \cdot (f (h' \cdot j))).
apply (@inc\_trans\_\_\_((f \# \cdot f) \cdot (j \# \cdot j))).
rewrite comp_assoc cap_comm -(@comp_assoc _ _ _ f).
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}r.
apply comp\_inc\_compat\_ab\_b.
apply H0.
apply f_equal.
apply inc_-def1 in H2.
by [rewrite -H2].
Qed.
```

Lemma 185 (surjection_conjugate) Let $e: A \rightarrow C$ be a surjection. Then,

$$\frac{f:A\to B\ \{e\cdot e^\sharp\sqsubseteq f\cdot f^\sharp\}}{h:C\to B}\ \frac{f=e\cdot h}{h=e^\sharp\cdot f}$$

```
Lemma surjection\_conjugate \{A \ B \ C : eqType\} \{e : Rel \ A \ C\}:
 surjection\_r \ e \rightarrow
 conjugate A B C B (fun f : Rel A B \Rightarrow ((e \cdot e \#) (f \cdot f \#)) \land function\_r f)
 (\mathbf{fun}\ h: Rel\ C\ B \Rightarrow function\_r\ h)\ (\mathbf{fun}\ h: Rel\ C\ B \Rightarrow e\ \cdot\ h)\ (\mathbf{fun}\ f: Rel\ A\ B \Rightarrow e\ \#
• f).
Proof.
elim.
elim \Rightarrow H H0 H1.
split.
move \Rightarrow alpha.
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
assert (function_r (e \# \bullet alpha)).
split.
apply (@inc\_trans \_ \_ \_ \_ H1).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H3).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H_4).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot ((alpha \cdot alpha \#) \cdot alpha))).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_a'b H2).
rewrite comp\_assoc -comp\_assoc.
apply (comp_inc_compat_ab_a H_4).
split.
apply H5.
apply Logic.eq_sym.
apply function_inc.
split.
apply H3.
apply H_4.
apply function_comp.
split.
apply H.
apply H0.
apply H5.
```

```
rewrite -comp\_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H.
move \Rightarrow beta.
elim \Rightarrow H2 H3.
assert (function_r (e \cdot beta)).
split.
apply (@inc\_trans \_ \_ \_ \_ H).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab H2).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ e).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H0).
split.
split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp_inc_compat_b_ab H2).
apply H_4.
rewrite -comp\_assoc.
replace (e \# \cdot e) with (Id \ C).
apply comp_{-}id_{-}l.
apply inc\_antisym.
rewrite /total_r in H1.
rewrite inv_invol in H1.
apply H1.
apply H0.
Qed.
  Lemma 186 (surjection_conjugate_corollary) Let e: A \rightarrow C be a surjection and
  f: A \to B be a function. Then,
                            e \cdot e^{\sharp} \sqsubseteq f \cdot f^{\sharp} \Leftrightarrow (\exists! h : C \to B, f = e \cdot h).
Lemma surjection\_conjugate\_corollary \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{e : Rel \ A \ C\}:
 surjection\_r \ e \rightarrow function\_r \ f \rightarrow
 ((e \cdot e \#) \quad (f \cdot f \#) \leftrightarrow \exists ! \ h : Rel \ C \ B, function\_r \ h \land f = e \cdot h).
Proof.
move \Rightarrow H H0.
move: (@surjection\_conjugate \_ B \_ \_ H).
elim \Rightarrow H1 H2.
```

```
split; move \Rightarrow H3.
\exists (e \# \cdot f).
split.
move: (H1 f (conj H3 H0)).
elim \Rightarrow H4 H5.
split.
apply H_4.
by [rewrite H5].
move \Rightarrow h.
elim \Rightarrow H4 H5.
rewrite H5 -comp\_assoc.
replace (e \# \cdot e) with (Id \ C).
apply comp_{-}id_{-}l.
rewrite /surjection_r/function_r/total_r in H.
rewrite inv\_invol in H.
apply inc\_antisym.
apply H.
apply H.
elim H3 \Rightarrow h.
elim.
elim \Rightarrow H4 H5 H6.
rewrite H5 comp_inv comp_assoc -(@comp_assoc _ _ _ h).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H_4.
Qed.
```

Lemma 187 (subid_conjugate) Subidentity $u \sqsubseteq id_A$ corresponds $\rho: I \to A$. That is,

$$\frac{\rho: I \multimap A}{u: A \multimap A \ \{u \sqsubseteq id_A\}} \ \frac{\rho = \nabla_{IA} \cdot u}{u = id_A \sqcap \nabla_{AI} \cdot \rho}.$$

```
Lemma subid\_conjugate \ \{A: eqType\}: \\ conjugate \ i \ A \ A \ True\_r \ (\texttt{fun} \ u: Rel \ A \ A \Rightarrow u \ Id \ A) \\ (\texttt{fun} \ u: Rel \ A \ A \Rightarrow i \ A \cdot u) \ (\texttt{fun} \ rho: Rel \ i \ A \Rightarrow Id \ A \quad (A \ i \cdot rho)). \\ \textbf{Proof.} \\ \textbf{split.} \\ \textbf{move} \Rightarrow alpha \ H. \\ \textbf{split.} \\ \textbf{apply} \ cap\_l. \\ \textbf{apply} \ inc\_antisym. \\ \textbf{apply} \ (@inc\_trans\_\_\_ (i \ A \cdot (A \ i \cdot alpha))). \\ \textbf{apply} \ comp\_inc\_compat\_ab\_ab'. \\ \end{cases}
```

```
apply cap_{-}r.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc_alpha_universal.
rewrite -(@inv\_universal\ i\ A).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (dedekind1)).
rewrite comp\_id\_r cap\_comm cap\_universal.
apply inc\_reft.
move \Rightarrow beta H.
split.
by [].
apply inc\_antisym.
rewrite cap_comm -comp_assoc lemma_for_tarski2.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite comp_id_l cap_comm cap_universal.
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_inc_move inv_id.
apply H.
apply inc\_cap.
split.
apply H.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_b\_ab.
rewrite lemma_for_tarski2.
apply inc\_alpha\_universal.
Qed.
```

```
Lemma 188 (subid_conjugate_corollary1) Let u, v : A \rightarrow A and u, v \sqsubseteq id_A. Then,
```

$$\nabla_{IA} \cdot u = \nabla_{IA} \cdot v \Rightarrow u = v.$$

```
Lemma subid\_conjugate\_corollary1 \{A: eqType\} \{u\ v: Rel\ A\ A\}: u \ Id\ A \to v \ Id\ A \to i\ A \cdot u = i\ A \cdot v \to u = v. Proof.

move \Rightarrow H\ H0\ H1.

move : (@subid\_conjugate\ A).

elim \Rightarrow H2\ H3.

move : (H3\ u\ H).

elim \Rightarrow H4\ H5.

rewrite -H5.

move : (H3\ v\ H0).

elim \Rightarrow H6\ H7.
```

```
rewrite -H7.
apply f_equal.
apply f_equal.
apply H1.
Qed.
```

Lemma 189 (subid_conjugate_corollary2) Let $\rho, \rho' : I \to A$. Then,

$$id_A \sqcap \nabla_{AI} \cdot \rho = id_A \sqcap \nabla_{AI} \cdot \rho' \Rightarrow \rho = \rho'.$$

```
Lemma subid\_conjugate\_corollary2 \{A : eqType\} \{rho \ rho' : Rel \ i \ A\}:
 Id A
               A \ i \cdot rho) = Id \ A \quad (A \ i \cdot rho') \rightarrow rho = rho'.
Proof.
move \Rightarrow H.
move: (@subid\_conjugate\ A).
elim \Rightarrow H0 \ H1.
move: (H0 \ rho \ I).
elim \Rightarrow H2 H3.
rewrite -H3.
move: (H0 \ rho' \ I).
elim \Rightarrow H4 H5.
rewrite -H5.
apply f_equal.
apply H.
Qed.
End main.
```

Chapter 9

Library Domain

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Require Import Dedekind.

Require Import Logic\_FunctionalExtensionality.

Module main\ (def:Relation).

Import def.

Module Basic\_Lemmas:=Basic\_Lemmas.main\ def.

Module Relation\_Properties:=Relation\_Properties.main\ def.

Module Functions\_Mappings:=Functions\_Mappings.main\ def.

Module Dedekind:=Dedekind.main\ def.

Import Basic\_Lemmas\ Relation\_Properties\ Functions\_Mappings\ Dedekind.
```

9.1 定義域の定義

```
関係 \alpha: A \to B に対して、その定義域 (関係) |\alpha|: A \to A は、
```

$$|\alpha| = \alpha \cdot \alpha^{\sharp} \sqcap id_A$$

で表される. また、Coq では以下のように表すことにする.

定義域の性質 $\overline{9.2}$

基本的な性質 9.2.1

Lemma 190 (domain_another_def) Let $\alpha : A \rightarrow B$. Then,

$$[\alpha] = \alpha \cdot \nabla_{BA} \cap id_A.$$

```
Lemma domain\_another\_def \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 domain \ alpha = (alpha \cdot B A)
```

Proof.

apply $inc_antisym$.

apply $cap_inc_compat_r$.

apply $comp_inc_compat_ab_ab$ '.

apply $inc_alpha_universal$.

apply inc_-cap .

split.

apply (@inc_trans _ _ _ (dedekind1)).

apply $comp_inc_compat_ab_ab$ '.

rewrite cap_comm comp_id_r cap_universal.

apply inc_reft .

apply cap_r .

Qed.

Lemma 191 (domain_inv) Let $\alpha : A \rightarrow B$. Then,

$$\lfloor \alpha \rfloor^{\sharp} = \lfloor \alpha \rfloor.$$

Lemma $domain_inv \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:$ $(domain \ alpha) \# = domain \ alpha.$

Proof.

apply dedekind_id1.

apply cap_r .

Qed.

Lemma 192 (domain_comp_alpha1, domain_comp_alpha2) Let $\alpha : A \rightarrow B$. Then,

$$[\alpha] \cdot \alpha = \alpha \wedge \alpha^{\sharp} \cdot [\alpha] = \alpha^{\sharp}.$$

Lemma $domain_comp_alpha1 \{A B : eqType\} \{alpha : Rel A B\}:$ $(domain \ alpha) \cdot alpha = alpha.$

Proof.

```
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_{-}r.
rewrite / domain.
rewrite cap\_comm.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (dedekind2)).
rewrite comp\_id\_l cap\_idem.
apply inc_refl.
Qed.
Lemma domain\_comp\_alpha2 {A B : eqType} {alpha : Rel A B}:
 alpha \# \bullet (domain \ alpha) = alpha \#.
Proof.
rewrite -domain_inv -comp_inv.
apply f_equal.
apply domain\_comp\_alpha1.
Qed.
  Lemma 193 (domain_inc_compat) Let \alpha, \alpha' : A \rightarrow B. Then,
                                       \alpha \sqsubseteq \alpha' \Rightarrow |\alpha| \sqsubseteq |\alpha'|.
Lemma domain\_inc\_compat \{A \ B : eqType\} \{alpha \ alpha' : Rel \ A \ B\}:
           alpha' \rightarrow domain \ alpha domain alpha'.
Proof.
move \Rightarrow H.
apply cap\_inc\_compat\_r.
apply comp_inc_compat.
apply H.
apply (@inc_inv_{-} - H).
Qed.
  Lemma 194 (domain_total) Let \alpha : A \rightarrow B. Then,
                                      "\alpha is total" \Leftrightarrow |\alpha| = id_A.
Lemma domain\_total \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 total\_r \ alpha \leftrightarrow domain \ alpha = Id \ A.
Proof.
split; move \Rightarrow H.
rewrite / domain.
rewrite cap\_comm.
apply Logic.eq_sym.
apply inc\_def1.
```

```
apply H. apply inc\_def1. rewrite /domain in H. by [rewrite cap\_comm H]. Qed.
```

```
Lemma 195 (domain_inc_id) Let u : A \rightarrow A. Then,
```

```
u \sqsubseteq id_A \Leftrightarrow \lfloor u \rfloor = u.
```

```
Lemma domain\_inc\_id \{A: eqType\} \{u: Rel\ A\ A\}: u Id\ A \leftrightarrow domain\ u = u. Proof. split; move \Rightarrow H. rewrite /domain. rewrite (dedekind\_id1\ H) (dedekind\_id2\ H). apply inc\_def1 in H. by [rewrite -H]. rewrite -H. apply cap\_r. Qed.
```

9.2.2 合成と定義域

```
Lemma 196 (comp_domain1, comp_domain2) Let \alpha: A \rightarrow B and \beta: B \rightarrow C. Then,
```

```
[\alpha \cdot \beta] = [\alpha \cdot [\beta]] \sqsubseteq [\alpha].
```

```
Lemma comp\_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 domain (alpha • beta)
                              domain alpha.
Proof.
rewrite /domain.
rewrite comp_-inv.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot (beta \cdot (beta \# \cdot alpha \#)) \quad alpha \#))
                                                                                          Id(A)).
replace (((alpha \cdot beta) \cdot (beta \# \cdot alpha \#)) Id A) with ((((alpha \cdot beta) \cdot
(\mathtt{beta} \ \# \ \bullet \ alpha \ \#)) \qquad Id \ A)
                                     Id\ A).
apply cap\_inc\_compat\_r.
rewrite comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite comp_{-}id_{-}r.
apply inc\_reft.
by [rewrite cap_assoc cap_idem].
apply cap\_inc\_compat\_r.
```

```
apply comp_inc_compat_ab_ab'.
apply cap_{-}r.
Qed.
Lemma comp\_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 domain (alpha \cdot beta) = domain (alpha \cdot domain beta).
Proof.
apply inc\_antisym.
replace (domain (alpha • beta)) with (domain ((alpha • domain beta) • beta)).
apply comp\_domain1.
by [rewrite comp_assoc domain_comp_alpha1].
apply (@inc\_trans \_ \_ \_ (domain (alpha \cdot (beta \cdot beta \#)))).
apply domain\_inc\_compat.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
rewrite - comp_assoc.
apply comp_domain1.
Qed.
```

Lemma 197 (comp_domain3) Let $\alpha: A \rightarrow B$ be a relation and $\beta: B \rightarrow C$ be a total relation. Then,

$$\lfloor \alpha \cdot \beta \rfloor = \lfloor \alpha \rfloor.$$

```
Lemma comp\_domain3 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}: total\_r \ beta \rightarrow domain \ (alpha \cdot beta) = domain \ alpha.

Proof.

move \Rightarrow H.

apply inc\_antisym.

apply comp\_domain1.

rewrite /domain.

rewrite comp\_inv \ comp\_assoc \cdot (@comp\_assoc \_ \_ \_ beta).

apply cap\_inc\_compat\_r.

apply comp\_inc\_compat\_ab\_ab.

apply (comp\_inc\_compat\_b\_ab \ H).

Qed.
```

Lemma 198 (comp_domain4) Let $\alpha : A \rightarrow B$ and $\beta : B \rightarrow C$. Then,

$$\lfloor \alpha^{\sharp} \rfloor \sqsubseteq \lfloor \beta \rfloor \Rightarrow \lfloor \alpha \cdot \beta \rfloor = \lfloor \alpha \rfloor.$$

Lemma $comp_domain4$ {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}: domain (alpha #) domain beta \rightarrow domain (alpha • beta) = domain alpha. Proof.

```
move \Rightarrow H.
apply inc\_antisym.
apply comp\_domain1.
rewrite / domain.
rewrite -(@domain_comp_alpha1 _ _ (alpha #)) comp_inv comp_assoc -(@comp_assoc _ _
_ _ beta).
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans \_ \_ \_ \_ H).
apply cap_{-}l.
Qed.
  Lemma 199 (comp_domain5) Let \alpha : A \rightarrow B be a univalent relation and \beta : B \rightarrow C.
  Then,
                                  |\alpha^{\sharp}| \sqsubset |\beta| \Leftrightarrow |\alpha \cdot \beta| = |\alpha|.
Lemma comp\_domain5 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow
                          domain beta \leftrightarrow domain (alpha \cdot beta) = domain alpha).
 (domain (alpha \#))
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply (comp\_domain4\ H0).
rewrite / domain.
rewrite inv\_invol.
apply cap\_inc\_compat\_r.
replace (alpha \# \cdot alpha) with (alpha \# \cdot (domain (alpha \cdot beta) \cdot alpha)).
rewrite / domain.
rewrite comp_{-}inv.
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot (((alpha \cdot beta) \cdot (beta \# \cdot alpha \#)) \cdot alpha))).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp_assoc comp_assoc -comp_assoc -(@comp_assoc _ _ _ beta).
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_b H)).
apply (comp\_inc\_compat\_ab\_a\ H).
by [rewrite H0 domain_comp_alpha1].
Qed.
```

```
Lemma 200 (comp_domain6) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                        \alpha \cdot |\beta| \sqsubseteq |\alpha \cdot \beta| \cdot \alpha.
Lemma comp\_domain6 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 (alpha • domain beta)
                                (domain (alpha \cdot beta) \cdot alpha).
Proof.
apply (@inc\_trans \_ \_ \_ \_ (@comp\_cap\_distr\_l \_ \_ \_ \_ )).
rewrite cap\_comm.
replace (alpha • Id B) with (Id A • alpha).
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite cap_comm -comp_assoc comp_assoc -comp_inv.
apply inc_refl.
by [rewrite comp_{-}id_{-}l comp_{-}id_{-}r].
Qed.
  Lemma 201 (comp_domain7) Let \alpha : A \to B be a univalent relation and \beta : B \to C.
  Then,
                                        \alpha \cdot |\beta| = |\alpha \cdot \beta| \cdot \alpha.
Lemma comp_domain7 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow alpha \cdot domain \ \mathsf{beta} = domain \ (alpha \cdot \mathsf{beta}) \cdot alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply comp\_domain6.
apply (@inc\_trans \_ \_ \_ \_ (@comp\_cap\_distr\_r \_ \_ \_ \_ )).
rewrite comp_id_l comp_inv comp_assoc comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
apply (fun H' \Rightarrow cap\_inc\_compat H' H).
rewrite comp_assoc -comp_assoc.
apply (comp\_inc\_compat\_ab\_a\ H).
Qed.
  Lemma 202 (comp_domain8) Let u: A \rightarrow A, \alpha: A \rightarrow B and u \sqsubseteq id_A. Then,
                                          |u \cdot \alpha| = u \cdot |\alpha|.
Lemma comp\_domain8 \{A \ B : eqType\} \{u : Rel \ A \ A\} \{alpha : Rel \ A \ B\}:
       Id A \rightarrow domain (u \cdot alpha) = u \cdot domain alpha.
Proof.
```

```
move \Rightarrow H.

apply inc\_antisym.

rewrite -(@cap\_idem\_\_ (domain (u \cdot alpha))).

rewrite (dedekind\_id3 \ H).

apply cap\_inc\_compat.

apply (@inc\_trans\_\_\_\_ (comp\_domain1)).

apply domain\_inc\_id \ in \ H.

rewrite H.

apply inc\_refl.

apply domain\_inc\_compat.

apply (comp\_inc\_compat\_ab\_b \ H).

apply (apple \ cap\_r).

apply (apple \ comp\_inc\_compat\_ab\_b \ H).

Apply (apple \ comp\_inc\_compat\_ab\_b \ H).
```

9.2.3 その他の性質

```
Lemma 203 (cap_domain) Let \alpha, \alpha' : A \to B. Then, \lfloor \alpha \sqcap \alpha' \rfloor = \alpha \cdot \alpha'^{\sharp} \sqcap id_A.
```

```
Lemma cap\_domain \{A \ B : eqType\} \{alpha \ alpha' : Rel \ A \ B\}:
 domain (alpha
                    alpha') = (alpha \cdot alpha' \#) \quad Id A.
Proof.
apply inc\_antisym.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat.
apply cap_{-}l.
apply inc_{-}inv.
apply cap_{-}r.
rewrite -(@cap\_idem \_ \_ (Id A)) - cap\_assoc.
apply cap\_inc\_compat\_r.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
rewrite inv_invol comp_id_l comp_id_r -inv_cap_distr (@cap_comm _ _ alpha').
apply inc\_reft.
Qed.
```

```
B) and P: predicate. Then,
                                   |\sqcup_{P(\alpha)} f(\alpha)| = \sqcup_{P(\alpha)} |f(\alpha)|.
Lemma cupP\_domain\_distr {A B C D : eqType} {f : Rel\ C\ D \rightarrow Rel\ A\ B} {P : Rel\ C\ D
\rightarrow Prop:
 domain ( _{\{P\}} f) = _{\{P\}} (fun \ alpha : Rel \ C \ D \Rightarrow domain (f \ alpha)).
Proof.
rewrite / domain.
rewrite inv\_cupP\_distr\_comp\_cupP\_distr\_l\_cap\_cupP\_distr\_r.
apply cupP_-eq.
move \Rightarrow alpha H.
rewrite - cap_domain - cap_domain.
apply f_equal.
rewrite cap_{-}idem.
apply inc\_antisym.
apply cap_{-}r.
apply inc\_cap.
split.
move: alpha H.
apply inc\_cupP.
apply inc_refl.
apply inc\_reft.
Qed.
Lemma cup\_domain\_distr {A B : eqType} {alpha alpha' : Rel A B}:
 domain (alpha
                     alpha') = domain \ alpha
                                                  domain alpha'.
Proof.
rewrite cup\_to\_cupP (@cup\_to\_cupP _ _ _ _ id).
apply cupP\_domain\_distr.
Qed.
  Lemma 205 (domain_universal1) Let \alpha : A \rightarrow B. Then,
                                       |\alpha| \cdot \nabla_{AC} = \alpha \cdot \nabla_{BC}.
Lemma domain\_universal1 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\}:
 domain \ alpha \cdot A \ C = alpha \cdot B \ C.
Proof.
apply inc\_antisym.
apply @inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot
                                                        A(C)).
apply comp\_inc\_compat\_ab\_a'b.
```

Lemma 204 (cupP_domain_distr, cup_domain_distr) Let $f: (C \rightarrow D) \rightarrow (A \rightarrow D)$

CHAPTER 9. LIBRARY DOMAIN

```
apply cap_{-}l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ ((domain alpha \cdot alpha) \cdot B C)).
rewrite domain_comp_alpha1.
apply inc\_reft.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
  Lemma 206 (domain_universal2) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                    \alpha \cdot |\beta| = \alpha \sqcap \nabla_{AC} \cdot \beta^{\sharp}.
Lemma domain\_universal2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 alpha \cdot domain beta = alpha \quad (A C \cdot beta \#).
apply inc\_antisym.
apply inc\_cap.
split.
apply comp\_inc\_compat\_ab\_a.
apply cap_{-}r.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite -inv_universal -comp_inv -domain_universal1.
rewrite comp_inv inv_universal domain_inv cap_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite cap_comm cap_universal domain_inv.
apply comp\_inc\_compat\_ab\_a.
apply cap_r.
Qed.
```

```
Lemma 207 (domain_lemma1) Let \alpha, \beta : A \rightarrow B and \beta is univalent. Then,
```

$$\alpha \sqsubseteq \beta \land |\alpha| = |\beta| \Rightarrow \alpha = \beta.$$

Lemma domain_lemma1 {A B : eqType} {alpha beta : Rel A B}:


```
beta \rightarrow domain \ alpha = domain \ beta \rightarrow alpha = beta.
 univalent_r beta \rightarrow alpha
Proof.
move \Rightarrow H H0 H1.
apply inc\_antisym.
apply H0.
rewrite -(@domain_comp_alpha1 _ _ beta) -H1.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_a'b.
apply @inc_inv_{-1} = H0.
Qed.
  Lemma 208 (domain_lemma2a, domain_lemma2b) Let \alpha : A \rightarrow B and \beta : A \rightarrow B
  C. Then,
                        |\alpha| \sqsubset |\beta| \Leftrightarrow \alpha \cdot \nabla_{BB} \sqsubseteq \beta \cdot \nabla_{CB} \Leftrightarrow \alpha \sqsubseteq \beta \cdot \beta^{\sharp} \cdot \alpha.
Lemma domain\_lemma2a \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 domain alpha
                     domain beta \leftrightarrow (alpha \cdot B B)
                                                                (beta • CB).
Proof.
split; move \Rightarrow H.
rewrite -(@domain_comp_alpha1 _ _ alpha) comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_a'b H)).
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_a'b (cap\_l))).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ (domain ((beta • beta #) • alpha))).
apply domain_inc_compat.
apply (@inc\_trans \_ \_ \_ (alpha (beta \cdot CB))).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cap\_inc\_compat\_l H)).
replace (alpha \quad (alpha \quad B \quad B)) with ((alpha \quad Id \quad B) \quad (alpha \quad \bullet \quad B)
                                                                                          B B)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (comp\_cap\_distr\_l)).
rewrite cap_universal comp_id_r.
apply inc\_reft.
by [rewrite comp_{-}id_{-}r].
rewrite cap_comm comp_assoc.
apply @inc\_trans \_ \_ \_ \_ (dedekind1).
rewrite cap_comm cap_universal.
apply inc\_reft.
```

CHAPTER 9. LIBRARY DOMAIN

```
rewrite comp_assoc.
apply comp_domain1.
Qed.
Lemma domain\_lemma2b {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
                   domain \ beta \leftrightarrow alpha \quad ((beta \cdot beta \#) \cdot alpha).
 domain alpha
Proof.
split; move \Rightarrow H.
apply domain\_lemma2a in H.
apply (@inc_trans _ _ _ (alpha
                                  (beta • C(B))).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cap\_inc\_compat\_l H)).
replace (alpha \quad (alpha \quad B \quad B)) with ((alpha \quad Id \quad B) \quad (alpha \quad B \quad B)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (comp\_cap\_distr\_l)).
rewrite cap_universal comp_id_r.
apply inc\_reft.
by |rewrite comp_{-}id_{-}r|.
rewrite cap_comm comp_assoc.
apply (@inc\_trans\_\_\_\_ (dedekind1)).
rewrite cap_comm cap_universal.
apply inc_refl.
apply domain\_inc\_compat in H.
apply (@inc\_trans \_ \_ \_ \_ H).
rewrite comp\_assoc.
apply comp_domain1.
Qed.
```

Lemma 209 (domain_corollary1) In below figure,

"\alpha and \beta are total" \land \alpha^\psi \cdot \beta \subseteq \rho^\psi \cdot \tau \infty \cdot \alpha^\psi \cdot \tau \infty \cdot \alpha^\psi \cdot \alpha^\psi \cdot \tau \cdot \alpha^\psi \cdot \alpha^\psi \cdot \tau \cdot \alpha^\psi \cdot \tau \cdot \alpha^\psi \cdot \tau \cdot \alpha^\psi \cdot \alpha \cdot \alpha^\psi \cdot \tau \cdot \alpha^\psi \cdot \alpha \cdot \alph




```
Lemma domain\_corollary1 {X \ Y \ Z \ T : eqType} 
{alpha : Rel \ X \ Y} {beta : Rel \ X \ Z} {rho : Rel \ T \ Y} {tau : Rel \ T \ Z}: 
total\_r \ alpha \rightarrow total\_r \ beta \rightarrow (alpha \ \# \ \ beta) \quad (rho \ \# \ \ \ tau) \rightarrow 
total\_r \ ((alpha \ \ \ \ rho \ \#) \quad (beta \ \ \ tau \ \#)). 
Proof. 
move \Rightarrow H \ H0 \ H1. 
move : (comp\_inc\_compat \ H \ H0) \Rightarrow H2.
```

```
rewrite comp\_id\_l -comp\_assoc (@comp\_assoc _ _ _ alpha) in H2.
rewrite /total_r.
replace (Id\ X) with (((alpha \cdot (rho \# \cdot tau)) \cdot beta \#)
                                                                 Id\ X).
rewrite -comp_assoc comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
rewrite comp_id_l comp_id_r comp_inv comp_inv inv_invol inv_invol.
rewrite inv_cap_distr comp_inv comp_inv inv_invol inv_invol (@cap_comm _ _ (tau •
beta \#)).
apply inc\_reft.
apply Logic.eq_sym.
rewrite cap\_comm.
apply inc\_def1.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp\_inc\_compat\_ab\_ab' H1).
Qed.
```

Lemma 210 (domain_corollary2) In below figure,

"\alpha and \beta are univalent" \land \rho \cdot \rho^\pm \pi \tau \cdot \tau^\pm = id_T \Rightarrow "\alpha \cdot \rho^\pm \pi \Bar \delta^\pm is univalent".



```
Lemma domain\_corollary2 {X \ Y \ Z \ T : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ X \ Z} {rho : Rel \ T \ Y} {tau : Rel \ T \ Z}: univalent\_r \ alpha \rightarrow univalent\_r \ beta \rightarrow (rho \cdot rho \#) \ (tau \cdot tau \#) = Id \ T \rightarrow univalent\_r \ ((alpha \cdot rho \#) \ (beta \cdot tau \#)).

Proof.

move \Rightarrow H \ H0 \ H1.

rewrite /univalent\_r.

rewrite -H1 \ inv\_cap\_distr.

apply (@inc\_trans\_\_\_\_\_ \ (comp\_cap\_distr\_l)).

apply cap\_inc\_compat.

apply (@inc\_trans\_\_\_\_\_ \ (comp\_cap\_distr\_r)).

apply (@inc\_trans\_\_\_\_\_ \ (cap\_l)).

rewrite comp\_inv \ inv\_invol \ -comp\_assoc \ (@comp\_assoc \ \_\_\_\_ \ rho).

apply comp\_inc\_compat\_ab\_a \ b.

apply comp\_inc\_compat\_ab\_a \ b.
```

CHAPTER 9. LIBRARY DOMAIN

```
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).

apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).

rewrite comp\_inv \ inv\_invol \ -comp\_assoc \ (@comp\_assoc \_ \_ \_ tau).

apply comp\_inc\_compat\_ab\_a'b.

apply (comp\_inc\_compat\_ab\_a \ H0).

Qed.
```

9.2.4 矩形関係

 $\alpha: A \rightarrow B \ \mathcal{N}$

$$\alpha \cdot \nabla_{BA} \cdot \alpha \sqsubseteq \alpha$$

を満たすとき, α は 矩形関係 (rectangular relation) であると言われる.

```
Definition rectangular \{A \ B : eqType\} (alpha : Rel \ A \ B) := ((alpha \cdot B \ A) \cdot alpha) alpha.
```

Lemma 211 (rectangular_inv) Let $\alpha : A \to B$ is a rectangular relation, then α^{\sharp} is also a rectangular relation.

```
Lemma rectangular_inv \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
rectangular alpha \rightarrow rectangular \ (alpha \#).

Proof.
move \Rightarrow H.
apply inv\_inc\_move.
rewrite comp\_inv \ comp\_inv \ inv\_invol \ inv\_universal -comp\_assoc.
apply H.
Qed.
```

Lemma 212 (rectangular_capP, rectangular_cap) Let $f(\alpha)$ is always a rectangular relation and P: predicate, then $\sqcap_{P(\beta)} f(\beta)$ is also a rectangular relation.

```
Lemma rectangular_capP {A B C D : eqType} {f : Rel C D \rightarrow Rel A B} {P : Rel C D \rightarrow Prop}: (\forall alpha : Rel C D, P alpha \rightarrow rectangular (f alpha)) \rightarrow rectangular (_{P} f). Proof. move \Rightarrow H. rewrite /rectangular. apply (@inc_trans _ _ _ ( _{P} (fun alpha : Rel C D \Rightarrow (f alpha \cdot B A) \cdot f alpha))). apply (@inc_trans _ _ _ _ (comp_capP_distr_l)). apply inc_capP. move \Rightarrow beta H0.
```

```
apply @inc\_trans\_\_\_ ((( _{P} f) \cdot B A) \cdot f beta)).
move: beta H0.
apply inc\_capP.
apply inc\_reft.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a'b.
move: H0.
apply inc\_capP.
apply inc\_reft.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (H beta H0)).
move : beta H0.
apply inc\_capP.
apply inc\_reft.
Qed.
Lemma rectangular\_cap \{A B : eqType\} \{alpha beta : Rel A B\}:
 rectangular \ alpha \rightarrow rectangular \ beta \rightarrow rectangular \ (alpha
Proof.
move \Rightarrow H H0.
rewrite (@cap\_to\_capP\_\_\_\_\_id).
apply rectangular_capP.
move \Rightarrow gamma.
case \Rightarrow H1; rewrite H1.
apply H.
apply H\theta.
Qed.
  Lemma 213 (rectangular_comp) Let \alpha : A \rightarrow B, \beta : B \rightarrow C and \alpha or \beta is a
  rectangular relation, then \alpha \cdot \beta is also a rectangular relation.
Lemma rectangular\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 rectangular\ alpha\ \lor\ rectangular\ beta \to rectangular\ (alpha\ •\ beta).
Proof.
rewrite / rectangular.
case; move \Rightarrow H.
rewrite - comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
```

CHAPTER 9. LIBRARY DOMAIN

End main.

```
apply inc\_alpha\_universal.
rewrite comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
rewrite - comp\_assoc - comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
  Lemma 214 (rectangular_unit) Let \alpha : A \rightarrow B. Then,
                    "\alpha is rectangular" \Leftrightarrow \exists \mu : I \to A, \exists \rho : I \to B, \alpha = \rho^{\sharp} \cdot \mu.
Lemma rectangular\_unit \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 rectangular alpha \leftrightarrow \exists (mu : Rel \ i \ A)(rho : Rel \ i \ B), \ alpha = mu \# \cdot rho.
Proof.
split; move \Rightarrow H.
\exists (i B \cdot alpha \#).
      i A \cdot alpha).
rewrite comp_inv inv_invol inv_universal.
rewrite -comp_assoc (@comp_assoc _ _ _ alpha) lemma_for_tarski2.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (relation\_rel\_inv\_rel)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply H.
elim H \Rightarrow mu.
elim \Rightarrow rho H0.
rewrite H0.
rewrite / rectangular.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_a.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
```

Chapter 10

Library Residual

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Require Import Dedekind.

Require Import Domain.

Require Import Logic\_FunctionalExtensionality.

Module main\ (def:Relation).

Import def.

Module Basic\_Lemmas:=Basic\_Lemmas.main\ def.

Module Relation\_Properties:=Relation\_Properties.main\ def.

Module Functions\_Mappings:=Functions\_Mappings.main\ def.

Module Dedekind:=Dedekind.main\ def.

Module Domain:=Domain.main\ def.

Import Basic\_Lemmas\ Relation\_Properties\ Functions\_Mappings\ Dedekind\ Domain.
```

10.1 剰余合成関係の性質

10.1.1 基本的な性質

```
Lemma 215 (double_residual) Let \alpha: A \to B, \ \beta: B \to C \ and \ \gamma: C \to D. Then \alpha \rhd (\beta \rhd \gamma) = (\alpha \cdot \beta) \rhd \gamma.
```

```
Lemma double\_residual {A \ B \ C \ D : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C} {gamma : Rel \ C \ D}: alpha (beta gamma) = (alpha \cdot beta) gamma.

Proof.

apply inc\_lower.
```

```
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_residual.
rewrite comp_inv comp_assoc.
rewrite -inc_residual -inc_residual.
apply H.
rewrite inc_residual inc_residual.
rewrite -comp_assoc -comp_inv.
apply inc\_residual.
apply H.
Qed.
  Lemma 216 (residual_to_complement) Let \alpha : A \to B and \beta : B \to C. Then
                                     \alpha \triangleright \beta = (\alpha \cdot \beta^{-})^{-}.
Lemma residual_to_complement {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
          beta = (alpha \cdot beta \hat{)} \hat{.}
 alpha
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
rewrite bool_lemma2 complement_invol cap_comm.
apply inc\_antisym.
apply (@inc\_trans\_\_\_\_ (dedekind1)).
replace (beta \hat{} (alpha \# • gamma)) with ( B C).
rewrite comp\_empty\_r.
apply inc_refl.
apply Logic.eq_sym.
rewrite cap\_comm.
apply bool_lemma2.
apply inc\_residual.
apply H.
apply inc\_empty\_alpha.
apply inc\_residual.
apply bool_lemma2.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite inv\_invol.
                     (alpha \cdot beta \hat{}) with (AC).
replace (gamma
rewrite comp\_empty\_r.
apply inc\_reft.
apply Logic.eq_sym.
```

```
rewrite -(@complement_invol_{--}(alpha \cdot beta^*)). apply bool_lemma2. apply H. apply inc_lempty_alpha. Qed.
```

Lemma 217 (inv_residual_inc) Let $\alpha : A \to B$ and $\beta : B \to C$. Then $\alpha^{\sharp} \cdot (\alpha \rhd \beta) \sqsubseteq \beta$.

Lemma $inv_residual_inc$ {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}: alpha # • (alpha beta) beta.

Proof.

apply $inc_residual$.

apply inc_reft .

Qed.

Lemma 218 (inc_residual_inv) Let $\alpha : A \rightarrow B$ and $\gamma : A \rightarrow C$. Then

$$\gamma \sqsubseteq \alpha \rhd \alpha^{\sharp} \cdot \gamma.$$

Lemma $inc_residual_inv$ { $A \ B \ C : eqType$ } { $alpha : Rel \ A \ B$ } { $gamma : Rel \ A \ C$ }: $gamma \quad (alpha \quad (alpha \ \# \quad gamma)$).

Proof.

apply $inc_residual$.

apply $inc_refl.$

Qed.

Lemma 219 (id_inc_residual) Let $\alpha : A \rightarrow B$. Then

$$id_A \sqsubseteq \alpha \rhd \alpha^{\sharp}$$
.

Lemma $id_inc_residual$ { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }: $Id \ A \ (alpha \ alpha \ \#)$. Proof.

apply $inc_residual$.

rewrite $comp_{-}id_{-}r$.

apply inc_reft .

Qed.

Lemma 220 (residual_universal) Let $\alpha : A \rightarrow B$. Then

$$\alpha \triangleright \nabla_{BC} = \nabla_{AC}$$
.

```
Lemma residual\_universal {A B C : eqType} {alpha : Rel A B}: alpha B C = A C. Proof. apply inc\_antisym. apply inc\_alpha\_universal. apply inc\_residual. apply inc\_alpha\_universal. Qed.
```

10.1.2 単調性と分配法則

```
Lemma 221 (residual_inc_compat) Let \alpha, \alpha' : A \to B and \beta, \beta' : B \to C. Then \alpha' \sqsubseteq \alpha \land \beta \sqsubseteq \beta' \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha' \rhd \beta'.
```

```
Lemma residual\_inc\_compat
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ B\} \{beta \ beta' : Rel \ B \ C\}:
             alpha \rightarrow beta \qquad beta' \rightarrow (alpha \qquad beta)
 alpha'
                                                                   (alpha')
Proof.
move \Rightarrow H H0.
apply inc\_residual.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
move: (@inc\_refl\_\_(alpha)
                                     beta)) \Rightarrow H1.
apply inc\_residual in H1.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply H.
Qed.
```

```
Lemma 222 (residual_inc_compat_l) Let \alpha : A \to B and \beta, \beta' : B \to C. Then \beta \sqsubseteq \beta' \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha \rhd \beta'.
```

```
Lemma residual\_inc\_compat\_l {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta beta' : Rel \ B \ C}: beta beta' \to (alpha \ beta) (alpha \ beta').

Proof.

move \Rightarrow H.

apply (@residual\_inc\_compat \ \_ \_ \_ \_ \_  (@inc\_refl \ \_ \_) H).

Qed.
```

```
\alpha' \sqsubseteq \alpha \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha' \rhd \beta.
Lemma residual\_inc\_compat\_r
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ B\} \{beta : Rel \ B \ C\}:
              alpha \rightarrow (alpha)
                                                (alpha')
 alpha'
                                    beta)
Proof.
move \Rightarrow H.
apply (@residual\_inc\_compat\_\_\_\_\_H (@inc\_refl\_\_\_)).
Qed.
  Lemma 224 (residual_capP_distr_l, residual_cap_distr_l) Let \alpha : A \rightarrow B, f :
  (D \rightarrow E) \rightarrow (B \rightarrow C) and P: predicate. Then
                                   \alpha \rhd (\sqcap_{P(\beta)} f(\beta)) = \sqcap_{P(\beta)} (\alpha \rhd f(\beta)).
Lemma residual\_capP\_distr\_l {A B C D E : eqType}
 \{alpha : Rel \ A \ B\} \{f : Rel \ D \ E \rightarrow Rel \ B \ C\} \{P : Rel \ D \ E \rightarrow Prop\}:
             ( \{P\} f) = \{P\} \text{ (fun beta : } Rel \ D \ E \Rightarrow alpha
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply inc\_residual.
move : beta H0.
apply inc\_capP.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply inc\_residual.
move: beta H0.
apply inc\_capP.
apply H.
Qed.
Lemma residual\_cap\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
 alpha
             (beta \quad gamma) = (alpha)
                                                   beta)
                                                               (alpha
                                                                            gamma).
```

Lemma 223 (residual_inc_compat_r) Let $\alpha, \alpha' : A \to B$ and $\beta : B \to C$. Then

Proof.

```
rewrite cap\_to\_capP (@cap\_to\_capP\_\_\_\_\_id).
apply residual\_capP\_distr\_l.
Qed.
  Lemma 225 (residual_cupP_distr_r, residual_cup_distr_r) Let f:(D \to E) \to
  (A \rightarrow B), \beta: B \rightarrow C \text{ and } P: \text{ predicate. Then}
                                (\sqcup_{P(\alpha)} f(\alpha)) \rhd \beta = \sqcap_{P(\alpha)} (f(\alpha) \rhd \beta).
Lemma residual\_cupP\_distr\_r {A B C D E : eqType}
 \{ beta : Rel \ B \ C \} \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \{ P : Rel \ D \ E \rightarrow Prop \} :
               beta = _{-}\{P\} (fun \ alpha : Rel \ D \ E \Rightarrow f \ alpha
Proof.
apply inc_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow alpha H0.
apply inc\_residual.
move: alpha H0.
apply inc\_cupP.
rewrite -comp\_cupP\_distr\_r -inv\_cupP\_distr.
apply inc\_residual.
apply H.
apply inc\_residual.
rewrite inv\_cupP\_distr\ comp\_cupP\_distr\_r.
apply inc\_cupP.
move \Rightarrow alpha H0.
apply inc\_residual.
move: alpha H0.
apply inc\_capP.
apply H.
Qed.
Lemma residual\_cup\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
            beta) qamma = (alpha \quad qamma)
                                                          (beta gamma).
 (alpha
Proof.
rewrite (@cup\_to\_cupP\_\_\_\_\_id) (@cap\_to\_capP\_\_\_\_\_\_(fun x \Rightarrow x)
                                                                                             qamma)).
apply residual_cupP_distr_r.
Qed.
```

10.1.3 剰余合成と関数

```
Lemma 226 (total_residual) Let \alpha: A \to B be a total relation and \beta: B \to C. Then
                                           \alpha \rhd \beta \sqsubseteq \alpha \cdot \beta.
Lemma total_residual {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total\_r \ alpha \rightarrow (alpha
                             beta) (alpha \cdot beta).
Proof.
move \Rightarrow H.
apply (@inc_trans _ _ _ ((alpha • alpha #) • (alpha
                                                                beta))).
apply (comp\_inc\_compat\_b\_ab\ H).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inv_residual_inc.
Qed.
  Lemma 227 (univalent_residual) Let \alpha : A \to B be a univalent relation and \beta :
  B \rightarrow C. Then
                                           \alpha \cdot \beta \sqsubseteq \alpha \rhd \beta.
Lemma univalent\_residual \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}:
 univalent_r \ alpha \rightarrow (alpha \cdot beta) \quad (alpha
                                                         beta).
Proof.
move \Rightarrow H.
apply (@inc_trans _ _ _ (@inc_residual_inv _ _ alpha _)).
apply residual_inc_compat_l.
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
  Lemma 228 (function_residual1) Let \alpha : A \to B be a function and \beta : B \to C.
  Then
                                           \alpha \triangleright \beta = \alpha \cdot \beta.
Lemma function\_residual1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
function_r \ alpha \rightarrow alpha
                                beta = alpha • beta.
Proof.
elim \Rightarrow H H0.
apply inc\_antisym.
apply (total\_residual\ H).
apply (univalent_residual H0).
Qed.
```

Qed.

```
Lemma 229 (residual_id) Let \alpha : A \rightarrow B. Then
                                              id_A \rhd \alpha = \alpha.
Lemma residual\_id {A B : eqType} {alpha : Rel A B}:
           alpha = alpha.
 Id A
Proof.
move: (@function\_residual1 \_ \_ \_ (Id A) alpha (@id\_function A)) \Rightarrow H.
rewrite comp_{-}id_{-}l in H.
apply H.
Qed.
  Lemma 230 (universal_residual) Let \alpha : A \to B. Then
                                              \nabla_{AA} \triangleright \alpha \sqsubseteq \alpha.
Lemma universal\_residual \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
              alpha
                         alpha.
     A A
Proof.
apply (@inc_trans _ _ _ (Id A
                                         alpha)).
apply residual_inc_compat_r.
apply inc\_alpha\_universal.
rewrite residual_id.
apply inc_refl.
Qed.
  Lemma 231 (function_residual2) Let \alpha: A \to B be a function, \beta: B \to C and
  \gamma: C \rightarrow D. Then
                                       \alpha \cdot (\beta \triangleright \gamma) = (\alpha \cdot \beta) \triangleright \gamma.
Lemma function\_residual2
 \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ C \ D\}:
 function\_r \ alpha \rightarrow alpha \cdot (beta \ gamma) = (alpha \cdot beta) \ gamma.
Proof.
move \Rightarrow H.
rewrite -(@function_residual1 _ _ _ _ H).
apply double_residual.
```

Lemma 232 (function_residual3) Let $\alpha:A \rightarrow B, \ \beta:B \rightarrow C$ be relations and $\gamma:D \rightarrow C$ be a function. Then

$$(\alpha \rhd \beta) \cdot \gamma^{\sharp} = \alpha \rhd (\beta \cdot \gamma^{\sharp}).$$

```
Lemma function_residual3
     \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ D \ C\}: \{gamma : Rel \ D \ C\}
    function\_r\ gamma \rightarrow (alpha \ beta) \cdot gamma \# = alpha \ (beta \cdot gamma \#).
Proof.
move \Rightarrow H.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H0.
apply inc\_residual.
rewrite -(@function\_move2\_\_\_\_\_H).
rewrite comp\_assoc.
apply inc\_residual.
rewrite (@function_move2 _ _ _ _ H).
apply H0.
rewrite -(@function\_move2\_\_\_\_\_H).
apply inc\_residual.
rewrite -comp_-assoc.
rewrite (@function\_move2\_\_\_\_\_H).
apply inc\_residual.
apply H0.
Qed.
```

Lemma 233 (function_residual4) Let $\alpha:A \rightarrow B, \ \gamma:C \rightarrow D$ be relations and $\beta:B \rightarrow C$ be a function. Then

$$\alpha \cdot \beta \rhd \gamma = \alpha \rhd \beta \cdot \gamma.$$

```
Lemma function_residual4  \{A \ B \ C \ D : eqType\} \ \{alpha : Rel \ A \ B\} \ \{beta : Rel \ B \ C\} \ \{gamma : Rel \ C \ D\}: function\_r \ beta \rightarrow (alpha \cdot beta) \quad gamma = alpha \quad (beta \cdot gamma).  Proof.  move \Rightarrow H.  rewrite -double\_residual.  by [rewrite (function\_residual1 \ H)]. Qed.
```

10.2 Galois 同値とその系

```
Lemma 234 (galois) Let \alpha: A \rightarrow B, \beta: B \rightarrow C and \gamma: A \rightarrow C. Then
                                         \gamma \sqsubseteq \alpha \rhd \beta \Leftrightarrow \alpha \sqsubseteq \gamma \rhd \beta^{\sharp}.
Lemma galois \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ A \ C\}:
 qamma
               (alpha
                            beta) \leftrightarrow alpha
                                                    (qamma)
                                                                   beta \#).
Proof.
split; move \Rightarrow H.
apply inc\_residual.
apply inv\_inc\_move.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inv_inc_invol.
rewrite comp_{-}inv \ inv_{-}invol.
apply inc\_residual.
apply H.
Qed.
  Lemma 235 (galois_corollary1) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then
                                              \alpha \sqsubset (\alpha \rhd \beta) \rhd \beta^{\sharp}.
Lemma galois_corollary1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
             ((alpha
                          beta) beta \#).
 alpha
Proof.
rewrite -galois.
apply inc\_reft.
Qed.
  Lemma 236 (galois_corollary2) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then
                                        ((\alpha \rhd \beta) \rhd \beta^{\sharp}) \rhd \beta = \alpha \rhd \beta.
Lemma galois_corollary2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 ((alpha
                         \mathtt{beta} \ \#) \qquad \mathtt{beta} = \mathit{alpha}
               beta)
Proof.
apply inc\_antisym.
apply residual_inc_compat_r.
```

```
apply galois\_corollary1.

move: (@galois\_corollary1\_\_\_(alpha beta) (beta #)) \Rightarrow H.

rewrite inv\_invol in H.

apply H.


Qed.
```

Lemma 237 (galois_corollary3) Let $\alpha : A \to B$ and $\beta : B \to C$. Then $\alpha = (\alpha \rhd \beta) \rhd \beta^{\sharp} \Leftrightarrow \exists \gamma : A \to C, \alpha = \gamma \rhd \beta^{\sharp}.$

```
Lemma galois\_corollary3 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}: alpha = (alpha \quad beta) \quad beta \# \leftrightarrow (\exists \ gamma : Rel A C, alpha = gamma \quad beta \#) Proof. split; move \Rightarrow H. \exists \ (alpha \quad beta). apply H. elim H \Rightarrow gamma \ H0. rewrite H0. move : (@galois\_corollary2 \ \_ \ \_ \ gamma \ (beta \#)) \Rightarrow H1. rewrite inv\_invol in H1. by [rewrite H1]. Qed.
```

10.3 その他の性質

この節では、特記が無い限り、記号は以下の図式に従って割り振られるものとする.



Lemma 238 (residual_property1)

$$(\alpha \rhd \beta) \cdot \gamma \sqsubseteq \alpha \rhd \beta \cdot \gamma.$$

```
Lemma residual\_property1 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}: ((alpha \ beta) \cdot gamma) \ (alpha \ (beta \cdot gamma)).
Proof.
```

Lemma 239 (residual_property2)

$$(\alpha \rhd \beta) \cdot (\beta^{\sharp} \rhd \eta) \sqsubseteq \alpha \rhd \eta.$$

```
Lemma residual\_property2 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{eta : Rel \ Y \ V\}: ((alpha beta) \cdot (beta \# eta)) \quad (alpha eta).

Proof.

apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).

apply residual\_inc\_compat\_l.

move : (@inv\_residual\_inc \_ \_ (beta \#) eta).

by [rewrite \ inv\_invol].

Qed.
```

Lemma 240 (residual_property3)

$$\alpha \rhd \beta \sqsubseteq \alpha \cdot \eta \rhd \eta^{\sharp} \cdot \beta.$$

```
{\color{red}\textbf{Lemma}}\ residual\_property3
```

Lemma 241 (residual_property4a, residual_property4b)

$$(\alpha \rhd \beta) \cdot \gamma \sqsubseteq (\alpha \rhd \beta \cdot \gamma) \sqcap \nabla_{XZ} \cdot \gamma \sqsubseteq (\alpha \rhd \beta \cdot \gamma) \cdot \gamma^{\sharp} \cdot \gamma.$$

```
Lemma residual\_property4a \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}:
```

```
beta) • gamma)
                                                (beta • qamma))
                                                                             X Z \cdot qamma)).
 ((alpha
                                    ((alpha
Proof.
rewrite -(@cap_universal _ _ (alpha
                                            beta)).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_r.
apply residual_property1.
Qed.
Lemma residual_property4b
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}:
             (beta \cdot gamma)) \quad (XZ \cdot gamma)) \quad ((alpha
                                                                             (beta \cdot gamma)) \cdot
(gamma \# \bullet gamma)).
Proof.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite cap_comm cap_universal comp_assoc.
apply inc\_reft.
Qed.
  Lemma 242 (residual_property5) Let \tau be a univalent relation. Then,
                             (\alpha \rhd \beta) \cdot \tau^{\sharp} = (\alpha \rhd \beta \cdot \tau^{\sharp}) \sqcap \nabla_{XZ} \cdot \tau^{\sharp}.
Lemma residual\_property5
 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{tau : Rel \ V \ Z\}:
 univalent_r tau \rightarrow
           beta) • tau \# = (alpha \quad (beta • tau \#)) \quad (XZ • tau \#).
 (alpha
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite -(@cap_universal _ _ (alpha
apply (@inc\_trans \_ \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_r.
apply residual_property1.
rewrite cap\_comm.
apply (@inc_trans _ _ _ (dedekind2)).
rewrite cap_comm cap_universal inv_invol.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply residual_inc_compat_l.
rewrite comp_{-}assoc.
apply (comp\_inc\_compat\_ab\_a\ H).
Qed.
```

Lemma 243 (residual_property6)

$$\alpha \rhd (\gamma^{\sharp} \rhd \beta^{\sharp})^{\sharp} = (\gamma^{\sharp} \rhd (\alpha \rhd \beta)^{\sharp})^{\sharp}.$$

```
Lemma residual_property6
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}:
          (gamma \ \# \ beta \ \#) \ \# = (gamma \ \#)
 alpha
                                                       (alpha
                                                                   beta) #) #.
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_move.
apply inc\_residual.
rewrite -comp_inv comp_assoc.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_invol.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_move.
rewrite comp_inv inv_invol inv_invol comp_assoc.
apply inc\_residual.
apply inv\_inc\_invol.
rewrite comp_{-}inv.
apply inc\_residual.
apply inv\_inc\_move.
apply H.
Qed.
```

Lemma 244 (residual_property7a, residual_property7b)

$$\alpha \rhd (\beta \Rightarrow \beta') \sqsubseteq (\alpha \cdot \beta \Rightarrow \alpha \cdot \beta') \sqsubseteq \alpha \rhd (\beta \Rightarrow \alpha^{\sharp} \cdot \alpha \cdot \beta').$$

Lemma $residual_property7a$ { $X \ Y \ Z : eqType$ } { $alpha : Rel \ X \ Y$ } {beta $beta' : Rel \ Y \ Z$ }: (alpha (beta » beta')) (($alpha \cdot beta'$)) » ($alpha \cdot beta'$)).

Proof. move $\Rightarrow H$.

```
apply inc_-rpc.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp_inc_compat_ab_ab'.
rewrite cap\_comm.
apply inc_-rpc.
apply inv\_residual\_inc.
Lemma residual\_property7b {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta beta' : Rel \ Y \ Z}:
 ((alpha \cdot beta) * (alpha \cdot beta')) (alpha \cdot (beta * (alpha # \cdot (alpha \cdot beta')))).
Proof.
rewrite inc_residual inc_rpc.
apply @inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite inv_invol -inc_rpc.
apply inc_refl.
Qed.
  Lemma 245 (residual_property8) Let \alpha be a univalent relation. Then,
                                  \alpha \rhd (\beta \Rightarrow \beta') = (\alpha \cdot \beta \Rightarrow \alpha \cdot \beta').
Lemma residual\_property8 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta \ beta' : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \quad (beta \gg beta') = (alpha \cdot beta) \gg (alpha \cdot beta').
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply residual_property7a.
apply (@inc_trans _ _ _ residual_property7b).
apply residual_inc_compat_l.
apply rpc\_inc\_compat\_l.
rewrite - comp_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
  Lemma 246 (residual_property9) Let \alpha be a univalent relation. Then,
                                     \alpha \rhd \beta = (\alpha \cdot \nabla_{YZ} \Rightarrow \alpha \cdot \beta).
Lemma residual\_property9 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \quad beta = (alpha \cdot Y Z) \otimes (alpha \cdot beta).
```

```
by [rewrite -(residual\_property8\ H) rpc\_universal\_alpha]. Qed.
```

Lemma 247 (residual_property10) Let α be a univalent relation. Then,

$$\alpha \cdot \beta = |\alpha| \cdot (\alpha \triangleright \beta).$$

```
Lemma residual\_property10 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
 univalent_r \ alpha \rightarrow alpha \cdot beta = domain \ alpha \cdot (alpha
Proof.
move \Rightarrow H.
apply inc\_antisym.
replace (alpha • beta) with (domain alpha • (alpha • beta)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite inc\_residual -comp\_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
by [rewrite -comp_assoc domain_comp_alpha1].
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot (alpha beta))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inv\_residual\_inc.
Qed.
```

Lemma 248 (residual_property11)

$$(\alpha \cdot \beta \Rightarrow \delta) \sqsubseteq \alpha \rhd (\beta \Rightarrow \alpha^{\sharp} \cdot \delta).$$

```
Lemma residual\_property11 \{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{delta : Rel \ X \ Z\}: \ ((alpha \cdot beta) \otimes delta) \ (alpha \ (beta \otimes (alpha \# \cdot delta))).

Proof.

apply inc\_residual.

apply inc\_rpc.

apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).

rewrite inv\_invol.

apply inc\_pc.

apply inc\_pc.

apply inc\_pc.

apply inc\_pc.

apply inc\_pc.
```

```
Lemma 249 (residual_property12a, residual_property12b) Let u \sqsubseteq id_X. Then,
```

```
u \rhd \alpha = u \cdot \nabla_{XY} \Rightarrow \alpha = u \rhd u \cdot \alpha.
```

```
Lemma residual\_property12a {X Y : eqType} {u : Rel X X} {alpha : Rel X Y}:
      Id X \to u
                    alpha = (u \cdot X Y) * alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
assert (univalent_r u).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_{-}id.
apply (@inc_inv_{-} - H).
rewrite (residual_property9 H0).
apply rpc\_inc\_compat\_l.
apply (comp\_inc\_compat\_ab\_b\ H).
apply (@inc_trans _ _ _ residual_property11).
apply residual_inc_compat_l.
rewrite rpc_universal_alpha.
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_{-}id.
apply (@inc_inv_{-} - H).
Qed.
Lemma residual\_property12b {X Y : eqType} {u : Rel X X} {alpha : Rel X Y}:
      Id X \rightarrow u
                    alpha = u \quad (u \cdot alpha).
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite (residual_property12a H).
apply (@inc_trans _ _ _ residual_property11).
apply residual_inc_compat_l.
rewrite rpc\_universal\_alpha.
apply comp\_inc\_compat\_ab\_a'b.
rewrite (dedekind_id1 H).
apply inc_refl.
apply residual\_inc\_compat\_l.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
```

Lemma 250 (residual_property13)

```
(\alpha \cdot \nabla_{YZ} \sqcap \delta) \rhd \gamma = (\alpha \cdot \nabla_{YW} \Rightarrow (\delta \rhd \gamma)).
```

```
Lemma residual_property13
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{gamma : Rel \ Z \ W\} \{delta : Rel \ X \ Z\}:
                                 qamma = (alpha \cdot Y W) \gg (delta)
 ((alpha \cdot Y Z)  delta)
Proof.
apply inc\_antisym.
rewrite inc\_rpc inc\_residual.
remember (((alpha \cdot Y Z) \quad delta) \quad gamma) \text{ as } sigma1.
apply (@inc\_trans \_ \_ \_ (((alpha \cdot Y Z) delta) \# \cdot sigma1)).
                                                                       (alpha \cdot
apply @inc\_trans \_ \_ \_ (((alpha \cdot Y Z) delta) \# \cdot (sigma1)
                                                                                      Y
W)))).
assert ((delta \# \cdot (sigma1 \quad (alpha \cdot Y W))) (delta \# \cdot sigma1)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
apply inc_{-}def1 in H.
rewrite H.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite (@inv_cap_distr _ _ _ delta) cap_comm.
apply cap\_inc\_compat\_r.
rewrite inv\_cap\_distr.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply @inc\_trans \_ \_ \_ \_ (cap\_r).
rewrite comp\_inv comp\_inv -comp\_assoc (@inv\_universal Y Z).
apply comp_inc_compat_ab_a'b.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
rewrite Hegsigma1.
apply inc\_residual.
apply inc\_reft.
rewrite inc\_residual.
apply @inc\_trans \_ \_ \_ (delta \# \cdot ((alpha \cdot Y W)))
                                                             sigma2))).
apply (@inc\_trans\_\_\_(((alpha \cdot YZ) \quad delta) \# \cdot ((alpha \cdot YW) \quad sigma2))).
assert ((((alpha \cdot YZ) delta) # sigma2) (delta # sigma2)).
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply cap_r.
```

```
apply inc\_def1 in H.
rewrite H.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm inv_invol.
apply cap\_inc\_compat\_r.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot Y Z) \cdot (delta \# \cdot sigma2))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply cap_r.
rewrite Hegsigma2.
rewrite -inc_residual cap_comm -inc_rpc.
apply inc\_reft.
Qed.
  Lemma 251 (residual_property14) Let \nabla_{XX} \cdot \alpha \sqsubseteq \alpha. Then,
                                      \nabla_{XX} \cdot (\alpha \rhd \beta) \sqsubset \alpha \rhd \beta.
Lemma residual\_property14 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
 (XX \cdot alpha) \quad alpha \rightarrow (XX \cdot (alpha))
                                                             beta))
                                                                          (alpha
                                                                                      beta).
Proof.
move \Rightarrow H.
apply (@inc\_trans \_ \_ \_ ( XX \cdot ( XX \quad (alpha \quad beta)))).
apply comp\_inc\_compat\_ab\_ab'.
rewrite double_residual.
apply (residual\_inc\_compat\_r\ H).
rewrite -inv_universal -inc_residual inv_universal.
apply inc\_reft.
Qed.
  Lemma 252 (residual_property15) Let \beta \cdot \nabla_{ZZ} \subseteq \beta. Then,
                                      (\alpha \rhd \beta) \cdot \nabla_{ZZ} \sqsubset \alpha \rhd \beta.
Lemma residual\_property15 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
            ZZ) beta \rightarrow ((alpha \text{ beta}) \cdot ZZ) (alpha)
Proof.
```

```
\begin{array}{l} \mathsf{move} \Rightarrow H. \\ \mathsf{apply} \; (@inc\_trans \_ \_ \_ \_ \; (residual\_property1)). \\ \mathsf{apply} \; (residual\_inc\_compat\_l \; H). \\ \\ \mathsf{Qed}. \end{array}
```

Lemma 253 (residual_property16)

```
id_X \sqsubseteq \alpha \rhd \alpha^{\sharp} \land (\alpha \rhd \alpha^{\sharp}) \cdot (\alpha \rhd \alpha^{\sharp}) \sqsubseteq \alpha \rhd \alpha^{\sharp}.
```

```
Lemma residual\_property16 {X \ Y : eqType} {alpha : Rel \ X \ Y}: Id \ X \quad (alpha \quad alpha \ \#) \land ((alpha \quad alpha \ \#) \quad (alpha \quad alpha \ \#)) \quad (alpha \quad alpha \ \#). Proof. split. rewrite inc\_residual \ comp\_id\_r. apply inc\_refl. move : (@residual\_property2 \ \_ \ \_ \ \_ \ alpha \ (alpha \ \#) \ (alpha \ \#)) \Rightarrow H. rewrite inv\_invol in H. apply H. Qed.
```

```
Lemma 254 (residual_property17) Let P(y) := "y : I \rightarrow Y \text{ is a function"}. Then,
```

```
\sqcup_{P(y)} y^{\sharp} \cdot y = id_Y \Rightarrow \alpha \rhd \beta = \sqcap_{P(y)} (\alpha \cdot y^{\sharp} \cdot \nabla_{IZ} \Rightarrow \alpha \cdot y^{\sharp} \cdot y \cdot \beta).
```

```
Lemma residual\_property17 \{X \ Y \ Z : eqType\}
 \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{P : Rel \ i \ Y \rightarrow Prop\}:
 P = (\mathbf{fun} \ y : Rel \ i \ Y \Rightarrow function_r \ y) \rightarrow
    \{P\} (fun y: Rel \ i \ Y \Rightarrow y \# \cdot y) = Id \ Y \rightarrow
 alpha beta = _{-}\{P\} (fun y: Rel i Y \Rightarrow
  ((alpha \cdot y \#) \cdot i Z) \gg ((alpha \cdot y \#) \cdot (y \cdot beta))).
Proof.
move \Rightarrow H H0.
replace (alpha
                      beta) with ((alpha • Id Y)
rewrite -H0 comp_cupP_distr_l residual_cupP_distr_r.
apply capP_{-}eq.
move \Rightarrow y H1.
rewrite H in H1.
rewrite -comp_assoc (function_residual4 H1).
apply residual_property9.
rewrite /univalent_r.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
```

by [rewrite $comp_{-}id_{-}r$]. Qed.

10.4 順序の関係と左剰余合成

10.4.1 max, sup, min, inf

 $\xi: X \to X$ を集合 X における順序と見なしたときの、関係 $\rho: V \to X$ の 最大値 (\max) 、上限 (\sup) 、最小値 (\min) 、下限 (\inf) はそれぞれ、以下のように定義される.

- $max(\rho, \xi) := \rho \sqcap (\rho \rhd \xi)$
- $sup(\rho, \xi) := (\rho \rhd \xi) \sqcap ((\rho \rhd \xi) \rhd \xi^{\sharp})$
- $min(\rho, \xi) := \rho \sqcap (\rho \rhd \xi^{\sharp}) (= max(\rho, \xi^{\sharp}))$
- $inf(\rho, \xi) := (\rho \triangleright \xi^{\sharp}) \sqcap ((\rho \triangleright \xi^{\sharp}) \triangleright \xi) (= sup(\rho, \xi^{\sharp}))$

```
Definition max \{ V \mid X : eqType \} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
             (rho
 := rho
                      xi).
Definition sup \{V \mid X : eqType\} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
                      ((rho
 := (rho)
                                xi)
                                     xi \#).
Definition min \{V \mid X : eqType\} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
 := rho
             (rho
                      xi \#).
Definition inf \{V \mid X : eqType\} (rho: Rel V X) (xi: Rel X X)
 := (rho
              xi \#
                       ((rho
                                  xi \#
                                              xi).
```

```
Lemma 255 (max_inc_sup) Let \rho: V \to X and \xi: X \to X. Then, max(\rho, \xi) \sqsubseteq sup(\rho, \xi).
```

```
Lemma max\_inc\_sup {V \ X : eqType} {rho : Rel \ V \ X} {xi : Rel \ X \ X}: max \ rho \ xi sup \ rho \ xi.

Proof.

rewrite /max/sup.

rewrite cap\_comm.

apply cap\_inc\_compat\_l.

apply galois\_corollary1.

Qed.
```

```
Lemma 256 (min_inc_inf) Let \rho: V \to X and \xi: X \to X. Then,
                                      min(\rho, \xi) \sqsubseteq inf(\rho, \xi).
Lemma min\_inc\_inf {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 min rho xi
                 inf rho xi.
Proof.
rewrite /min/inf.
rewrite cap\_comm.
apply cap\_inc\_compat\_l.
move: (@galois\_corollary1\_\_\_rho(xi\#)) \Rightarrow H.
rewrite inv\_invol in H.
apply H.
Qed.
  Lemma 257 (inf_to_sup) Let \rho: V \to X and \xi: X \to X. Then,
                                   inf(\rho,\xi) = sup(\rho \triangleright \xi^{\sharp}, \xi).
Lemma inf\_to\_sup {V X : eqType} {rho : Rel\ V\ X} {xi : Rel\ X\ X}:
 inf \ rho \ xi = sup \ (rho
                             xi \#) xi.
Proof.
rewrite /sup/inf.
rewrite cap\_comm.
move: (@galois\_corollary2 \_ \_ \_ rho (xi \#)) \Rightarrow H.
rewrite inv\_invol in H.
by [rewrite H].
Qed.
  Lemma 258 (sup_to_inf) Let \rho: V \to X and \xi: X \to X. Then,
                                    sup(\rho, \xi) = inf(\rho \triangleright \xi, \xi).
Lemma sup\_to\_inf {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 sup \ rho \ xi = inf \ (rho
                             xi) xi.
Proof.
rewrite /sup/inf.
rewrite cap\_comm.
by [rewrite qalois_corollary2].
Qed.
```

```
Lemma 259 (residual_inc_sup1, residual_inc_sup2) Let \rho: V \to X and \xi: X \to X
  X. Then,
                                 sup(\rho, \xi) \sqsubseteq \rho \rhd \xi \sqsubseteq sup(\rho, \xi) \rhd \xi.
Lemma residual\_inc\_sup1 { V X : eqType} { rho : Rel V X} { xi : Rel X X}:
                 (rho
 sup rho xi
                          xi).
Proof.
apply cap_{-}l.
Qed.
Lemma residual\_inc\_sup2 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                  ((sup \ rho \ xi)
 (rho
          xi
                                     xi).
Proof.
rewrite qalois.
apply cap_{-}r.
Qed.
  Lemma 260 (max_inc_xi_cap) Let \rho: V \to X and \xi: X \to X. Then,
                                 (max(\rho,\xi))^{\sharp} \cdot max(\rho,\xi) \sqsubseteq \xi \sqcap \xi^{\sharp}.
Lemma max\_inc\_xi\_cap {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (max \ rho \ xi \ \# \ \bullet \ max \ rho \ xi) (xi)
                                               xi \#).
Proof.
rewrite /max.
rewrite inv\_cap\_distr.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat.
apply inc\_residual.
apply cap_{-}r.
apply inv\_inc\_move.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply residual\_inc\_compat\_r.
apply cap_{-}l.
Qed.
  Lemma 261 (sup_inc_xi_cap) Let \rho: V \to X and \xi: X \to X. Then,
                                  (sup(\rho,\xi))^{\sharp} \cdot sup(\rho,\xi) \sqsubseteq \xi \sqcap \xi^{\sharp}.
Lemma sup\_inc\_xi\_cap {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (sup \ rho \ xi \ \# \ \bullet \ sup \ rho \ xi) (xi)
                                             xi \#).
```

```
CHAPTER 10. LIBRARY RESIDUAL
Proof.
move: (@max\_inc\_xi\_cap\_\_ (rho
                                           xi) (xi \#).
rewrite /max/sup.
by [rewrite inv_invol (@cap_comm _ _ xi)].
Qed.
  Lemma 262 (transitive_sup1) Let \rho: V \to X, \xi: X \to X and \xi \cdot \xi \sqsubseteq \xi. Then,
                                  sup(\rho, \xi) \cdot (\xi \sqcap \xi^{\sharp}) = sup(\rho, \xi).
Lemma transitive\_sup1 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                xi \rightarrow sup \ rho \ xi \cdot (xi \quad xi \#) = sup \ rho \ xi.
 (xi \cdot xi)
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite /sup.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply (residual\_inc\_compat\_l\ H).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply residual_inc_compat_l.
rewrite -comp_inv inv_inc_move inv_invol.
apply H.
apply (@inc_trans _ _ _ _ (relation_rel_inv_rel)).
rewrite comp_{-}assoc.
apply (comp\_inc\_compat\_ab\_ab' sup\_inc\_xi\_cap).
Qed.
  Lemma 263 (transitive_sup2) Let \rho: V \to X, \xi: X \to X and \xi \cdot \xi \sqsubseteq \xi. Then,
                                sup(\rho, \xi) \cdot \xi = |sup(\rho, \xi)| \cdot (\rho \triangleright \xi).
Lemma transitive\_sup2 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                xi \rightarrow sup \ rho \ xi \cdot xi = domain (sup \ rho \ xi) \cdot (rho
 (xi \cdot xi)
Proof.
move \Rightarrow H.
apply inc\_antisym.
replace (sup rho xi · xi) with (domain (sup rho xi) · (sup rho xi · xi)).
```

apply $comp_inc_compat_ab_ab$ '.

```
apply @inc\_trans \_ \_ \_ ((rho
                                    xi) \cdot xi).
apply (comp\_inc\_compat\_ab\_a'b cap\_l).
apply (@inc_trans _ _ _ _ (residual_property1) (residual_inc_compat_l H)).
by [rewrite -comp_assoc domain_comp_alpha1].
apply (@inc_trans _ _ _ (domain (sup rho xi) • (sup rho xi)
                                                                     xi))).
apply comp_inc_compat_ab_ab'.
apply galois.
apply cap_r.
rewrite / domain.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp\_assoc.
apply comp_inc_compat_ab_ab'.
apply inc\_residual.
apply inc_refl.
Qed.
  Lemma 264 (domain_sup_inc) Let \rho: V \to X and \xi: X \to X. Then,
                                |sup(\rho,\xi)| \cdot \rho \sqsubseteq sup(\rho,\xi) \cdot \xi^{\sharp}.
Lemma domain\_sup\_inc {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (domain (sup rho xi) \cdot rho) (sup rho xi \cdot xi \#).
Proof.
apply (@inc\_trans \_ \_ \_ (domain (sup rho xi) \cdot (sup rho xi xi \#))).
apply comp\_inc\_compat\_ab\_ab'.
rewrite - galois.
apply cap_l.
rewrite / domain.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_residual.
apply inc\_reft.
Qed.
  Lemma 265 (sup_function) Let \rho: V \to X, \xi: X \to X be relations and f: W \to V
  be a function. Then,
                                 f \cdot sup(\rho, \xi) = sup(f \cdot \rho, \xi).
```

Lemma $sup_function \{V \ W \ X : eqType\} \{rho : Rel \ V \ X\} \{xi : Rel \ X \ X\} \{f : Rel \ W \ V\}:$

```
function_r f \to f • sup rho xi = \sup (f \cdot rho) xi.
Proof.
move \Rightarrow H.
rewrite /sup.
rewrite (function_cap_distr_l H).
by [rewrite (function_residual2 H) (function_residual2 H) (function_residual2 H)].
Qed.
  Lemma 266 (max_univalent) Let \rho: V \to X, \xi: X \to X be relations and \varphi: W \to X
  V be a univalent relation. Then,
                                  \varphi \cdot max(\rho, \xi) = max(\varphi \cdot \rho, \xi).
Lemma max\_univalent \{ V \ W \ X : eqType \}
 \{rho: Rel\ V\ X\}\ \{xi: Rel\ X\ X\}\ \{phi: Rel\ W\ V\}:
 univalent_r \ phi \rightarrow phi \cdot max \ rho \ xi = max \ (phi \cdot rho) \ xi.
Proof.
move \Rightarrow H.
rewrite /max.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat\_l.
apply (@inc\_trans \_ \_ \_ \_ (univalent\_residual H)).
rewrite double_residual.
apply inc\_reft.
```

10.4.2 左剰余合成

apply $cap_inc_compat_l$.

apply inc_reft .

Qed.

apply (@inc_trans _ _ _ _ (dedekind1)).

rewrite -inc_residual double_residual.

apply $comp_inc_compat_ab_ab$ '.

```
関係 \alpha: X \to Y, \beta: Y \to Z に対し、左剰余合成を \alpha \triangleleft \beta := (\beta^\sharp \rhd \alpha^\sharp)^\sharp で定義する.
```

```
Definition leftres \{X \ Y \ Z : eqType\} (alpha : Rel X Y) (beta : Rel Y Z) := (beta \# alpha \#) \#.
```

Lemma 267 (inc_leftres) Let $\alpha: X \to Y, \ \beta: Y \to Z \ and \ \delta: X \to Z$. Then, $\delta \sqsubseteq \alpha \lhd \beta \Leftrightarrow \delta \cdot \beta^{\sharp} \sqsubseteq \alpha.$

Lemma $inc_leftres$ {X Y Z : eqType}

 $\{alpha: Rel\ X\ Y\}\ \{beta: Rel\ Y\ Z\}\ \{delta: Rel\ X\ Z\}:$

delta $leftres \ alpha \ beta \leftrightarrow (delta \cdot beta \#) \ alpha$

Proof.

rewrite / leftres.

by [rewrite inv_inc_move inc_residual -comp_inv inv_inc_move inv_invol]. Qed.

Lemma 268 (residual_leftres_assoc) Let $\alpha: X \to Y$, $\beta: Y \to Z$ and $\gamma: Z \to W$. Then,

$$(\alpha \rhd \beta) \lhd \gamma = \alpha \rhd (\beta \lhd \gamma).$$

Lemma $residual_leftres_assoc$ { W X Y Z : eqType}

 $\{alpha: Rel\ X\ Y\}\ \{beta: Rel\ Y\ Z\}\ \{gamma: Rel\ Z\ W\}:$

leftres (alpha beta) gamma = alpha leftres beta gamma.

Proof.

apply inc_lower .

 $move \Rightarrow delta$.

by [rewrite inc_leftres inc_residual -comp_assoc -inc_leftres -inc_residual]. Qed.

End main.

Chapter 11

Library Schroder

```
Require Import Basic_Notations.

Require Import Basic_Lemmas.

Require Import Relation_Properties.

Require Import Functions_Mappings.

Require Import Dedekind.

Require Import Residual.

Require Import Logic.FunctionalExtensionality.

Module main (def: Relation).

Import def.

Module Basic_Lemmas := Basic_Lemmas.main def.

Module Relation_Properties := Relation_Properties.main def.

Module Functions_Mappings := Functions_Mappings.main def.


Module Dedekind := Dedekind.main def.

Module Residual := Residual.main def.

Import Basic_Lemmas Relation_Properties Functions_Mappings Dedekind Residual.
```

11.1 Schröder 圏の性質

この節では、特記が無い限り、記号は以下の図式に従って割り振られるものとする.



Lemma 269 (schroder_equivalence1, schroder_equivalence2)

```
\alpha \cdot \beta \sqsubseteq \delta \Leftrightarrow \alpha^{\sharp} \cdot \delta^{-} \sqsubseteq \beta^{-} \Leftrightarrow \delta^{-} \cdot \beta^{\sharp} \sqsubseteq \alpha^{-}.
```

```
Lemma schroder\_equivalence1
 \{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{delta : Rel \ X \ Z\}:
 (alpha \cdot beta) delta \leftrightarrow (alpha \# \cdot delta \hat{}) beta \hat{}.
Proof.
split; move \Rightarrow H.
rewrite bool_lemma2 complement_invol.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply bool\_lemma2 in H.
rewrite cap_comm inv_invol H comp_empty_r.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite bool_lemma2.
apply inc\_antisym.
apply (@inc_trans _ _ _ (dedekind1)).
apply bool\_lemma2 in H.
rewrite cap_comm -(@complement_invol _ _ beta) H comp_empty_r.
apply inc\_reft.
apply inc\_empty\_alpha.
Qed.
Lemma schroder\_equivalence2
 \{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{delta : Rel \ X \ Z\}:
 (alpha \cdot beta) \quad delta \leftrightarrow (delta \cdot beta \#)
Proof.
split; move \Rightarrow H.
rewrite bool_lemma2 complement_invol.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply bool\_lemma2 in H.
rewrite cap_comm inv_invol H comp_empty_l.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite bool_lemma2.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply bool\_lemma2 in H.
rewrite cap_comm -(@complement_invol _ _ alpha) H comp_empty_l.
apply inc\_reft.
```

apply inc_empty_alpha . Qed.

Lemma 270 (function_inv_complement) Let α and τ be functions. Then,

$$(\alpha \cdot \beta \cdot \tau^{\sharp})^{-} = \alpha \cdot \beta^{-} \cdot \tau^{\sharp}.$$

```
Lemma function_inv_complement
 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{tau : Rel \ V \ Z\}:
 function\_r \ alpha \rightarrow function\_r \ tau \rightarrow
 ((alpha \cdot beta) \cdot tau \#) = (alpha \cdot beta) \cdot tau \#.
Proof.
move \Rightarrow H H0.
apply inc\_antisym.
rewrite bool_lemma1 complement_invol.
apply inc\_antisym.
rewrite -comp_cup_distr_r -comp_cup_distr_l complement_classic.
apply (@inc\_trans \_ \_ \_ (((alpha \cdot alpha \#) \cdot X V) \cdot (tau \cdot tau \#))).
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot X V)).
apply comp\_inc\_compat\_b\_ab.
apply H.
apply comp\_inc\_compat\_a\_ab.
apply H0.
rewrite -comp_assoc (@comp_assoc _ _ _ alpha) (@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
rewrite bool_lemma2 complement_invol.
apply inc\_antisym.
rewrite -(function_cap_distr H H0) cap_comm cap_complement_empty comp_empty_r comp_empty_l.
apply inc\_reft.
apply inc\_empty\_alpha.
Qed.
```

Lemma 271 (schroder_univalent1) Let α be a univalent relation and $\beta \subseteq \beta'$. Then,

$$\alpha \cdot (\beta' \sqcap \beta^{-}) = \alpha \cdot \beta' \sqcap (\alpha \cdot \beta)^{-}.$$

```
Lemma schroder\_univalent1

\{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta \ beta' : Rel \ Y \ Z\}: \ univalent\_r \ alpha \rightarrow beta \ beta' \rightarrow \ alpha \cdot (beta' \ beta') = (alpha \cdot beta') \ (alpha \cdot beta) ^.
```

```
Proof.
move \Rightarrow H H0.
apply (@cap_cup_unique _ _ (alpha • beta)).
                               (alpha \cdot (beta' beta'))) with (XZ).
replace ((alpha • beta)
rewrite (@cap_comm _ _ (alpha • beta')) -cap_assoc.
by [rewrite cap_complement_empty cap_comm cap_empty].
apply inc\_antisym.
apply inc\_empty\_alpha.
                                                ((alpha \cdot beta') \quad (alpha \cdot beta^{\circ}))).
apply @inc\_trans \_ \_ \_ ((alpha \cdot beta))
apply cap\_inc\_compat\_l.
apply comp\_cap\_distr\_l.
                                               (alpha • beta ^)).
replace (XZ) with ((alpha \cdot beta)
apply cap\_inc\_compat\_l.
apply cap_r.
apply inc\_antisym.
move: (@univalent\_residual \_ \_ \_ \_ beta H) \Rightarrow H1.
rewrite -inc\_rpc.
rewrite residual_to_complement in H1.
apply H1.
apply inc\_empty\_alpha.
apply inc_{-}def2 in H0.
rewrite -comp_cup_distr_l cup_cap_distr_l.
rewrite -H0 complement_classic cap_universal.
rewrite cup\_cap\_distr\_l -comp\_cup\_distr\_l.
by [rewrite -H0 complement_classic cap_universal].
Qed.
  Lemma 272 (schroder_univalent2) Let \alpha be a univalent relation. Then,
                                 \alpha \cdot \beta^- = \alpha \cdot \nabla_{YZ} \sqcap (\alpha \cdot \beta)^-.
Lemma schroder\_univalent2 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \cdot beta \hat{} = (alpha \cdot Y Z)
                                                                  (alpha \cdot beta) ^.
Proof.
move \Rightarrow H.
move: (@schroder_univalent1 _ _ alpha beta ( Y Z) H (@inc_alpha_universal _ _ ))
\Rightarrow H0.
rewrite cap_comm cap_universal in H0.
apply H0.
Qed.
```

CHAPTER 11. LIBRARY SCHRODER

Lemma 273 (schroder_univalent3) Let α be a univalent relation. Then,

$$(\alpha \cdot \beta)^- = (\alpha \cdot \nabla_{YZ})^- \sqcup \alpha \cdot \beta^-.$$

Lemma $schroder_univalent3$ { $X \ Y \ Z : eqType$ } { $alpha : Rel \ X \ Y$ } {beta : $Rel \ Y \ Z$ }: $univalent_r \ alpha \rightarrow (alpha \cdot beta) ^ = (alpha \cdot Y \ Z) ^ (alpha \cdot beta ^).$ Proof.

move $\Rightarrow H$.

rewrite ($schroder_univalent2 \ H$).

rewrite $cup_cap_distr_l \ cup_comm \ complement_classic \ cap_comm \ cap_universal$.

apply inc_def2 .

apply inc_def2 .

apply $comp_inc_compat_r$.

apply $comp_inc_compat_ab_ab$.

apply $inc_alpha_universal$.

Qed.

Lemma 274 (schroder_univalent4) Let α be a univalent relation. Then,

$$\alpha \rhd \beta = (\alpha \cdot \nabla_{YZ})^- \sqcup \alpha \cdot \beta.$$

Lemma $schroder_univalent4$ { $X \ Y \ Z : eqType$ } { $alpha : Rel \ X \ Y$ } {beta : $Rel \ Y \ Z$ }: $univalent_r \ alpha \rightarrow alpha$ beta = $(alpha \cdot Y \ Z) \cdot (alpha \cdot beta)$.

Proof.

move $\Rightarrow H$.

rewrite $(residual_property9 \ H)$.

apply $Logic.eq_sym$.

 $\verb"apply" cup_to_rpc".$

Qed.

Lemma 275 (schroder_universal) Let $\nabla_{XZ} \cdot \nabla_{ZW} = \nabla_{XW}$. Then,

$$(\alpha \cdot \nabla_{YZ})^- \cdot \nabla_{ZW} = (\alpha \cdot \nabla_{YW})^-.$$

Lemma $schroder_universal\ \{W\ X\ Y\ Z\ : eqType\}\ \{alpha\ :\ Rel\ X\ Y\}:$ $(XZ \cdot ZW) = XW \rightarrow (alpha \cdot YZ) \cdot ZW = (alpha \cdot YW) \cdot .$ Proof. $move \Rightarrow H.$ $apply\ (@cap_cup_unique__(alpha \cdot YW)).$ $rewrite\ cap_complement_empty\ cap_comm.$ $apply\ (@inc_trans____(dedekind2)).$ $apply\ (@inc_trans____(((alpha \cdot YZ) \cdot (alpha \cdot YZ)) \cdot ZW)).$

CHAPTER 11. LIBRARY SCHRODER

```
apply comp\_inc\_compat\_ab\_a'b.
apply cap\_inc\_compat\_l.
rewrite comp\_assoc.
apply comp_inc_compat_ab_ab'.
apply inc\_alpha\_universal.
rewrite cap_comm cap_complement_empty comp_empty_l.
apply inc_refl.
apply inc\_empty\_alpha.
rewrite complement_classic.
apply inc\_antisym.
apply inc\_alpha\_universal.
rewrite -H -(@complement_classic _ _ (alpha • Y Z)) comp_cup_distr_r.
apply cup\_inc\_compat\_r.
rewrite comp_-assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc_alpha_universal.
Qed.
```

Lemma 276 (residual_inv)

Qed.

$$(\alpha \rhd \beta)^{\sharp} = \beta^{-\sharp} \rhd \alpha^{-\sharp}.$$

```
Lemma residual\_inv {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}: (alpha \ beta) # = (beta \hat{}) # (alpha \hat{}) #.

Proof.

rewrite residual\_to\_complement \ residual\_to\_complement.

by [rewrite -inv\_complement \ complement\_invol \ inv\_complement \ comp\_inv].
```

Lemma 277 (residual_cupP_distr_l, residual_cup_distr_l) Let α be a univalent relation, $f:(V \to W) \to (Y \to Z)$ and $\exists \beta, P(\beta)$. Then,

$$\alpha \rhd (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \rhd f(\beta)).$$

```
Lemma residual\_cupP\_distr\_l {V \ W \ X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {f : Rel \ V \ W \to Rel \ Y \ Z} {P : Rel \ V \ W \to Prop}: univalent\_r \ alpha \to (\exists \ beta' : Rel \ V \ W, \ P \ beta') \to alpha \quad ( \ \_\{P\} \ f) = \ \_\{P\} \ (fun \ beta : Rel \ V \ W \Rightarrow alpha \quad f \ beta).

Proof.

move \Rightarrow H.

elim \Rightarrow beta' \ H0.

rewrite (schroder\_univalent4 \ H) \ comp\_cupP\_distr\_l.

replace (\ \_\{P\} \ (fun \ beta : Rel \ V \ W \Rightarrow alpha \quad f \ beta)) with (\ \_\{P\} \ (fun \ beta : Rel \ V \ W \Rightarrow alpha \quad f \ beta))
```

CHAPTER 11. LIBRARY SCHRODER

```
Rel\ V\ W \Rightarrow (alpha\ \cdot
                               YZ)^{\hat{}}
                                            (alpha • f beta))).
apply (@cap\_cup\_unique\_\_(alpha \cdot
                                                 (Y Z)).
rewrite cap_cup_distr_l cap_cupP_distr_l cap_complement_empty cup_comm cup_empty.
rewrite cap\_cupP\_distr\_l.
apply cupP_-eq.
move \Rightarrow qamma\ H1.
by [rewrite cap_cup_distr_l cap_complement_empty cup_comm cup_empty].
rewrite -cup_assoc complement_classic cup_comm cup_universal.
rewrite -(@complement_invol _ _ (alpha •
apply bool_lemma1.
rewrite complement_invol.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot Y Z) ^ (alpha \cdot f beta'))).
apply cup_{-}l.
move: beta' H0.
apply inc\_cupP.
apply inc_reft.
apply cupP_{-}eq.
move \Rightarrow qamma\ H1.
by [rewrite (schroder_univalent4 H)].
Qed.
Lemma residual\_cup\_distr\_l
 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta \ beta' : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow
                       beta') = (alpha)
                                                      (alpha
 alpha
            (beta
                                             beta)
                                                                    beta').
Proof.
move \Rightarrow H.
rewrite cup\_to\_cupP (@cup\_to\_cupP _ _ _ _ id).
apply (residual\_cupP\_distr\_l\ H).
∃ beta.
by [left].
Qed.
  Lemma 278 (residual_capP_distr_r, residual_cap_distr_r) Let f: (Y \rightarrow Z) \rightarrow
  (I \rightarrow X) and \exists \alpha, P(\alpha). Then,
                                (\sqcap_{P(\alpha)} f(\alpha)^{\sharp}) \rhd \rho = \sqcup_{P(\alpha)} (f(\alpha)^{\sharp} \rhd \rho).
Lemma residual\_capP\_distr\_r
 \{X \ Y \ Z : eqType\} \ \{rho : Rel \ i \ X\} \ \{f : Rel \ Y \ Z \rightarrow Rel \ i \ X\} \ \{P : Rel \ Y \ Z \rightarrow Prop\}:
 (\exists alpha' : Rel Y Z, P alpha') \rightarrow
 ( _{P} (fun \ alpha : Rel \ Y \ Z \Rightarrow f \ alpha \#)) \quad rho = _{P} (fun \ alpha : Rel \ Y \ Z \Rightarrow f \ alpha \#))
f alpha \#
             rho).
```

```
Proof.
elim \Rightarrow alpha' H.
rewrite residual_to_complement.
\texttt{rewrite -} (@complement\_invol \_ \_ ( \_\{P\} (\texttt{fun} \ alpha : Rel \ Y \ Z \Rightarrow f \ alpha \ \# \quad rho))).
apply f_equal.
rewrite de_{-}morgan3.
replace (fun alpha: Rel Y Z \Rightarrow (f alpha # rho) \hat{}) with (fun alpha: Rel Y Z \Rightarrow f
alpha \# \cdot rho \hat{}).
apply inc\_antisym.
apply comp\_capP\_distr\_r.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
apply (@inc\_trans \_ \_ \_ ((( \_{P} (fun alpha : Rel Y Z \Rightarrow f alpha \# \cdot rho ^))) \cdot (fun alpha : Rel Y Z \Rightarrow f alpha \# \cdot rho ^))
alpha' \# \cdot rho ^) \#) \cdot (f \ alpha' \# \cdot rho ^))).
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_ab'.
move: alpha'H.
apply inc\_capP.
rewrite inv\_capP\_distr.
apply inc_refl.
move: alpha' H.
apply inc\_capP.
apply inc\_reft.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (comp\_capP\_distr\_r)).
apply inc\_capP.
move \Rightarrow beta H\theta.
apply (@inc\_trans \_ \_ \_ ((f beta \# \cdot rho ^) \cdot ((f alpha' \# \cdot rho ^) \# \cdot f alpha' \#))).
move : beta H0.
apply inc\_capP.
apply inc\_reft.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite residual_to_complement complement_invol].
Qed.
End main.
```

Chapter 12

Library Sum_Product

```
Require Import Basic_Notations.
Require Import Basic_Lemmas.
Require Import Relation_Properties.
Require Import Functions_Mappings.
Require Import Dedekind.
Require Import Conjugate.
Require Import Domain.
Require Import Logic. Indefinite Description.
Module main (def : Relation).
Import def.
Module\ Basic\_Lemmas := Basic\_Lemmas.main\ def.
Module\ Relation\_Properties := Relation\_Properties.main\ def.
Module\ Functions\_Mappings := Functions\_Mappings.main\ def.
Module Dedekind := Dedekind.main def.
Module Conjugate := Conjugate.main def.
Module Domain := Domain.main \ def.
Import Basic_Lemmas Relation_Properties Functions_Mappings Dedekind Conjugate Do-
main.
```

12.1 関係の直和

12.1.1 入射対,関係直和の定義

入射対の存在公理 (Axiom 23) で入射対が存在することまでは仮定済みなので、実際に入射対 $j:A \rightarrow A+B, k:B \rightarrow A+B$ を定義する関数を定義する.

```
Definition sum_r (A B : eqType):

\{x : (Rel \ A \ (sum_eqType \ A \ B)) \times (Rel \ B \ (sum_eqType \ A \ B)) \mid
```

```
(fst \ x) \cdot (fst \ x) \# = Id \ A \wedge (snd \ x) \cdot (snd \ x) \# = Id \ B \wedge
 (fst \ x) \cdot (snd \ x) \# =
                            A B \wedge
 ((fst \ x) \# \cdot (fst \ x))
                           ((snd \ x) \# \cdot (snd \ x)) = Id (sum\_eqType \ A \ B).
apply constructive_indefinite_description.
elim (@pair_of_inclusions \ A \ B) \Rightarrow j.
elim \Rightarrow k H.
\exists (j,k).
simpl.
apply H.
Defined.
Definition inl_r (A B : eqType) := fst (sval (sum_r A B)).
Definition inr_r (A B : eqType) := snd (sval (sum_r A B)).
  またこの定義による入射対が、入射対としての性質 (Axiom 23) + \alpha を満たしていること
  も事前に証明しておく.
Lemma inl\_id \{A B : eqType\}: inl\_r A B \cdot inl\_r A B \# = Id A.
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inr_id \{A B : eqType\}: inr_r A B \cdot inr_r A B \# = Id B.
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inl\_inr\_empty \{A B : eqType\}: inl\_r A B \cdot inr\_r A B \# = 1
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inr\_inl\_empty \{A B : eqType\}: inr\_r A B \cdot inl\_r A B \# = 1
                                                                           B A.
Proof.
apply inv_invol2.
rewrite comp_inv inv_invol inv_empty.
apply inl\_inr\_empty.
Qed.
Lemma inl\_inr\_cup\_id \{A \ B : eqType\}:
 (inl\_r \ A \ B \ \# \ \cdot \ inl\_r \ A \ B) = Id \ (sum\_eqType \ A \ B).
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inl\_function \{A B : eqType\}: function\_r (inl\_r A B).
Proof.
move: (proj2\_sig\ (sum\_r\ A\ B)).
```

```
\mathtt{elim} \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2.
split.
rewrite /total_{-}r.
rewrite H.
apply inc_refl.
rewrite /univalent_r.
rewrite -H2.
apply cup_{-}l.
Qed.
Lemma inr\_function \{A \ B : eqType\}: function\_r (inr\_r \ A \ B).
Proof.
move: (proj2\_sig\ (sum\_r\ A\ B)).
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2.
split.
rewrite /total_{-}r.
rewrite H0.
apply inc_refl.
rewrite /univalent_r.
rewrite -H2.
apply cup_r.
Qed.
  さらに \alpha:A \to C と \beta:B \to C の関係直和 \alpha \perp \beta:A+B \to C を, \alpha \perp \beta:=j^{\sharp} \cdot \alpha \sqcup k^{\sharp} \cdot \beta
```

で定義する.

```
Definition Rel\_sum \{A \ B \ C : eqType\} \ (alpha : Rel \ A \ C) \ (beta : Rel \ B \ C):=
 (inl\_r \ A \ B \ \# \ \bullet \ alpha) \quad (inr\_r \ A \ B \ \# \ \bullet \ \mathsf{beta}).
```

12.1.2関係直和の性質

```
Lemma 279 (sum_inc_compat) Let \alpha, \alpha' : A \to C and \beta, \beta' : B \to C. Then,
                                                 \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \bot \beta \sqsubseteq \alpha' \bot \beta'.
```

```
Lemma sum\_inc\_compat
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ C\} \{beta \ beta' : Rel \ B \ C\}:
             alpha' \rightarrow \mathtt{beta}
                                    beta' \rightarrow Rel\_sum \ alpha \ beta Rel\_sum \ alpha' \ beta'.
 alpha
Proof.
```

```
CHAPTER 12. LIBRARY SUM_PRODUCT
move \Rightarrow H H0.
apply cup\_inc\_compat.
apply (comp\_inc\_compat\_ab\_ab' H).
apply (comp\_inc\_compat\_ab\_ab' H0).
Qed.
  Lemma 280 (sum_inc_compat_l) Let \alpha : A \to C and \beta, \beta' : B \to C. Then,
                                         \beta \sqsubseteq \beta' \Rightarrow \alpha \bot \beta \sqsubseteq \alpha \bot \beta'.
Lemma sum\_inc\_compat\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta \ beta' : Rel \ B \ C\}:
           beta' \rightarrow Rel\_sum \ alpha \ beta
                                                 Rel_sum alpha beta'.
Proof.
move \Rightarrow H.
apply (sum\_inc\_compat (@inc\_refl \_ \_ alpha) H).
Qed.
  Lemma 281 (sum_inc_compat_r) Let \alpha, \alpha' : A \to C and \beta : B \to C. Then,
                                         \alpha \sqsubseteq \alpha' \Rightarrow \alpha \bot \beta \sqsubseteq \alpha' \bot \beta.
Lemma sum\_inc\_compat\_r
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ C\} \{beta : Rel B \ C\}:
            alpha' \rightarrow Rel\_sum \ alpha \ beta Rel\_sum \ alpha' \ beta.
Proof.
move \Rightarrow H.
apply (sum_inc_compat H (@inc_refl _ beta)).
Qed.
  Lemma 282 (total_sum) Let \alpha: A \rightarrow C and \beta: B \rightarrow C are total relations, then
  \alpha \perp \beta is also a total relation.
Lemma total\_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 total\_r \ alpha \rightarrow total\_r \ beta \rightarrow total\_r \ (Rel\_sum \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /total_r/Rel_sum.
\texttt{rewrite-} inl\_inr\_cup\_id\ inv\_cup\_distr\ comp\_cup\_distr\_l\ comp\_cup\_distr\_r\ comp\_cup\_distr\_r.
rewrite comp_inv comp_inv inv_invol inv_invol.
apply cup\_inc\_compat.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cup\_l)).
```

rewrite comp_assoc -(@comp_assoc _ _ _ alpha).

```
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cup\_r)).
rewrite comp_assoc -(@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H0).
Qed.
  Lemma 283 (univalent_sum) Let \alpha : A \rightarrow C and \beta : B \rightarrow C are univalent relations,
  then \alpha \perp \beta is also a univalent relation.
Lemma univalent\_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (Rel\_sum \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /univalent_r/Rel_sum.
rewrite inv_cup_distr comp_cup_distr_l comp_cup_distr_r comp_cup_distr_r.
\verb"rewrite" comp\_inv comp\_inv inv\_invol inv\_invol.
rewrite comp_assoc -(@comp_assoc _ _ _ (inl_r A B)) inl_id comp_id_l.
\verb"rewrite" comp\_assoc -(@comp\_assoc \_ \_ \_ \_ (inr\_r \ A \ B)) \ inr\_inl\_empty \ comp\_empty\_l
comp\_empty\_r cup\_empty.
rewrite-cup_assoc comp_assoc -(@comp_assoc _ _ _ (inl_r A B)) inl_inr_empty comp_empty_l
comp\_empty\_r cup\_empty.
rewrite comp\_assoc -(@comp\_assoc _ _ _ (inr\_r A B)) inr\_id comp\_id\_l.
apply inc_-cup.
split.
apply H.
apply H0.
Qed.
  Lemma 284 (function_sum) Let \alpha: A \to C and \beta: B \to C are functions, then \alpha \perp \beta
  is also a function.
Lemma function_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 function\_r \ alpha \rightarrow function\_r \ \mathsf{beta} \rightarrow function\_r \ (Rel\_sum \ alpha \ \mathsf{beta}).
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (total_sum H H1).
apply (univalent_sum H0 H2).
Qed.
```


Lemma 285 (sum_conjugate) Let $\alpha: A \rightarrow C$, $\beta: B \rightarrow C$ and $\gamma: A+B \rightarrow C$ be relations, $j: A \rightarrow A+B$ and $k: B \rightarrow A+B$ be inclusions. Then,

$$j \cdot \gamma = \alpha \wedge k \cdot \gamma = \beta \Leftrightarrow \gamma = \alpha \perp \beta.$$

```
Lemma sum\_conjugate
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\} \{gamma : Rel \ (sum\_eqType \ A \ B)\}
C}:
 inl_r A B \cdot gamma = alpha \wedge inr_r A B \cdot gamma = beta \leftrightarrow
 gamma = Rel\_sum \ alpha \ beta.
Proof.
split; move \Rightarrow H.
elim H \Rightarrow H0 \ H1.
rewrite -(@comp\_id\_l\_\_\_gamma).
rewrite -inl_inr_cup_id comp_cup_distr_r comp_assoc comp_assoc.
by [rewrite H0 H1].
split.
rewrite H comp\_cup\_distr\_l -comp\_assoc -comp\_assoc.
rewrite inl_id inl_inr_empty comp_id_l comp_empty_l.
by [rewrite cup\_empty].
rewrite H comp\_cup\_distr\_l -comp\_assoc -comp\_assoc.
rewrite inr_id inr_inl_empty comp_id_l comp_empty_l.
by [rewrite cup\_comm\ cup\_empty].
Qed.
```

Lemma 286 (sum_comp) In below figure,

$$(\alpha \perp \beta)^{\sharp} \cdot (\gamma \perp \delta) = \alpha^{\sharp} \cdot \gamma \sqcup \beta^{\sharp} \cdot \delta.$$



```
Lemma sum\_comp {W \ X \ Y \ Z : eqType} {alpha : Rel \ Y \ X} {beta : Rel \ Z \ X} {gamma : Rel \ Y \ W} {delta : Rel \ Z \ W}: (Rel\_sum \ alpha \ beta) # • Rel\_sum \ gamma \ delta = (alpha \ \# \ \bullet \ gamma) (beta # • delta).

Proof.
rewrite /Rel\_sum.
```

```
rewrite inv\_cup\_distr\ comp\_cup\_distr\_l\ comp\_cup\_distr\_r\ comp\_cup\_distr\_r. rewrite comp\_inv\ comp\_inv\ inv\_invol\ inv\_invol. apply f\_equal2. rewrite comp\_assoc\ -(@comp\_assoc\ -(= (inl\_r\ Y\ Z))\ inl\_id\ comp\_id\_l. by [rewrite comp\_assoc\ -(@comp\_assoc\ -(= (inl\_r\ Y\ Z))\ inr\_inl\_empty\ comp\_empty\_l\ comp\_empty\_r\ cup\_empty]. rewrite <math>comp\_assoc\ -(@comp\_assoc\ -(= (inl\_r\ Y\ Z))\ inl\_inr\_empty\ comp\_empty\_l\ comp\_empty\_r\ cup\_comm\ cup\_empty. by [rewrite comp\_assoc\ -(@comp\_assoc\ -(= (inr\_r\ Y\ Z))\ inr\_id\ comp\_id\_l]. Qed.
```

12.1.3 分配法則

```
Lemma 287 (sum_cap_distr_l) Let \alpha: A \to C and \beta, \beta': B \to C. Then, \alpha \bot (\beta \sqcap \beta') \sqsubseteq (\alpha \bot \beta) \sqcap (\alpha \bot \beta').
```

```
Lemma sum\_cap\_distr\_l {A \ B \ C : eqType} {alpha : Rel \ A \ C} {beta beta' : Rel \ B \ C}: Rel\_sum \ alpha (beta beta') (Rel\_sum \ alpha beta Rel\_sum \ alpha \ beta'). Proof. rewrite -cup\_cap\_distr\_l. apply cup\_inc\_compat\_l. apply comp\_cap\_distr\_l. Qed.
```

```
Lemma 288 (sum_cap_distr_r) Let \alpha, \alpha' : A \to C and \beta : B \to C. Then, (\alpha \sqcap \alpha') \bot \beta \sqsubseteq (\alpha \bot \beta) \sqcap (\alpha' \bot \beta).
```

```
Lemma sum\_cap\_distr\_r {A \ B \ C : eqType} {alpha \ alpha' : Rel \ A \ C} {beta : Rel \ B \ C}: Rel\_sum \ (alpha \ alpha') beta (Rel\_sum \ alpha \ beta \ Rel\_sum \ alpha' \ beta). Proof. rewrite -cup\_cap\_distr\_r. apply cup\_inc\_compat\_r. apply comp\_cap\_distr\_l. Qed.
```

Lemma 289 (sum_cup_distr_l) Let $\alpha : A \rightarrow C$ and $\beta, \beta' : B \rightarrow C$. Then,

$$\alpha \perp (\beta \sqcup \beta') = (\alpha \perp \beta) \sqcup (\alpha \perp \beta').$$

Lemma $sum_cup_distr_l$

 $\{A\ B\ C: eqType\}\ \{alpha: Rel\ A\ C\}\ \{beta\ beta': Rel\ B\ C\}:$

Rel_sum alpha (beta beta') = Rel_sum alpha beta Rel_sum alpha beta'.

Proof.

rewrite $-cup_assoc$ (@ $cup_comm__$ (Rel_sum alpha beta)) $-cup_assoc$. by [rewrite cup_idem cup_assoc $-comp_cup_distr_l$]. Qed.

Lemma 290 (sum_cup_distr_r) Let $\alpha, \alpha' : A \rightarrow C$ and $\beta : B \rightarrow C$. Then,

$$(\alpha \sqcup \alpha') \bot \beta = (\alpha \bot \beta) \sqcup (\alpha' \bot \beta).$$

Lemma $sum_cup_distr_r$

 $\{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ C\} \{$ beta : Rel $B \ C\}$:

Rel_sum (alpha alpha') beta = (Rel_sum alpha beta Rel_sum alpha' beta).

Proof.

rewrite cup_assoc (@ $cup_comm__$ (inr_r A B # • beta)) cup_assoc . by [rewrite cup_idem - cup_assoc - $comp_cup_distr_l$]. Qed.

Lemma 291 (comp_sum_distr_r) Let $\alpha: A \rightarrow C, \ \beta: B \rightarrow C \ and \ \gamma: C \rightarrow D.$ Then,

$$(\alpha \bot \beta) \cdot \gamma = \alpha \cdot \gamma \bot \beta \cdot \gamma.$$

Lemma $comp_sum_distr_r$

 $\{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ C\} \{$ beta : $Rel \ B \ C\} \{gamma : Rel \ C \ D\}$: $(Rel_sum \ alpha \ beta) \cdot gamma = Rel_sum \ (alpha \cdot gamma) \ (beta \cdot gamma).$

Proof.

by [rewrite $comp_cup_distr_r$ $comp_assoc$ $comp_assoc$]. Qed.

12.2 関係の直積

12.2.1 射影対,関係直積の定義

Definition $prod_r$ (A B : eqType):

射影対の存在公理 (Axiom 24) で射影対が存在することまでは仮定済みなので、実際に射影対 $p:A\times B\to A, k:A\times B\to B$ を定義する関数を定義する.

 $\{x: (Rel\ (prod_eqType\ A\ B)\ A) \times (Rel\ (prod_eqType\ A\ B)\ B) \mid$

```
(fst \ x) \# \cdot (snd \ x) = A B \wedge
 ((fst\ x)\ \cdot\ (fst\ x)\ \#)\ ((snd\ x)\ \cdot\ (snd\ x)\ \#)=Id\ (prod\_eqType\ A\ B)\ \land
 univalent_r (fst \ x) \land univalent_r (snd \ x).
apply constructive_indefinite_description.
elim (@pair_of_projections \ A \ B) \Rightarrow p.
elim \Rightarrow q H.
\exists (p,q).
simpl.
apply H.
Defined.
Definition fst_r (A B : eqType) := fst (sval (prod_r A B)).
Definition snd_r (A B : eqType):= snd (sval (prod_r A B)).
  またこの定義による射影対が、射影対としての性質 (Axiom 24) + \alpha を満たしていること
  も事前に証明しておく.
Lemma fst\_snd\_universal \{A B : eqType\}: fst\_r A B \# \bullet snd\_r A B =
                                                                                A B.
Proof.
apply (proj2\_sig\ (prod\_r\ A\ B)).
Lemma snd\_fst\_universal\ \{A\ B: eqType\}:\ snd\_r\ A\ B\ \#\ \bullet\ fst\_r\ A\ B=
Proof.
apply inv_invol2.
rewrite comp_inv inv_invol inv_universal.
apply fst\_snd\_universal.
Qed.
Lemma fst\_snd\_cap\_id \{A B : eqType\}:
 (fst_r \ A \ B \cdot fst_r \ A \ B \#) \quad (snd_r \ A \ B \cdot snd_r \ A \ B \#) = Id (prod_eqType \ A \ B).
Proof.
apply (proj2\_sig\ (prod\_r\ A\ B)).
Lemma fst\_function \{A B : eqType\}: function\_r (fst\_r A B).
Proof.
```

```
move: (proj2\_sig\ (prod\_r\ A\ B)).
elim \Rightarrow H.
elim \Rightarrow H0 \ H1.
split.
rewrite /total_{-}r.
rewrite -H0.
apply cap_{-}l.
apply H1.
Qed.
Lemma snd\_function \{A \ B : eqType\}: function\_r (snd\_r \ A \ B).
move: (proj2\_sig\ (prod\_r\ A\ B)).
elim \Rightarrow H.
elim \Rightarrow H0 \ H1.
split.
rewrite /total_r.
rewrite -H0.
apply cap_{-}r.
apply H1.
Qed.
  さらに \alpha:A \to B と \beta:A \to C の関係直積 \alpha \top \beta:A \to B \times C を, \alpha \top \beta:=\alpha \cdot p^\sharp \sqcap \beta \cdot q^\sharp
```

さらに $\alpha:A \to B \succeq \beta:A \to C$ の関係直積 $\alpha \top \beta:A \to B \times C$ を, $\alpha \top \beta:=\alpha \cdot p^\sharp \sqcap \beta \cdot q^\sharp$ で定義する.

```
Definition Rel\_prod \{A \ B \ C : eqType\} \ (alpha : Rel \ A \ B) \ (beta : Rel \ A \ C) := (alpha \cdot fst\_r \ B \ C \ \#) \ (beta \cdot snd\_r \ B \ C \ \#).
```

12.2.2 関係直積の性質

```
Lemma 292 (prod_inc_compat) Let \alpha, \alpha' : A \to B and \beta, \beta' : A \to C. Then, \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \top \beta \sqsubseteq \alpha' \top \beta'.
```

```
Lemma prod\_inc\_compat {A \ B \ C : eqType} {alpha \ alpha' : Rel \ A \ B} {beta beta' : Rel \ A \ C}: alpha \ alpha' \rightarrow beta \ beta' \rightarrow Rel\_prod \ alpha \ beta \ Rel\_prod \ alpha' \ beta'. Proof.

move \Rightarrow H \ H0.

apply cap\_inc\_compat.

apply (comp\_inc\_compat\_ab\_a'b \ H).

apply (comp\_inc\_compat\_ab\_a'b \ H0).

Qed.
```

```
Lemma 293 (prod_inc_compat_l) Let \alpha : A \rightarrow B and \beta, \beta' : A \rightarrow C. Then,
                                         \beta \sqsubseteq \beta' \Rightarrow \alpha \top \beta \sqsubseteq \alpha \top \beta'.
Lemma prod\_inc\_compat\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ beta' : Rel \ A \ C\}:
            beta' \rightarrow Rel\_prod\ alpha\ beta'. Rel\_prod\ alpha\ beta'.
 beta
Proof.
move \Rightarrow H.
apply (prod_inc_compat (@inc_refl _ alpha) H).
Qed.
  Lemma 294 (prod_inc_compat_r) Let \alpha, \alpha' : A \to B and \beta : A \to C. Then,
                                         \alpha \sqsubseteq \alpha' \Rightarrow \alpha \top \beta \sqsubseteq \alpha' \top \beta.
Lemma prod\_inc\_compat\_r
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
             alpha' \rightarrow Rel\_prod\ alpha\ beta Rel\_prod\ alpha'\ beta.
 alpha
Proof.
move \Rightarrow H.
apply (prod\_inc\_compat \ H \ (@inc\_refl \_ \_ beta)).
Qed.
  Lemma 295 (total_prod) Let \alpha: A \rightarrow B and \beta: A \rightarrow C are total relations, then
  \alpha \top \beta is also a total relation.
Lemma total\_prod \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 total\_r \ alpha \rightarrow total\_r \ beta \rightarrow total\_r \ (Rel\_prod \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite domain_total cap_domain cap_comm.
apply Logic.eq_sym.
apply inc\_def1.
apply (@inc\_trans \_ \_ \_ \_ H).
rewrite comp_inv inv_invol comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot (beta \cdot beta \#))).
apply (comp\_inc\_compat\_a\_ab\ H0).
rewrite -comp_assoc -comp_assoc fst_snd_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
```

Lemma 296 (univalent_prod) Let $\alpha : A \to B$ and $\beta : A \to C$ are univalent relations, then $\alpha \top \beta$ is also a univalent relation.

```
Lemma univalent\_prod \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (Rel\_prod \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /univalent_r/Rel_prod.
rewrite inv_cap_distr comp_inv inv_invol comp_inv inv_invol.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
rewrite -fst\_snd\_cap\_id.
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
rewrite comp\_assoc -(@comp\_assoc _ _ _ _ beta).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H0).
Qed.
```

Lemma 297 (function_prod) Let $\alpha : A \to B$ and $\beta : A \to C$ are functions, then $\alpha \top \beta$ is also a function.

```
Lemma function_prod {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}: function_r alpha \rightarrow function_r beta \rightarrow function_r (Rel_prod alpha beta). Proof. elim \Rightarrow H H0. elim \Rightarrow H1 H2. split. apply (total_prod H H1). apply (univalent_prod H0 H2). Qed.
```

```
Lemma 298 (prod_fst_surjection) Let p: B \times C \to B be a projection. Then, "p is a surjection" \Leftrightarrow \forall D, \nabla_{BD} = \nabla_{BC} \cdot \nabla_{CD}.
```

Lemma $prod_fst_surjection \{B \ C : eqType\}:$

```
surjection_r (fst_r B C) \leftrightarrow \forall D : eqType,
                                                      B D =
                                                                  B C \cdot
                                                                                C D.
Proof.
split; move \Rightarrow H.
move \Rightarrow D.
elim H \Rightarrow H0 \ H1.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((fst\_r \ B \ C \# \cdot (fst\_r \ B \ C \#) \#) \cdot B \ D)).
apply (comp\_inc\_compat\_b\_ab\ H1).
rewrite inv\_invol.
apply (@inc\_trans \_ \_ \_ (((fst\_r \ B \ C \# \cdot snd\_r \ B \ C) \cdot (snd\_r \ B \ C \# \cdot fst\_r \ B \ C))
   B(D)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc - (@comp\_assoc \_ \_ \_ \_ (snd\_r \ B \ C)).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_b\_ab.
apply snd_-function.
rewrite (@comp_assoc _ _ _ _ ( B D)).
apply comp\_inc\_compat.
apply inc_alpha_universal.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
split.
apply fst_function.
rewrite /total_r.
rewrite -(@cap\_universal\_\_(Id\ B))(H\ B)-(@fst\_snd\_universal\ B\ C)\ cap\_comm\ comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite comp_{-}id_{-}r.
apply cap_{-}r.
Qed.
  Lemma 299 (prod_snd_surjection) Let q: B \times C \to C be a projection. Then,
                          "q is a surjection" \Leftrightarrow \forall D, \nabla_{CD} = \nabla_{CB} \cdot \nabla_{BD}.
Lemma prod\_snd\_surjection \{B \ C : eqType\}:
 surjection\_r (snd\_r \ B \ C) \leftrightarrow \forall \ D : eqType, \qquad C \ D =
                                                                 C B \cdot
                                                                                BD.
Proof.
split; move \Rightarrow H.
move \Rightarrow D.
elim H \Rightarrow H0 \ H1.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((snd\_r \ B \ C \# \cdot (snd\_r \ B \ C \#) \#) \cdot C \ D)).
```

```
apply (comp_inc_compat_b_ab H1).
rewrite inv_invol.
apply (@inc\_trans \_ \_ \_ (((snd\_r B C \# \cdot fst\_r B C) \cdot (fst\_r B C \# \cdot snd\_r B C)) \cdot
   (CD)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc -(@comp\_assoc _ _ _ _ (fst\_r B C)).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_b\_ab.
apply fst\_function.
rewrite (@comp_assoc _ _ _ _ ( C D)).
apply comp\_inc\_compat.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
split.
apply snd_-function.
rewrite /total_r.
rewrite - (@cap\_universal\_\_(Id\ C))(H\ C) - (@snd\_fst\_universal\ B\ C)\ cap\_comm\ comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp_inc_compat_ab_ab'.
rewrite comp_{-}id_{-}r.
apply cap_r.
Qed.
 Lemma 300 (prod_fst_domain1) Let p: B \times C \rightarrow B be a projection, \alpha: A \rightarrow B
  and \beta: A \rightarrow C. Then,
                                     (\alpha \top \beta) \cdot p = |\beta| \cdot \alpha.
Lemma prod_fst_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • fst\_r\ B\ C=domain\ beta • alpha.
Proof.
rewrite (@comp_inv_inv A A) domain_inv.
rewrite domain_universal2 inv_cap_distr comp_inv inv_invol inv_invol inv_universal.
rewrite -snd_fst_universal.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
rewrite comp_assoc comp_assoc.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_a.
apply fst\_function.
rewrite cap\_comm -comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
```

```
rewrite cap\_comm.
apply inc\_reft.
Qed.
  Lemma 301 (prod_fst_domain2) Let p: B \times C \rightarrow B be a projection, \alpha: A \rightarrow B
  and \beta: A \rightarrow C. Then,
                                  (\alpha \top \beta) \cdot p = \alpha \Leftrightarrow |\alpha| \sqsubset |\beta|.
Lemma prod_fst_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • fst\_r\ B\ C=alpha\leftrightarrow domain\ alpha
                                                                         domain beta.
Proof.
rewrite prod_fst_domain1.
split; move \Rightarrow H.
apply domain_lemma2b.
assert ((domain beta • alpha)
                                       ((beta \cdot beta \#) \cdot alpha)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite H in H0.
apply H0.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_r.
apply (@inc_trans _ _ _ (domain alpha • alpha)).
rewrite domain_comp_alpha1.
apply inc\_reft.
apply (comp\_inc\_compat\_ab\_a'b\ H).
Qed.
  Lemma 302 (prod_snd_domain1) Let q: B \times C \to C be a projection, \alpha: A \to B
  and \beta: A \rightarrow C. Then,
                                       (\alpha \top \beta) \cdot q = |\alpha| \cdot \beta.
Lemma prod_snd_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ {\tt beta}) • snd\_r\ B\ C=domain\ alpha • {\tt beta}.
Proof.
rewrite (@comp_inv_inv A A) domain_inv.
rewrite domain_universal2 inv_cap_distr comp_inv inv_invol inv_invol inv_universal.
rewrite -fst\_snd\_universal.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
rewrite comp_assoc comp_assoc cap_comm.
apply cap\_inc\_compat\_r.
```

apply $comp_inc_compat_ab_a$.

apply $snd_-function$.

```
rewrite cap\_comm -comp\_assoc.
apply dedekind2.
Qed.
  Lemma 303 (prod_snd_domain2) Let q: B \times C \to C be a projection, \alpha: A \to B
  and \beta: A \rightarrow C. Then,
                                  (\alpha \top \beta) \cdot q = \beta \Leftrightarrow |\beta| \sqsubseteq |\alpha|.
Lemma prod_snd_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • snd\_r\ B\ C = beta \leftrightarrow domain\ beta
                                                                      domain\ alpha.
Proof.
rewrite prod_snd_domain1.
split; move \Rightarrow H.
apply domain_lemma2b.
                                       ((alpha \cdot alpha \#) \cdot beta)).
assert ((domain alpha • beta)
apply comp\_inc\_compat\_ab\_a'b.
apply cap_l.
rewrite H in H0.
apply H0.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_{-}r.
apply (@inc_trans _ _ _ (domain beta • beta)).
rewrite domain_comp_alpha1.
apply inc\_reft.
apply (comp\_inc\_compat\_ab\_a'b\ H).
Qed.
  Lemma 304 (prod_to_cap) Let \alpha : A \rightarrow B and \beta : A \rightarrow C. Then,
                                      |\alpha \top \beta| = |\alpha| \sqcap |\beta|.
Lemma prod\_to\_cap \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 domain (Rel_prod alpha beta) = domain alpha
                                                        domain beta.
Proof.
replace (domain (Rel_prod alpha beta)) with (domain (Rel_prod alpha beta • snd_r B
C)).
rewrite prod_snd_domain1 comp_domain8.
apply dedekind_id3.
apply cap_r.
apply cap_r.
apply cap_r.
```

```
apply comp\_domain3. apply snd\_function. Qed.
```

Lemma 305 (prod_conjugate1) Let $\alpha: A \to B$ and $\beta: A \to C$ be functions, $p: B \times C \to B$ and $q: B \times C \to C$ be projections. Then,

$$(\alpha \top \beta) \cdot p = \alpha \wedge (\alpha \top \beta) \cdot q = \beta.$$

Lemma $prod_conjugate1$ { $A \ B \ C : eqType$ } { $alpha : Rel \ A \ B$ } {beta : $Rel \ A \ C$ }: $function_r \ alpha \rightarrow function_r \ \mathtt{beta} \rightarrow$ $Rel_prod\ alpha\ {\tt beta}\ {\tt \cdot}\ fst_r\ B\ C=alpha\ \land\ Rel_prod\ alpha\ {\tt beta}\ {\tt \cdot}\ snd_r\ B\ C={\tt beta}.$ Proof. move $\Rightarrow H H0$. split. rewrite prod_fst_domain1. elim $H0 \Rightarrow H1 \ H2$. apply $inc_{-}def1$ in H1. rewrite / domain. by [rewrite $cap_comm - H1 \ comp_id_l$]. rewrite prod_snd_domain1. elim $H \Rightarrow H1 H2$. apply inc_def1 in H1. rewrite / domain. by [rewrite $cap_comm - H1 \ comp_id_l$]. Qed.

Lemma 306 (prod_conjugate2) Let $\gamma: A \to B \times C$ be a function, $p: B \times C \to B$ and $q: B \times C \to C$ be projections. Then,

$$(\gamma \cdot p) \top (\gamma \cdot q) = \gamma.$$

Lemma $prod_conjugate2 \{A \ B \ C : eqType\} \{gamma : Rel \ A \ (prod_eqType \ B \ C)\}:$

 $\begin{array}{l} \textit{function_r gamma} \rightarrow \textit{Rel_prod (gamma • fst_r B C) (gamma • snd_r B C)} = \textit{gamma}. \\ \textbf{Proof.} \\ \textbf{move} \Rightarrow \textit{H.} \\ \textbf{rewrite } / \textit{Rel_prod.} \\ \textbf{rewrite } \textit{comp_assoc comp_assoc - (function_cap_distr_l \ H)}. \\ \textbf{by [rewrite } \textit{fst_snd_cap_id comp_id_r]}. \\ \textbf{Qed.} \end{array}$

projections. Then,

```
\frac{\alpha: A \to B}{u \sqsubseteq id_{A \times B}} \frac{\alpha = p^{\sharp} \cdot u \cdot q}{u = |p \cdot \alpha \sqcap q|}.
Lemma diagonal\_conjugate \{A B : eqType\} \{alpha : Rel A B\}:
 conjugate \ A \ B \ (prod\_eqType \ A \ B) \ (prod\_eqType \ A \ B)
 True_r (\mathbf{fun} \ u \Rightarrow u)
                             Id (prod\_eqType A B))
 (fun u \Rightarrow (fst_r A B \# \cdot u) \cdot snd_r A B)
                                                         snd_r A B).
 (\mathbf{fun} \ alpha \Rightarrow domain \ ((fst_r \ A \ B \cdot alpha))
Proof.
split.
move \Rightarrow alpha0 H.
split.
apply cap_r.
rewrite cap\_domain.
apply inc\_antisym.
apply (@inc\_trans\_\_\_((fst\_r\ A\ B\ \#\ \cdot\ ((fst\_r\ A\ B\ \bullet\ alpha0)\ \bullet\ snd\_r\ A\ B\ \#))\ \bullet\ snd\_r
A B)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
rewrite comp_assoc comp_assoc -(@comp_assoc _ _ _ _ (fst_r A B #)).
apply (@inc\_trans \_ \_ \_ ((fst\_r \ A \ B \ \# \cdot fst\_r \ A \ B) \cdot alpha0)).
apply comp\_inc\_compat\_ab\_a.
apply snd_-function.
apply comp\_inc\_compat\_ab\_b.
apply fst\_function.
apply @inc\_trans \_ \_ \_ (alpha0)
                                           ((fst\_r \ A \ B \ \# \ \cdot \ Id \ (prod\_eqType \ A \ B)) \ \cdot \ snd\_r \ A
B))).
{\tt rewrite}\ comp\_id\_r\ fst\_snd\_universal\ cap\_universal.
apply inc\_reft.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm inv_invol comp_assoc.
apply inc\_reft.
move \Rightarrow u H.
split.
by [].
```

Lemma 307 (diagonal_conjugate) Let $p: B \times C \rightarrow B$ and $q: B \times C \rightarrow C$ be


```
replace ((fst_r \ A \ B \cdot ((fst_r \ A \ B \# \cdot u) \cdot snd_r \ A \ B)) \quad snd_r \ A \ B) with (u \cdot snd_r \ A \ B)
A B).
apply domain\_inc\_id in H.
move: (@snd\_function \ A \ B) \Rightarrow H0.
elim H0 \Rightarrow H1 \ H2.
by [rewrite (comp\_domain3\ H1)\ H].
rewrite comp_assoc -comp_assoc.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((u \cdot snd\_r A B) \quad snd\_r A B)).
apply inc_-cap.
split.
apply inc_refl.
apply (comp\_inc\_compat\_ab\_b\ H).
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_b\_ab.
apply fst_-function.
apply (@inc_trans _ _ _ _ (dedekind2)).
apply comp\_inc\_compat\_ab\_b.
rewrite -fst\_snd\_cap\_id.
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_ab'.
apply inc_{-}inv.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
```

12.2.3 鋭敏性

この節の補題は以下の 1 つのみだが、証明が異様に長いため単独の節を設ける.

Lemma 308 (sharpness) In below figure,

$$\alpha \cdot \gamma^\sharp \sqcap \beta \cdot \delta^\sharp = (\alpha \cdot p^\sharp \sqcap \beta \cdot q^\sharp) \cdot (p \cdot \gamma^\sharp \sqcap q \cdot \delta^\sharp).$$



Lemma $sharpness \{ W \ X \ Y \ Z : eqType \}$ $\{ alpha : Rel \ X \ Y \} \{ beta : Rel \ X \ Z \} \{ gamma : Rel \ W \ Y \} \{ delta : Rel \ W \ Z \}:$

```
(alpha \cdot gamma \#)
                                 (beta • delta #) =
 ((alpha \cdot fst_r Y Z \#)
                                     (beta • snd_r Y Z \#))
  • ((fst_r \ Y \ Z \ \bullet \ gamma \ \#) \ (snd_r \ Y \ Z \ \bullet \ delta \ \#)).
Proof.
apply inc\_antisym.
move: (rationality \_ \_ alpha) \Rightarrow H.
move: (rationality \_ \_ beta) \Rightarrow H0.
move: (rationality \_ \_ (gamma \#)) \Rightarrow H1.
move: (rationality \_ \_ (delta \#)) \Rightarrow H2.
elim H \Rightarrow R.
elim \Rightarrow f\theta.
elim \Rightarrow g\theta H3.
elim H\theta \Rightarrow R\theta.
elim \Rightarrow f1.
elim \Rightarrow g1 H_4.
elim H1 \Rightarrow R1.
elim \Rightarrow h\theta.
elim \Rightarrow k0 H5.
elim H2 \Rightarrow R2.
elim \Rightarrow h1.
elim \Rightarrow k1 H6.
move: (rationality \_ \_ (g0 \cdot h0 \#)) \Rightarrow H7.
move: (rationality \_ \_ (g1 \cdot h1 \#)) \Rightarrow H8.
move: (rationality _ _ ((alpha • gamma #)
                                                             (beta \cdot delta \#)) \Rightarrow H9.
elim H7 \Rightarrow R3.
elim \Rightarrow s\theta.
elim \Rightarrow t0 \ H10.
elim H8 \Rightarrow R4.
elim \Rightarrow s1.
elim \Rightarrow t1 \ H11.
elim H9 \Rightarrow R5.
elim \Rightarrow x.
elim \Rightarrow z H12.
assert (alpha \cdot gamma \# = (f0 \# \cdot (s0 \# \cdot t0)) \cdot k0).
replace alpha with (f0 \# \cdot g0).
replace (gamma \#) with (h0 \# \cdot k0).
rewrite -comp\_assoc (@comp\_assoc\_\_\_ (f0 \#)).
apply f_{-}equal2.
apply f_equal.
apply H10.
by ||.
apply Logic.eq_sym.
```

```
apply H5.
apply Logic.eq_sym.
apply H3.
assert (beta • delta \# = (f1 \# \bullet (s1 \# \bullet t1)) \bullet k1).
replace beta with (f1 \# \bullet g1).
replace (delta \#) with (h1 \# \cdot k1).
rewrite -comp\_assoc (@comp\_assoc\_\_\_ (f1 \#)).
apply f_{-}equal2.
apply f_equal.
apply H11.
by [].
apply Logic.eq_sym.
apply H6.
apply Logic.eq_sym.
apply H_4.
assert (t\theta \cdot h\theta = s\theta \cdot g\theta).
apply function_inc.
apply function_comp.
apply H10.
apply H5.
apply function_comp.
apply H10.
apply H3.
apply (@inc\_trans \_ \_ \_ (s\theta \cdot ((s\theta \# \cdot t\theta) \cdot h\theta))).
rewrite comp_assoc -comp_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H10.
apply comp\_inc\_compat\_ab\_ab'.
replace (s\theta \# \cdot t\theta) with (g\theta \cdot h\theta \#).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H5.
apply H10.
assert (t1 \cdot h1 = s1 \cdot g1).
apply function_inc.
apply function_comp.
apply H11.
apply H6.
apply function_comp.
apply H11.
apply H_4.
apply (@inc\_trans \_ \_ \_ (s1 \cdot ((s1 \# \cdot t1) \cdot h1))).
```

```
rewrite comp_assoc -comp_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H11.
apply comp\_inc\_compat\_ab\_ab'.
replace (s1 \# \cdot t1) with (g1 \cdot h1 \#).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply H6.
apply H11.
remember ((x \cdot (s0 \cdot f0) \#) (z \cdot (t0 \cdot k0) \#)) as m0.
remember ((x \cdot (s1 \cdot f1) \#) (z \cdot (t1 \cdot k1) \#)) as m1.
assert (total_r \ m0).
rewrite Hegm0.
apply domain_corollary1.
apply H12.
apply H12.
replace (x \# \cdot z) with ((alpha \cdot gamma \#) (beta \cdot delta \#)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_inv H13 -comp_assoc comp_assoc.
apply inc\_reft.
apply H12.
assert (total_r \ m1).
rewrite Heqm1.
apply domain_corollary1.
apply H12.
apply H12.
replace (x \# \cdot z) with ((alpha \cdot gamma \#) \pmod{beta} \cdot delta \#)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
rewrite comp_inv H14 -comp_assoc comp_assoc.
apply inc\_reft.
apply H12.
remember (m0 \cdot (s0 \cdot g0)) as n0.
remember (m1 \cdot (s1 \cdot g1)) as n1.
assert (total_r \ n\theta).
rewrite Hegn\theta.
apply (total_comp H17).
apply total_comp.
apply H10.
apply H3.
assert (total_r \ n1).
rewrite Hegn1.
apply (total_comp H18).
```

```
apply total\_comp.
apply H11.
apply H_4.
assert (total_r ((n0 \cdot fst_r Y Z \#) (n1 \cdot snd_r Y Z \#))).
apply (domain_corollary1 H19 H20).
rewrite fst\_snd\_universal.
apply inc\_alpha\_universal.
assert ((x \# \cdot n\theta))
                         alpha).
replace alpha with (f0 \# \cdot g0).
rewrite Hegn0\ Hegm0.
apply (@inc\_trans \_ \_ \_ (((x \# \cdot x) \cdot f0 \#) \cdot ((s0 \# \cdot s0) \cdot g0))).
rewrite comp_assoc comp_assoc.
apply comp_inc_compat_ab_ab'.
rewrite -comp_assoc -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply comp_inc_compat_ab_a'b.
rewrite comp_assoc -comp_inv.
apply cap_{-}l.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_b.
apply H12.
apply comp\_inc\_compat\_ab\_b.
apply H10.
apply Logic.eq_sym.
apply H3.
assert ((x \# \cdot n1)  beta).
replace beta with (f1 \# \bullet g1).
rewrite Heqn1 Heqm1.
apply (@inc\_trans \_ \_ \_ (((x \# \cdot x) \cdot f1 \#) \cdot ((s1 \# \cdot s1) \cdot g1))).
rewrite comp\_assoc comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp_assoc -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc -comp_inv.
apply cap_{-}l.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_b.
apply H12.
apply comp\_inc\_compat\_ab\_b.
apply H11.
apply Logic.eq_sym.
```

```
apply H_4.
assert ((n0 \# \cdot z)
                      qamma \#).
replace (gamma \#) with (h0 \# \cdot k0).
rewrite Heqn0 Heqm0 -H15 comp_inv comp_inv inv_cap_distr.
apply (@inc\_trans\_\_\_((h0 \# \cdot (t0 \# \cdot t0)) \cdot (k0 \cdot (z \# \cdot z)))).
rewrite -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp_assoc (@comp_inv _ _ z) inv_invol.
apply cap_{-}r.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a.
apply H10.
apply comp\_inc\_compat\_ab\_a.
apply H12.
apply Logic.eq_sym.
apply H5.
assert ((n1 \# \cdot z)  delta \#).
replace (delta \#) with (h1 \# \cdot k1).
rewrite Heqn1 Heqm1 -H16 comp_inv comp_inv inv_cap_distr.
apply (@inc\_trans \_ \_ \_ ((h1 \# \cdot (t1 \# \cdot t1)) \cdot (k1 \cdot (z \# \cdot z)))).
rewrite -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp_assoc (@comp_inv _ _ z) inv_invol.
apply cap_r.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a.
apply H11.
apply comp\_inc\_compat\_ab\_a.
apply H12.
apply Logic.eq_sym.
apply H6.
replace ((alpha \cdot gamma \#) (beta \cdot delta \#)) with (x \# \cdot z).
apply (@inc\_trans \_ \_ \_ ((x \# \cdot (((n0 \cdot fst\_r Y Z \#) (n1 \cdot snd\_r Y Z \#)) \cdot (((n0 \cdot fst\_r Y Z \#)))))
• fst_r \ Y \ Z \ \#) (n1 \cdot snd_r \ Y \ Z \ \#))) \ \#)) \cdot z)).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp_inc_compat_a_ab H21).
```

```
rewrite -comp_assoc comp_assoc.
apply comp\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_ab\_a'b H22).
rewrite - comp_assoc.
apply (comp\_inc\_compat\_ab\_a'b H23).
rewrite inv_cap_distr comp_inv comp_inv inv_invol inv_invol.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat.
rewrite comp_{-}assoc.
apply (comp_inc_compat_ab_ab' H24).
rewrite comp_-assoc.
apply (comp_inc_compat_ab_ab' H25).
apply Logic.eq_sym.
apply H12.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite -comp_assoc (@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a.
apply fst\_function.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
rewrite -comp_assoc (@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a.
apply snd_-function.
Qed.
```

12.2.4 分配法則

```
Lemma 309 (prod_cap_distr_l) Let \alpha: A \to B and \beta, \beta': A \to C. Then, \alpha \top (\beta \sqcap \beta') = (\alpha \top \beta) \sqcap (\alpha \top \beta').
```

```
Lemma prod\_cap\_distr\_l {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta beta' : Rel \ A \ C}: Rel\_prod \ alpha (beta beta') = Rel\_prod \ alpha beta Rel\_prod \ alpha beta'. Proof.
```

rewrite $/Rel_prod$.
rewrite $-cap_assoc$ (@ cap_comm $_$ $_$ $_$ (alpha • fst_r B C #)) $-cap_assoc$ cap_idem cap_assoc .
apply f_equal.
apply $function_cap_distr_r$.
apply $snd_function$.
Qed.

Lemma 310 (prod_cap_distr_r) Let $\alpha, \alpha' : A \rightarrow B$ and $\beta : A \rightarrow C$. Then,

$$(\alpha \sqcap \alpha') \top \beta = (\alpha \top \beta) \sqcap (\alpha' \top \beta).$$

Lemma $prod_cap_distr_r$ { $A \ B \ C : eqType$ } { $alpha \ alpha' : Rel \ A \ B$ } {beta : $Rel \ A \ C$ }: $Rel_prod \ (alpha \ alpha')$ beta = $Rel_prod \ alpha$ beta $Rel_prod \ alpha'$ beta. Proof.

rewrite $/Rel_prod$.

rewrite cap_assoc (@ $cap_comm__$ (beta • snd_r B C #)) cap_assoc cap_idem - cap_assoc . apply (@f_equal $__$ (fun $x \Rightarrow$ @ $cap___x$ (beta • snd_r B C #))). apply $function_cap_distr_r$. apply $fst_function$.

appiy jst_junction

Qed.

Lemma 311 (prod_cup_distr_l) Let $\alpha : A \rightarrow B$ and $\beta, \beta' : A \rightarrow C$. Then,

$$\alpha \top (\beta \sqcup \beta') = (\alpha \top \beta) \sqcup (\alpha \top \beta').$$

Lemma $prod_cup_distr_l$ {A B C : eqType} {alpha : Rel A B} {beta beta' : Rel A C}: Rel_prod alpha (beta beta') = Rel_prod alpha beta Rel_prod alpha beta'.

Proof.

by [rewrite $-cap_cup_distr_l$ $-comp_cup_distr_r$]. Qed.

Lemma 312 (prod_cup_distr_r) Let $\alpha, \alpha' : A \to B$ and $\beta : A \to C$. Then,

$$(\alpha \sqcup \alpha') \top \beta = (\alpha \top \beta) \sqcup (\alpha' \top \beta).$$

Lemma $prod_cup_distr_r$ { $A \ B \ C : eqType$ } { $alpha \ alpha' : Rel \ A \ B$ } {beta : $Rel \ A \ C$ }: $Rel_prod \ (alpha \ alpha')$ beta = $Rel_prod \ alpha$ beta $Rel_prod \ alpha'$ beta. Proof.

by [rewrite $-cap_cup_distr_r$ $-comp_cup_distr_r$]. Qed.

```
Lemma 313 (comp_prod_distr_l) Let \alpha : A \rightarrow B, \beta : B \rightarrow C and \gamma : B \rightarrow D.
   Then,
                                            \alpha \cdot (\beta \top \gamma) \sqsubseteq \alpha \cdot \beta \top \alpha \cdot \gamma.
Lemma comp\_prod\_distr\_l
 \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ B \ D\}:
 alpha • Rel_prod beta gamma
                                             Rel\_prod (alpha \cdot beta) (alpha \cdot gamma).
Proof.
rewrite /Rel_prod.
rewrite comp_assoc comp_assoc.
apply comp\_cap\_distr\_l.
Qed.
  Lemma 314 (function_prod_distr_l) Let \alpha : A \rightarrow B be a function, \beta : B \rightarrow C and
  \gamma: B \rightarrow D. Then,
                                            \alpha \cdot (\beta \top \gamma) = \alpha \cdot \beta \top \alpha \cdot \gamma.
Lemma function\_prod\_distr\_l
 \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ B \ D\}:
 function\_r \ alpha \rightarrow alpha \cdot Rel\_prod \ \mathsf{beta} \ gamma = Rel\_prod \ (alpha \cdot \mathsf{beta}) \ (alpha \cdot \mathsf{beta})
qamma).
Proof.
move \Rightarrow H.
rewrite /Rel_prod.
rewrite comp_assoc comp_assoc.
apply (function\_cap\_distr\_l\ H).
Qed.
  Lemma 315 (comp_prod_universal) Let \alpha : A \rightarrow B, \beta : B \rightarrow C and \gamma : D \rightarrow E.
   Then.
                                      \alpha \cdot (\beta \top \nabla_{BD} \cdot \gamma) = \alpha \cdot \beta \top \nabla_{AD} \cdot \gamma.
Lemma comp\_prod\_universal
 \{A \ B \ C \ D \ E : eqType\} \ \{alpha : Rel \ A \ B\} \ \{beta : Rel \ B \ C\} \ \{gamma : Rel \ D \ E\}:
 alpha \cdot Rel\_prod \ \mathsf{beta} \ ( B \ D \cdot gamma) = Rel\_prod \ (alpha \cdot \mathsf{beta}) \ (
                                                                                                   AD \cdot qamma).
Proof.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_prod\_distr\_l)).
apply prod_inc_compat_l.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
```

```
rewrite /Rel_prod.
rewrite comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp_inc_compat_ab_ab'.
apply cap\_inc\_compat\_l.
rewrite comp_assoc comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
  Lemma 316 (fst_cap_snd_distr) Let u, v : A \times B \rightarrow A \times B and u, v \sqsubseteq id_{A \times B},
  p: B \times C \to B and q: B \times C \to C be projections. Then,
                               p^{\sharp} \cdot (u \sqcap v) \cdot q = p^{\sharp} \cdot u \cdot q \sqcap p^{\sharp} \cdot v \cdot q.
Lemma fst\_cap\_snd\_distr
 \{A \ B : eqType\} \{u \ v : Rel \ (prod\_eqType \ A \ B) \ (prod\_eqType \ A \ B)\}:
       Id (prod\_eqType \ A \ B) \rightarrow v \qquad Id (prod\_eqType \ A \ B) \rightarrow
 fst_r A B \# \cdot (u \quad v) \cdot snd_r A B =
 ((fst_r A B \# \cdot u) \cdot snd_r A B) ((fst_r A B \# \cdot v) \cdot snd_r A B).
Proof.
move \Rightarrow H H0.
apply inc\_antisym.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (comp\_cap\_distr\_r)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_cap\_distr\_l.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite -(dedekind_id3 H H0) -(@comp_assoc _ _ _ u) (@comp_assoc _ _ _ (fst_r A
B \# \cdot u v.
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_b.
rewrite comp_inv comp_inv inv_invol -fst_snd_cap_id.
apply cap\_inc\_compat.
rewrite comp_assoc (dedekind_id1 H).
apply (comp\_inc\_compat\_ab\_b\ H).
rewrite -comp_assoc (dedekind_id1 H0).
apply (comp\_inc\_compat\_ab\_a\ H0).
Qed.
```

End main.

Chapter 13

Library Point_Axiom

```
Require Import Basic\_Notations\_Set.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Require Import Dedekind.

Require Import Logic.IndefiniteDescription.

Module main (def:Relation).

Import def.

Module Basic\_Lemmas := Basic\_Lemmas.main \ def.

Module Relation\_Properties := Relation\_Properties.main \ def.

Module Functions\_Mappings := Functions\_Mappings.main \ def.

Module Dedekind := Dedekind.main \ def.

Import Basic\_Lemmas \ Relation\_Properties \ Functions\_Mappings \ Dedekind.
```

13.1 I-点

13.1.1 I-点の定義

Dedekind 圏における域 X の I-点 x とは, 関数 $x:I \to X$ のことであり, 記号 $x \in X$ によって表される. また関係 $\rho:I \to X$ と I-点 $x:I \to X$ に対して, 記号 $x \in \rho$ で $x \sqsubseteq \rho$ を表すものとする.

ちなみに I-点の定義 $x \in X$ は $x \in \nabla_{IX}$ と言い換えることも可能である.

```
Definition point\_inc\ \{X: eqType\}\ (x\ rho: Rel\ i\ X):=function\_r\ x \wedge x rho.
Definition point\ \{X: eqType\}\ (x: Rel\ i\ X):=point\_inc\ x\ (\ i\ X).
```

13.1.2 I-点の性質

```
Lemma 317 (point_property1) Let x, y \in X. Then,
                                         x = y \Leftrightarrow x \cdot y^{\sharp} = id_I.
Lemma point\_property1 \{X : eqType\} \{x \ y : Rel \ i \ X\}:
 point x \to point \ y \to (x = y \leftrightarrow x \cdot y \# = Id \ i).
Proof.
move \Rightarrow H H0.
split; move \Rightarrow H1.
apply inc\_antisym.
rewrite unit\_identity\_is\_universal.
apply inc\_alpha\_universal.
rewrite H1.
apply H0.
apply Logic.eq_sym.
apply function_inc.
apply H0.
apply H.
rewrite -(@comp\_id\_l\_\_y) -H1 comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
Qed.
  Lemma 318 (point_property2a, point_property2b) Let \rho: I \to X be a total re-
  lation. Then,
                                        \rho \cdot \rho^{\sharp} = \rho \cdot \nabla_{XI} = id_I.
Lemma point\_property2a \{X : eqType\} \{rho : Rel \ i \ X\}:
 total\_r \ rho \rightarrow rho \cdot rho \# = Id \ i.
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite unit\_identity\_is\_universal.
apply inc\_alpha\_universal.
apply H.
Qed.
Lemma point\_property2b \{X : eqType\} \{rho : Rel \ i \ X\}:
 total\_r \ rho \rightarrow rho \ \cdot \ rho \ \# = rho \ \cdot \ X \ i.
Proof.
move \Rightarrow H.
```

```
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
rewrite (point_property2a H) unit_identity_is_universal.
apply inc_alpha_universal.
Qed.
  Lemma 319 (point_property3) Let \rho: I \to X. Then,
                                    \exists x \in \rho \Rightarrow "\rho \text{ is total"} \land \rho \neq \phi_{IX}.
Lemma point\_property3 \{X : eqType\} \{rho : Rel \ i \ X\}:
 (\exists x : Rel \ i \ X, point\_inc \ x \ rho) \rightarrow total\_r \ rho \land rho \neq i
                                                                         i X.
Proof.
elim \Rightarrow x H.
assert (total_r rho).
elim H \Rightarrow H0 \ H1.
elim H0 \Rightarrow H2 \ H3.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply comp\_inc\_compat.
apply H1.
apply (@inc_inv_{-1} - H1).
split.
apply H0.
move \Rightarrow H1.
rewrite /total_r in H0.
rewrite H1 comp_-empty_-l in H0.
apply unit_identity_not_empty.
apply inc\_antisym.
apply H0.
apply inc\_empty\_alpha.
Qed.
  Lemma 320 (point_property4)
                               \exists x \in X \Rightarrow "\nabla_{IX} \text{ is total"} \land \nabla_{IX} \neq \phi_{IX}.
Lemma point\_property4 \{X : eqType\}:
 (\exists x : Rel \ i \ X, \ point \ x) \rightarrow total_r \ (i \ X) \land (i \ X) \neq
                                                                           i X.
Proof.
move \Rightarrow H.
apply (@point\_property3 \_ ( i X) H).
Qed.
```

この "点公理" を使えば、I-点に関する様々な定理や補題が導出できる.

13.2 I-点に関する諸公理

13.2.1 点公理

elim $H0 \Rightarrow x H1$.

```
Lemma 321 (point_axiom) Let \rho: I \to X. Then,
                                                 \rho = \sqcup_{x \in \rho} x.
Lemma lemma\_for\_PA \{X : eqType\} \{rho : Rel \ i \ X\}:
                i(X) \rightarrow False) \rightarrow False) \rightarrow rho = i(X).
Proof.
move \Rightarrow H.
case (@unit_empty_or_universal (rho • rho \#)) \Rightarrow H0.
apply inc\_antisym.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
rewrite H0 comp\_empty\_l.
apply inc\_reft.
apply inc\_empty\_alpha.
apply False_ind.
apply H.
move \Rightarrow H1.
rewrite H1 comp_-empty_-l in H0.
apply (unit_empty_not_universal H0).
Qed.
Lemma point\_axiom \{X : eqType\} \{rho : Rel \ i \ X\}:
            _{-}\{fun x: Rel \ i \ X \Rightarrow point\_inc \ x \ rho\} \ id.
 rho =
Proof.
apply inc\_antisym.
apply bool_lemma2.
assert ((\exists x : Rel \ i \ X, point\_inc \ x \ (( \_{fun} \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho) \ id)
( _{\text{-}}\{\text{fun } x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho\} \ id) \hat{\ })) \rightarrow False).
move \Rightarrow H.
move: (point\_property3 \ H) \Rightarrow H0.
apply H0.
apply cap\_complement\_empty.
assert ((\exists x : Rel \ i \ X, point\_inc \ x \ (rho) \ ( \_\{fun \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho\} \ id)
\hat{})) \rightarrow False).
move \Rightarrow H0.
apply H.
```

```
\exists x.
split.
apply H1.
apply inc\_cap.
split.
assert (point\_inc \ x \ rho).
split.
apply H1.
elim H1 \Rightarrow H2 \ H3.
apply inc\_cap in H3.
apply H3.
clear H1.
move: x H2.
apply inc\_cupP.
apply inc_refl.
elim H1 \Rightarrow H2 \ H3.
apply inc\_cap in H3.
apply H3.
apply lemma\_for\_PA.
move \Rightarrow H1.
apply H0.
apply axiom\_of\_choice.
rewrite /total_r.
remember (rho
                   ( _{\text{fun }} x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho \} \ id) ^) \text{ as } rho'.
case (@unit\_empty\_or\_universal\ (rho' \cdot rho' \#)) \Rightarrow H2.
apply False\_ind.
apply H1.
apply inc\_antisym.
apply (@inc_trans _ _ _ _ (relation_rel_inv_rel)).
rewrite H2 comp\_empty\_l.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite H2.
apply inc_alpha_universal.
apply inc\_cupP.
move \Rightarrow beta H.
apply H.
Qed.
  Lemma 322 (PA_corollary1)
```

 $\nabla_{IX} = \sqcup_{x \in X} x.$

```
Lemma PA\_corollary1 \{X: eqType\}: i\ X = \_\{point\}\ id. Proof. apply point\_axiom. Qed.
```

Lemma 323 (PA_corollary2)

```
id_X = \sqcup_{x \in X} x^{\sharp} \cdot x.
```

```
Lemma PA\_corollary2 \{X : eqType\}:
            \{point\}\ (\mathbf{fun}\ x: Rel\ i\ X \Rightarrow x\ \#\ \cdot\ x).
 Id X =
Proof.
rewrite -(@cap_universal _ _ (Id X)) -lemma_for_tarski2 PA_corollary1.
rewrite comp\_cupP\_distr\_l cap\_cupP\_distr\_l.
apply cupP_{-}eq.
move \Rightarrow alpha H.
apply inc\_antisym.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite comp_id_l cap_comm cap_universal.
apply inc_refl.
apply inc\_cap.
split.
apply H.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
```

Lemma 324 (PA_corollary3) Let $\alpha, \beta: X \to Y$. Then,

$$(\forall x \in X, x \cdot \alpha = x \cdot \beta) \Rightarrow \alpha = \beta.$$

```
Lemma PA\_corollary3 \{X \ Y : eqType\} \{alpha \ beta : Rel \ X \ Y\}: (\forall \ x : Rel \ i \ X, \ point \ x \to x \ \bullet \ alpha = x \ \bullet \ beta) \to alpha = beta. Proof. move \Rightarrow H. rewrite -(@comp\_id\_l\_\_\_alpha) - (@comp\_id\_l\_\_\_beta) \ PA\_corollary2. rewrite comp\_cupP\_distr\_r \ comp\_cupP\_distr\_r. apply cupP\_eq. move \Rightarrow gamma \ H0. by [rewrite comp\_assoc \ comp\_assoc \ (H \ gamma \ H0)]. Qed.
```

```
Lemma 325 (PA_corollary4) Let \alpha: X \to Y. Then,
                                 "\alpha is total" \Leftrightarrow \forall x \in X, "x \cdot \alpha is total".
Lemma PA\_corollary4 \{X \ Y : eqType\} \{alpha : Rel \ X \ Y\}:
 total\_r \ alpha \leftrightarrow \forall \ x : Rel \ i \ X, \ point \ x \rightarrow total\_r \ (x \cdot alpha).
Proof.
split; move \Rightarrow H.
move \Rightarrow x H0.
apply total\_comp.
apply H0.
apply H.
rewrite /total_r.
rewrite PA\_corollary2.
apply inc\_cupP.
move \Rightarrow x H0.
move: (H \times H0) \Rightarrow H1.
apply (@inc\_trans \_ \_ \_ ((x \# \cdot ((x \cdot alpha) \cdot (x \cdot alpha) \#)) \cdot x)).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp_inc_compat_a_ab H1).
rewrite comp_inv -comp_assoc -comp_assoc -comp_assoc.
rewrite comp\_assoc (@comp\_assoc\_\_\_\_ (x \# \cdot x)).
apply (@inc\_trans \_ \_ \_ ((x \# \cdot x) \cdot (alpha \cdot alpha \#))).
apply comp\_inc\_compat\_ab\_a.
apply H0.
apply comp\_inc\_compat\_ab\_b.
apply H\theta.
Qed.
  Lemma 326 (PA_corollary5) Let \alpha: X \to Y. Then,
                           "\alpha is univalent" \Leftrightarrow \forall x \in X, "x \cdot \alpha is univalent".
Lemma PA\_corollary5 \{X \ Y : eqType\} \{alpha : Rel \ X \ Y\}:
 univalent_r \ alpha \leftrightarrow \forall \ x : Rel \ i \ X, \ point \ x \rightarrow univalent_r \ (x \cdot alpha).
Proof.
split; move \Rightarrow H.
move \Rightarrow x H0.
apply univalent_comp.
apply H0.
apply H.
rewrite /univalent_r.
rewrite -(@comp\_id\_r\_\_(alpha \#)) PA_corollary2.
```

```
rewrite comp\_cupP\_distr\_l\ comp\_cupP\_distr\_r. apply inc\_cupP. move \Rightarrow x\ H0. move : (H\ x\ H0) \Rightarrow H1. rewrite -comp\_assoc\ -comp\_inv\ comp\_assoc. apply H1. Qed.
```

13.2.2 全域性公理

```
Lemma 327 (total_axiom) Let \rho: I \to X. Then, \rho \neq \phi_{IX} \Rightarrow id_I = \rho \cdot \rho^{\sharp}.
```

```
Lemma total\_axiom \{X: eqType\} \{rho: Rel \ i \ X\}: rho \neq i \ X \rightarrow Id \ i = rho \cdot rho \#.

Proof.

move \Rightarrow H.

case (@unit\_empty\_or\_universal (rho \cdot rho \#)) \Rightarrow H0.

apply False\_ind.

apply H.

apply inc\_antisym.

apply (@inc\_trans _ _ _ _ (relation\_rel\_inv\_rel)).

rewrite H0 comp\_empty\_l.

apply inc\_refl.

apply inc\_empty\_alpha.

by [rewrite H0 unit\_identity\_is\_universal].

Qed.
```

```
Lemma 328 (Tot_corollary1) Let \rho: I \to X and x \in X. Then,
```

$$\rho \sqsubseteq x \Rightarrow \rho = \phi_{IX} \lor \rho = x.$$

```
Lemma Tot\_corollary1 \{X: eqType\} \{rho\ x: Rel\ i\ X\}: point\ x \to rho = i\ X \lor rho = x. Proof.

move \Rightarrow H\ H0.

case (@unit\_empty\_or\_universal (rho\ \cdot\ rho\ \#)) \Rightarrow H1.

left.

apply inc\_antisym.

apply (@inc\_trans _ _ _ _ (relation\_rel\_inv\_rel)).

rewrite H1\ comp\_empty\_l.
```

```
apply inc\_reft.
apply inc\_empty\_alpha.
right.
apply inc\_antisym.
apply H0.
rewrite -(@comp\_id\_l\_\_x) unit_identity_is_universal -H1 comp_assoc.
apply (@inc\_trans \_ \_ \_ (rho \cdot (x \# \cdot x))).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc_inv_- - H0).
apply comp\_inc\_compat\_ab\_a.
apply H.
Qed.
  Lemma 329 (Tot_corollary2) Let x, y \in X. Then,
                                           x \neq y \Leftrightarrow x \cdot y^{\sharp} = \phi_{II}.
Lemma Tot\_corollary2 \{X : eqType\} \{x \ y : Rel \ i \ X\}:
 point \ x \rightarrow point \ y \rightarrow (x \neq y \leftrightarrow x \ \bullet \ y \ \# =
Proof.
move \Rightarrow H H0.
\texttt{assert} \; (x = y \leftrightarrow x \; \cdot \; y \; \# \neq x)
rewrite (point_property1 H H0).
split; move \Rightarrow H1.
rewrite H1.
apply unit_identity_not_empty.
case (@unit\_empty\_or\_universal\ (x \cdot y \#)) \Rightarrow H2.
apply False_ind.
apply (H1 H2).
by [rewrite H2 unit_identity_is_universal].
rewrite H1.
split; move \Rightarrow H2.
apply (lemma_for_PA H2).
move \Rightarrow H3.
apply (H3 H2).
Qed.
  Lemma 330 (Tot_corollary3) Let f: (I \rightarrow X) \rightarrow (I \rightarrow Y). Then,
                  (\forall x \in X, "f(x) \text{ is a function"}) \Rightarrow "\sqcup_{x \in X} x^{\sharp} \cdot f(x) \text{ is a function"}.
```

```
(\forall x : Rel \ i \ X, \ point \ x \rightarrow function_r \ (f \ x)) \rightarrow function_r \ (f \ x) \ (fun \ x : Rel \ i \ X)
\Rightarrow x \# \cdot f(x)).
Proof.
\mathtt{move} \Rightarrow H.
assert (\forall x : Rel \ i \ X, \ point \ x \rightarrow x \bullet ( \ _{\{point\}} \ (\mathbf{fun} \ x\theta : Rel \ i \ X \Rightarrow x\theta \ \# \bullet f \ x\theta)) =
f(x).
move \Rightarrow x H0.
assert (x \cdot x \# = Id \ i).
apply inc\_antisym.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
apply H0.
rewrite -(@comp_id_l - (f x)) - H1.
apply inc\_antisym.
rewrite comp\_cupP\_distr\_l.
apply inc\_cupP.
move \Rightarrow y H2.
rewrite -comp\_assoc.
case (@unit_empty_or_universal (x \cdot y \#) \Rightarrow H3.
rewrite H3 comp\_empty\_l.
apply inc\_empty\_alpha.
rewrite -unit_identity_is_universal in H3.
apply (point_property1 H0 H2) in H3.
rewrite H3.
apply inc\_reft.
rewrite comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
clear H1.
move: x H0.
apply inc\_cupP.
apply inc\_reft.
split.
rewrite PA_corollary4.
move \Rightarrow x H1.
rewrite (H0 \times H1).
apply (H \times H1).
rewrite PA\_corollary5.
move \Rightarrow x H1.
rewrite (H0 \ x \ H1).
apply (H \times H1).
Qed.
```

13.2.3 その他の公理

```
\rho \neq \phi_{IX} \Rightarrow \exists x \dot{\in} \rho. Lemma nonempty_axiom \{X : eqType\} \{rho : Rel \ i \ X\}: rho \neq i \ X \rightarrow \exists \ x : Rel \ i \ X, \ point_inc \ x \ rho. Proof.

move : (@axiom\_of\_choice\_\_ rho) \Rightarrow H.
move \Rightarrow H0.
apply H.
rewrite /total\_r.
rewrite /total\_axiom\ H0).
apply inc\_refl.
Qed.
```

Lemma 331 (nonempty_axiom) Let $\rho: I \to X$. Then,

Lemma 332 (axiom_of_subobjects2) Let $\rho: I \to X$. Then,

```
\exists S, \exists j: S \to X, \rho = \nabla_{IS} \cdot j \wedge j \cdot j^{\sharp} = id_{S}.
```

```
Lemma axiom\_of\_subobjects2 \{X : eqType\} \{rho : Rel \ i \ X\}:
 \exists (S : eqType)(j : Rel S X), rho = i S \cdot j \wedge j \cdot j \# = Id S.
Proof.
elim (@rationality \_ \_ rho) \Rightarrow R.
elim \Rightarrow f.
elim \Rightarrow g.
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2.
\exists R.
\exists q.
split.
rewrite H1.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans\_\_\_( i R \cdot (f \cdot f \#))).
apply comp\_inc\_compat\_a\_ab.
apply H.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
```

```
\verb"rewrite" unit\_identity\_is\_universal".
apply inc\_alpha\_universal.
rewrite -H2 cap_comm inc_def1.
assert ((f \# \cdot g)
                       rho).
rewrite H1.
apply inc_refl.
apply (function\_move1 \ H) in H3.
apply (@inc\_trans \_ \_ \_ ((f \cdot rho) \cdot (f \cdot rho) \#)).
apply comp_inc_compat.
apply H3.
apply (@inc_inv_{-1} - H3).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ rho).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
```

13.3 その他の補題

```
Lemma 333 (point_atomic) Let x \in X, then x is atomic.
```

```
Lemma point\_atomic\ \{X: eqType\}\ \{x: Rel\ i\ X\}:\ point\ x \to atomic\ x. Proof.

move \Rightarrow H.

split.

move: (@point\_property3\ X\ x) \Rightarrow H0.

apply H0.

\exists\ x.

split.

apply H.

apply H.

apply inc\_refl.

move \Rightarrow beta.

apply (Tot\_corollary1\ H).

Qed.
```

Lemma 334 (point_atomic2) Let $x \in X$ and $y \in Y$, then $x^{\sharp} \cdot y$ is atomic.

```
Lemma point\_atomic2 {X \ Y : eqType} {x : Rel \ i \ X} {y : Rel \ i \ Y}: point \ x \rightarrow point \ y \rightarrow atomic \ (x \ \# \ ^{\bullet} \ y).
Proof.
```

```
move \Rightarrow H H\overline{\theta}.
split.
move \Rightarrow H1.
assert (Id \ i = (x \cdot x \#) \cdot (y \cdot y \#)).
apply inc\_antisym.
rewrite -(@comp_{-}id_{-}l_{-} - (Id i)).
apply comp_inc_compat.
apply H.
apply H0.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
rewrite comp\_assoc -(@comp\_assoc _ _ _ _ (x \#)) in H2.
rewrite H1 comp_empty_l comp_empty_r in H2.
apply (unit_identity_not_empty H2).
move \Rightarrow beta H1.
case (@unit_empty_or_universal (( i X \cdot beta) · Y i)) \Rightarrow H2.
apply inc\_antisym.
replace (XY) with ((Xi)
                                       i i) \cdot
                                                    i Y).
rewrite -H2 -comp_assoc -comp_assoc unit_universal.
rewrite comp_assoc unit_universal.
apply (@inc\_trans\_\_\_(XX \cdot beta)).
apply comp\_inc\_compat\_b\_ab.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_a\_ab.
apply inc_alpha_universal.
by [rewrite comp\_empty\_r comp\_empty\_l].
apply inc\_empty\_alpha.
right.
apply inc\_antisym.
apply H1.
assert (beta \neq X Y).
move \Rightarrow H3.
rewrite H3 comp\_empty\_r comp\_empty\_l in H2.
apply (unit\_empty\_not\_universal\ H2).
apply @inc\_trans\_\_\_(x \# \cdot (x \cdot beta))).
apply comp\_inc\_compat\_ab\_ab'.
assert ((x \cdot beta) y).
apply (@inc\_trans \_ \_ \_ (x \cdot (x \# \cdot y))).
apply (comp\_inc\_compat\_ab\_ab' H1).
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
```

```
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
apply inc\_def1 in H1.
rewrite H1 in H3.
assert (x \# \cdot ((x \cdot beta) \quad y) \neq X Y).
move \Rightarrow H5.
apply H3.
apply inc\_antisym.
rewrite cap\_comm.
apply (@inc_trans _ _ _ _ (dedekind1)).
rewrite cap_comm inv_invol H5.
apply inc_refl.
apply inc\_empty\_alpha.
case (Tot\_corollary1\ H0\ H4) \Rightarrow H6.
rewrite H6 cap\_comm cap\_empty comp\_empty\_r in H5.
apply False\_ind.
by [apply H5].
rewrite H6.
apply inc_refl.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
apply H.
Qed.
End main.
```

Bibliography

[1] R. Affeldt and M. Hagiwara. Formalization of Shannon 's Theorems in SSReflect-Coq. In 3rd Conference on Interactive Theorem Proving, LNCS 7406, 233–249, 2012.