

# INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY

### LOGIC AND COMPUTATION PROJECT

# Coq Modules for Relational Calculus

(Ver.0.1)

Hisaharu TANAKA Saga University

Shuichi INOKUCHI Fukuoka Institute of Technology Toshiaki MATSUSHIMA Kyushu University

Yoshihiro MIZOGUCHI Kyushu University

October 20, 2015

Repository: svn://sakura.imi.kyushu-u.ac.jp/Coq/RelationalCalculus/branches/20150716/

## Contents

1	Libr	eary Basic_Notations	4
	1.1	このライブラリについて	4
2		eary Basic_Notations_Rel	5
	2.1	定義	5
	2.2	関数の定義	8
	2.3	関係の公理	8
		2.3.1 Dedekind <b>圏の公理</b>	8
		2.3.2 排中律	13
		2.3.3 単域	13
		2.3.4 弱選択公理	13
			14
			14
3	Libr		<b>1</b> 5
	3.1	定義	15
	3.2	関数の定義	16
	3.3	関係の公理	17
		3.3.1 Dedekind <b>圏の公理</b>	17
		3.3.2 排中律	25
		3.3.3 単域	26
			27
		• • • • • • •	27
			31
		SISTEMATICAL PROPERTY OF THE P	01
4	Libr	eary Basic_Lemmas	35
	4.1	束論に関する補題	35
		4.1.1 和関係, 共通関係	35
			44
			46
	4.2		48
	4.3	· ·	55
	4.4	Rool 代数に関する補題	59

5	<b>Libr</b> 5.1	ary Re	elation 算の基																	<b>62</b>
	5.2		异い巫 inc_con																	
	5.3		に関す																	
	5.4		関する																	80
	5.5		Tarski																	85
	0.0	T-%C	Idibiti			• •		•	 •	 •	•	 •	 •	 •	•	• •	•	•	 •	00
6	Libr	ary Fu																		90
	6.1	全域性	,一価性	ŧ, 写像	に関	する	補	題												90
	6.2	全射, 🖺	単射に関	する神	甫題															100
_	T	-																		
7		ary Do			· 88 <del>- }-</del>	フ <del>)</del> 날														109
	7.1	Dedeki																		
		Dedeki Dedeki																		
	7.3	Deaeki	na iorn	nuia C	一一一	关  行	† .	•	 •	 •	•	 •	 •	 •	•		•	•	 •	116
8	Libr	ary Ra	ational	itv																116
•	8.1		から導		系 .															116
9		ary Co																		118
	9.1		の定義																	
	9.2	共役の	例					•					 •					•		119
10	Libr	ary Do	amain																	128
10	10 1	定義域	Jiliaili の定義																	
		定義域																		
	10.2		基本的																	
			合成と																	131
			その他																	
			矩形関																	
		10.2.1	Λ <u>Ε/Ι</u> ΖΙ <del>Χ</del> Ι	101		• •		•	 •	 •	•	 •	 •	 •	•	• •	•	•	 •	110
11	Libr	ary Re	esidual																	<b>14</b> 4
	11.1	剰余合	成関係	の性質																144
		11.1.1	基本的	な性質																144
			単調性																	147
		11.1.3	剰余合	成と関	数 .															150
		Galois		_																153
		その他																		154
	11.4	順序の																		164
			max, s																	164
		11.4.2	左剰余	合成										 •						169
10	T:L.	C -	hnode-	2																171
14		ary Sc																		1/1

13 Library Sum_Product	179
13.1 関係の直和	 179
13.1.1 入射対,関係直和の定義	 179
13.1.2 関係直和の性質	 181
13.1.3 分配法則	 185
13.2 関係の直積	 186
13.2.1 射影対, 関係直積の定義	 186
13.2.2 関係直積の性質	 188
13.2.3 鋭敏性	 197
13.2.4 分配法則	 203
14 Library Point_Axiom	207
14.1 I-点	 207
14.1.1 I-点の定義	 207
14.1.2 I- <b>点の性質</b>	 207
14.2 I-点に関する諸公理	 210
14.2.1 点公理	 210
14.2.2 その他の公理	213

## Chapter 1

## Library Basic\_Notations

### 1.1 このライブラリについて

- このライブラリは河原康雄先生の"関係の理論 Dedekind 圏概説 -" をもとに制作されている.
- 現状サポートしているのは、
  - 1.4 節大半, 1.5 1.6 節全部
  - 2.1 2.3 節全部, 2.4 2.5 節大半, 2.6 節全部, 2.7 節大半, 2.8 節有理性
  - 4.2 4.3 節全部, 4.4 4.5 節大半, 4.6 節命題 4.6.1, 4.7 節大半, 4.9 節全部
  - 4.8 節は部分域公理を用いるので、そちらが終わり次第

といったところである.

● 関係論で話を進めたい場合は、下の行に Require Export Basic\_Notations\_Rel. を、集合論で話を進めたい場合は、Require Export Basic\_Notations\_Set. を記述する.

Require Export Basic\_Notations\_Set.

なお、証明の書き方が悪いと、まれに"関係論では証明が通ったのに、集合論では通らない"といったことも起こるようなので、ある程度注意しておく必要がある.

## Chapter 2

## Library Basic\_Notations\_Rel

Require Export ssreflect eqtype bigop.
Require Export Logic. Classical Facts.

Axiom prop\_extensionality\_ok: prop\_extensionality.

### 2.1 定義

- A, B を eqType として, A から B への関係の型を (Rel A B) と書き,  $A \to B \to Prop$  として定義する. 本文中では型 (Rel A B) を  $A \to B$  と書く.
- 関係  $\alpha:A\to B$  の逆関係  $\alpha^{\sharp}:B\to A$  は (inverse  $\alpha$ ) で, Coq では  $(\alpha$  #) と記述する.
- 2 つの関係  $\alpha:A \to B$ ,  $\beta:B \to C$  の合成関係  $\alpha \cdot \beta (\text{or } \alpha\beta):A \to C$  は  $(\text{composite } \alpha \beta)$  で,  $(\alpha \cdot \beta)$  と記述する.
- 剰余合成関係  $\alpha \triangleright \beta : A \rightarrow C$  は (residual  $\alpha \beta$ ) で,  $(\alpha \beta)$  と記述する.
- 恒等関係  $\mathrm{id}_A:A\to A$  は (identity A) で, (Id A) と記述する.
- 空関係  $\phi_{AB}: A \rightarrow B$  は (empty AB) で, (AB) と記述する.
- 全関係  $\nabla_{AB}: A \rightarrow B$  は (universal AB) で, (AB) と記述する.
- 2 つの関係  $\alpha:A\to B$ ,  $\beta:A\to B$  の和関係  $\alpha\sqcup\beta:A\to B$  は  $(\operatorname{cup}\ \alpha\ \beta)$  で,  $(\alpha\qquad\beta)$  と記述する.
- 共通関係  $\alpha \sqcap \beta : A \to B$  は (cap  $\alpha \beta$ ) で,  $(\alpha \quad \beta)$  と記述する.
- 相対擬補関係  $\alpha \Rightarrow \beta : A \rightarrow B$  は (rpc  $\alpha \beta$ ) で,  $(\alpha >> \beta)$  と記述する.
- 関係  $\alpha:A\to B$  の補関係  $\alpha^-:A\to B$  は (complement  $\alpha$ ) で, Coq では  $(\alpha ^\circ)$  と記述する.

	数式	Coq	Notation
逆関係	$lpha^{\sharp}$	(inverse $\alpha$ )	(α #)
合成関係	$\alpha \cdot \beta$	(composite $\alpha \beta$ )	$(\alpha \cdot \beta)$
剰余合成関係	$\alpha \rhd \beta$	$( exttt{residual} \ lpha \ eta)$	$(\alpha \qquad \beta)$
恒等関係	$\mathrm{id}_A$	(identity A)	$(\operatorname{Id} A)$
空関係	$\phi_{AB}$	$(\texttt{empty } A \ B)$	(AB)
全関係	$ abla_{AB}$	(universal AB)	(AB)
和関係	$\alpha \sqcup \beta$	$(\operatorname{cup} \ \alpha \ \beta)$	$(\alpha \qquad \beta)$
共通関係	$\alpha \sqcap \beta$	$(\operatorname{cap}\ \alpha\ eta)$	$(\alpha \qquad \beta)$
相対擬補関係	$\alpha \Rightarrow \beta$	$(\operatorname{\mathtt{rpc}}\ \alpha\ \beta)$	$(\alpha \gg \beta)$
補関係	$\alpha^{-}$	$(\texttt{complement} \ \alpha)$	(α <b>^</b> )
差関係	$\alpha - \beta$	(difference $\alpha \beta$ )	$(\alpha \beta)$
添字付和関係	$\sqcup_{P(\alpha)} f(\alpha)$	$(\operatorname{cupP} P f)$	$( -\{P\} f)$
添字付共通関係	$\sqcap_{P(\alpha)} f(\alpha)$	$(\mathtt{capP}\ P\ f)$	$( -\{P\} f)$

Table 2.1: 関係の表記について

- 2 つの関係  $\alpha:A\to B$ ,  $\beta:A\to B$  の差関係  $\alpha-\beta:A\to B$  は (difference  $\alpha$   $\beta$ ) で,  $(\alpha$  --  $\beta)$  と記述する.
- (cupP) と (capP) は添字付の和関係と共通関係であり、述語 P に対し、 $\{f(\alpha) \mid P(\alpha)\}$  の和関係、共通関係を表す.
- また, 1 点集合 I = {\*} は i と表記する.
- なお, 通常の共通関係, 和関係も添字付のもので表現することができるため, ここでは それを用いて表記する.
- 後で述べるように、剰余合成  $\alpha \triangleright \beta$  も  $(\alpha \cdot \beta^-)^-$  のように表現することは可能だが、"剰余合成が存在すれば、それは  $(\alpha \cdot \beta^-)^-$  に等しい" というレベルのものであるため、剰余合成に関する公理はやはり必要となる.

#### 表 2.1 に関係の表記についてまとめる.

```
Definition Rel(A B : eqType) := A \rightarrow B \rightarrow Prop.
```

Parameter inverse:  $(\forall A B : eqType, Rel A B \rightarrow Rel B A)$ .

Notation "a #" :=  $(inverse \_ \_ a)$  (at level 20).

Parameter composite:  $(\forall A B C : eqType, Rel A B \rightarrow Rel B C \rightarrow Rel A C)$ .

Notation "a'  $\cdot$  'b" :=  $(composite \_ \_ \_ a \ b)$  (at level 50).

Parameter residual:  $(\forall A B C : eqType, Rel A B \rightarrow Rel B C \rightarrow Rel A C)$ .

Notation "a', b" :=  $(residual \_ \_ \_ a \ b)$  (at level 50).

Parameter identity:  $(\forall A : eqType, Rel A A)$ .

Notation "'Id'" := identity.

```
Parameter empty: (\forall A B : egType, Rel A B).
Notation "' := empty.
Parameter universal: (\forall A B : eqType, Rel A B).
Notation "' := universal.
Parameter include: (\forall A B : eqType, Rel A B \rightarrow Rel A B \rightarrow Prop).
Notation "a', b" := (include \_ \_ a \ b) (at level 50).
Parameter cupP: (\forall A \ B \ C \ D : eqType, (Rel \ C \ D \rightarrow Prop) \rightarrow (Rel \ C \ D \rightarrow Rel \ A \ B) \rightarrow
Rel\ A\ B).
                \{ p' \}'  f'' := (cupP_{---} p f)  (at level 50).
Notation "'
Parameter capP: (\forall A B C D : eqType, (Rel C D \rightarrow Prop) \rightarrow (Rel C D \rightarrow Rel A B) \rightarrow
Rel\ A\ B).
                \{', p'\}', f'' := (capP_{-} - p_f) \text{ (at level 50)}.
Notation "'
Definition cup \{A B : eqType\} (alpha beta : Rel A B)
       _{\text{-}}\{\text{fun } gamma : Rel \ A \ B \Rightarrow gamma = alpha \lor gamma = \text{beta}\} \ id.
Notation "a', b" := (cup \ a \ b) (at level 50).
Definition cap \{A B : eqType\} (alpha beta : Rel A B)
       _{-}\{fun gamma : Rel \ A \ B \Rightarrow gamma = alpha \lor gamma = beta \} id.
Notation "a', b" := (cap \ a \ b) (at level 50).
Parameter rpc: (\forall A B : eqType, Rel A B \rightarrow Rel A B \rightarrow Rel A B).
Notation "a'»' b" := (rpc - a b) (at level 50).
Definition complement \{A \ B : eqType\} (alpha : Rel \ A \ B) := alpha \gg
                                                                                     A B.
Notation "a '^' := (complement \ a) (at level 20).
Definition difference \{A \ B : eqType\}\ (alpha \ \mathsf{beta} : Rel \ A \ B) := alpha \ \mathsf{beta} \ \hat{}.
Notation "a - b" := (difference a b) (at level 50).
Notation "'i'" := unit\_eqType.
```

### 2.2 関数の定義

 $\alpha:A\to B$  に対し、全域性 total\_r、一価性 univalent\_r、関数 function\_r、全射 surjective\_r、単射 injective\_r、全単射 bijection\_r を以下のように定義する.

- total\_r :  $id_A \sqsubseteq \alpha \cdot \alpha^{\sharp}$
- univalent\_r :  $\alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- function\_r :  $id_A \sqsubseteq \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- surjection\_r :  $id_A \sqsubseteq \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha = id_B$
- injection\_r :  $id_A = \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha \sqsubseteq id_B$
- bijection\_r :  $id_A = \alpha \cdot \alpha^{\sharp} \wedge \alpha^{\sharp} \cdot \alpha = id_B$

```
Definition total\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (Id \ A) (alpha \cdot alpha \#). Definition univalent\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (alpha \# \cdot alpha) (Id \ B). Definition function\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (total\_r \ alpha) \land (univalent\_r \ alpha). Definition surjection\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (function\_r \ alpha) \land (total\_r \ (alpha \#)). Definition injection\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (function\_r \ alpha) \land (univalent\_r \ (alpha \#)). Definition bijection\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (function\_r \ alpha) \land (total\_r \ (alpha \#)) \land (univalent\_r \ (alpha \#)).
```

### 2.3 関係の公理

今後の諸定理の証明は、原則以下の公理群、およびそれらから導かれる補題のみを用いて行っていくことにする.

### 2.3.1 Dedekind 圏の公理

```
Axiom 1 (comp_id_l, comp_id_r) Let \alpha : A \to B. Then, id_A \cdot \alpha = \alpha \cdot id_B = \alpha.
```

```
Definition axiom1a := \forall (A \ B : eqType)(alpha : Rel \ A \ B), Id \ A \cdot alpha = alpha.
Axiom comp\_id\_l : axiom1a.
Definition axiom1b := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha \cdot Id \ B = alpha.
Axiom comp\_id\_r : axiom1b.
```

**Axiom 2 (comp\_assoc)** Let  $\alpha : A \rightarrow B$ ,  $\beta : B \rightarrow C$ , and  $\gamma : C \rightarrow D$ . Then,

$$(\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).$$

Definition axiom2 :=

 $\forall (A \ B \ C \ D : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ C \ D),$ 

 $(alpha \cdot beta) \cdot gamma = alpha \cdot (beta \cdot gamma).$ 

Axiom  $comp\_assoc: axiom2$ .

**Axiom 3 (inc\_refl)** Let  $\alpha : A \rightarrow B$ . Then,

 $\alpha \sqsubseteq \alpha$ .

Definition  $axiom3 := \forall (A B : eqType)(alpha : Rel A B), alpha alpha.$ 

Axiom  $inc\_refl: axiom3$ .

**Axiom 4 (inc\_trans)** Let  $\alpha, \beta, \gamma : A \rightarrow B$ . Then,

 $\alpha \sqsubseteq \beta \land \beta \sqsubseteq \gamma \Rightarrow \alpha \sqsubseteq \gamma.$ 

Definition  $axiom4 := \forall (A B : eqType)(alpha beta gamma : Rel A B),$ 

alpha beta  $\rightarrow$  beta  $gamma \rightarrow alpha$  gamma.

Axiom  $inc\_trans : axiom4$ .

**Axiom 5 (inc\_antisym)** Let  $\alpha, \beta : A \rightarrow B$ . Then,

 $\alpha \sqsubseteq \beta \land \beta \sqsubseteq \alpha \Rightarrow \alpha = \beta.$ 

Definition  $axiom5 := \forall (A B : eqType)(alpha beta : Rel A B),$ 

alpha beta  $\rightarrow$  beta  $alpha \rightarrow alpha =$ beta.

Axiom  $inc\_antisym : axiom 5$ .

**Axiom 6 (inc\_empty\_alpha)** Let  $\alpha : A \rightarrow B$ . Then,

 $\phi_{AB} \sqsubseteq \alpha$ .

**Definition**  $axiom6 := \forall (A B : eqType)(alpha : Rel A B), A B alpha.$ 

Axiom  $inc\_empty\_alpha$ : axiom 6.

```
Axiom 7 (inc_alpha_universal) Let \alpha : A \rightarrow B. Then,
```

$$\alpha \sqsubseteq \nabla_{AB}$$
.

Definition  $axiom 7 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha Axiom <math>inc\_alpha\_universal : axiom 7$ .

### Axiom 8 (inc\_capP, inc\_cap)

- 1.  $\mathbf{inc\_capP}$ : Let  $\alpha: A \to B$ ,  $f: (C \to D) \to (A \to B)$  and P: predicate. Then,  $\alpha \sqsubseteq (\sqcap_{P(\beta)} f(\beta)) \Leftrightarrow \forall \beta: C \to D, P(\beta) \Rightarrow \alpha \sqsubseteq f(\beta).$
- 2. inc\_cap : Let  $\alpha, \beta, \gamma : A \rightarrow B$ . Then,

$$\alpha \sqsubset (\beta \sqcap \gamma) \Leftrightarrow \alpha \sqsubset \beta \land \alpha \sqsubset \gamma.$$

```
Definition axiom8a :=
 \forall (A B C D : eqType)
 (alpha: Rel\ A\ B)(f: Rel\ C\ D \rightarrow Rel\ A\ B)(P: Rel\ C\ D \rightarrow Prop),
            ( \{P\} f) \leftrightarrow \forall \text{ beta} : Rel \ C \ D, P \text{ beta} \rightarrow alpha
Axiom inc\_capP: axiom8a.
Definition axiom8b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
 alpha
            (beta
                        gamma) \leftrightarrow (alpha \quad beta) \wedge (alpha)
                                                                         qamma).
Lemma inc\_cap: axiom8b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
rewrite inc\_capP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

### Axiom 9 (inc\_cupP, inc\_cup)

- 1.  $\operatorname{inc\_cupP} : \operatorname{Let} \alpha : A \to B, \ f : (C \to D) \to (A \to B) \ and \ P : \ predicate. \ Then,$   $(\sqcup_{P(\beta)} f(\beta)) \sqsubseteq \alpha \Leftrightarrow \forall \beta : C \to D, P(\beta) \Rightarrow f(\beta) \sqsubseteq \alpha.$
- 2. inc\_cup: Let  $\alpha, \beta, \gamma : A \rightarrow B$ . Then,

$$(\beta \sqcup \gamma) \sqsubseteq \alpha \Leftrightarrow \beta \sqsubseteq \alpha \land \gamma \sqsubseteq \alpha.$$

```
Definition axiom9a :=
 \forall (A \ B \ C \ D : eqType)
 (alpha: Rel\ A\ B)(f: Rel\ C\ D \rightarrow Rel\ A\ B)(P: Rel\ C\ D \rightarrow Prop),
                   alpha \leftrightarrow \forall beta : Rel\ C\ D, P beta \rightarrow f beta
Axiom inc\_cupP: axiom9a.
Definition axiom9b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             gamma)
                           alpha \leftrightarrow (\texttt{beta} \quad alpha) \land (gamma)
Lemma inc\_cup: axiom9b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
rewrite inc\_cupP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

```
Axiom 10 (inc_rpc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
```

$$\alpha \sqsubseteq (\beta \Rightarrow \gamma) \Leftrightarrow (\alpha \sqcap \beta) \sqsubseteq \gamma.$$

Definition  $axiom10 := \forall (A \ B : eqType)(alpha \ beta \ gamma : Rel \ A \ B),$   $alpha \ (beta \gg gamma) \leftrightarrow (alpha \ beta) \ gamma.$ Axiom  $inc\_rpc : axiom10$ .

**Axiom 11 (inv\_invol)** Let  $\alpha : A \rightarrow B$ . Then,

$$(\alpha^{\sharp})^{\sharp} = \alpha.$$

Definition  $axiom11 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), (alpha \#) \# = alpha.$ Axiom  $inv\_invol : axiom11$ .

**Axiom 12 (comp\_inv)** Let  $\alpha : A \rightarrow B$  and  $\beta : B \rightarrow C$ . Then,

$$(\alpha \cdot \beta)^{\sharp} = \beta^{\sharp} \cdot \alpha^{\sharp}.$$

Definition  $axiom12 := \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C),$   $(alpha \cdot beta) \# = (beta \# \cdot alpha \#).$ 

Axiom  $comp_inv : axiom12$ .

**Axiom 13 (inc\_inv)** Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$\alpha \sqsubseteq \beta \Rightarrow \alpha^{\sharp} \sqsubseteq \beta^{\sharp}.$$

Definition axiom13 :=

 $\forall (A \ B : eqType)(alpha \ \mathsf{beta} : Rel \ A \ B), \ alpha \ \mathsf{beta} \to alpha \ \# \ \mathsf{beta} \ \#.$  Axiom  $inc\_inv : axiom13$ .

**Axiom 14 (dedekind)** Let  $\alpha: A \rightarrow B$ ,  $\beta: B \rightarrow C$ , and  $\gamma: A \rightarrow C$ . Then,

$$(\alpha \cdot \beta) \sqcap \gamma \sqsubseteq (\alpha \sqcap (\gamma \cdot \beta^{\sharp})) \cdot (\beta \sqcap (\alpha^{\sharp} \cdot \gamma)).$$

Definition axiom14 :=

 $\forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),$   $((alpha \cdot beta) \quad gamma)$ 

 $((alpha \quad (gamma \cdot beta \#)) \cdot (beta \quad (alpha \# \cdot gamma))).$ 

Axiom dedekind : axiom 14.

**Axiom 15 (inc\_residual)** Let  $\alpha: A \to B$ ,  $\beta: B \to C$ , and  $\gamma: A \to C$ . Then,

$$\gamma \sqsubseteq (\alpha \rhd \beta) \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.$$

Definition axiom15 :=

 $\forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)($ beta : Rel  $B \ C)(gamma : Rel \ A \ C),$ 

 $qamma \quad (alpha \quad beta) \leftrightarrow (alpha \# \cdot qamma) \quad beta.$ 

Axiom  $inc\_residual$ : axiom15.

### 2.3.2 排中律

Dedekind 圏の公理のほかに、以下の"排中律"を仮定すれば、与えられる圏は Schröder 圏となり、Bool 代数の性質も満たされる. ちなみに剰余合成は補関係から定義可能なので、本来 Schröder 圏には剰余合成に関する公理は存在しない.

**Axiom 16 (complement\_classic)** Let  $\alpha : A \rightarrow B$ . Then,

$$\alpha \sqcup \alpha^- = \nabla_{AB}$$

**Definition**  $axiom16 := \forall (A B : eqType)(alpha : Rel A B),$ 

 $alpha \quad alpha \quad = \quad A B.$ 

Axiom complement\_classic : axiom16.

### 2.3.3 単域

### 1 点集合 *I* が単域となるための条件は

$$\phi_{II} \neq id_I \wedge id_I = \nabla_{II} \wedge \forall X, \nabla_{XI} \cdot \nabla_{IX} = \nabla_{XX}$$

だが、Rel の定義から左 2 つは証明できるため、右の式だけ仮定する.

Axiom 17 (unit\_universal)

$$\nabla_{AI} \cdot \nabla_{IA} = \nabla_{AA}$$

Definition  $axiom 17 := \forall (A : eqType), A i \cdot i A = A A.$ 

Axiom unit\_universal: axiom17.

### 2.3.4 弱選択公理

この "弱選択公理" を仮定すれば、排中律と単域の存在 (厳密には全域性公理) を利用して 点公理を導出できる.

**Axiom 18 (weak\_axiom\_of\_choice)** Let  $\alpha: I \to A$  be a total relation. Then,

$$\exists \beta: I \to A, \beta \sqsubseteq \alpha.$$

Definition  $axiom18 := \forall (A : eqType)(alpha : Rel i A),$ 

 $total\_r \ alpha \rightarrow \exists \ \mathsf{beta} : Rel \ i \ A, function\_r \ \mathsf{beta} \land \ \mathsf{beta} \qquad alpha.$ 

Axiom  $weak\_axiom\_of\_choice : axiom18$ .

### 2.3.5 関係の有理性

集合論では色々インポートしながら頑張って証明したので、できればそちらもご覧ください.

**Axiom 19 (rationality)** *Let*  $\alpha : A \rightarrow B$ . *Then,* 

$$\exists R, \exists f: R \to A, \exists g: R \to B, \alpha = f^{\sharp} \cdot g \land f \cdot f^{\sharp} \sqcap g \cdot g^{\sharp} = id_R.$$

Definition  $axiom19 := \forall (A \ B : eqType)(alpha : Rel \ A \ B),$   $\exists (R : eqType)(f : Rel \ R \ A)(g : Rel \ R \ B),$   $function\_r \ f \land function\_r \ g \land alpha = f \ \# \ \bullet \ g \land ((f \ \bullet f \ \#) \ (g \ \bullet g \ \#)) = Id \ R.$ Axiom rationality : axiom19.

### 2.3.6 直和と直積

### 任意の直和に対して、入射対が存在することを仮定する.

**Axiom 20 (pair\_of\_inclusions)**  $\exists j : A \to A + B, \exists k : B \to A + B,$ 

$$j \cdot j^{\sharp} = id_A \wedge k \cdot k^{\sharp} = id_B \wedge j \cdot k^{\sharp} = \phi_{AB} \wedge j^{\sharp} \cdot j \sqcup k^{\sharp} \cdot k = id_{A+B}.$$

Definition axiom20 :=

 $\forall$   $(A \ B : eqType), \exists (j : Rel \ A \ (sum\_eqType \ A \ B))(k : Rel \ B \ (sum\_eqType \ A \ B)),$  $<math>j \cdot j \# = Id \ A \wedge k \cdot k \# = Id \ B \wedge j \cdot k \# = A \ B \wedge$   $(j \# \cdot j) \quad (k \# \cdot k) = Id \ (sum\_eqType \ A \ B).$ Axiom  $pair\_of\_inclusions : axiom20$ .

### 任意の直積に対して、射影対が存在することを仮定する.

**Axiom 21** (pair\_of\_projections)  $\exists p: A \times B \to A, \exists q: A \times B \to B,$ 

$$p^{\sharp} \cdot q = \nabla_{AB} \wedge p \cdot p^{\sharp} \cap q \cdot q^{\sharp} = id_{A \times B}.$$

Definition axiom21 :=

 $\forall$  (A B : eqType),  $\exists$  (p : Rel (prod\_eqType A B) A)(q : Rel (prod\_eqType A B) B),  $p \# \cdot q = A B \wedge (p \cdot p \#) \quad (q \cdot q \#) = Id (prod_eqType A B) \wedge univalent_r p \wedge univalent_r q$ .

Axiom  $pair\_of\_projections : axiom21$ .

## Chapter 3

## Library Basic\_Notations\_Set

```
Require Export ssreflect eqtype bigop.
Require Export Logic.ClassicalFacts.
Require Import Logic.FunctionalExtensionality.
Require Import Logic.Classical_Prop.
Require Import Logic.IndefiniteDescription.
Require Import Logic.ProofIrrelevance.

Axiom prop_extensionality_ok: prop_extensionality.
```

### 3.1 定義

この章では、関係を集合論的に定義した場合の定義、およびその定義で諸公理が成立することを示す。公理名や記号などは Basic\_Notations と同じものを使用するため、Basic\_Lemms 以降ではそれの代わりにこのライブラリをインポートすることもできる.

```
Definition Rel\ (A\ B: eqType) := A \to B \to Prop.

Definition inverse\ \{A\ B: eqType\}\ (alpha: Rel\ A\ B): Rel\ B\ A
:= (fun\ (b:B)(a:A) \Rightarrow alpha\ a\ b).

Notation "a #" := (inverse\ a) (at level 20).

Definition composite\ \{A\ B\ C: eqType\}\ (alpha: Rel\ A\ B) (beta: Rel\ B\ C): Rel\ A\ C
:= (fun\ (a:A)(c:C) \Rightarrow \exists\ b:B,\ alpha\ a\ b \land beta\ b\ c).

Notation "a' 'b" := (composite\ a\ b) (at level 50).

Definition residual\ \{A\ B\ C: eqType\}\ (alpha: Rel\ A\ B) (beta: Rel\ B\ C): Rel\ A\ C
:= (fun\ (a:A)(c:C) \Rightarrow \forall\ b:B,\ alpha\ a\ b \to beta\ b\ c).

Notation "a' 'b" := (residual\ a\ b) (at level 50).

Definition identity\ (A: eqType): Rel\ A\ A: = (fun\ a\ a\theta:A \Rightarrow a=a\theta).

Notation "'Id'" := identity.

Definition empty\ (A\ B: eqType): Rel\ A\ B: = (fun\ (a:A)(b:B) \Rightarrow False).

Notation "' := empty.

Definition universal\ (A\ B: eqType): Rel\ A\ B: = (fun\ (a:A)(b:B) \Rightarrow True).
```

```
Notation "' := universal.
Definition include \{A \ B : eqType\} (alpha \ beta : Rel \ A \ B) : Prop
 := (\forall (a: A)(b: B), alpha \ a \ b \rightarrow beta \ a \ b).
Notation "a', b" := (include \ a \ b) (at level 50).
Definition cupP \{A \ B \ C \ D : eqType\}\ (P : Rel \ C \ D \to Prop)\ (f : Rel \ C \ D \to Rel \ A \ B)
: Rel \ A \ B
 := (\mathbf{fun} \ (a : A)(b : B) \Rightarrow \exists \ alpha : Rel \ C \ D, P \ alpha \wedge (f \ alpha) \ a \ b).
Notation "' _{\{',p'\}'} f" := (cupP \ p \ f) (at level 50).
Definition cap P \{A \ B \ C \ D : eqType\} \ (P : Rel \ C \ D \rightarrow Prop) \ (f : Rel \ C \ D \rightarrow Rel \ A \ B) :
Rel A B
 := (\mathbf{fun} \ (a : A)(b : B) \Rightarrow \forall \ alpha : Rel \ C \ D, \ P \ alpha \rightarrow (f \ alpha) \ a \ b).
Notation "' _{-}{' p '}' f" := (capP \ p \ f) (at level 50).
Definition cup \{A B : eqType\} (alpha beta : Rel A B)
       \{\text{fun } gamma : Rel \ A \ B \Rightarrow gamma = alpha \lor gamma = \text{beta}\} \ id.
Notation "a', b" := (cup \ a \ b) (at level 50).
Definition cap \{A B : eqType\} (alpha beta : Rel A B)
       \{\text{fun } gamma : Rel \ A \ B \Rightarrow gamma = alpha \lor gamma = \text{beta}\} \ id.
Notation "a', b" := (cap \ a \ b) (at level 50).
Definition rpc \{A \ B : eqType\} (alpha \ beta : Rel \ A \ B) : Rel \ A \ B
 := (fun (a : A)(b : B) \Rightarrow alpha \ a \ b \rightarrow beta \ a \ b).
Notation "a'»' b" := (rpc \ a \ b) (at level 50).
Definition complement \{A \ B : eqType\} (alpha : Rel \ A \ B) := alpha \gg
                                                                                        A B.
Notation "a '^' := (complement \ a) (at level 20).
Definition difference \{A B : eqType\} \{alpha beta : Rel A B\} := alpha
                                                                                         beta ^.
Notation "a - b" := (difference \ a \ b) (at level 50).
Notation "'i' := unit\_eqType.
```

### 3.2 関数の定義

```
Definition total\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (Id \ A) (alpha \cdot alpha \#).

Definition univalent\_r {A \ B : eqType} (alpha : Rel \ A \ B) := (alpha \# \cdot alpha) (Id \ B).

Definition function\_r {A \ B : eqType} (alpha : Rel \ A \ B)

:= (function\_r \ alpha) \land (funivalent\_r \ alpha).

Definition function\_r \ alpha) \land (funivalent\_r \ alpha).
```

Lemma 1 (comp\_id\_l, comp\_id\_r) Let  $\alpha : A \rightarrow B$ . Then,

### 3.3 関係の公理

今後の諸定理の証明は、原則以下の公理群、およびそれらから導かれる補題のみを用いて行っていくことにする.

### 3.3.1 Dedekind 圏の公理

```
id_A \cdot \alpha = \alpha \cdot id_B = \alpha.
Definition axiom1a := \forall (A B : eqType)(alpha : Rel A B), Id A • alpha = alpha.
Lemma comp_{-}id_{-}l: axiom1a.
Proof.
move \Rightarrow A \ B \ alpha.
{\tt apply} \ functional\_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split.
elim \Rightarrow a\theta.
elim \Rightarrow H H0.
rewrite H.
apply H0.
move \Rightarrow H.
\exists a.
split.
by [].
apply H.
Qed.
Definition axiom1b := \forall (A B : eqType)(alpha : Rel A B), alpha • Id B = alpha.
Lemma comp\_id\_r: axiom1b.
Proof.
move \Rightarrow A \ B \ alpha.
apply functional_extensionality.
move \Rightarrow a.
apply functional\_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split.
elim \Rightarrow b\theta.
```

 $elim \Rightarrow H H0$ .

split.

```
rewrite -H0.
apply H.
move \Rightarrow H.
\exists b.
split.
apply H.
by [].
Qed.
  Lemma 2 (comp_assoc) Let \alpha: A \rightarrow B, \beta: B \rightarrow C, and \gamma: C \rightarrow D. Then,
                                             (\alpha \cdot \beta) \cdot \gamma = \alpha \cdot (\beta \cdot \gamma).
Definition axiom2 :=
 \forall (A \ B \ C \ D : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ C \ D),
 (alpha \cdot beta) \cdot gamma = alpha \cdot (beta \cdot gamma).
Lemma comp\_assoc: axiom2.
Proof.
move \Rightarrow A B C D alpha beta gamma.
apply functional_extensionality.
\mathtt{move} \Rightarrow a.
apply functional_extensionality.
move \Rightarrow d.
apply prop_extensionality_ok.
split.
elim \Rightarrow c.
elim \Rightarrow H H0.
elim H \Rightarrow b \ H1.
\exists b.
split.
apply H1.
\exists c.
split.
apply H1.
apply H0.
elim \Rightarrow b.
elim \Rightarrow H.
elim \Rightarrow c H0.
\exists c.
split.
\exists b.
```

```
CHAPTER 3. LIBRARY BASIC_NOTATIONS_SET
apply H.
apply H0.
apply H0.
Qed.
  Lemma 3 (inc_refl) Let \alpha : A \rightarrow B. Then,
                                                    \alpha \sqsubset \alpha.
Definition axiom3 := \forall (A B : eqType)(alpha : Rel A B), alpha
                                                                                       alpha.
Lemma inc\_refl: axiom3.
Proof.
by [rewrite / axiom3 / include].
Qed.
  Lemma 4 (inc_trans) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                         \alpha \sqsubseteq \beta \land \beta \sqsubseteq \gamma \Rightarrow \alpha \sqsubseteq \gamma.
Definition axiom4 := \forall (A B : eqType)(alpha beta gamma : Rel A B),
                                  gamma \rightarrow alpha
            \mathtt{beta} \to \mathtt{beta}
                                                           qamma.
Lemma inc\_trans: axiom4.
Proof.
move \Rightarrow A B alpha beta gamma H H0 a b H1.
apply (H0 - (H - H1)).
Qed.
  Lemma 5 (inc_antisym) Let \alpha, \beta : A \rightarrow B. Then,
                                         \alpha \sqsubseteq \beta \land \beta \sqsubseteq \alpha \Rightarrow \alpha = \beta.
Definition axiom5 := \forall (A B : eqType)(alpha beta : Rel A B),
            beta \rightarrow beta
                                  alpha \rightarrow alpha = beta.
Lemma inc\_antisym : axiom5.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ H \ H0.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split.
```

apply H.

apply H0.

Qed.

Lemma 6 (inc\_empty\_alpha) Let  $\alpha : A \rightarrow B$ . Then,

 $\phi_{AB} \sqsubseteq \alpha$ .

**Definition**  $axiom6 := \forall (A B : eqType)(alpha : Rel A B), A B alpha.$ 

Lemma  $inc\_empty\_alpha$ : axiom6.

Proof.

 $move \Rightarrow A \ B \ alpha \ a \ b.$ 

apply  $False\_ind$ .

Qed.

Lemma 7 (inc\_alpha\_universal) Let  $\alpha : A \rightarrow B$ . Then,

 $\alpha \sqsubseteq \nabla_{AB}$ .

**Definition**  $axiom 7 := \forall (A B : eqType)(alpha : Rel A B), alpha A B.$ 

Lemma  $inc\_alpha\_universal: axiom 7.$ 

Proof

move  $\Rightarrow A \ B \ alpha \ a \ b \ H$ .

apply I.

Qed.

### Lemma 8 (inc\_capP, inc\_cap)

1. inc\_capP: Let  $\alpha: A \to B$ ,  $f: (C \to D) \to (A \to B)$  and P: predicate. Then,

$$\alpha \sqsubseteq (\sqcap_{P(\beta)} f(\beta)) \Leftrightarrow \forall \beta : C \to D, P(\beta) \Rightarrow \alpha \sqsubseteq f(\beta).$$

2. inc\_cap : Let  $\alpha, \beta, \gamma : A \rightarrow B$ . Then,

$$\alpha \sqsubseteq (\beta \sqcap \gamma) \Leftrightarrow \alpha \sqsubseteq \beta \land \alpha \sqsubseteq \gamma.$$

Definition axiom8a :=

 $\forall (A \ B \ C \ D : eqType)$ 

 $(alpha: Rel\ A\ B)(f: Rel\ C\ D \rightarrow Rel\ A\ B)(P: Rel\ C\ D \rightarrow Prop),$ 

alpha ( $\{P\} f$ )  $\leftrightarrow \forall$  beta:  $Rel\ C\ D$ , P beta  $\to alpha$  f beta.

Lemma  $inc\_capP$ : axiom8a.

Proof.

move  $\Rightarrow A B C D alpha f P$ .

 $split; move \Rightarrow H.$ 

```
move \Rightarrow beta H0 \ a \ b H1.
apply (H - H1 - H0).
move \Rightarrow a \ b \ H0 beta H1.
apply (H - H1 - H0).
Qed.
Definition axiom8b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
           (beta
                      qamma) \leftrightarrow (alpha)
                                               beta) \land (alpha)
                                                                     qamma).
Lemma inc\_cap: axiom8b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ qamma.
rewrite inc\_capP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
```

### Lemma 9 (inc\_cupP, inc\_cup)

- 1.  $inc\_cupP : Let \alpha : A \rightarrow B, f : (C \rightarrow D) \rightarrow (A \rightarrow B) \text{ and } P : predicate. Then,$  $(\sqcup_{P(\beta)} f(\beta)) \sqsubseteq \alpha \Leftrightarrow \forall \beta : C \to D, P(\beta) \Rightarrow f(\beta) \sqsubseteq \alpha.$
- 2. inc\_cup: Let  $\alpha, \beta, \gamma : A \rightarrow B$ . Then,

$$(\beta \sqcup \gamma) \sqsubseteq \alpha \Leftrightarrow \beta \sqsubseteq \alpha \land \gamma \sqsubseteq \alpha.$$

alpha.

```
Definition axiom9a :=
 \forall (A \ B \ C \ D : eqType)
 (alpha: Rel \ A \ B)(f: Rel \ C \ D \rightarrow Rel \ A \ B)(P: Rel \ C \ D \rightarrow Prop),
                   alpha \leftrightarrow \forall beta : Rel\ C\ D,\ P\ beta \rightarrow f\ beta
Lemma inc\_cupP: axiom9a.
Proof.
move \Rightarrow A B C D alpha f P.
split.
move \Rightarrow H beta H0 a b H1.
apply H.
∃ beta.
split.
apply H0.
apply H1.
```

```
move \Rightarrow H \ a \ b.
elim \Rightarrow beta.
elim \Rightarrow H0 \ H1.
apply (H \text{ beta } H0 - H1).
Definition axiom9b := \forall (A B : eqType)(alpha beta gamma : Rel A B),
                             alpha \leftrightarrow (\texttt{beta})
                                                     alpha) \wedge (qamma)
             gamma)
Lemma inc\_cup: axiom9b.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ qamma.
rewrite inc\_cupP.
split; move \Rightarrow H.
split; apply H.
by [left].
by [right].
move \Rightarrow delta H\theta.
case H0 \Rightarrow H1; rewrite H1; apply H.
Qed.
  Lemma 10 (inc_rpc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                        \alpha \sqsubseteq (\beta \Rightarrow \gamma) \Leftrightarrow (\alpha \sqcap \beta) \sqsubseteq \gamma.
Definition axiom 10 := \forall (A B : eqType)(alpha beta gamma : Rel A B),
             (beta \gg gamma) \leftrightarrow (alpha)
                                                  beta)
                                                                qamma.
Lemma inc\_rpc: axiom10.
Proof.
move \Rightarrow A \ B \ alpha \ beta \ gamma.
split; move \Rightarrow H.
move \Rightarrow a \ b \ H0.
apply H.
apply H0.
by [left].
apply H0.
by [right].
move \Rightarrow a \ b \ H0 \ H1.
apply H.
move \Rightarrow delta.
case \Rightarrow H2; rewrite H2.
apply H0.
apply H1.
Qed.
```

```
Lemma 11 (inv_invol) Let \alpha : A \rightarrow B. Then,
                                                    (\alpha^{\sharp})^{\sharp} = \alpha.
Definition axiom11 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), (alpha \#) \# = alpha.
Lemma inv\_invol: axiom11.
Proof.
by [move \Rightarrow A \ B \ alpha].
Qed.
  Lemma 12 (comp_inv) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                               (\alpha \cdot \beta)^{\sharp} = \beta^{\sharp} \cdot \alpha^{\sharp}.
Definition axiom12 := \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C),
 (alpha \cdot beta) \# = (beta \# \cdot alpha \#).
Lemma comp_inv : axiom12.
Proof.
move \Rightarrow A B C alpha beta.
apply functional_extensionality.
move \Rightarrow c.
apply functional_extensionality.
move \Rightarrow a.
apply prop_extensionality_ok.
split; elim \Rightarrow b.
elim \Rightarrow H H0.
\exists b.
split.
apply H0.
apply H.
elim \Rightarrow H H0.
\exists b.
split.
apply H0.
apply H.
Qed.
  Lemma 13 (inc_inv) Let \alpha, \beta : A \rightarrow B. Then,
```

 $\alpha \sqsubseteq \beta \Rightarrow \alpha^{\sharp} \sqsubseteq \beta^{\sharp}.$ 

Definition axiom13 :=

 $\forall (A B : eqType)(alpha beta : Rel A B), alpha$  $\mathtt{beta} o alpha \ \#$ beta #.

```
Lemma inc_inv : axiom13.
move \Rightarrow A \ B \ alpha \ beta \ H \ b \ a \ H\theta.
apply (H - H0).
Qed.
  Lemma 14 (dedekind) Let \alpha: A \rightarrow B, \beta: B \rightarrow C, and \gamma: A \rightarrow C. Then,
                                 (\alpha \cdot \beta) \sqcap \gamma \sqsubseteq (\alpha \sqcap (\gamma \cdot \beta^{\sharp})) \cdot (\beta \sqcap (\alpha^{\sharp} \cdot \gamma)).
Definition axiom14 :=
 \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),
 ((alpha • beta)
                           gamma)
                    (gamma \cdot beta \#)) \cdot (beta
                                                                (alpha \# \bullet gamma))).
     ((alpha
Lemma dedekind: axiom 14.
Proof.
move \Rightarrow A B C alpha beta gamma a c H.
assert (\exists b : B, alpha \ a \ b \land beta \ b \ c).
apply H.
by [left].
elim H\theta \Rightarrow b.
\mathtt{elim} \Rightarrow \mathit{H1}\ \mathit{H2}.
\exists b.
repeat split.
move \Rightarrow delta H3.
case H3 \Rightarrow H4; rewrite H4.
apply H1.
unfold id.
\exists c.
split.
apply H.
by [right].
apply H2.
move \Rightarrow delta H3.
case H3 \Rightarrow H4; rewrite H4.
apply H2.
unfold id.
\exists a.
split.
apply H1.
apply H.
by [right].
Qed.
```

```
Lemma 15 (inc_residual) Let \alpha: A \to B, \beta: B \to C, and \gamma: A \to C. Then, \gamma \sqsubseteq (\alpha \rhd \beta) \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.
```

```
Definition axiom15 :=
 \forall (A \ B \ C : eqType)(alpha : Rel \ A \ B)(beta : Rel \ B \ C)(gamma : Rel \ A \ C),
                          \texttt{beta}) \leftrightarrow (alpha \# \bullet gamma)
Lemma inc_residual: axiom15.
Proof.
move \Rightarrow A B C alpha beta gamma.
split; move \Rightarrow H.
move \Rightarrow b c.
elim \Rightarrow a H0.
apply (H \ a).
apply H0.
apply H0.
move \Rightarrow a \ c \ H0 \ b \ H1.
apply H.
\exists a.
split.
apply H1.
apply H0.
Qed.
```

### 3.3.2 排中律

Dedekind 圏の公理のほかに、以下の"排中律"を仮定すれば、与えられる圏は Schröder 圏となり、Bool 代数の性質も満たされる. ちなみに剰余合成は補関係から定義可能なので、本来 Schröder 圏には剰余合成に関する公理は存在しない.

Lemma 16 (complement\_classic) Let  $\alpha : A \rightarrow B$ . Then,

$$\alpha \sqcup \alpha^- = \nabla_{AB}$$

```
Definition axiom16 := \forall (A \ B : eqType)(alpha : Rel \ A \ B), alpha \quad alpha \ \hat{\ } = A \ B.
Lemma complement\_classic : axiom16.
Proof.
move \Rightarrow A \ B \ alpha.
apply functional\_extensionality.
move \Rightarrow a.
apply functional\_extensionality.
```

```
move \Rightarrow b.

apply prop\_extensionality\_ok.

split; move \Rightarrow H.

apply I.

case (classic\ (alpha\ a\ b)) \Rightarrow H0.

\exists\ alpha.

split.

by [left].

apply H0.

\exists\ (\mathbf{fun}\ (a0:A)\ (b0:B) \Rightarrow alpha\ a0\ b0 \rightarrow False).

split.

by [right].

apply H0.

Qed.
```

### 3.3.3 単域

### 1点集合 / が単域となるための条件は

$$\phi_{II} \neq id_I \wedge id_I = \nabla_{II} \wedge \forall X, \nabla_{XI} \cdot \nabla_{IX} = \nabla_{XX}$$

だが、Rel の定義から左2 つは証明できるため、右の式だけ仮定する.

Lemma 17 (unit\_universal)

$$\nabla_{AI} \cdot \nabla_{IA} = \nabla_{AA}$$

```
Definition axiom 17 := \forall (A : eqType),
                                                                           A A.
                                                   A i \cdot i A =
Lemma unit\_universal : axiom17.
Proof.
move \Rightarrow A.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow a\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
apply I.
\exists tt.
by [].
Qed.
```

### 3.3.4 弱選択公理

この "弱選択公理" を仮定すれば、排中律と単域の存在 (厳密には全域性公理) を利用して 点公理を導出できる.

Lemma 18 (weak\_axiom\_of\_choice) Let  $\alpha: I \to A$  be a total relation. Then,

$$\exists \beta: I \to A, \beta \sqsubseteq \alpha.$$

```
Definition axiom18 := \forall (A : eqType)(alpha : Rel i A),
 total_r \ alpha \rightarrow \exists \ \mathtt{beta} : Rel \ i \ A, function_r \ \mathtt{beta} \land \mathtt{beta}
                                                                                     alpha.
Lemma weak_axiom_of_choice: axiom18.
Proof.
move \Rightarrow A \ alpha.
rewrite / function_r / total_r / univalent_r / identity / include / composite / inverse.
move \Rightarrow H.
move: (H tt tt (Logic.eq_refl tt)).
elim \Rightarrow a H0.
\exists (\mathbf{fun} (\_: i)(a\theta : A) \Rightarrow a = a\theta).
repeat split.
move \Rightarrow tt \ tt0 \ H1.
by [\exists a].
move \Rightarrow a\theta \ a1.
elim \Rightarrow tt0.
elim \Rightarrow H1 H2.
by [rewrite -H1 -H2].
induction a\theta.
move \Rightarrow a0 H1.
rewrite -H1.
apply H0.
Qed.
```

### 3.3.5 関係の有理性

集合の選択公理 (Logic.IndefiniteDescription) や証明の一意性 (Logic.ProofIrrelevance) を仮定すれば、集合論上ならごり押しで証明できる. 旧ライブラリの頃は無理だと諦めて Axiom を追加していたが、Standard Library のインポートだけで解けた. 正直びっくり.

Lemma 19 (rationality) Let  $\alpha : A \rightarrow B$ . Then,

$$\exists R, \exists f: R \to A, \exists g: R \to B, \alpha = f^{\sharp} \cdot g \land f \cdot f^{\sharp} \sqcap g \cdot g^{\sharp} = id_R.$$

この付近は、ごり押しのための補題. 命題の真偽を選択公理で bool 値に変換したり、部分集合の元から上位集合の元を生成する sval (proj1\_sig) の単射性を示したりしている.

```
Lemma is\_true\_inv0: \forall P: Prop, \exists b: bool, P \leftrightarrow is\_true b.
Proof.
move \Rightarrow P.
case (classic P); move \Rightarrow H.
\exists true.
split; move \Rightarrow H0.
by ||.
apply H.
\exists false.
split; move \Rightarrow H0.
apply False_ind.
apply (H H\theta).
discriminate H0.
Definition is\_true\_inv : Prop \rightarrow bool.
move \Rightarrow P.
move: (is\_true\_inv0 \ P) \Rightarrow H.
apply constructive\_indefinite\_description in H.
apply H.
Defined.
Lemma is\_true\_id : \forall P : Prop, is\_true (is\_true\_inv P) \leftrightarrow P.
Proof.
move \Rightarrow P.
unfold is\_true\_inv.
move: (constructive\_indefinite\_description (fun b : bool \Rightarrow P \leftrightarrow is\_true b) (is\_true\_inv0)
(P)) \Rightarrow x\theta.
apply (@sig\_ind\ bool\ (fun\ b \Rightarrow (P \leftrightarrow is\_true\ b))\ (fun\ y \Rightarrow is\_true\ (let\ (x,\_) := y\ in\ x)
\leftrightarrow P)).
\mathtt{move} \Rightarrow x\ H.
apply iff_-sym.
apply H.
Qed.
Lemma sval\_inv : \forall (A : Type)(P : A \rightarrow Prop)(x : sig P)(a : A), a = sval x \rightarrow \exists (H : P a),
x = exist P a H.
Proof.
move \Rightarrow A P x a H0.
rewrite H0.
\exists (proj2\_sig \ x).
apply (@sig\_ind \ A \ P \ (fun \ y \Rightarrow y = exist \ P \ (sval \ y) \ (proj2\_sig \ y))).
```

```
move \Rightarrow a\theta H.
by [simpl].
Qed.
Lemma sval\_injective : \forall (A : Type)(P : A \rightarrow Prop)(x \ y : siq \ P), sval \ x = sval \ y \rightarrow x = y.
Proof.
move \Rightarrow A P x y H.
move: (sval\_inv \ A \ P \ y \ (sval \ x) \ H).
elim \Rightarrow H0 \ H1.
rewrite H1.
assert (H0 = proj2\_siq x).
apply proof_irrelevance.
rewrite H2.
apply (@siq\_ind \ A \ P \ (fun \ y \Rightarrow y = exist \ P \ (sval \ y) \ (proj2\_siq \ y))).
move \Rightarrow a0 H3.
by [simpl].
Qed.
Definition axiom 19 := \forall (A B : eqType)(alpha : Rel A B),
 \exists (R : eqType)(f : Rel R A)(g : Rel R B),
 function\_r\ f \land function\_r\ g \land alpha = f \# \bullet g \land ((f \bullet f \#) \quad (g \bullet g \#)) = Id\ R.
Lemma rationality: axiom19.
Proof.
move \Rightarrow A \ B \ alpha.
\verb"rewrite" / function\_r/total\_r/univalent\_r/cap/capP/identity/composite/inverse/include.
\exists (sig\_eqType (fun \ x : prod\_eqType \ A \ B \Rightarrow is\_true\_inv (alpha (fst \ x) (snd \ x)))).
\exists (\mathbf{fun} \ x \ a \Rightarrow a = (fst \ (sval \ x))).
\exists (\mathbf{fun} \ x \ b \Rightarrow b = (snd \ (sval \ x))).
simpl.
repeat split.
move \Rightarrow x \ x\theta \ H.
\exists (fst (sval x)).
repeat split.
by [rewrite H].
move \Rightarrow a \ a\theta.
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
move \Rightarrow x \ x\theta \ H.
\exists (snd (sval x)).
repeat split.
by [rewrite H].
move \Rightarrow b \ b\theta.
```

```
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split; move \Rightarrow H.
assert (is\_true (is\_true\_inv (alpha (fst (a,b)) (snd (a,b))))).
simpl.
apply is_true_id.
apply H.
\exists (exist (fun \ x \Rightarrow (is\_true \ (is\_true\_inv \ (alpha \ (fst \ x) \ (snd \ x))))) (a,b) \ H0).
by simpl.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 H1.
apply is\_true\_id.
apply (@sig\_ind (A × B) (fun x \Rightarrow is\_true (is\_true\_inv (alpha (fst x) (snd x)))) (fun x
\Rightarrow is\_true\ (is\_true\_inv\ (alpha\ (fst\ (sval\ x))\ (snd\ (sval\ x)))))).
simpl.
by [move \Rightarrow x\theta].
apply functional_extensionality.
move \Rightarrow y.
apply functional_extensionality.
move \Rightarrow y\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
apply sval_injective.
move: (H \text{ (fun } a \ c : \{x : A \times B \mid is\_true \ (is\_true\_inv \ (alpha \ (fst \ x) \ (snd \ x)))\} \Rightarrow \exists \ b :
A, b = fst (sval \ a) \land b = fst (sval \ c)) (or\_introl \ Logic.eq\_reft)).
move: (H \text{ (fun } a \text{ } c \text{ : } \{x \text{ : } A \times B \mid is\_true \text{ (} is\_true\_inv \text{ (} alpha \text{ (} fst \text{ } x \text{) (} snd \text{ } x \text{))})\} \Rightarrow \exists b \text{ : }
B, b = snd (sval \ a) \land b = snd (sval \ c)) (or\_intror \ Logic.eq\_reft)).
unfold id.
clear H.
elim \Rightarrow b.
elim \Rightarrow H H0.
elim \Rightarrow a.
elim \Rightarrow H1 H2.
rewrite (surjective_pairing (sval y0)) -H0 -H2 H H1.
apply surjective_pairing.
```

```
rewrite H.

move \Rightarrow beta H\theta.

case H\theta \Rightarrow H1; rewrite H1; unfold id.

\exists \ (fst \ (sval \ y\theta)).

repeat split.

\exists \ (snd \ (sval \ y\theta)).

repeat split.

Qed.
```

### 3.3.6 直和と直積

```
任意の直和に対して、入射対が存在することを仮定する。
Lemma 20 (pair_of_inclusions) \exists j:A \to A+B, \exists k:B \to A+B, j\cdot j^\sharp=id_A \wedge k\cdot k^\sharp=id_B \wedge j\cdot k^\sharp=\phi_{AB} \wedge j^\sharp\cdot j\sqcup k^\sharp\cdot k=id_{A+B}.
Definition axiom20:=
```

```
Definition axiom20 :=
      \forall (A B : eqType), \exists (j : Rel A (sum\_eqType A B))(k : Rel B (sum\_eqType A B)),
    j \cdot j \# = Id \ A \wedge k \cdot k \# = Id \ B \wedge j \cdot k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = Id \ B \wedge j \circ k \# = I
                                                                                                                                                                                                                                                                                                                       A B \wedge
      (j \# \cdot j) (k \# \cdot k) = Id (sum\_eqType A B).
Lemma pair\_of\_inclusions: axiom20.
Proof.
move \Rightarrow A B.
\exists (fun (a : A)(x : sum\_eqType A B) \Rightarrow x = inl a).
\exists (\mathbf{fun} \ (b : B)(x : sum\_eqType \ A \ B) \Rightarrow x = inr \ b).
repeat split.
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow a\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
by [injection H1].
\exists (inl a).
repeat split.
by [rewrite H].
apply functional_extensionality.
move \Rightarrow b.
```

```
apply functional_extensionality.
move \Rightarrow b\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
by [injection H1].
\exists (inr \ b).
repeat split.
by [rewrite H].
apply functional_extensionality.
move \Rightarrow a.
apply functional_extensionality.
move \Rightarrow b.
apply prop_extensionality_ok.
split; move \Rightarrow H.
elim H \Rightarrow x.
elim \Rightarrow H0 \ H1.
rewrite H0 in H1.
discriminate H1.
apply False_ind.
apply H.
apply functional_extensionality.
move \Rightarrow x.
apply functional_extensionality.
move \Rightarrow x\theta.
apply prop_extensionality_ok.
split.
elim \Rightarrow alpha.
elim \Rightarrow H0 \ H1.
case H0 \Rightarrow H2; rewrite H2 in H1.
elim H1 \Rightarrow a.
elim \Rightarrow H3 H4.
by [rewrite H3 H4].
elim H1 \Rightarrow b.
elim \Rightarrow H3 H4.
by [rewrite H3 H4].
assert ((\exists a : A, x = inl \ a) \lor (\exists b : B, x = inr \ b)).
move: x.
apply sum_{-}ind.
move \Rightarrow a.
```

```
left.
by [\exists a].
move \Rightarrow b.
right.
by [\exists b].
case H.
elim \Rightarrow a H0 H1.
\exists (fun x \ x\theta \Rightarrow \exists \ a\theta : A, (x = inl \ a\theta \land x\theta = inl \ a\theta)).
split.
by [left].
\exists a.
by [rewrite -H1 H0].
elim \Rightarrow b H0 H1.
\exists (fun x \ x\theta \Rightarrow \exists \ b\theta : B, (x = inr \ b\theta \land x\theta = inr \ b\theta)).
split.
by [right].
\exists b.
by [rewrite -H1 H0].
Qed.
  任意の直積に対して、射影対が存在することを仮定する.
  Lemma 21 (pair_of_projections) \exists p: A \times B \to A, \exists q: A \times B \to B,
                                     p^{\sharp} \cdot q = \nabla_{AB} \wedge p \cdot p^{\sharp} \sqcap q \cdot q^{\sharp} = id_{A \times B}.
Definition axiom21 :=
 \forall (A B : eqType), \exists (p : Rel (prod\_eqType A B) A)(q : Rel (prod\_eqType A B) B),
 p \# \bullet q = A B \land (p \bullet p \#) \quad (q \bullet q \#) = Id (prod\_eqType A B) \land univalent\_r p
```

```
Definition axiom21 := \forall (A \ B : eqType), \exists (p : Rel (prod_eqType \ A \ B) \ A)(q : Rel (prod_eqType \ A \ B) \ p \# \bullet q = A \ B \land (p \bullet p \#) \ (q \bullet q \#) = Id (prod_eqType \ A \ B) \land un \land univalent\_r \ q.
Lemma pair\_of\_projections : axiom21.
Proof.

move \Rightarrow A \ B.
\exists (fun \ (x : prod_eqType \ A \ B)(a : A) \Rightarrow a = (fst \ x)).
\exists (fun \ (x : prod_eqType \ A \ B)(b : B) \Rightarrow b = (snd \ x)).
split.
apply functional\_extensionality.
move \Rightarrow a.
apply functional\_extensionality.
move \Rightarrow b.
apply functional\_extensionality\_ok.
split; functional\_extensionality\_ok.
split; functional\_extensionality\_ok.
```

```
\exists (a,b).
by [simpl].
split.
apply functional_extensionality.
move \Rightarrow x.
apply functional_extensionality.
move \Rightarrow x\theta.
apply prop_extensionality_ok.
split; move \Rightarrow H.
move: (H (fun a c: prod_eqType A B \Rightarrow \exists b: A, b = fst a \land b = fst c) (or_introl
Logic.eq\_refl).
move: (H \text{ (fun } a \ c : prod\_eqType } A \ B \Rightarrow \exists \ b : B, \ b = snd \ a \land b = snd \ c) \ (or\_intror)
Logic.eq\_refl).
unfold id.
clear H.
elim \Rightarrow b.
elim \Rightarrow H H0.
elim \Rightarrow a.
elim \Rightarrow H1 H2.
rewrite (surjective_pairing x0) -H0 -H2 H H1.
apply surjective_pairing.
rewrite H.
move \Rightarrow alpha H0.
case H0 \Rightarrow H1; rewrite H1; unfold id.
\exists (fst \ x\theta).
repeat split.
\exists (snd x\theta).
repeat split.
split.
move \Rightarrow a \ a\theta.
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
move \Rightarrow b \ b\theta.
elim \Rightarrow x.
elim \Rightarrow H H0.
by [rewrite H H\theta].
Qed.
```

## Chapter 4

## Library Basic\_Lemmas

```
Require Import Basic_Notations.
Require Import Logic.Classical_Prop.
```

### 4.1 束論に関する補題

### 4.1.1 和関係, 共通関係

```
Lemma 22 (cap_l) Let \alpha, \beta : A \rightarrow B. Then,
                                               \alpha \sqcap \beta \sqsubseteq \alpha.
Lemma cap_l {A B : eqType} {alpha beta : Rel A B}: (alpha
                                                                                       alpha.
                                                                           beta)
Proof.
assert ((alpha
                      beta)
                                 (alpha
                                            beta)).
apply inc\_reft.
apply inc\_cap in H.
apply H.
Qed.
  Lemma 23 (cap_r) Let \alpha, \beta : A \rightarrow B. Then,
                                               \alpha \sqcap \beta \sqsubseteq \beta.
Lemma cap_r \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: (alpha
                                                                            beta)
                                                                                       beta.
Proof.
                               (alpha
assert ((alpha
                     beta)
                                            beta)).
apply inc_refl.
apply inc\_cap in H.
apply H.
```

Qed.

```
Lemma 24 (cup_l) Let \alpha, \beta : A \rightarrow B. Then,
                                                 \alpha \sqsubseteq \alpha \sqcup \beta.
Lemma cup_l \{A B : eqType\} \{alpha \text{ beta} : Rel A B\}: alpha
                                                                              (alpha
                                                                                          beta).
assert ((alpha
                      beta)
                                  (alpha
                                              beta)).
apply inc\_reft.
apply inc\_cup in H.
apply H.
Qed.
  Lemma 25 (cup_r) Let \alpha, \beta : A \rightarrow B. Then,
                                                 \beta \sqsubseteq \alpha \sqcup \beta.
Lemma cup_r \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: beta
                                                                              (alpha
                                                                                          beta).
Proof.
assert ((alpha
                      beta)
                                (alpha
                                              beta)).
apply inc_refl.
apply inc\_cup in H.
apply H.
Qed.
  Lemma 26 (inc_def1) Let \alpha, \beta : A \rightarrow B. Then,
                                           \alpha = \alpha \sqcap \beta \Leftrightarrow \alpha \sqsubseteq \beta.
Lemma inc\_def1 {A B : eqType} {alpha beta : Rel A B}:
 alpha = alpha
                    \mathtt{beta} \leftrightarrow alpha
                                            beta.
Proof.
split; move \Rightarrow H.
                     (alpha
assert (alpha
                                 beta)).
rewrite -H.
apply inc\_reft.
apply inc\_cap in H0.
apply H0.
apply inc\_antisym.
apply inc_-cap.
split.
apply inc\_reft.
```

```
apply H.
apply cap_{-}l.
Qed.
  Lemma 27 (inc_def2) Let \alpha, \beta : A \rightarrow B. Then,
                                        \beta = \alpha \sqcup \beta \Leftrightarrow \alpha \sqsubset \beta.
Lemma inc\_def2 {A B : eqType} {alpha beta : Rel A B}:
                   \mathtt{beta} \leftrightarrow alpha
 beta = alpha
                                         beta.
Proof.
split; move \Rightarrow H.
assert ((alpha)
                     beta)
                               beta).
rewrite -H.
apply inc_refl.
apply inc\_cup in H0.
apply H0.
apply inc\_antisym.
assert ((alpha
                     beta)
                              (alpha  beta)).
apply inc\_reft.
apply cup_r.
apply inc_-cup.
split.
apply H.
apply inc_refl.
Qed.
  Lemma 28 (cap_assoc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                     (\alpha \sqcap \beta) \sqcap \gamma = \alpha \sqcap (\beta \sqcap \gamma).
Lemma cap\_assoc {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha
            beta)
                       gamma = alpha
                                            (beta
                                                         qamma).
Proof.
apply inc\_antisym.
rewrite inc\_cap.
split.
apply (inc_trans _ _ _ (alpha
                                     beta)).
apply cap_{-}l.
apply cap_{-}l.
rewrite inc_-cap.
split.
apply (inc_trans _ _ _ (alpha
                                     beta)).
```

```
apply cap_{-}l.
apply cap_{-}r.
apply cap_{-}r.
rewrite inc\_cap.
split.
rewrite inc_-cap.
split.
apply cap_{-}l.
apply (inc_trans _ _ _ (beta
                                   gamma)).
apply cap_{-}r.
apply cap_{-}l.
apply (inc_trans _ _ _ (beta
                                   gamma)).
apply cap_r.
apply cap_{-}r.
Qed.
  Lemma 29 (cup_assoc) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                   (\alpha \sqcup \beta) \sqcup \gamma = \alpha \sqcup (\beta \sqcup \gamma).
Lemma cup\_assoc\ \{A\ B: eqType\}\ \{alpha\ beta\ gamma: Rel\ A\ B\}:
                      gamma = alpha
 (alpha
            beta)
                                           (beta
                                                      qamma).
Proof.
apply inc\_antisym.
rewrite inc\_cup.
split.
rewrite inc\_cup.
split.
apply cup_l.
apply (inc_trans _ _ _ (beta
                                   qamma)).
apply cup_{-}l.
apply cup_r.
apply (inc_trans _ _ _ (beta
                                    qamma)).
apply cup_{-}r.
apply cup_r.
rewrite inc\_cup.
split.
apply (inc_trans _ _ _ (alpha
                                   beta)).
apply cup_{-}l.
apply cup_{-}l.
rewrite inc_-cup.
split.
apply (inc_trans _ _ _ (alpha
                                   beta)).
```

```
apply cup_r.
apply cup_{-}l.
apply cup_{-}r.
Qed.
  Lemma 30 (cap_comm) Let \alpha, \beta : A \rightarrow B. Then,
                                          \alpha \sqcap \beta = \beta \sqcap \alpha.
Lemma cap\_comm {A B : eqType} {alpha beta : Rel A B}: alpha
                                                                            beta = beta
                                                                                               alpha.
Proof.
apply inc\_antisym.
rewrite inc\_cap.
split.
apply cap_{-}r.
apply cap_{-}l.
rewrite inc\_cap.
split.
apply cap_r.
apply cap_{-}l.
Qed.
  Lemma 31 (cup_comm) Let \alpha, \beta : A \rightarrow B. Then,
                                          \alpha \sqcup \beta = \beta \sqcup \alpha.
Lemma cup\_comm {A B : eqType} {alpha beta : Rel A B}: alpha beta = beta
                                                                                               alpha.
Proof.
apply inc\_antisym.
rewrite inc_-cup.
split.
apply cup_r.
apply cup_{-}l.
rewrite inc\_cup.
split.
apply cup_r.
apply cup_{-}l.
Qed.
  Lemma 32 (cup_cap_abs) Let \alpha, \beta : A \rightarrow B. Then,
```

 $\alpha \sqcup (\alpha \sqcap \beta) = \alpha.$ 

```
Lemma cup\_cap\_abs {A B : eqType} {alpha beta : Rel A B}:
                      beta) = alpha.
 alpha
           (alpha
Proof.
move: (@cap_l - alpha beta) \Rightarrow H.
apply inc\_def2 in H.
by [rewrite cup\_comm - H].
Qed.
  Lemma 33 (cap_cup_abs) Let \alpha, \beta : A \rightarrow B. Then,
                                         \alpha \sqcap (\alpha \sqcup \beta) = \alpha.
Lemma cap\_cup\_abs {A B : eqType} {alpha beta : Rel A B}:
 alpha
           (alpha
                      beta) = alpha.
Proof.
move: (@cup_l - alpha beta) \Rightarrow H.
apply inc\_def1 in H.
by [rewrite -H].
Qed.
  Lemma 34 (cap_idem) Let \alpha : A \rightarrow B. Then,
                                            \alpha \sqcap \alpha = \alpha.
Lemma cap\_idem \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha
                                                                      alpha = alpha.
Proof.
apply inc\_antisym.
apply cap_{-}l.
apply inc\_cap.
split; apply inc\_refl.
Qed.
  Lemma 35 (cup_idem) Let \alpha : A \rightarrow B. Then,
                                            \alpha \sqcup \alpha = \alpha.
Lemma cup\_idem \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha
                                                                      alpha = alpha.
Proof.
apply inc\_antisym.
apply inc\_cup.
split; apply inc\_refl.
apply cup_{-}l.
Qed.
```

```
Lemma 36 (cap_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                                      \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha' \sqcap \beta'.
Lemma cap_inc_compat {A B : eqType} {alpha alpha' beta beta' : Rel A B}:
              alpha' \rightarrow \mathtt{beta} \qquad beta' \rightarrow (alpha)
                                                                 beta)
                                                                              (alpha'
                                                                                              beta').
Proof.
move \Rightarrow H H0.
rewrite -inc_{-}def1.
apply inc\_def1 in H.
apply inc\_def1 in H0.
rewrite cap\_assoc -(@cap\_assoc _ _ beta).
rewrite (@cap\_comm\_\_beta).
rewrite cap\_assoc -(@cap\_assoc _ _ alpha).
by [rewrite -H - H\theta].
Qed.
  Lemma 37 (cap_inc_compat_l) Let \alpha, \beta, \beta' : A \rightarrow B. Then,
                                             \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha \sqcap \beta'.
Lemma cap\_inc\_compat\_l {A B : eqType} {alpha beta beta' : Rel A B}:
             beta' \rightarrow (alpha \quad beta) \quad (alpha)
                                                                 beta').
 beta
Proof.
move \Rightarrow H.
apply (@cap_inc_compat _ _ _ _ (@inc_reft _ _ alpha) H).
Qed.
  Lemma 38 (cap_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                             \alpha \sqsubseteq \alpha' \Rightarrow \alpha \sqcap \beta \sqsubseteq \alpha' \sqcap \beta.
Lemma cap\_inc\_compat\_r {A B : eqType} {alpha \ alpha' \ beta : Rel A B}:
 alpha
              alpha' \rightarrow (alpha \quad beta)
                                                    (alpha')
                                                                     beta).
Proof.
move \Rightarrow H.
apply (@cap_inc_compat _ _ _ H (@inc_reft _ beta)).
Qed.
  Lemma 39 (cup_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                                      \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha' \sqcup \beta'.
```

```
Lemma cup_inc_compat {A B : eqType} {alpha alpha' beta beta' : Rel A B}:
            alpha' \rightarrow beta \qquad beta' \rightarrow (alpha)
 alpha
                                                          beta)
                                                                     (alpha')
                                                                                  beta').
Proof.
move \Rightarrow H H0.
rewrite -inc_-def2.
apply inc_{-}def2 in H.
apply inc\_def2 in H0.
rewrite cup\_assoc -(@cup\_assoc _ _ beta).
rewrite (@cup\_comm\_\_ beta).
rewrite cup\_assoc -(@cup\_assoc _ _ alpha).
by [rewrite -H - H\theta].
Qed.
  Lemma 40 (cup_inc_compat_l) Let \alpha, \beta, \beta' : A \rightarrow B. Then,
                                        \beta \sqsubseteq \beta' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha \sqcup \beta'.
Lemma cup\_inc\_compat\_l {A B : eqType} {alpha beta beta' : Rel A B}:
           beta' \rightarrow (alpha \quad beta)
                                           (alpha
 beta
                                                          beta').
Proof.
move \Rightarrow H.
apply (@cup\_inc\_compat\_\_\_\_\_ (@inc\_refl\_\_ alpha) H).
Qed.
  Lemma 41 (cup_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                        \alpha \sqsubseteq \alpha' \Rightarrow \alpha \sqcup \beta \sqsubseteq \alpha' \sqcup \beta.
Lemma cup\_inc\_compat\_r {A B : eqType} {alpha \ alpha' \ beta : Rel A B}:
            alpha' \rightarrow (alpha)
                                              (alpha')
 alpha
                                   beta)
                                                             beta).
Proof.
move \Rightarrow H.
apply (@cup\_inc\_compat\_\_\_\_\_H (@inc\_refl\_\_beta)).
Qed.
  Lemma 42 (cap_empty) Let \alpha : A \rightarrow B. Then,
                                              \alpha \sqcap \phi_{AB} = \phi_{AB}.
Lemma cap\_empty \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha
                                                                                 A B =
                                                                                              A B.
Proof.
apply inc\_antisym.
apply cap_{-}r.
```

apply  $inc\_empty\_alpha$ . Qed.

**Lemma 43 (cup\_empty)** Let  $\alpha : A \rightarrow B$ . Then,

 $\alpha \sqcup \phi_{AB} = \alpha$ .

Lemma  $cup\_empty$   $\{A \ B : eqType\}$   $\{alpha : Rel \ A \ B\}$ : alpha  $A \ B = alpha$ . Proof.

apply  $inc\_antisym$ .

apply  $inc\_cup$ .

split.

apply  $inc\_reft$ .

apply  $inc\_empty\_alpha$ .

apply  $cup_{-}l$ .

Qed.

Lemma 44 (cap\_universal) Let  $\alpha : A \rightarrow B$ . Then,

 $\alpha \sqcap \nabla_{AB} = \alpha.$ 

Proof.

apply  $inc\_antisym$ .

apply  $cap_{-}l$ .

apply  $inc\_cap$ .

split.

apply  $inc\_reft$ .

apply  $inc\_alpha\_universal$ .

Qed.

Lemma 45 (cup\_universal) Let  $\alpha : A \rightarrow B$ . Then,

 $\alpha \sqcup \nabla_{AB} = \nabla_{AB}.$ 

Lemma  $cup\_universal\ \{A\ B: eqType\}\ \{alpha: Rel\ A\ B\}: alpha$   $A\ B=A\ B.$ 

Proof.

apply  $inc\_antisym$ .

apply  $inc_-cup$ .

split.

apply  $inc_alpha_universal$ .

apply  $inc\_reft$ .

apply  $cup_r$ .

Qed.

```
Lemma 46 (inc_lower) Let \alpha, \beta : A \rightarrow B. Then,
                                    \alpha = \beta \Leftrightarrow (\forall \gamma : A \to B, \gamma \sqsubseteq \alpha \Leftrightarrow \gamma \sqsubseteq \beta).
Lemma inc\_lower {A \ B : eqType} {alpha \ beta : Rel \ A \ B}:
 alpha = beta \leftrightarrow (\forall \ gamma : Rel \ A \ B, \ gamma)
                                                                        alpha \leftrightarrow qamma
                                                                                                      beta).
Proof.
split; move \Rightarrow H.
move \Rightarrow qamma.
by [rewrite H].
apply inc\_antisym.
rewrite -H.
apply inc\_reft.
rewrite H.
apply inc_refl.
Qed.
  Lemma 47 (inc_upper) Let \alpha, \beta : A \rightarrow B. Then,
                                    \alpha = \beta \Leftrightarrow (\forall \gamma : A \rightarrow B, \alpha \sqsubseteq \gamma \Leftrightarrow \beta \sqsubseteq \gamma).
Lemma inc\_upper \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 alpha = beta \leftrightarrow (\forall \ gamma : Rel \ A \ B, \ alpha
                                                                    gamma \leftrightarrow \mathtt{beta}
                                                                                                   gamma).
Proof.
split; move \Rightarrow H.
move \Rightarrow gamma.
by [rewrite H].
apply inc\_antisym.
rewrite H.
apply inc\_reft.
rewrite -H.
apply inc_refl.
Qed.
             分配法則
4.1.2
```

Lemma 48 (cap\_cup\_distr\_l) Let 
$$\alpha, \beta, \gamma : A \to B$$
. Then,  

$$\alpha \sqcap (\beta \sqcup \gamma) = (\alpha \sqcap \beta) \sqcup (\alpha \sqcap \gamma).$$

```
Lemma cap\_cup\_distr\_l {A \ B : eqType} {alpha \ beta \ qamma : Rel \ A \ B}:
                     qamma) = (alpha)
 alpha
           (beta
                                             beta)
                                                       (alpha
                                                                   qamma).
Proof.
apply inc\_upper.
move \Rightarrow delta.
split; move \Rightarrow H.
rewrite cap\_comm (@cap\_comm\_\_\_ gamma).
apply inc_-cup.
rewrite -inc\_rpc -inc\_rpc.
apply inc_-cup.
rewrite inc\_rpc cap\_comm.
apply H.
rewrite cap\_comm -inc\_rpc.
apply inc\_cup.
rewrite inc\_rpc inc\_rpc.
apply inc_-cup.
rewrite cap_comm (@cap_comm _ _ gamma).
apply H.
Qed.
  Lemma 49 (cap_cup_distr_r) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                (\alpha \sqcup \beta) \sqcap \gamma = (\alpha \sqcap \gamma) \sqcup (\beta \sqcap \gamma).
Lemma cap\_cup\_distr\_r {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
 (alpha
            beta)
                      qamma = (alpha
                                             qamma)
                                                          (beta
                                                                     qamma).
Proof.
rewrite (@cap\_comm\_\_(alpha beta)) (@cap\_comm\_\_alpha) (@cap\_comm\_\_beta).
apply cap\_cup\_distr\_l.
Qed.
  Lemma 50 (cup_cap_distr_l) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                \alpha \sqcup (\beta \sqcap \gamma) = (\alpha \sqcup \beta) \sqcap (\alpha \sqcup \gamma).
Lemma cup\_cap\_distr\_l {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
 alpha
           (beta
                     gamma) = (alpha)
                                            beta)
                                                       (alpha
                                                                   qamma).
Proof.
rewrite cap\_cup\_distr\_l.
rewrite (@cap_comm _ _ (alpha beta)) cap_cup_abs (@cap_comm _ _ (alpha beta)).
rewrite cap\_cup\_distr\_l.
rewrite -cup_assoc (@cap_comm _ _ gamma) cup_cap_abs.
by [rewrite cap\_comm].
```

Qed.

```
Lemma 51 (cup_cap_distr_r) Let \alpha, \beta, \gamma : A \to B. Then, (\alpha \sqcap \beta) \sqcup \gamma = (\alpha \sqcup \gamma) \sqcap (\beta \sqcup \gamma).
```

Lemma  $cup\_cap\_distr\_r$  { $A \ B : eqType$ } { $alpha \ beta \ gamma : Rel \ A \ B$ }: ( $alpha \ beta$ )  $gamma = (alpha \ gamma)$  (beta gamma).

Proof.

rewrite ( $@cup\_comm\_\_(alpha \ beta)$ ) ( $@cup\_comm\_\_alpha$ ) ( $@cup\_comm\_\_beta$ ). apply  $cup\_cap\_distr\_l$ .

Qed.

```
Lemma 52 (cap_cup_unique) Let \alpha, \beta, \gamma : A \to B. Then, \alpha \sqcap \beta = \alpha \sqcap \gamma \land \alpha \sqcup \beta = \alpha \sqcup \gamma \Rightarrow \beta = \gamma.
```

```
Lemma cap\_cup\_unique {A \ B : eqType} {alpha \ beta \ gamma : Rel \ A \ B}:
alpha \ beta = alpha \ gamma \rightarrow alpha \ beta = alpha \ gamma \rightarrow beta = gamma.

Proof.

move \Rightarrow H \ H0.

rewrite -(@cap\_cup\_abs\_\_\_beta \ alpha) \ cup\_comm \ H0.

rewrite cap\_cup\_distr\_l.

rewrite cap\_cup\_distr\_l.

rewrite -cap\_cup\_distr\_r.

rewrite H0 \ cap\_comm \ cup\_comm.

apply cap\_cup\_abs.

Qed.
```

# 4.1.3 原子性

空関係でない  $\alpha: A \rightarrow B$  が、任意の  $\beta: A \rightarrow B$  について

$$\beta \sqsubseteq \alpha \Rightarrow \beta = \phi_{AB} \lor \beta = \alpha$$

を満たすとき,  $\alpha$  は原子的 (atomic) であると言われる.

```
Definition atomic \{A \ B : eqType\}\ (alpha : Rel \ A \ B) := alpha \neq A \ B \land (\forall \ beta : Rel \ A \ B, \ beta = alpha) \rightarrow beta = A \ B \lor beta = alpha).
```

```
\alpha \sqcap \beta = \phi_{AB}.
Lemma atomic\_cap\_empty {A B : eqType} {alpha beta : Rel A B}:
 atomic\ alpha 
ightarrow atomic\ beta 
ightarrow alpha 
eq beta 
ightarrow alpha
                                                                                       A B.
Proof.
move \Rightarrow H H0.
apply or_{-}to_{-}imply.
case (classic (alpha
                            beta =
                                          (A B)); move \Rightarrow H1.
right.
apply H1.
left.
move \Rightarrow H2.
apply H2.
apply inc\_antisym.
apply inc\_def1.
elim H \Rightarrow H3 H4.
case (H4 (alpha
                        beta) (cap_l); move \Rightarrow H5.
apply False_ind.
apply (H1 \ H5).
by [rewrite H5].
apply inc\_def1.
elim H0 \Rightarrow H3 H4.
case (H4 (alpha
                         beta) (cap_r); move \Rightarrow H5.
apply False_ind.
apply (H1 \ H5).
by [rewrite cap\_comm\ H5].
Qed.
  Lemma 54 (atomic_cup) Let \alpha, \beta, \gamma : A \rightarrow B and \alpha is atomic. Then,
                                      \alpha \sqsubseteq \beta \sqcup \gamma \Rightarrow \alpha \sqsubseteq \beta \vee \alpha \sqsubseteq \gamma.
Lemma atomic\_cup \{A \ B : eqType\} \{alpha \ beta \ qamma : Rel \ A \ B\}:
 atomic\ alpha \rightarrow alpha
                                 (beta
                                            qamma) \rightarrow alpha
                                                                      beta \vee alpha
                                                                                           qamma.
Proof.
move \Rightarrow H H0.
apply inc\_def1 in H0.
rewrite cap\_cup\_distr\_l in H0.
elim H \Rightarrow H1 H2.
rewrite H0 in H1.
                                   A B \vee alpha
assert (alpha
                      beta \neq
                                                    qamma \neq
                                                                        A B).
```

**Lemma 53 (atomic\_cap\_empty)** Let  $\alpha, \beta : A \rightarrow B$  are atomic and  $\alpha \neq \beta$ . Then,

```
apply not\_and\_or.
elim \Rightarrow H3 H4.
rewrite H3 H4 in H1.
apply H1.
by [rewrite cup\_empty].
case H3; move \Rightarrow H4.
left.
apply inc\_def1.
case (H2 (alpha
                     beta) (cap_{-}l); move \Rightarrow H5.
apply False_ind.
apply (H4 H5).
by [rewrite H5].
right.
apply inc\_def1.
case (H2 (alpha
                     gamma) (cap_l); move \Rightarrow H5.
apply False_ind.
apply (H4 H5).
by [rewrite H5].
Qed.
```

# 4.2 Heyting 代数に関する補題

```
Lemma 55 (rpc_universal) Let \alpha:A\to B. Then, (\alpha\Rightarrow\alpha)=\nabla_{AB}. Lemma rpc_universal \{A\ B:eqType\} \{alpha:Rel\ A\ B\}:(alpha \gg alpha)=A\ B. Proof. apply inc_ulower. move \Rightarrow gamma. split; move \Rightarrow H. apply inc_ulowersal. Qed. Lemma 56 (rpc_r) Let \alpha,\beta:A\to B. Then,
```

Lemma  $rpc_r \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: (alpha \ beta)$  beta = beta.

 $(\alpha \Rightarrow \beta) \sqcap \beta = \beta.$ 

```
Proof.
assert (beta
                    (alpha \gg beta)).
apply inc\_rpc.
apply cap_l.
apply inc\_def1 in H.
by [rewrite cap\_comm - H].
Qed.
  Lemma 57 (inc_def3) Let \alpha, \beta : A \rightarrow B. Then,
                                      (\alpha \Rightarrow \beta) = \nabla_{AB} \Leftrightarrow \alpha \sqsubseteq \beta.
Lemma inc\_def3 {A B : eqType} {alpha beta : Rel A B}:
 (alpha \gg beta) =
                         A B \leftrightarrow alpha
Proof.
split; move \Rightarrow H.
rewrite -(@rpc_universal _ _ alpha) in H.
assert ((alpha \gg alpha) (alpha \gg beta)).
rewrite H.
apply inc_refl.
apply inc\_rpc in H0.
rewrite rpc_{-}r in H0.
apply H0.
apply inc\_antisym.
apply inc\_alpha\_universal.
rewrite -(@rpc_universal _ _ alpha).
apply inc\_rpc.
rewrite rpc_-r.
apply H.
Qed.
  Lemma 58 (rpc_l) Let \alpha, \beta : A \rightarrow B. Then,
                                        \alpha \sqcap (\alpha \Rightarrow \beta) = \alpha \sqcap \beta.
Lemma rpc_l {A B : eqType} {alpha beta : Rel A B}:
            (alpha \gg beta) = alpha
 alpha
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_cap.
apply inc\_cap in H.
```

```
split.
apply H.
elim H \Rightarrow H0 \ H1.
apply inc\_rpc in H1.
rewrite -(@cap_idem _ _ gamma).
apply (inc_trans _ _ _ (gamma
                                             alpha)).
apply cap\_inc\_compat.
apply inc_refl.
apply H0.
apply H1.
apply inc\_cap.
apply inc\_cap in H.
split.
apply H.
apply inc\_rpc.
apply (inc\_trans \_ \_ \_ gamma).
apply cap_l.
apply H.
Qed.
  Lemma 59 (rpc_inc_compat) Let \alpha, \alpha', \beta, \beta' : A \rightarrow B. Then,
                                \alpha' \sqsubseteq \alpha \land \beta \sqsubseteq \beta' \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha' \Rightarrow \beta').
Lemma rpc\_inc\_compat {A B : eqType} {alpha \ alpha' \ beta \ beta' : Rel \ A \ B}:
 alpha'
              alpha \rightarrow \mathtt{beta}
                                  beta' \rightarrow (alpha \gg beta') (alpha' \gg beta').
Proof.
move \Rightarrow H H0.
apply inc\_rpc.
apply (@inc_trans _ _ _ ((alpha » beta)
                                                         alpha)).
apply (@cap\_inc\_compat\_l\_\_\_\_\_H).
rewrite cap\_comm \ rpc\_l.
apply @inc_trans_{-} - beta).
apply cap_{-}r.
apply H0.
Qed.
  Lemma 60 (rpc_inc_compat_l) Let \alpha, \beta, \beta' : A \rightarrow B. Then,
                                      \beta \sqsubset \beta' \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha \Rightarrow \beta').
```

```
Proof.
move \Rightarrow H.
apply (@rpc\_inc\_compat\_\_\_\_\_ (@inc\_refl\_\_alpha) H).
Qed.
  Lemma 61 (rpc_inc_compat_r) Let \alpha, \alpha', \beta : A \rightarrow B. Then,
                                    \alpha' \sqsubseteq \alpha \Rightarrow (\alpha \Rightarrow \beta) \sqsubseteq (\alpha' \Rightarrow \beta).
Lemma rpc\_inc\_compat\_r {A B : eqType} {alpha \ alpha' \ beta : Rel A B}:
             alpha \rightarrow (alpha \gg beta) (alpha' \gg beta).
Proof.
move \Rightarrow H.
apply (@rpc_inc_compat _ _ _ _ H (@inc_refl _ _ beta)).
Qed.
  Lemma 62 (rpc_universal_alpha) Let \alpha : A \rightarrow B. Then,
                                              \nabla_{AB} \Rightarrow \alpha = \alpha.
Lemma rpc\_universal\_alpha {A B : eqType} {alpha : Rel A B}:
                                                                               A B \gg alpha = alpha.
Proof.
apply inc_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_rpc in H.
rewrite cap\_universal in H.
apply H.
apply inc\_rpc.
rewrite cap_universal.
apply H.
Qed.
  Lemma 63 (rpc_lemma1) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                    (\alpha \Rightarrow \beta) \sqsubseteq ((\alpha \sqcap \gamma) \Rightarrow (\beta \sqcap \gamma)).
Lemma rpc\_lemma1 {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha \gg beta)
                       ((alpha
                                   gamma) \gg (beta
                                                             gamma)).
Proof.
apply inc_rpc.
rewrite - cap_assoc (@cap_comm _ _ alpha).
rewrite rpc_{-}l.
```

```
apply cap\_inc\_compat\_r.
apply cap_{-}r.
Qed.
  Lemma 64 (rpc_lemma2) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                (\alpha \Rightarrow \beta) \sqcap (\alpha \Rightarrow \gamma) = (\alpha \Rightarrow (\beta \sqcap \gamma)).
Lemma rpc\_lemma2 {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha \gg beta)
                      (alpha \gg gamma) = alpha \gg (beta)
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
rewrite inc\_rpc.
apply inc\_cap in H.
apply inc\_cap.
rewrite -inc\_rpc -inc\_rpc.
apply H.
apply inc_-cap.
rewrite inc\_rpc inc\_rpc.
apply inc_-cap.
rewrite -inc_-rpc.
apply H.
Qed.
  Lemma 65 (rpc_lemma3) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                            (\alpha \Rightarrow \beta) \sqcap (\beta \Rightarrow \gamma) \sqsubset ((\alpha \sqcup \beta) \Rightarrow (\beta \sqcap \gamma)).
Lemma rpc\_lemma3 {A B : eqType} {alpha beta gamma : Rel A B}:
 ((alpha \gg beta)
                        (beta » gamma))
                                                 ((alpha
                                                            beta) » (beta
                                                                                   gamma)).
Proof.
apply inc_-rpc.
rewrite cap\_cup\_distr\_l.
rewrite cap_comm -cap_assoc rpc_l.
rewrite (@cap_assoc _ _ _ beta) (@cap_comm _ _ (beta » gamma)) -cap_assoc rpc_r.
rewrite cap_assoc rpc_l.
apply inc\_cup.
split.
apply cap_r.
apply inc\_reft.
Qed.
```

```
Lemma 66 (rpc_lemma4) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                     (\alpha \Rightarrow \beta) \sqcap (\beta \Rightarrow \gamma) \sqsubseteq (\alpha \Rightarrow \gamma).
Lemma rpc\_lemma4 {A B : eqType} {alpha beta gamma : Rel A B}:
 ((alpha » beta) (beta » gamma))
                                                   (alpha \gg qamma).
Proof.
apply (@inc\_trans \_ \_ \_ ((alpha beta) » (beta))
                                                                   qamma))).
apply rpc\_lemma3.
apply rpc\_inc\_compat.
apply cup_{-}l.
apply cap_r.
Qed.
  Lemma 67 (rpc_lemma5) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                      \alpha \Rightarrow (\beta \Rightarrow \gamma) = (\alpha \sqcap \beta) \Rightarrow \gamma.
Lemma rpc\_lemma5 {A B : eqType} {alpha beta gamma : Rel A B}:
 alpha \gg (beta \gg gamma) = (alpha \implies beta) \gg gamma.
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_rpc.
rewrite -cap\_assoc.
rewrite -inc\_rpc -inc\_rpc.
apply H.
rewrite inc\_rpc inc\_rpc.
rewrite cap\_assoc.
apply inc\_rpc.
apply H.
Qed.
  Lemma 68 (rpc_lemma6) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                 \alpha \Rightarrow (\beta \Rightarrow \gamma) \sqsubseteq (\alpha \Rightarrow \beta) \Rightarrow (\alpha \Rightarrow \gamma).
Lemma rpc\_lemma6 {A B : eqType} {alpha beta gamma : Rel A B}:
 (alpha \gg (beta \gg gamma)) ((alpha \gg beta) \gg (alpha \gg gamma)).
Proof.
rewrite inc\_rpc inc\_rpc.
rewrite cap_assoc (@cap_comm _ _ alpha).
```

```
rewrite rpc_-l.
rewrite -cap_assoc (@cap_comm _ _ alpha).
rewrite rpc_{-}l.
rewrite cap\_assoc (@cap\_comm _ _ _ beta).
rewrite rpc_{-}l.
rewrite -cap_-assoc.
apply cap_r.
Qed.
  Lemma 69 (rpc_lemma7) Let \alpha, \beta, \gamma, \delta : A \rightarrow B and \beta \sqsubseteq \alpha \sqsubseteq \gamma. Then,
             (\alpha \sqcap \delta = \beta) \land (\alpha \sqcup \delta = \gamma) \Leftrightarrow (\gamma \sqsubseteq \alpha \sqcup (\alpha \Rightarrow \beta)) \land (\delta = \gamma \sqcap (\alpha \Rightarrow \beta)).
Lemma rpc\_lemma? \{A \ B : eqType\} \{alpha \ beta \ gamma \ delta : Rel \ A \ B\}:
beta
          alpha \rightarrow alpha
                                qamma \rightarrow (alpha)
                                                        delta = beta \land alpha
                                                                                        delta = qamma
                              (alpha \gg beta)) \land delta = gamma
                                                                          (alpha \gg beta)).
 \leftrightarrow qamma
                 (alpha
Proof.
move \Rightarrow H H0.
split; elim; move \Rightarrow H1 H2; split.
rewrite -H2.
apply cup\_inc\_compat\_l.
apply inc_-rpc.
rewrite cap_comm H1.
apply inc_refl.
rewrite -H2.
rewrite cap\_cup\_distr\_r\ rpc\_l.
assert (delta
                     (alpha \gg beta).
apply inc\_rpc.
rewrite cap_comm H1.
apply inc\_reft.
apply inc\_def1 in H3.
rewrite -H3 -H1.
rewrite -cap_assoc cap_idem.
by [rewrite cap_comm cup_comm cup_cap_abs].
rewrite H2.
rewrite (@cap_comm _ _ gamma) -cap_assoc rpc_l.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ (beta gamma)).
apply cap\_inc\_compat\_r.
apply cap_r.
apply cap_{-}l.
move: (@inc\_trans \_ \_ \_ \_ H H0) \Rightarrow H3.
apply inc\_def1 in H.
```

```
apply inc\_def1 in H3.

rewrite cap\_comm in H.

rewrite -H -H3.

apply inc\_reft.

rewrite H2.

rewrite cup\_cap\_distr\_t.

apply inc\_def2 in H0.

rewrite -H0.

apply inc\_def1 in H1.

by [rewrite -H1].

Qed.
```

# 4.3 補関係に関する補題

# Lemma 70 (complement\_universal)

$$\nabla_{AB}^{-} = \phi_{AB}$$
.

Lemma 71 (complement\_alpha\_universal) Let  $\alpha : A \rightarrow B$ . Then,

$$\alpha^- = \nabla_{AB} \Leftrightarrow \alpha = \phi_{AB}.$$

```
Lemma complement\_alpha\_universal\ \{A\ B: eqType\}\ \{alpha: Rel\ A\ B\}:
 alpha \hat{} =
               A B \leftrightarrow alpha =
                                  A B.
Proof.
split; move \Rightarrow H.
apply inc\_antisym.
rewrite -(@cap_universal _ _ alpha) cap_comm.
apply inc_rpc.
rewrite -H.
apply inc\_reft.
apply inc\_empty\_alpha.
apply inc\_antisym.
apply inc\_alpha\_universal.
apply inc\_rpc.
rewrite cap_comm cap_universal.
rewrite H.
```

apply  $inc\_reft$ .

Qed.

# Lemma 72 (complement\_empty)

$$\phi_{AB}^{-} = \nabla_{AB}$$
.

Lemma  $complement\_empty \{A \ B : eqType\}: A \ B \hat{\ } = A \ B.$ 

Proof.

by [apply complement\_alpha\_universal].

Qed.

# Lemma 73 (complement\_invol\_inc) Let $\alpha : A \rightarrow B$ . Then,

$$\alpha \sqsubseteq (\alpha^{-})^{-}$$
.

Lemma  $complement\_invol\_inc$  { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }: alpha (alpha ^) ^. Proof.

apply  $inc\_rpc$ .

rewrite  $cap\_comm$ .

apply  $inc\_rpc$ .

apply  $inc\_reft$ .

Qed.

## Lemma 74 (cap\_complement\_empty) Let $\alpha : A \rightarrow B$ . Then,

$$\alpha \sqcap \alpha^- = \phi_{AB}$$
.

Lemma  $cap\_complement\_empty$  {A B : eqType} {alpha : Rel A B}:

alpha alpha  $^{\circ} = A B.$ 

Proof.

apply  $inc\_antisym$ .

rewrite  $cap\_comm$ .

apply  $inc\_rpc$ .

apply  $inc\_reft$ .

apply  $inc\_empty\_alpha$ .

Qed.

# Lemma 75 (complement\_invol) Let $\alpha : A \rightarrow B$ . Then,

$$(\alpha^{-})^{-}=\alpha$$
.

```
Proof.

rewrite -(@cap_universal _ _ ((alpha ^) ^)).

rewrite -(@complement_classic _ _ alpha).

rewrite cap_cup_distr_l.

rewrite (@cap_comm _ _ _ (alpha ^)) cap_complement_empty.

rewrite cup_empty cap_comm.

apply Logic.eq_sym.

apply inc_def1.

apply complement_invol_inc.

Qed.
```

**Lemma 76 (complement\_move)** Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$\alpha = \beta^- \Leftrightarrow \alpha^- = \beta$$
.

```
Lemma complement\_move \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: alpha = beta ^ \leftrightarrow alpha ^ = beta.

Proof.

split; move \Rightarrow H.

by [rewrite H \ complement\_invol].

by [rewrite -H \ complement\_invol].

Qed.
```

Lemma 77 (contraposition) Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$(\alpha \Rightarrow \beta) = (\beta^- \Rightarrow \alpha^-).$$

```
Lemma contraposition {A B : eqType} {alpha beta : Rel A B}: alpha » beta = beta ^ » alpha ^.

Proof.

apply inc_antisym.

apply inc_rpc.

apply rpc_lemma4.

replace (alpha » beta) with ((alpha ^) ^ » (beta ^) ^).

apply inc_rpc.

apply rpc_lemma4.

by [rewrite complement_invol complement_invol].

Qed.
```

Lemma 78 (de\_morgan1) Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$(\alpha \sqcup \beta)^- = \alpha^- \sqcap \beta^-.$$

```
Lemma de\_morgan1 {A B : eqType} {alpha beta : Rel A B}:
           beta) \hat{} = alpha \hat{}
 (alpha
                                beta ^.
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_cap.
rewrite inc\_rpc inc\_rpc.
apply inc\_cup.
rewrite -cap\_cup\_distr\_l.
apply inc\_rpc.
apply H.
apply inc\_rpc.
rewrite cap\_cup\_distr\_l.
apply inc\_cup.
rewrite -inc\_rpc -inc\_rpc.
apply inc\_cap.
apply H.
Qed.
 Lemma 79 (de_morgan2) Let \alpha, \beta : A \rightarrow B. Then,
                                    (\alpha \sqcap \beta)^- = \alpha^- \sqcup \beta^-.
Lemma de\_morgan2 {A B : eqType} {alpha beta : Rel A B}:
           beta) \hat{} = alpha \hat{}
 (alpha
                                beta ^.
by [rewrite -complement_move de_morgan1 complement_invol complement_invol].
Qed.
 Lemma 80 (cup_to_rpc) Let \alpha, \beta : A \rightarrow B. Then,
                                     \alpha^- \sqcup \beta = (\alpha \Rightarrow \beta).
beta = alpha \gg beta.
Proof.
apply inc\_antisym.
apply inc\_rpc.
rewrite cap\_cup\_distr\_r cap\_comm.
rewrite cap_complement_empty cup_comm cup_empty.
apply cap_{-}l.
```

rewrite -(@cap\_universal \_ \_ (alpha » beta)) cap\_comm.

```
rewrite -(@complement\_classic\_\_alpha). rewrite cap\_cup\_distr\_r cup\_comm. apply cup\_inc\_compat. apply cap\_l. rewrite rpc\_l. apply cap\_r. Qed.
```

Lemma 81 (beta\_contradiction) Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$(\alpha \Rightarrow \beta) \sqcap (\alpha \Rightarrow \beta^{-}) = \alpha^{-}.$$

```
Lemma beta_contradiction \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}: (alpha \ beta) (alpha \ beta \ ) = alpha \ .
Proof.

rewrite -cup\_to\_rpc -cup\_to\_rpc.

rewrite -cup\_cap\_distr\_l.

by [rewrite \ cap\_complement\_empty \ cup\_empty].
Qed.
```

# 4.4 Bool 代数に関する補題

```
Lemma 82 (bool_lemma1) Let \alpha, \beta : A \rightarrow B. Then,
```

$$\alpha \sqsubseteq \beta \Leftrightarrow \nabla_{AB} = \alpha^- \sqcup \beta.$$

```
Lemma bool_lemma1 {A B : eqType} {alpha beta : Rel A B}: alpha beta \leftrightarrow A B = alpha beta.

Proof.

split; move \Rightarrow H.

apply inc\_antisym.

rewrite -(@complement\_classic _ _ alpha) cup\_comm.

apply cup\_inc\_compat\_l.

apply H.

apply inc\_alpha\_universal.

apply inc\_alpha\_universal.

apply inc\_def3.

rewrite H.

apply (Logic.eq\_sym cup\_to\_rpc).

Qed.
```

```
Lemma 83 (bool_lemma2) Let \alpha, \beta : A \rightarrow B. Then,
                                    \alpha \sqsubseteq \beta \Leftrightarrow \alpha \sqcap \beta^- = \phi_{AB}.
Lemma bool_lemma2 {A B : eqType} {alpha beta : Rel A B}:
           beta \leftrightarrow alpha
                              beta ^ =
Proof.
split; move \Rightarrow H.
rewrite -(@cap_universal _ _ (alpha
                                            beta ^)).
apply bool\_lemma1 in H.
rewrite H.
rewrite cap\_cup\_distr\_l.
rewrite (@cap_comm _ _ alpha) cap_assoc cap_complement_empty cap_empty.
rewrite cap_comm -cap_assoc cap_complement_empty cap_comm cap_empty.
by [rewrite cup\_empty].
rewrite -(@cap_universal _ alpha).
rewrite -(@complement_classic _ _ beta).
rewrite cap\_cup\_distr\_l.
rewrite H cup\_empty.
apply cap_r.
Qed.
  Lemma 84 (bool_lemma3) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                    \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \alpha \sqcap \beta^- \sqsubseteq \gamma.
Lemma bool_lemma3 {A B : eqType} {alpha beta gamma : Rel A B}:
           (beta
                     gamma) \leftrightarrow (alpha)
 alpha
                                             beta ^)
                                                          qamma.
Proof.
split; move \Rightarrow H.
apply (@inc_trans _ _ _ ((beta
                                                    beta ^)).
                                       gamma)
apply cap\_inc\_compat\_r.
apply H.
rewrite cap\_cup\_distr\_r.
rewrite cap_complement_empty cup_comm cup_empty.
apply cap_{-}l.
apply (@inc_trans _ _ _ (beta
                                      (alpha
                                                 beta ^))).
rewrite cup\_cap\_distr\_l.
rewrite complement_classic cap_universal.
apply cup_r.
apply cup\_inc\_compat\_l.
apply H.
Qed.
```

```
Lemma 85 (bool_lemma4) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                        \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \beta^- \sqsubseteq \alpha^- \sqcup \gamma.
Lemma bool\_lemma4 {A B : eqType} {alpha beta gamma : Rel A B}:
                      gamma) \leftrightarrow beta \hat{} (alpha \hat{} gamma).
            (beta
Proof.
rewrite bool_lemma3.
rewrite cap\_comm.
apply iff_-sym.
                         alpha) with (beta ^ (alpha ^) ^).
replace (beta ^
apply bool_lemma3.
by [rewrite complement_invol].
Qed.
  Lemma 86 (bool_lemma5) Let \alpha, \beta, \gamma : A \rightarrow B. Then,
                                    \alpha \sqsubseteq \beta \sqcup \gamma \Leftrightarrow \nabla_{AB} = \alpha^- \sqcup \beta \sqcup \gamma.
Lemma bool_lemma5 {A B : eqType} {alpha beta gamma : Rel A B}:
            (beta
                        gamma) \leftrightarrow A B = (alpha \hat{ } beta)
 alpha
Proof.
rewrite bool_lemma1.
by [rewrite cup\_assoc].
Qed.
```

# Chapter 5

# Library Relation\_Properties

```
Require Import Basic\_Notations.
Require Import Basic\_Lemmas.
Require Import Logic.FunctionalExtensionality.
Require Import Logic.Classical\_Prop.
```

# 5.1 関係計算の基本的な性質

```
Lemma 87 (RelAB_unique) \phi_{AB} = \nabla_{AB} \Leftrightarrow \forall \alpha, \beta : A \to B, \alpha = \beta. Lemma RelAB_unique {AB: eqType}:
```

```
Lemma RelAB\_unique \{A B : eqType\}:
              A B \leftrightarrow (\forall alpha beta : Rel A B, alpha = beta).
   A B =
Proof.
split; move \Rightarrow H.
move \Rightarrow alpha beta.
replace beta with (
                         A B).
apply inc\_antisym.
rewrite H.
apply inc\_alpha\_universal.
apply inc\_empty\_alpha.
apply inc\_antisym.
apply inc\_empty\_alpha.
rewrite H.
apply inc\_alpha\_universal.
apply H.
Qed.
```

apply  $(H0 \ a \ b)$ .

Qed.

# Lemma 88 (either\_empty) $\phi_{AB} = \nabla_{AB} \Leftrightarrow A = \emptyset \lor B = \emptyset.$ Lemma either\_empty $\{A \ B : eqType\}$ : $A \ B =$ $A \ B \leftrightarrow (A \rightarrow False) \lor (B \rightarrow False).$ Proof. rewrite $RelAB\_unique$ . $split; move \Rightarrow H.$ case $(classic (\exists \_: A, True)).$ $elim \Rightarrow a H0.$ right. $move \Rightarrow b$ . remember (fun ( $\_: A$ ) ( $\_: B$ ) $\Rightarrow True$ ) as T. remember (fun ( $\_: A$ ) ( $\_: B$ ) $\Rightarrow$ False) as F. move: $(H \ T \ F) \Rightarrow H1$ . assert $(T \ a \ b = F \ a \ b)$ . by [rewrite H1]. rewrite HeqT HeqF in H2. rewrite -H2. apply I. move $\Rightarrow H0$ . left. $move \Rightarrow a$ . apply H0. $\exists a.$ apply I. move $\Rightarrow alpha$ beta. assert $(A \rightarrow B \rightarrow False)$ . move $\Rightarrow a \ b$ . case H; move $\Rightarrow H\theta$ . apply $(H0 \ a)$ . apply $(H0\ b)$ . apply functional\_extensionality. $move \Rightarrow a$ . $apply functional\_extensionality.$ move $\Rightarrow b$ . apply False\_ind.

```
Lemma 89 (unit_empty_not_universal)
```

```
\phi_{II} \neq \nabla_{II}.
```

```
Lemma unit\_empty\_not\_universal: i i \neq i i. Proof.

move \Rightarrow H.

apply either\_empty in H.

case H; move \Rightarrow H0.

apply (H0 \ tt).

apply (H0 \ tt).

Qed.
```

# Lemma 90 (unit\_empty\_or\_universal) Let $\alpha: I \rightarrow I$ . Then,

$$\alpha = \phi_{II} \vee \alpha = \nabla_{II}$$
.

```
Lemma unit\_empty\_or\_universal \{alpha : Rel \ i \ i\}: alpha = i \ i \lor alpha =
                                                                                           i i.
assert (\forall beta : Rel\ i\ i, beta = (fun (_ _ : i) \Rightarrow True) \lor beta = (fun (_ _ : i) \Rightarrow False)).
move \Rightarrow beta.
case (classic (beta tt tt)); move \Rightarrow H.
left.
apply functional_extensionality.
\verb"induction" x.
apply functional_extensionality.
induction x.
apply prop_extensionality_ok.
split; move \Rightarrow H0.
apply I.
apply H.
right.
apply functional_extensionality.
induction x.
apply functional_extensionality.
induction x.
apply prop_extensionality_ok.
split.
apply H.
apply False_ind.
assert ((fun \_ : i \Rightarrow True) \neq (fun \_ : i \Rightarrow False)).
move \Rightarrow H0.
remember (fun \_ : i \Rightarrow True) as T.
```

```
remember (fun \_ : i \Rightarrow False) as F.
assert (T tt tt = F tt tt).
by [rewrite H\theta].
rewrite HeqT HeqF in H1.
rewrite -H1.
apply I.
case (H(i)); move \Rightarrow H1.
case (H(i)); move \Rightarrow H2.
apply False_ind.
apply unit_empty_not_universal.
by [rewrite H1 H2].
case (H \ alpha); move \Rightarrow H3.
left.
by [rewrite H3 H1].
right.
by [rewrite H3 H2].
case (H(i)); move \Rightarrow H2.
case (H \ alpha); move \Rightarrow H3.
right.
by [rewrite H3 H2].
left.
by [rewrite H3 H1].
apply False_ind.
apply unit_empty_not_universal.
by [rewrite H1 H2].
Qed.
```

## Lemma 91 (unit\_identity\_is\_universal)

```
id_I = \nabla_{II}.
```

```
Lemma unit\_identity\_is\_universal: Id\ i = i\ i.

Proof.

case (@unit\_empty\_or\_universal\ (Id\ i)); move \Rightarrow H.

apply False\_ind.

assert (Id\ i\ (i\ i\ \#\ i\ i)).

rewrite H.

apply inc\_empty\_alpha.

apply inc\_empty\_alpha.

apply inc\_residual\ in\ H0.

rewrite inv\_invol\ comp\_id\_r\ in\ H0.

apply unit\_empty\_not\_universal.

apply inc\_antisym.

apply inc\_empty\_alpha.
```

```
apply H\theta.
```

Qed.

# Lemma 92 (unit\_identity\_not\_empty)

 $id_I \neq \phi_{II}$ .

Lemma  $unit\_identity\_not\_empty: Id \ i \neq i$ . Proof.

move  $\Rightarrow H$ .

apply unit\_empty\_not\_universal.

rewrite -H.

apply  $unit\_identity\_is\_universal$ .

Qed.

**Lemma 93 (cupP\_False)** Let  $f:(C \rightarrow D) \rightarrow (A \rightarrow B)$  and  $P(\alpha) :=$  "False". Then,

$$\sqcup_{P(\alpha)} f(\alpha) = \phi_{AB}.$$

Proof.

apply  $inc\_antisym$ .

apply  $inc\_cupP$ .

 $move \Rightarrow beta$ .

apply False\_ind.

apply  $inc\_empty\_alpha$ .

Qed.

**Lemma 94 (capP\_False)** Let  $f:(C \rightarrow D) \rightarrow (A \rightarrow B)$  and  $P(\alpha) :=$  "False". Then,

$$\sqcap_{P(\alpha)} f(\alpha) = \nabla_{AB}.$$

Lemma capP-False  $\{A \ B \ C \ D : eqType\} \ \{f : Rel \ C \ D \rightarrow Rel \ A \ B\}:$ \_ $\{\mathbf{fun} \ \_: Rel \ C \ D \Rightarrow False\} \ f = A \ B.$ 

Proof.

apply  $inc\_antisym$ .

apply  $inc_alpha_universal$ .

apply  $inc\_capP$ .

 $move \Rightarrow beta.$ 

apply False\_ind.

Qed.

```
(\forall \alpha: C \to D, P(\alpha) \Rightarrow f(\alpha) = g(\alpha)) \Rightarrow \sqcup_{P(\alpha)} f(\alpha) = \sqcup_{P(\alpha)} g(\alpha).
Lemma cupP_-eq {A B C D : eqType}
 \{f g : Rel \ C \ D \rightarrow Rel \ A \ B\} \ \{P : Rel \ C \ D \rightarrow Prop\}:
 (\forall alpha : Rel \ C \ D, P \ alpha \rightarrow f \ alpha = g \ alpha) \rightarrow -\{P\} \ f = -\{P\} \ g.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc\_cupP.
move \Rightarrow beta H0.
rewrite (H - H0).
move : beta H0.
apply inc\_cupP.
apply inc_refl.
apply inc\_cupP.
move \Rightarrow beta H\theta.
rewrite -(H - H\theta).
move : beta H0.
apply inc\_cupP.
apply inc\_reft.
Qed.
  Lemma 96 (capP_eq) Let f, g: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                    (\forall \alpha : C \to D, P(\alpha) \Rightarrow f(\alpha) = g(\alpha)) \Rightarrow \sqcap_{P(\alpha)} f(\alpha) = \sqcap_{P(\alpha)} g(\alpha).
Lemma capP_{-}eq \{A \ B \ C \ D : eqType\}
 \{f g : Rel \ C \ D \rightarrow Rel \ A \ B\} \ \{P : Rel \ C \ D \rightarrow Prop\}:
 (\forall alpha : Rel \ C \ D, P \ alpha \rightarrow f \ alpha = g \ alpha) \rightarrow \_\{P\} \ f = \_\{P\} \ q.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc\_capP.
move \Rightarrow beta H\theta.
rewrite -(H - H0).
move : beta H0.
apply inc\_capP.
apply inc_refl.
```

**Lemma 95 (cupP\_eq)** Let  $f, g: (C \rightarrow D) \rightarrow (A \rightarrow B)$  and P: predicate. Then,

apply  $inc\_capP$ . move  $\Rightarrow$  beta  $H\theta$ .

```
rewrite (H - H\theta).
move: beta H0.
apply inc\_capP.
apply inc\_reft.
Qed.
  Lemma 97 (cap_cupP_distr_l) Let \alpha: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                   \alpha \sqcap (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \sqcap f(\beta)).
Lemma cap\_cupP\_distr\_l \{A \ B \ C \ D : eqType\}
 \{alpha: Rel\ A\ B\}\ \{f: Rel\ C\ D \rightarrow Rel\ A\ B\}\ \{P: Rel\ C\ D \rightarrow Prop\}:
 alpha
            ( _{P} f) = _{P} (fun beta : Rel \ C \ D \Rightarrow alpha f beta).
Proof.
apply inc\_upper.
move \Rightarrow qamma.
split; move \Rightarrow H.
apply inc\_cupP.
move \Rightarrow beta H\theta.
apply (@inc_trans _ _ _ (alpha
                                            _{-}\{P\}\ f)).
apply cap\_inc\_compat\_l.
move: H0.
apply inc\_cupP.
apply inc\_reft.
apply H.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow (alpha f beta)
                                                                           qamma).
apply inc\_cupP.
apply H.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow f beta (alpha \gg gamma)).
move \Rightarrow beta H1.
rewrite inc\_rpc\ cap\_comm.
apply (H0 - H1).
rewrite cap\_comm -inc\_rpc.
apply inc\_cupP.
apply H1.
Qed.
  Lemma 98 (cap_cupP_distr_r) Let \beta: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                   (\sqcup_{P(\alpha)} f(\alpha)) \sqcap \beta = \sqcup_{P(\alpha)} (f(\alpha) \sqcap \beta).
```

```
\{ beta : Rel \ A \ B \} \{ f : Rel \ C \ D \rightarrow Rel \ A \ B \} \{ P : Rel \ C \ D \rightarrow Prop \} :
                              \{P\} (fun alpha : Rel C D \Rightarrow f alpha
 (-\{P\}f)
                beta =
Proof.
rewrite cap\_comm.
replace (fun alpha: Rel\ C\ D \Rightarrow f\ alpha beta) with (fun alpha: Rel\ C\ D \Rightarrow beta
f alpha).
apply cap\_cupP\_distr\_l.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite cap\_comm].
Qed.
  Lemma 99 (cup_capP_distr_l) Let \alpha: A \rightarrow B, f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P:
  predicate. Then,
                                 \alpha \sqcup (\sqcap_{P(\beta)} f(\beta)) = \sqcap_{P(\beta)} (\alpha \sqcup f(\beta)).
Lemma cup\_capP\_distr\_l \{A \ B \ C \ D : eqType\}
 \{alpha: Rel\ A\ B\}\ \{f: Rel\ C\ D \rightarrow Rel\ A\ B\}\ \{P: Rel\ C\ D \rightarrow Prop\}:
            ( _{P} f) = _{P} (fun beta : Rel C D \Rightarrow alpha f beta).
 alpha
Proof.
apply inc\_lower.
move \Rightarrow qamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply (@inc\_trans \_ \_ \_ (alpha \_ \{P\} f)).
apply H.
apply cup\_inc\_compat\_l.
move: H0.
apply inc\_capP.
apply inc\_reft.
rewrite bool_lemma3.
assert (\forall beta : Rel\ C\ D,\ P\ beta \rightarrow gamma (alpha
                                                                       f beta)).
apply inc\_capP.
apply H.
apply inc\_capP.
move \Rightarrow beta H1.
rewrite -bool_lemma3.
apply (H0 - H1).
Qed.
```

```
predicate. Then,
                                  (\sqcap_{P(\alpha)} f(\alpha)) \sqcup \beta = \sqcap_{P(\alpha)} (f(\alpha) \sqcup \beta).
Lemma cup\_capP\_distr\_r \{A \ B \ C \ D : eqType\}
 \{ beta : Rel \ A \ B \} \{ f : Rel \ C \ D \rightarrow Rel \ A \ B \} \{ P : Rel \ C \ D \rightarrow Prop \} :
                 beta = _{-}\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha
Proof.
rewrite cup\_comm.
replace (fun alpha : Rel \ C \ D \Rightarrow f \ alpha
                                                      beta) with (fun alpha : Rel \ C \ D \Rightarrow beta
f alpha).
apply cup\_capP\_distr\_l.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite cup\_comm].
Qed.
  Lemma 101 (de_morgan3) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                      (\sqcup_{P(\alpha)} f(\alpha))^- = (\sqcap_{P(\alpha)} f(\alpha)^-).
Lemma de\_morgan3
 \{A \ B \ C \ D : eqType\} \{f : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Prop\}:
 ( -\{P\} f)^{\circ} = -\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha^{\circ}).
Proof.
apply inc\_lower.
move \Rightarrow gamma.
rewrite inc\_capP.
split; move \Rightarrow H.
move \Rightarrow beta H0.
rewrite bool_lemma1 -de_morgan2 complement_move complement_universal.
apply bool\_lemma2 in H.
apply inc\_antisym.
apply inc\_empty\_alpha.
rewrite - H complement_invol.
apply cap\_inc\_compat\_l.
move: H0.
apply inc\_cupP.
apply inc_refl.
rewrite bool_lemma2 complement_invol.
rewrite cap\_cupP\_distr\_l.
apply inc\_antisym.
```

**Lemma 100 (cup\_capP\_distr\_r)** Let  $\beta: A \rightarrow B$ ,  $f: (C \rightarrow D) \rightarrow (A \rightarrow B)$  and P:

```
apply inc\_cupP.
move \Rightarrow beta H\theta.
rewrite -inc_-rpc.
apply (H - H\theta).
apply inc\_empty\_alpha.
Qed.
  Lemma 102 (de_morgan4) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                       (\sqcap_{P(\alpha)} f(\alpha))^- = (\sqcup_{P(\alpha)} f(\alpha)^-).
Lemma de\_morgan4
 \{A \ B \ C \ D : eqType\} \{f : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Prop\}:
 ( -\{P\} f)^{=} -\{P\} (fun \ alpha : Rel \ C \ D \Rightarrow f \ alpha^{=}).
Proof.
rewrite -complement_move de_morgan3.
replace (fun alpha : Rel \ C \ D \Rightarrow (f \ alpha \ \hat{}) \ \hat{}) with f.
by ||.
{\tt apply} \ functional\_extensionality.
move \Rightarrow x.
by [rewrite complement_invol].
Qed.
  Lemma 103 (cup_to_cupP) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                        f(\alpha) \sqcup f(\beta) = \sqcup_{\gamma = \alpha \lor \gamma = \beta} f(\gamma).
Lemma cup\_to\_cupP
 \{A \ B \ C \ D : eqType\} \{alpha \ \mathsf{beta} : Rel \ C \ D\} \{f : Rel \ C \ D \to Rel \ A \ B\}:
                f \text{ beta}) = -\{\text{fun } gamma : Rel \ C \ D \Rightarrow gamma = alpha \lor gamma = \text{beta}\}
 (f alpha
f.
Proof.
apply inc\_upper.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_cupP.
apply inc\_cup in H.
move \Rightarrow gamma\ H0.
case H0 \Rightarrow H1.
rewrite H1.
apply H.
rewrite H1.
apply H.
```

```
apply inc\_cup.
assert (\forall gamma : Rel \ C \ D, gamma = alpha \lor gamma = beta \rightarrow f \ gamma
apply inc\_cupP.
apply H.
split.
apply (H0 \ alpha).
by [left].
apply (H0 \text{ beta}).
by [right].
Qed.
  Lemma 104 (cap_to_capP) Let f:(C \rightarrow D) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                     f(\alpha) \sqcap f(\beta) = \sqcap_{\gamma = \alpha \lor \gamma = \beta} f(\gamma).
Lemma cap\_to\_capP
 \{A \ B \ C \ D : eqType\} \{alpha \ \mathsf{beta} : Rel \ C \ D\} \{f : Rel \ C \ D \to Rel \ A \ B\}:
               f \text{ beta}) = -\{\text{fun } gamma : Rel \ C \ D \Rightarrow gamma = alpha \lor gamma = \text{beta}\}
f.
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inc\_capP.
apply inc\_cap in H.
move \Rightarrow gamma H0.
case H0 \Rightarrow H1.
rewrite H1.
apply H.
rewrite H1.
apply H.
apply inc\_cap.
assert (\forall gamma : Rel \ C \ D, gamma = alpha \lor gamma = beta \rightarrow delta f gamma).
apply inc\_capP.
apply H.
split.
apply (H0 \ alpha).
by [left].
apply (H0 \text{ beta}).
by [right].
Qed.
```

# 5.2 comp\_inc\_compat と派生補題

```
Lemma 105 (comp_inc_compat_ab_ab') Let \alpha : A \to B and \beta, \beta' : B \to C. Then,
                                          \beta \sqsubseteq \beta' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha \cdot \beta'.
Lemma comp\_inc\_compat\_ab\_ab'
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ beta' : Rel \ B \ C\}:
            beta' \rightarrow (alpha \cdot beta') (alpha · beta').
Proof.
move \Rightarrow H.
replace (alpha \cdot beta) with ((alpha \#) \# \cdot beta).
apply inc\_residual.
apply (@inc\_trans \_ \_ \_ beta').
apply H.
apply inc\_residual.
rewrite inv_{-}invol.
apply inc\_reft.
by [rewrite inv\_invol].
Qed.
  Lemma 106 (comp_inc_compat_ab_a'b) Let \alpha, \alpha' : A \to B and \beta : B \to C. Then,
                                          \alpha \sqsubseteq \alpha' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha' \cdot \beta.
Lemma comp_inc_compat_ab_a'b
 \{A \ B \ C : eqType\} \ \{alpha \ alpha' : Rel \ A \ B\} \ \{beta : Rel \ B \ C\}:
             alpha' \rightarrow (alpha \cdot beta) \quad (alpha' \cdot beta).
 alpha
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ (alpha • beta)).
rewrite -(@inv_invol _ _ (alpha' • beta)).
apply inc_{-}inv.
rewrite comp_{-}inv \ comp_{-}inv.
apply comp\_inc\_compat\_ab\_ab'.
apply inc_-inv.
apply H.
Qed.
```

```
Lemma 107 (comp_inc_compat) Let \alpha, \alpha' : A \to B and \beta, \beta' : B \to C. Then,
                                     \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \cdot \beta \sqsubseteq \alpha' \cdot \beta'.
Lemma comp\_inc\_compat
 \{A \ B \ C : eqType\} \ \{alpha \ alpha' : Rel \ A \ B\} \ \{beta \ beta' : Rel \ B \ C\}:
                                     beta' \rightarrow (alpha \cdot beta') \quad (alpha' \cdot beta').
 alpha
             alpha' \rightarrow \mathtt{beta}
Proof.
move \Rightarrow H H0.
apply (@inc_trans _ _ _ (alpha' • beta)).
apply (@comp\_inc\_compat\_ab\_a'b\_\_\_\_\_H).
apply (@comp\_inc\_compat\_ab\_ab'\_\_\_\__H0).
Qed.
  Lemma 108 (comp_inc_compat_ab_a) Let \alpha : A \to B and \beta : B \to B. Then,
                                            \beta \sqsubseteq id_B \Rightarrow \alpha \cdot \beta \sqsubseteq \beta.
Lemma comp\_inc\_compat\_ab\_a {A \ B : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ B}:
            Id B \rightarrow (alpha \cdot beta)
                                                alpha.
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_ab'\_\_\_alpha\_\_H) \Rightarrow H0.
rewrite comp_{-}id_{-}r in H0.
apply H0.
Qed.
  Lemma 109 (comp_inc_compat_a_ab) Let \alpha : A \rightarrow B and \beta : B \rightarrow B. Then,
                                            id_B \sqsubseteq \beta \Rightarrow \beta \sqsubseteq \alpha \cdot \beta.
Lemma comp_inc_compat_a_ab {A B : eqType} {alpha : Rel A B} {beta : Rel B B}:
 Id\ B
            beta \rightarrow alpha
                                   (alpha \cdot beta).
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_ab'\_\_\_alpha\_\_H) \Rightarrow H0.
rewrite comp_{-}id_{-}r in H0.
apply H0.
Qed.
```

```
Lemma 110 (comp_inc_compat_ab_b) Let \alpha : A \rightarrow A and \beta : A \rightarrow B. Then,
                                             \alpha \sqsubseteq id_A \Rightarrow \alpha \cdot \beta \sqsubseteq \beta.
Lemma comp\_inc\_compat\_ab\_b {A \ B : eqType} {alpha : Rel \ A \ A} {beta : Rel \ A \ B}:
             Id A \rightarrow (alpha \cdot beta)
                                                  beta.
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_a"b\_\_\_\_\_ beta H) \Rightarrow H0.
rewrite comp_{-}id_{-}l in H0.
apply H0.
Qed.
  Lemma 111 (comp_inc_compat_b_ab) Let \alpha : A \rightarrow A and \beta : A \rightarrow B. Then,
                                             id_A \sqsubseteq \alpha \Rightarrow \beta \sqsubseteq \alpha \cdot \beta.
Lemma comp\_inc\_compat\_b\_ab {A \ B : eqType} {alpha : Rel \ A \ A} {beta : Rel \ A \ B}:
             alpha \rightarrow \mathtt{beta}
 Id A
                                   (alpha \cdot beta).
Proof.
move \Rightarrow H.
move: (@comp\_inc\_compat\_ab\_a"b\_\_\_\_\_ beta H) \Rightarrow H0.
rewrite comp_{-}id_{-}l in H0.
apply H0.
Qed.
          逆関係に関する補題
5.3
  Lemma 112 (inv_move) Let \alpha : A \rightarrow B and \beta : B \rightarrow A. Then,
                                                \alpha = \beta^{\sharp} \Leftrightarrow \alpha^{\sharp} = \beta.
Lemma inv\_move \{A \ B : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ A\}:
 alpha = \mathtt{beta} \ \# \leftrightarrow alpha \ \# = \mathtt{beta}.
Proof.
split; move \Rightarrow H.
by [rewrite H inv_invol].
```

by [rewrite -H inv\_invol].

Qed.

```
Lemma 113 (comp_inv_inv) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                            \alpha \cdot \beta = (\beta^{\sharp} \cdot \alpha^{\sharp})^{\sharp}.
alpha • beta = (beta # • alpha #) #.
Proof.
apply inv_move.
apply comp_{-}inv.
Qed.
  Lemma 114 (inv_inc_move) Let \alpha : A \rightarrow B and \beta : B \rightarrow A. Then,
                                            \alpha \sqsubseteq \beta^{\sharp} \Leftrightarrow \alpha^{\sharp} \sqsubseteq \beta.
Lemma inv\_inc\_move \{A \ B : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ A\}:
            beta \# \leftrightarrow alpha \#
                                        beta.
Proof.
split; move \Rightarrow H.
rewrite -(@inv_invol_{-} beta).
apply inc_{-}inv.
apply H.
rewrite -(@inv_invol _ _ alpha).
apply inc_{-}inv.
apply H.
Qed.
  Lemma 115 (inv_invol2) Let \alpha, \beta : A \rightarrow B. Then,
                                            \alpha^{\sharp} = \beta^{\sharp} \Rightarrow \alpha = \beta.
Lemma inv\_invol2 {A B : eqType} {alpha beta : Rel A B}:
 alpha \# = \mathtt{beta} \# \to alpha = \mathtt{beta}.
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ alpha) -(@inv_invol _ _ beta).
apply f_equal.
apply H.
Qed.
```

Lemma 116 (inv\_inc\_invol) Let  $\alpha, \beta : A \rightarrow B$ . Then,

```
\alpha^{\sharp} \sqsubseteq \beta^{\sharp} \Rightarrow \alpha \sqsubseteq \beta.
Lemma inv\_inc\_invol {A B : eqType} {alpha beta : Rel A B}:
               \mathtt{beta} \ \# \to alpha
 alpha \#
                                       beta.
Proof.
move \Rightarrow H.
rewrite -(@inv_invol _ _ alpha) -(@inv_invol _ _ beta).
apply inc_inv.
apply H.
Qed.
  Lemma 117 (inv_cupP_distr, inv_cup_distr) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and
  P: predicate. Then,
                                     (\sqcup_{P(\alpha)} f(\alpha))^{\sharp} = (\sqcup_{P(\alpha)} f(\alpha)^{\sharp}).
Lemma inv\_cupP\_distr {A B C D : eqType} {f : Rel C D \rightarrow Rel A B} {P : Rel C D \rightarrow
Prop}:
 ( _{P} f) # = ( _{P} (fun alpha : Rel C D \Rightarrow f alpha #)).
Proof.
apply inc\_antisym.
rewrite -inv\_inc\_move.
apply inc\_cupP.
assert (\forall beta : Rel\ C\ D,\ P\ beta \to f\ beta # _{{}^{-}}{P}\ (fun\ alpha:\ Rel\ C\ D \Rightarrow f
alpha \#)).
apply inc\_cupP.
apply inc_refl.
move \Rightarrow beta H\theta.
rewrite inv\_inc\_move.
apply (H - H\theta).
apply inc\_cupP.
move \Rightarrow beta H0.
apply inc_{-}inv.
move: H0.
apply inc\_cupP.
apply inc\_reft.
Qed.
Lemma inv\_cup\_distr {A B : eqType} {alpha beta : Rel A B}:
             beta) \# = alpha \# beta \#.
 (alpha
Proof.
by [rewrite cup\_to\_cupP -inv\_cupP\_distr -cup\_to\_cupP].
```

Qed.

```
Lemma 118 (inv_capP_distr, inv_cap_distr) Let f: (C \rightarrow D) \rightarrow (A \rightarrow B) and P
  : predicate. Then,
                                       (\sqcap_{P(\alpha)} f(\alpha))^{\sharp} = (\sqcap_{P(\alpha)} f(\alpha)^{\sharp}).
Lemma inv\_capP\_distr {A B C D : eqType} {f : Rel\ C\ D \rightarrow Rel\ A\ B} {P : Rel\ C\ D \rightarrow Rel\ A\ B}
 ( -\{P\} f) \# = ( -\{P\} (\mathbf{fun} \ alpha : Rel \ C \ D \Rightarrow f \ alpha \#)).
Proof.
apply inc\_antisym.
apply inc\_capP.
move \Rightarrow beta H.
apply inc_{-}inv.
move: H.
apply inc\_capP.
apply inc\_reft.
rewrite inv\_inc\_move.
apply inc\_capP.
assert (\forall beta : Rel\ C\ D, P beta \rightarrow _{{P}} (fun\ alpha : Rel\ C\ D \Rightarrow f\ alpha\ \#)
                                                                                                                f
beta \#).
apply inc\_capP.
apply inc\_reft.
move \Rightarrow beta H\theta.
rewrite -inv\_inc\_move.
apply (H - H\theta).
Qed.
Lemma inv\_cap\_distr \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 (alpha
              beta) \# = alpha \# beta \#.
Proof.
by [rewrite cap_to_capP -inv_capP_distr -cap_to_capP].
Qed.
  Lemma 119 (rpc_inv_distr) Let \alpha, \beta : A \rightarrow B. Then,
                                           (\alpha \Rightarrow \beta)^{\sharp} = \alpha^{\sharp} \Rightarrow \beta^{\sharp}.
Lemma rpc\_inv\_distr {A B : eqType} {alpha beta : Rel A B}:
 (alpha \gg beta) \# = alpha \# \gg beta \#.
Proof.
apply inc_lower.
move \Rightarrow gamma.
```

$$\begin{split} & \text{split; move} \Rightarrow H. \\ & \text{apply } inc\_rpc. \\ & \text{rewrite } inv\_inc\_move \ inv\_cap\_distr \ inv\_invol. \\ & \text{rewrite } -inc\_rpc \ -inv\_inc\_move. \\ & \text{apply } H. \\ & \text{rewrite } inv\_inc\_move \ inc\_rpc. \\ & \text{rewrite } -(@inv\_invol \ \_ \ alpha) \ -inv\_cap\_distr \ -inv\_inc\_move. \\ & \text{apply } inc\_rpc. \\ & \text{apply } inc\_rpc. \\ & \text{apply } H. \\ & \text{Qed.} \end{split}$$

# Lemma 120 (inv\_empty)

$$\phi_{AB}^{\sharp} = \phi_{BA}.$$

Lemma  $inv\_empty$  { $A \ B : eqType$ }:  $A \ B \# = B \ A$ . Proof.

apply  $inc\_antisym$ .

rewrite  $-inv\_inc\_move$ .

apply  $inc\_empty\_alpha$ .

apply  $inc\_empty\_alpha$ .

Qed.

# Lemma 121 (inv\_universal)

$$abla_{AB}^{\sharp} = 
abla_{BA}.$$

Lemma  $inv\_universal\ \{A\ B: eqType\}: A\ B\ \#=B\ A.$ 

Proof.

apply  $inc\_antisym$ .

apply  $inc\_alpha\_universal$ .

rewrite  $inv\_inc\_move$ .

apply  $inc\_alpha\_universal$ .

Qed.

## Lemma 122 (inv\_id)

$$id_A^{\sharp} = id_A.$$

Proof

replace  $(Id \ A \ \#)$  with  $((Id \ A \ \#) \ \# \cdot Id \ A \ \#)$ .

by [rewrite -comp\_inv comp\_id\_l inv\_invol].

by [rewrite inv\_invol comp\_id\_l].

Qed.

Lemma 123 (inv\_complement) Let  $\alpha : A \rightarrow B$ . Then,

$$(\alpha^{-})^{\sharp} = (\alpha^{\sharp})^{-}.$$

Lemma  $inv\_complement \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: (alpha ^) \# = (alpha \#) ^.$ apply  $inc\_antisym$ . apply  $inc\_rpc$ . rewrite  $-inv\_cap\_distr$ . rewrite cap\_comm -inv\_inc\_move inv\_empty. rewrite cap\_complement\_empty. apply  $inc\_reft$ . rewrite  $inv\_inc\_move$ . apply  $inc\_rpc$ . alpha) with  $(((alpha \#) \hat{}) \# (alpha \#) \#)$ . replace (((alpha #) ^) #  $\verb"rewrite" - inv\_cap\_distr".$ rewrite cap\_comm -inv\_inc\_move inv\_empty. rewrite cap\_complement\_empty. apply  $inc\_reft$ . by [rewrite  $inv_{-}invol$ ]. Qed.

Lemma 124 (inv\_difference\_distr) Let  $\alpha, \beta : A \rightarrow B$ . Then,

$$(\alpha - \beta)^{\sharp} = \alpha^{\sharp} - \beta^{\sharp}.$$

Lemma  $inv\_difference\_distr$  { $A \ B : eqType$ } { $alpha \ beta : Rel \ A \ B$ }: (alpha - beta)  $\# = alpha \ \# - beta \ \#$ .

Proof.

rewrite  $inv\_cap\_distr$ .

by [rewrite  $inv\_complement$ ].

Qed.

# 5.4 合成に関する補題

**Lemma 125 (comp\_cupP\_distr\_l, comp\_cup\_distr\_l)** Let  $\alpha: A \rightarrow B$ ,  $f: (D \rightarrow E) \rightarrow (B \rightarrow C)$  and P: predicate. Then,

$$\alpha \cdot (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \cdot f(\beta)).$$

Lemma  $comp\_cupP\_distr\_l$  {A B C D E : eqType}

```
\{alpha: Rel\ A\ B\}\ \{f: Rel\ D\ E \rightarrow Rel\ B\ C\}\ \{P: Rel\ D\ E \rightarrow Prop\}:
 alpha \cdot ( _{P} f) = _{P} (fun beta : Rel D E \Rightarrow (alpha \cdot f beta)).
Proof.
apply inc\_upper.
move \Rightarrow gamma.
split; move \Rightarrow H.
rewrite -(@inv\_invol\_\_alpha) in H.
apply inc\_residual in H.
apply inc\_cupP.
assert (\forall beta : Rel\ D\ E, P beta \rightarrow f beta (alpha\ \#\ qamma)).
apply inc\_cupP.
apply H.
move \Rightarrow beta H1.
rewrite -(@inv_invol _ _ alpha).
apply inc\_residual.
apply (H0 - H1).
rewrite -(@inv_invol _ _ alpha).
apply inc\_residual.
apply inc\_cupP.
assert (\forall beta : Rel\ D\ E, P beta \rightarrow (alpha • f beta)
                                                                       qamma).
apply inc\_cupP.
apply H.
move \Rightarrow beta H1.
apply inc_residual.
rewrite inv\_invol.
apply (H0 - H1).
Qed.
Lemma comp\_cup\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
 alpha • (beta
                      gamma) = (alpha \cdot beta)
                                                          (alpha \cdot gamma).
by [rewrite cup\_to\_cupP -comp\_cupP\_distr\_l -cup\_to\_cupP].
Qed.
  Lemma 126 (comp_cupP_distr_r, comp_cup_distr_r) Let f:(D \rightarrow E) \rightarrow (A \rightarrow E)
  B), \beta: B \rightarrow C and P: predicate. Then,
                                  (\sqcup_{P(\alpha)} f(\alpha)) \cdot \beta = \sqcup_{P(\alpha)} (f(\alpha) \cdot \beta).
Lemma comp\_cupP\_distr\_r {A B C D E : eqType}
 \{ beta : Rel \ B \ C \} \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \{ P : Rel \ D \ E \rightarrow Prop \} :
 ( \{P\} f) \cdot beta = \{P\} (fun \ alpha : Rel \ D \ E \Rightarrow (f \ alpha \cdot beta)).
```

```
Proof.
replace (fun alpha: Rel D E \Rightarrow f alpha • beta) with (fun alpha: Rel D E \Rightarrow (beta #
• f alpha #) #).
rewrite -inv\_cupP\_distr.
rewrite -comp\_cupP\_distr\_l.
rewrite -inv\_cupP\_distr.
rewrite comp_{-}inv.
by [rewrite inv_invol inv_invol].
apply functional_extensionality.
move \Rightarrow x.
rewrite comp_{-}inv.
by [rewrite inv_invol inv_invol].
Qed.
Lemma comp\_cup\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
            beta) • gamma = (alpha • gamma)
 (alpha
                                                          (beta • gamma).
Proof.
by [rewrite (@cup\_to\_cupP\_\_\_\_\_\_id) comp\_cupP\_distr\_r -cup\_to\_cupP].
Qed.
  Lemma 127 (comp_capP_distr) Let \alpha: A \to B, \gamma: C \to D, f: (E \to F) \to (B \to F)
  C) and P: predicate. Then,
                             \alpha \cdot (\sqcap_{P(\beta)} f(\beta)) \cdot \gamma \sqsubseteq \sqcap_{P(\beta)} (\alpha \cdot f(\beta) \cdot \gamma).
Lemma comp\_capP\_distr {A B C D E F : eqType}
 {alpha : Rel \ A \ B} \{gamma : Rel \ C \ D\}
 \{f: Rel\ E\ F \to Rel\ B\ C\}\ \{P: Rel\ E\ F \to \operatorname{Prop}\}:
 (alpha \cdot ( _{\{P\}} f)) \cdot gamma
       \{P\} (fun beta : Rel\ E\ F \Rightarrow ((alpha \cdot f beta) \cdot gamma)).
Proof.
apply inc\_capP.
move \Rightarrow beta H.
apply comp\_inc\_compat\_ab\_a'b.
apply comp_inc_compat_ab_ab'.
move: H.
apply inc\_capP.
apply inc\_reft.
Qed.
```

```
Lemma 128 (comp_capP_distr_l, comp_cap_distr_l) Let \alpha: A \rightarrow B, f: (D \rightarrow B)
  E) \rightarrow (B \rightarrow C) and P: predicate. Then,
                                    \alpha \cdot (\sqcap_{P(\beta)} f(\beta)) \sqsubseteq \sqcap_{P(\beta)} (\alpha \cdot f(\beta)).
Lemma comp\_capP\_distr\_l \{A \ B \ C \ D \ E : eqType\}
 \{alpha: Rel\ A\ B\}\ \{f: Rel\ D\ E \rightarrow Rel\ B\ C\}\ \{P: Rel\ D\ E \rightarrow Prop\}:
                              \{P\} (fun beta : Rel\ D\ E \Rightarrow (alpha \cdot f\ beta)).
 (alpha \cdot (-\{P\}f))
Proof.
move: (@comp\_capP\_distr\_\_\_\_ alpha (Id C) f P) \Rightarrow H.
rewrite comp_{-}id_{-}r in H.
replace (fun beta: Rel\ D\ E \Rightarrow (alpha \cdot f beta) \cdot Id\ C) with (fun beta: Rel\ D\ E \Rightarrow
(alpha \cdot f beta)) in H.
apply H.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite comp_{-}id_{-}r].
Qed.
Lemma comp\_cap\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
 (alpha • (beta
                       qamma))
                                       ((alpha \cdot beta) (alpha \cdot gamma)).
Proof.
rewrite cap\_to\_capP (@cap\_to\_capP\_\_\_\_\_id).
apply comp\_capP\_distr\_l.
Qed.
  Lemma 129 (comp_capP_distr_r, comp_cap_distr_r) Let \beta: B \rightarrow C, f: (D \rightarrow C)
  (E) \rightarrow (A \rightarrow B) and P: predicate. Then,
                                    (\sqcap_{P(\alpha)} f(\alpha)) \cdot \beta \sqsubseteq \sqcap_{P(\alpha)} (f(\alpha) \cdot \beta).
{\tt Lemma}\ comp\_capP\_distr\_r
 \{A \ B \ C \ D \ E : eqType\} \ \{ beta : Rel \ B \ C \} \ \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \ \{ P : Rel \ D \ E \rightarrow Rel \ A \ B \} \} 
Prop}:
 (( _{P} f) \cdot beta) _{P} (fun alpha : Rel D E \Rightarrow (f alpha · beta)).
Proof.
move: (@comp\_capP\_distr\_\_\_\_\_(Id\ A)\ beta\ f\ P) \Rightarrow H.
rewrite comp_{-}id_{-}l in H.
replace (fun alpha: Rel D E \Rightarrow (Id A • f alpha) • beta) with (fun alpha: Rel D E
\Rightarrow f \ alpha \cdot beta) \ in \ H.
apply H.
apply functional_extensionality.
```

```
move \Rightarrow x.
by [rewrite comp_{-}id_{-}l].
Qed.
Lemma comp\_cap\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
             beta) • qamma) ((alpha • qamma) (beta • qamma)).
Proof.
rewrite (@cap\_to\_capP\_\_\_\_\_id) (@cap\_to\_capP\_\_\_\_\_\_(fun x \Rightarrow x \cdot gamma)).
apply comp\_capP\_distr\_r.
Qed.
  Lemma 130 (comp_empty_l, comp_empty_r) Let \alpha : A \rightarrow B, \beta : B \rightarrow C. Then,
                                    \alpha \cdot \phi_{BC} = \phi_{AB} \cdot \beta = \phi_{AC}.
Lemma comp\_empty\_r \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\}: alpha  • B \ C =
apply inc\_antisym.
rewrite -(@inv_invol _ _ alpha).
apply inc_residual.
apply inc\_empty\_alpha.
apply inc\_empty\_alpha.
Qed.
Lemma comp\_empty\_l \{A \ B \ C : eqType\} \{ beta : Rel \ B \ C \}: A \ B \cdot beta = A \ C.
Proof.
rewrite -(@inv_invol_{-} ( AB \cdot beta)).
rewrite -inv_move comp_inv inv_empty inv_empty.
apply comp\_empty\_r.
Qed.
  Lemma 131 (comp_either_empty) Let \alpha : A \rightarrow B, \beta : B \rightarrow C. Then,
                               \alpha = \phi_{AB} \vee \beta = \phi_{BC} \Rightarrow \alpha \cdot \beta = \phi_{AC}.
Lemma comp_either_empty {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
              A \ B \lor \mathtt{beta} = B \ C \to alpha \bullet \mathtt{beta} = A \ C.
 alpha =
Proof.
case; move \Rightarrow H.
rewrite H.
apply comp\_empty\_l.
rewrite H.
apply comp\_empty\_r.
```

Qed.

```
\alpha \cdot \beta \neq \phi_{AC} \Rightarrow \alpha \neq \phi_{AB} \wedge \beta \neq \phi_{BC}. Lemma comp\_neither\_empty \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}: alpha \cdot beta \neq A \ C \rightarrow alpha \neq A \ B \wedge beta \neq B \ C. Proof. move \Rightarrow H. split; move \Rightarrow H0. apply H. rewrite H0. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_l. apply comp\_empty\_r. Qed.
```

Lemma 132 (comp\_neither\_empty) Let  $\alpha : A \to B$ ,  $\beta : B \to C$ . Then,

# 5.5 単域と Tarski の定理

```
Lemma 133 (lemma_for_tarski1) Let \alpha : A \to B and \alpha \neq \phi_{AB}. Then, \nabla_{IA} \cdot \alpha \cdot \nabla_{BI} = id_I.
```

```
Lemma lemma\_for\_tarski1 \{A B : eqType\} \{alpha : Rel A B\}:
             A B \rightarrow ((i A \cdot alpha) \cdot B i) = Id i.
 alpha \neq
Proof.
move \Rightarrow H.
              i A \cdot alpha) \cdot B i \neq i i.
assert (((
move \Rightarrow H0.
apply H.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((Ai \cdot ((iA \cdot alpha) \cdot Bi)) \cdot iB)).
rewrite comp_assoc comp_assoc unit_universal.
rewrite -comp_assoc -comp_assoc unit_universal.
apply (@inc\_trans \_ \_ \_ ((Id A \cdot alpha) \cdot Id B)).
rewrite comp\_id\_l comp\_id\_r.
apply inc\_reft.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
```

```
apply inc\_alpha\_universal.

rewrite H0\ comp\_empty\_r\ comp\_empty\_l.

apply inc\_refl.

apply inc\_empty\_alpha.

case (@unit\_empty\_or\_universal (( i A · alpha) · B i)); move \Rightarrow H1.

apply False\_ind.

apply (H0\ H1).

rewrite unit\_identity\_is\_universal.

apply H1.

Qed.
```

# Lemma 134 (lemma\_for\_tarski2)

$$\nabla_{AI} \cdot \nabla_{IB} = \nabla_{AB}$$
.

Lemma  $lemma\_for\_tarski2$  {A B : eqType}: A  $i \cdot i B = i$ A B. Proof. apply  $inc\_antisym$ . apply  $inc\_alpha\_universal$ . apply  $@inc\_trans\_\_\_$  (  $AA \cdot$ A B)). apply  $(@inc\_trans \_ \_ \_ (Id A \cdot A B))$ . rewrite  $comp_{-}id_{-}l$ . apply  $inc_refl.$ apply  $comp\_inc\_compat\_ab\_a'b$ . apply  $inc\_alpha\_universal$ . rewrite -(@unit\_universal A) comp\_assoc. apply  $comp\_inc\_compat\_ab\_ab$ '. apply  $inc\_alpha\_universal$ . Qed.

**Lemma 135 (tarski)** Let  $\alpha : A \rightarrow B$  and  $\alpha \neq \phi_{AB}$ . Then,

$$\nabla_{AA} \cdot \alpha \cdot \nabla_{BB} = \nabla_{AB}.$$

```
Lemma tarski {A \ B : eqType} {alpha : Rel \ A \ B}: alpha \neq A \ B \rightarrow ((A \ A \ \cdot \ alpha) \ \cdot B \ B) = A \ B.

Proof.

move \Rightarrow H.

rewrite -(@unit\_universal \ A) -(@unit\_universal \ B).

move : (@lemma\_for\_tarski1 \ \_ \ alpha \ H) \Rightarrow H0.

rewrite -comp\_assoc \ (@comp\_assoc \ \_ \ \_ \ \_ \ (A \ i)) \ (@comp\_assoc \ \_ \ \_ \ \_ \ (A \ i)).

rewrite H0 \ comp\_id\_r.

apply lemma\_for\_tarski2.
```

Qed.

```
Lemma 136 (comp_universal1) Let B \neq \emptyset. Then,
```

$$\nabla_{AB} \cdot \nabla_{BC} = \nabla_{AC}.$$

```
Lemma comp\_universal\ \{A\ B\ C: eqType\}: B \rightarrow A\ B \cdot B\ C =
                                                                          A C.
Proof.
move \Rightarrow b.
replace (AB) with (AB \cdot BB).
rewrite -(@lemma_for_tarski2 A B) -(@lemma_for_tarski2 B C).
rewrite (@comp_assoc _ _ _ ( A i)) (@comp_assoc _ _ _ ( A i)) -(@comp_assoc _
---(Bi).
rewrite lemma\_for\_tarski1.
rewrite comp_{-}id_{-}l.
apply lemma_for_tarski2.
apply not\_eq\_sym.
move \Rightarrow H.
apply either\_empty in H.
case H; move \Rightarrow H0.
apply (H0\ b).
apply (H0\ b).
apply inc\_antisym.
apply inc_alpha_universal.
apply (@inc\_trans \_ \_ \_ ( AB \cdot Id B)).
rewrite comp_{-}id_{-}r.
apply inc\_reft.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
```

# Lemma 137 (comp\_universal2)

$$\nabla_{IA}^{\sharp} \cdot \nabla_{IB} = \nabla_{AB}.$$

Lemma  $comp\_universal2$  { $A \ B : eqType$ }:  $i \ A \ \# \bullet$   $i \ B = A \ B$ . Proof. rewrite  $inv\_universal$ . apply  $lemma\_for\_tarski2$ . Qed.

# Lemma 138 (empty\_equivalence1, empty\_equivalence2, empty\_equivalence3)

```
A = \emptyset \Leftrightarrow \nabla_{IA} = \phi_{IA} \Leftrightarrow \nabla_{AA} = \phi_{AA} \Leftrightarrow id_A = \phi_{AA}.
```

```
Lemma empty_equivalence1 \{A : eqType\}: (A \rightarrow False) \leftrightarrow i A =
Proof.
move: (@either\_empty \ i \ A) \Rightarrow H.
split; move \Rightarrow H0.
apply Logic.eq_sym.
apply H.
right.
apply H0.
apply Logic.eq\_sym in H0.
apply H in H0.
case H0.
move \Rightarrow H1 H2.
apply H1.
apply tt.
by [].
Qed.
Lemma empty\_equivalence2 \{A : eqType\}: (A \rightarrow False) \leftrightarrow
                                                                  A A =
                                                                               A A.
Proof.
move: (@either\_empty\ A\ A) \Rightarrow H.
split; move \Rightarrow H0.
apply Logic.eq_sym.
apply H.
left.
apply H0.
apply Logic.eq_sym in H0.
apply H in H\theta.
case H0.
by [].
by [].
Lemma empty_equivalence3 \{A: eqType\}: (A \rightarrow False) \leftrightarrow Id A = AA.
split; move \Rightarrow H.
assert ( AA =
                        A A).
apply empty_equivalence2.
apply H.
apply RelAB\_unique.
apply Logic.eq_sym.
```

```
apply H0.
assert ( AA = AA).
by [rewrite -(@comp\_id\_r\_\_ ( AA)) H comp\_empty\_r].
apply either\_empty in H0.
case H0.
by [].
by [].
```

# Chapter 6

# Library Functions\_Mappings

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Logic.FunctionalExtensionality.
```

# 6.1 全域性,一価性,写像に関する補題

```
Lemma 139 (id_function) id_A: A \rightarrow A is a function.

Lemma id\_function \{A: eqType\}: function\_r \ (Id\ A).

Proof.

rewrite /function\_r/total\_r/univalent\_r.

rewrite inv\_id\ comp\_id\_l.

split.

apply inc\_refl.

apply inc\_refl.

Qed.
```

```
Lemma 140 (unit_function) \nabla_{AI}: A \rightarrow I is a function.
```

```
Lemma unit\_function \{A : eqType\}: function\_r (A i).

Proof.

rewrite /function\_r/total\_r/univalent\_r.

rewrite inv\_universal lemma\_for\_tarski2 unit\_identity\_is\_universal.

split.

apply inc\_alpha\_universal.

apply inc\_alpha\_universal.

Qed.
```

```
Lemma 141 (total_comp) Let \alpha: A \to B and \beta: B \to C be total relations, then
  \alpha \cdot \beta is also a total relation.
Lemma total\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 total_r \ alpha \rightarrow total_r \ beta \rightarrow total_r \ (alpha \cdot beta).
Proof.
rewrite /total_r.
move \Rightarrow H H0.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H0.
Qed.
  Lemma 142 (univalent_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be univalent relations,
  then \alpha \cdot \beta is also a univalent relation.
Lemma univalent_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (alpha \cdot beta).
Proof.
rewrite /univalent_r.
move \Rightarrow H H0.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ (alpha #)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_ab\_b.
apply H.
Qed.
  Lemma 143 (function_comp) Let \alpha: A \to B and \beta: B \to C be functions, then \alpha \cdot \beta
  is also a function.
Lemma function\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 function\_r \ alpha \rightarrow function\_r \ beta \rightarrow function\_r \ (alpha \ \cdot \ beta).
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (total\_comp\ H\ H1).
apply (univalent\_comp\ H0\ H2).
Qed.
```

```
Lemma 144 (total_comp2) Let \alpha: A \to B, \beta: B \to C and \alpha \cdot \beta be a total relation,
  then \alpha is also a total relation.
Lemma total\_comp2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total_r (alpha \cdot beta) \rightarrow total_r alpha.
Proof.
move \Rightarrow H.
apply inc\_def1 in H.
rewrite comp\_inv cap\_comm comp\_assoc in H.
rewrite /total_r.
rewrite H.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ )).
apply comp\_inc\_compat.
apply cap_{-}l.
rewrite comp_{-}id_{-}r.
apply cap_r.
Qed.
  Lemma 145 (univalent_comp2) Let \alpha: A \rightarrow B, \beta: B \rightarrow C, \alpha \cdot \beta be a univalent
  relation and \alpha^{\sharp} be a total relation, then \beta is a univalent relation.
Lemma univalent\_comp2 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 univalent_r (alpha \cdot beta) \rightarrow total_r (alpha \#) \rightarrow univalent_r beta.
Proof.
move \Rightarrow H H0.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp_inc_compat_ab_ab'.
rewrite /total_r in H0.
rewrite inv_{-}invol in H0.
apply (comp\_inc\_compat\_b\_ab\ H0).
Qed.
  Lemma 146 (total_inc) Let \alpha : A \to B be a total relation and \alpha \sqsubseteq \beta, then \beta is also
  a total relation.
Lemma total\_inc {A B : eqType} {alpha beta : Rel A B}:
 total\_r \ alpha \rightarrow alpha \quad beta \rightarrow total\_r \ beta.
Proof.
move \Rightarrow H H0.
apply @inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat.
apply H0.
```

```
apply (@inc_inv_{-} - H0). Qed.
```

**Lemma 147 (univalent\_inc)** Let  $\alpha : A \to B$  be a univalent relation and  $\beta \sqsubseteq \alpha$ , then  $\beta$  is also a univalent relation.

```
Lemma univalent\_inc {A \ B : eqType} {alpha \ beta : Rel \ A \ B}: univalent\_r \ alpha \rightarrow beta \quad alpha \rightarrow univalent\_r \ beta. Proof.

move \Rightarrow H \ H0.

apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' \ H).

apply comp\_inc\_compat.

apply (@inc\_inv _ _ _ H0).

apply H0.

Qed.
```

**Lemma 148 (function\_inc)** Let  $\alpha, \beta : A \to B$  be functions and  $\alpha \sqsubseteq \beta$ . Then,

$$\alpha = \beta$$
.

```
Lemma function\_inc \{A \ B : eqType\} \{alpha \ beta : Rel \ A \ B\}:
 function\_r \ alpha \rightarrow function\_r \ \mathtt{beta} \rightarrow alpha
                                                       beta \rightarrow alpha = beta.
Proof.
move \Rightarrow H H0 H1.
apply inc\_antisym.
apply H1.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot beta)).
apply comp\_inc\_compat\_b\_ab.
apply H.
move: (@inc_inv_- - H1) \Rightarrow H2.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot beta \#) \cdot beta)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply H2.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
Qed.
```

Lemma 149 (total\_universal) If  $\nabla_{IB}$  be a total relation, then

$$\nabla_{AB} \cdot \nabla_{BC} = \nabla_{AC}$$
.

```
Lemma total\_universal \{A \ B \ C : eqType\}:
 total_r ( i B) \rightarrow
                         AB \cdot BC =
Proof.
move \Rightarrow H.
rewrite -(@lemma_for_tarski2 A B) -(@lemma_for_tarski2 B C).
rewrite comp\_assoc -(@comp\_assoc _ _ _ _ ( i B)).
replace ( i B •
                           B i) with (Id i).
rewrite comp_{-}id_{-}l.
apply lemma_for_tarski2.
apply inc\_antisym.
rewrite /total_r in H.
rewrite inv\_universal in H.
apply H.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
  Lemma 150 (function_rel_inv_rel) Let \alpha : A \to B be function. Then,
                                             \alpha \cdot \alpha^{\sharp} \cdot \alpha = \alpha
Lemma function\_rel\_inv\_rel \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 function\_r \ alpha \rightarrow (alpha \cdot alpha \#) \cdot alpha = alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H.
apply comp\_inc\_compat\_b\_ab.
apply H.
Qed.
  Lemma 151 (function_capP_distr) Let f: A \to B, g: D \to C be functions, \theta:
  (E \rightarrow F) \rightarrow (B \rightarrow C) and P: predicate. Then,
                              f \cdot (\sqcap_{P(\theta)} \theta(\alpha)) \cdot g^{\sharp} = \sqcap_{P(\alpha)} (f \cdot \theta(\alpha) \cdot g^{\sharp}).
Lemma function\_capP\_distr {A B C D E F : eqType}
 \{f: Rel\ A\ B\}\ \{g: Rel\ D\ C\}\ \{theta: Rel\ E\ F \to Rel\ B\ C\}\ \{P: Rel\ E\ F \to Prop\}:
 function_r f \rightarrow function_r g \rightarrow
 (f \cdot (-\{P\} \ theta)) \cdot g \# =
    _{-}\{P\} (fun alpha : Rel E F \Rightarrow (f • theta alpha) • g \#).
```

```
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
apply inc\_antisym.
apply comp\_capP\_distr.
apply (@inc\_trans \_ \_ \_ (((f \cdot f \#) \cdot \_ \{P\} (fun \ alpha : Rel \ E \ F \Rightarrow (f \cdot theta \ alpha)))
• g \# )) • (g • g \# ))).
apply (@inc\_trans \_ \_ \_ ((f \cdot f \#) \cdot ( \_\{P\} (fun \ alpha : Rel \ E \ F \Rightarrow (f \cdot theta \ alpha)))
• g \#)))).
apply (comp\_inc\_compat\_b\_ab\ H).
apply (comp\_inc\_compat\_a\_ab\ H1).
rewrite (@comp\_assoc\_\_\_\_ (f \#)) comp\_assoc\_(@comp\_assoc\_\_\_\_ g) - comp\_assoc.
apply comp_inc_compat_ab_a'b.
apply comp_inc_compat_ab_ab'.
apply (@inc_trans _ _ _ ( _{P} (fun alpha : Rel E F \Rightarrow (f # • ((f • theta alpha) • q
\#)) \cdot g))).
apply comp\_capP\_distr.
replace (fun alpha: Rel E F \Rightarrow (f # • ((f • theta alpha) • g #)) • g) with (fun alpha
: Rel\ E\ F \Rightarrow ((f \# \bullet f) \bullet theta\ alpha) \bullet (g \# \bullet g)).
apply inc\_capP.
move \Rightarrow beta H3.
apply (@inc\_trans\_\_\_((f \# \cdot f) \cdot theta beta)).
apply (@inc\_trans \_ \_ \_ (((f \# \cdot f) \cdot theta \ beta) \cdot (g \# \cdot g))).
move: beta H3.
apply inc\_capP.
apply inc\_reft.
apply (comp\_inc\_compat\_ab\_a\ H2).
apply (comp\_inc\_compat\_ab\_b\ H0).
apply functional_extensionality.
move \Rightarrow x.
by rewrite comp_assoc comp_assoc comp_assoc comp_assoc.
Qed.
  Lemma 152 (function_cap_distr, function_cap_distr_l, function_cap_distr_r)
  Let f: A \to B, q: D \to C be functions and \alpha, \beta: B \to C. Then,
                              f \cdot (\alpha \sqcap \beta) \cdot q^{\sharp} = (f \cdot \alpha \cdot q^{\sharp}) \sqcap (f \cdot \beta \cdot q^{\sharp}).
Lemma function\_cap\_distr
 \{A \ B \ C \ D : eqType\} \ \{f : Rel \ A \ B\} \ \{alpha \ \mathsf{beta} : Rel \ B \ C\} \ \{g : Rel \ D \ C\}: \}
```

 $(f \cdot (alpha \quad beta)) \cdot q \# = ((f \cdot alpha) \cdot q \#) \quad ((f \cdot beta) \cdot q \#).$ 

 $function_r f \rightarrow function_r g \rightarrow$ 

Proof.

```
\overline{\text{rewrite } (@cap\_to\_capP\_\_\_\_\_id) \ (@cap\_to\_capP\_\_\_\_\_\_(\mathbf{fun} \ x \Rightarrow (f \cdot x) \cdot g)}
\#)).
apply function\_capP\_distr.
Qed.
Lemma function\_cap\_distr\_l
 \{A \ B \ C : eqType\} \ \{f : Rel \ A \ B\} \ \{alpha \ beta : Rel \ B \ C\}:
 function_r f \rightarrow
                  \mathtt{beta}) = (f \cdot alpha) \quad (f \cdot \mathtt{beta}).
 f \cdot (alpha)
Proof.
move: (@id\_function\ C) \Rightarrow H.
move \Rightarrow H0.
apply (@function\_cap\_distr\_\_\_\_f \ alpha \ beta) in H.
rewrite inv_id comp_id_r comp_id_r comp_id_r in H.
apply H.
apply H0.
Qed.
Lemma function\_cap\_distr\_r
 \{B\ C\ D: eqType\}\ \{alpha\ \mathsf{beta}: Rel\ B\ C\}\ \{g: Rel\ D\ C\}:
 function_r g \rightarrow
              beta) • q \# = (alpha • q \#) (beta • q \#).
 (alpha
Proof.
move: (@id\_function B) \Rightarrow H.
move \Rightarrow H0.
apply (@function\_cap\_distr\_\_\_\_ alpha beta g) in H.
rewrite comp_{-}id_{-}l comp_{-}id_{-}l comp_{-}id_{-}l in H.
apply H.
apply H0.
Qed.
  Lemma 153 (function_move1) Let \alpha : A \rightarrow B be a function, \beta : B \rightarrow C and
  \gamma: A \rightarrow C. Then,
                                           \gamma \sqsubseteq \alpha \cdot \beta \Leftrightarrow \alpha^{\sharp} \cdot \gamma \sqsubseteq \beta.
Lemma function\_move1 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C} {gamma : Rel \ A \ B} {gamma : Rel \ B \ C} {gamma : Rel \ B \ C}
Rel\ A\ C:
 function\_r \ alpha \rightarrow (gamma \ (alpha \cdot beta) \leftrightarrow (alpha \# \cdot gamma) \ beta).
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply @inc\_trans \_ \_ \_ ((alpha \# \cdot alpha) \cdot beta)).
rewrite comp\_assoc.
apply (comp\_inc\_compat\_ab\_ab' H0).
```

```
apply comp\_inc\_compat\_ab\_b.
apply H.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot gamma)).
apply comp\_inc\_compat\_b\_ab.
apply H.
rewrite comp_-assoc.
apply (comp\_inc\_compat\_ab\_ab' H0).
Qed.
  Lemma 154 (function_move2) Let \beta: B \rightarrow C be a function, \alpha: A \rightarrow B and
  \gamma: A \rightarrow C. Then,
                                           \alpha \cdot \beta \sqsubseteq \gamma \Leftrightarrow \alpha \sqsubseteq \gamma \cdot \beta^{\sharp}.
Lemma function_move2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C} {qamma :
Rel\ A\ C:
 function\_r \ \mathsf{beta} \to ((alpha \cdot \mathsf{beta}) \quad gamma \leftrightarrow alpha \quad (gamma \cdot \mathsf{beta} \#)).
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot beta) \cdot beta \#)).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_a\_ab.
apply H.
apply (comp\_inc\_compat\_ab\_a'b H0).
apply (@inc\_trans \_ \_ \_ ((gamma \cdot beta \#) \cdot beta)).
apply (comp\_inc\_compat\_ab\_a'b\ H0).
rewrite comp_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H.
Qed.
  Lemma 155 (function_rpc_distr) Let f: A \rightarrow B, g: D \rightarrow C be functions and
  \alpha, \beta: B \rightarrow C. Then,
                               f \cdot (\alpha \Rightarrow \beta) \cdot q^{\sharp} = (f \cdot \alpha \cdot q^{\sharp}) \Rightarrow (f \cdot \beta \cdot q^{\sharp}).
Lemma function\_rpc\_distr
 \{A \ B \ C \ D : eqType\} \{f : Rel \ A \ B\} \{alpha \ beta : Rel \ B \ C\} \{g : Rel \ D \ C\}:
 function_r f \rightarrow function_r g \rightarrow
 (f \cdot (alpha \otimes beta)) \cdot g \# = ((f \cdot alpha) \cdot g \#) \otimes ((f \cdot beta) \cdot g \#).
Proof.
move \Rightarrow H H0.
apply inc_lower.
```

```
move \Rightarrow qamma.
split; move \Rightarrow H1.
apply inc\_rpc.
apply (function_move2 H0).
apply (function\_move1 \ H).
apply (@inc\_trans\_\_\_ (((f \# \cdot gamma) \cdot g) ((f \# \cdot ((f \cdot alpha) \cdot g \#)) \cdot g))).
rewrite -comp\_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (@comp\_cap\_distr\_r \_ \_ \_ \_ \_)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_cap\_distr\_l.
apply (function_move2 H0) in H1.
apply (function_move1 H) in H1.
rewrite -inc_rpc comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ H1).
apply rpc\_inc\_compat\_r.
rewrite comp_assoc comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ (alpha \cdot (g \# \cdot g))).
apply comp\_inc\_compat\_ab\_b.
apply H.
apply comp\_inc\_compat\_ab\_a.
apply H0.
apply (function_move2 H0).
apply (function\_move1 \ H).
apply inc\_rpc.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
apply (@inc\_trans \_ \_ \_ (f \# \cdot ((gamma \cdot g) ((f \#) \# \cdot alpha)))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_l.
rewrite inv_invol.
apply (@inc\_trans \_ \_ \_ ((f \# \cdot (gamma ((f \cdot alpha) \cdot g \#))) \cdot g)).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc\_trans \_ \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
apply (function_move2 H0).
apply (function\_move1 \ H).
rewrite -inc_rpc -comp_assoc.
apply H1.
Qed.
```

Then,

Lemma 156 (function\_inv\_rel1, function\_inv\_rel2) Let  $f : A \to B$  be a function.

```
f^{\sharp} \cdot f = id_B \cap f^{\sharp} \cdot \nabla_{AA} \cdot f = id_B \cap \nabla_{BA} \cdot f.
Lemma function\_inv\_rel1 \{A B : eqType\} \{f : Rel A B\}:
 function_r f \to f \# \cdot f = Id B \quad ((f \# \cdot A A) \cdot f).
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply inc\_cap.
split.
apply H.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_a\_ab.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ (Id B (B A \cdot f))).
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite cap\_comm.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite comp_id_l comp_id_r cap_comm inv_universal.
rewrite cap_universal cap_universal.
apply inc\_reft.
Qed.
Lemma function_inv_rel2 \{A \ B : eqType\} \{f : Rel \ A \ B\}:
function\_r f \rightarrow f \# \cdot f = Id B \quad (BA \cdot f).
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite (@function_inv_rel1 _ _ _ H).
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite cap\_comm.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite comp_id_l comp_id_r cap_comm inv_universal.
rewrite cap_universal cap_universal.
apply inc_refl.
Qed.
```

function,  $\mu: C \to A$  and  $\rho: C \to B$ . Then,

```
(\mu \sqcap \rho \cdot f^{\sharp}) \cdot f = \mu \cdot f \sqcap \rho \wedge \rho \cdot f^{\sharp} \cdot f = \nabla_{CA} \cdot f \sqcap \rho.
Lemma function_dedekind1
 \{A\ B\ C: eqType\}\ \{f: Rel\ A\ B\}\ \{mu: Rel\ C\ A\}\ \{rho: Rel\ C\ B\}:
 function_r f \rightarrow (mu \quad (rho \cdot f \#)) \cdot f = (mu \cdot f)
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H.
apply (@inc\_trans \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
Qed.
Lemma function_dedekind2 \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{rho : Rel \ C \ B\}:
 function_r f \rightarrow (rho \cdot f \#) \cdot f = (CA \cdot f)
Proof.
move \Rightarrow H.
move: (@function\_dedekind1 \_ \_ \_ f ( CA) rho H) \Rightarrow H0.
rewrite cap\_comm\ cap\_universal\ in\ H0.
apply H0.
Qed.
```

Lemma 157 (function\_dedekind1, function\_dedekind2) Let  $f: A \rightarrow B$  be a

# 6.2 全射, 単射に関する補題

```
Lemma 158 (surjection_comp) Let \alpha: A \rightarrow B and \beta: B \rightarrow C be surjections, then \alpha \cdot \beta is also a surjection.
```

```
Lemma surjection\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}: surjection\_r \ alpha \rightarrow surjection\_r \ beta \rightarrow surjection\_r \ (alpha \ ^{\bullet} \ beta).

Proof.
rewrite /surjection\_r.
elim \Rightarrow H \ H0.
elim \Rightarrow H1 \ H2.
split.
```

```
apply (function_comp H H1).
rewrite comp_{-}inv.
apply (total\_comp\ H2\ H0).
Qed.
  Lemma 159 (injection_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be injections, then
  \alpha \cdot \beta is also an injection.
Lemma injection\_comp {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C}:
 injection\_r \ alpha \rightarrow injection\_r \ beta \rightarrow injection\_r \ (alpha \cdot beta).
Proof.
rewrite /injection_r.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (function_comp H H1).
rewrite comp_{-}inv.
apply (univalent\_comp\ H2\ H0).
Qed.
  Lemma 160 (bijection_comp) Let \alpha : A \rightarrow B and \beta : B \rightarrow C be bijections, then
  \alpha \cdot \beta is also a bijection.
Lemma bijection_comp \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}:
 bijection\_r \ alpha \rightarrow bijection\_r \ beta \rightarrow bijection\_r \ (alpha \cdot beta).
Proof.
rewrite /bijection_r.
elim \Rightarrow H.
elim \Rightarrow H0 \ H1.
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
split.
apply (function\_comp\ H\ H2).
rewrite comp_{-}inv.
split.
apply (total\_comp\ H3\ H0).
apply (univalent\_comp\ H4\ H1).
Qed.
  Lemma 161 (surjection_unique1) Let e: A \rightarrow B be a surjection, f: A \rightarrow C be a
```

```
function and e \cdot e^{\sharp} \sqsubseteq f \cdot f^{\sharp}, then there exists a unique function g : B \to C s.t. f = eg.
```

Lemma  $surjection\_unique1 \{A \ B \ C : eqType\} \{e : Rel \ A \ B\} \{f : Rel \ A \ C\}:$ 

```
surjection\_r \ e \rightarrow function\_r \ f \rightarrow (e \cdot e \#)
                                                    (f \cdot f \#) \rightarrow
 (\exists ! \ g : Rel \ B \ C, function\_r \ g \land f = e \cdot g).
Proof.
rewrite /surjection\_r/function\_r/total\_r/univalent\_r.
elim.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 H3 H4.
\exists (e \# \cdot f).
repeat split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ f).
apply (@inc\_trans \_ \_ \_ \_ H1).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H2.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
apply (@inc\_trans\_\_\_(f \# \cdot ((f \cdot f \#) \cdot f))).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_a'b H_4).
rewrite comp\_assoc -comp\_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H3).
apply function_inc.
split.
apply H2.
apply H3.
split.
rewrite /total_r.
rewrite comp_inv comp_inv inv_invol.
rewrite -(@comp\_assoc\_\_\_\_e) (@comp\_assoc\_\_\_\_e) (@comp\_assoc\_\_\_\_f)
-(@comp\_assoc\_\_\_f).
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_a\_ab.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply (comp\_inc\_compat\_a\_ab\ H).
rewrite /univalent_r.
rewrite comp_inv comp_inv inv_invol.
rewrite (@comp\_assoc\_\_\_\_e) -(@comp\_assoc\_\_\_e) comp_assoc -(@comp\_assoc
_ _ _ f).
apply (@inc\_trans \_ \_ \_ (f \# \cdot (((f \cdot f \#) \cdot (f \cdot f \#)) \cdot f))).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat.
```

```
apply H_4.
apply H_4.
rewrite comp\_assoc (@comp\_assoc _ _ _ _ f) -(@comp\_assoc _ _ _ _ (f \#)) -(@comp\_assoc
\_\_\_\_(f \#)) (@comp\_assoc\_\_\_\_(f \#)) - (@comp\_assoc\_\_\_(f \#)).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H3).
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_b\_ab\ H).
move \Rightarrow q.
elim.
elim \Rightarrow H5 \ H6 \ H7.
replace q with (e \# \cdot (e \cdot q)).
apply f_equal.
apply H\gamma.
rewrite -comp\_assoc.
apply inc\_antisym.
apply (comp\_inc\_compat\_ab\_b\ H0).
rewrite inv\_invol in H1.
apply (comp\_inc\_compat\_b\_ab\ H1).
Qed.
  Lemma 162 (surjection_unique2) Let e: A \rightarrow B be a surjection, f: A \rightarrow C be a
  function and e \cdot e^{\sharp} = f \cdot f^{\sharp}, then function e^{\sharp} f is an injection.
Lemma surjection\_unique2 \{A \ B \ C : eqType\} \{e : Rel \ A \ B\} \{f : Rel \ A \ C\}:
 surjection\_r \ e \rightarrow function\_r \ f \rightarrow (e \cdot e \#) = (f \cdot f \#) \rightarrow injection\_r \ (e \# \cdot f).
Proof.
rewrite /surjection_r/injection_r/function_r/total_r/univalent_r.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 \ H3 \ H4.
repeat split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ f).
apply (@inc\_trans \_ \_ \_ \_ H1).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H2.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
rewrite H_4.
rewrite comp_assoc -comp_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
```

```
apply (comp_inc_compat_ab_a H3).
rewrite inv_invol comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
rewrite -H4.
rewrite comp_assoc -comp_assoc.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp\_inc\_compat\_ab\_a.
apply H0.
Qed.
  Lemma 163 (injection_unique1) Let m: B \rightarrow A be an injection, f: C \rightarrow A be a
  function and f^{\sharp} \cdot f \sqsubseteq m^{\sharp} \cdot m, then there exists a unique function q: C \to B s.t. f = qm.
Lemma injection\_unique1 \{A \ B \ C : eqType\} \{m : Rel \ B \ A\} \{f : Rel \ C \ A\}:
 injection\_r \ m \rightarrow function\_r \ f \rightarrow (f \# \bullet f) \ (m \# \bullet m) \rightarrow
 (\exists ! \ g : Rel \ C \ B, function\_r \ g \land f = g \bullet m).
rewrite /injection_r/function_r/total_r/univalent_r.
elim.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 H3 H4.
\exists (f \cdot m \#).
repeat split.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ m).
apply (@inc\_trans\_\_\_(f \cdot ((f \# \cdot f) \cdot f \#))).
rewrite comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply (comp\_inc\_compat\_a\_ab\ H2).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_a'b\ H_4).
rewrite comp\_inv \ comp\_assoc \ -(@comp\_assoc \ \_ \ \_ \ \_ \ f).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
rewrite comp\_assoc.
apply Logic.eq_sym.
apply function_inc.
split.
rewrite /total_r.
rewrite comp_inv comp_inv inv_invol.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply comp\_inc\_compat.
apply (@inc\_trans\_\_\_(f \cdot (f \# \cdot f))).
rewrite -comp\_assoc.
```

```
apply (comp\_inc\_compat\_b\_ab\ H2).
apply (comp_inc_compat_ab_ab' H4).
apply (@inc\_trans\_\_\_((f \# \bullet f) \bullet f \#)).
rewrite comp_assoc.
apply (comp\_inc\_compat\_a\_ab H2).
apply (comp_inc_compat_ab_a'b H_4).
rewrite /univalent_r.
rewrite comp_inv comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
apply (comp\_inc\_compat\_ab\_a\ H0).
split.
apply H2.
apply H3.
apply (comp\_inc\_compat\_ab\_a\ H0).
move \Rightarrow g.
elim.
elim \Rightarrow H5 \ H6 \ H7.
rewrite H7 comp\_assoc.
apply inc\_antisym.
rewrite inv_invol in H1.
apply (comp\_inc\_compat\_ab\_a\ H1).
apply (comp\_inc\_compat\_a\_ab\ H).
Qed.
  Lemma 164 (injection_unique2) Let m: B \rightarrow A be an injection, f: C \rightarrow A be a
  function and f^{\sharp} \cdot f = m^{\sharp} \cdot m, then function f \cdot m^{\sharp} is a surjection.
Lemma injection\_unique2 {A B C : eqType} {m : Rel B A} {f : Rel C A}:
 injection\_r \ m \rightarrow function\_r \ f \rightarrow (f \# \bullet f) = (m \# \bullet m) \rightarrow surjection\_r \ (f \bullet m \#).
Proof.
rewrite /surjection_r/injection_r/function_r/total_r/univalent_r.
elim.
elim \Rightarrow H H0 H1.
elim \Rightarrow H2 \ H3 \ H4.
repeat split.
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ m).
apply (@inc\_trans \_ \_ \_ (f \cdot ((f \# \cdot f) \cdot f \#))).
rewrite comp_assoc -comp_assoc.
apply @inc_trans_H = H2.
apply (comp\_inc\_compat\_a\_ab\ H2).
apply comp\_inc\_compat\_ab\_ab'.
```

```
rewrite H4.
apply inc\_reft.
rewrite comp\_inv \ comp\_assoc \ -(@comp\_assoc \ \_ \ \_ \ \_ \ f).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
rewrite inv_invol comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ _ f).
apply (@inc\_trans \_ \_ \_ \_ H).
apply comp\_inc\_compat\_ab\_ab'.
rewrite H4 comp_assoc.
apply (comp\_inc\_compat\_a\_ab\ H).
Qed.
  Lemma 165 (bijection_inv) Let \alpha: A \to B, \beta: B \to A, \alpha \cdot \beta = id_A and \beta \cdot \alpha = id_B,
  then \alpha and \beta are bijections and \beta = \alpha^{\sharp}.
Lemma bijection_inv {A B : eqType} {alpha : Rel A B} {beta : Rel B A}:
 alpha • beta = Id\ A \rightarrow beta • alpha = Id\ B \rightarrow bijection\_r\ alpha \land bijection\_r\ beta \land
beta = alpha \#.
Proof.
move \Rightarrow H H0.
move: (@id_function A) \Rightarrow H1.
move: (@id\_function B) \Rightarrow H2.
assert (bijection_r \ alpha \land bijection_r \ beta).
assert (total_r \ alpha \land total_r \ (alpha \#) \land total_r \ beta \land total_r \ (beta \#)).
repeat split.
apply (@total\_comp2 \_ \_ \_ \_ beta).
rewrite H.
apply H1.
apply (@total\_comp2\_\_\_\_ (beta \#)).
rewrite - comp_inv H0 inv_id.
apply H2.
apply (@total\_comp2\_\_\_\_alpha).
rewrite H0.
apply H2.
apply (@total\_comp2\_\_\_\_(alpha \#)).
rewrite -comp_inv H inv_id.
apply H1.
repeat split.
apply H3.
apply (@univalent_comp2 _ _ beta).
rewrite H0.
apply H2.
```

```
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ (beta \#)).
rewrite -comp\_inv \ H \ inv\_id.
apply H1.
rewrite inv_-invol.
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ alpha).
rewrite H.
apply H1.
apply H3.
apply H3.
apply (@univalent\_comp2 \_ \_ \_ (alpha \#)).
rewrite -comp_inv H0 inv_id.
apply H2.
rewrite inv\_invol.
apply H3.
split.
apply H3.
split.
apply H3.
rewrite -(@comp\_id\_r\_\_\_beta) -(@comp\_id\_l\_\_\_(alpha \#)).
rewrite -H0 comp\_assoc.
apply f_equal.
apply inc\_antisym.
apply H3.
rewrite comp_inv_inv -inv_inc_move inv_id.
apply H3.
Qed.
 Lemma 166 (bijection_inv_corollary) Let \alpha : A \rightarrow B be a bijection, then \alpha^{\sharp} is also
  a bijection.
Lemma bijection_inv_corollary \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 bijection\_r \ alpha \rightarrow bijection\_r \ (alpha \#).
Proof.
move: (@bijection\_inv \_ \_ alpha (alpha \#)) \Rightarrow H.
rewrite /bijection\_r/function\_r/total\_r/univalent\_r in H0.
rewrite inv\_invol in H0.
apply H.
```

## CHAPTER 6. LIBRARY FUNCTIONS\_MAPPINGS

```
apply inc\_antisym. apply H0. apply H0. apply inc\_antisym. apply H0. apply H0. Qed.
```

# Chapter 7

# Library Dedekind

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.
```

## 7.1 Dedekind formula に関する補題

```
Lemma 167 (dedekind1) Let \alpha: A \to B, \beta: B \to C and \gamma: A \to C. Then \alpha \cdot \beta \sqcap \gamma \sqsubseteq \alpha \cdot (\beta \sqcap \alpha^{\sharp} \cdot \gamma).
Lemma dedekind1  \{A \ B \ C: eqType\} \ \{alpha: Rel \ A \ B\} \ \{beta: Rel \ B \ C\} \ \{gamma: Rel \ A \ C\}: ((alpha \bullet beta) \ gamma) \ (alpha \bullet (beta \ (alpha \# \bullet gamma))).
Proof. apply (@inc\_trans \_ \_ \_ \_ \_ (@dedekind \_ \_ \_ \_ \_)). apply comp\_inc\_compat\_ab\_a'b. apply cap\_l. Qed.
```

```
Lemma 168 (dedekind2) Let \alpha: A \to B, \ \beta: B \to C \ and \ \gamma: A \to C. Then \alpha \cdot \beta \sqcap \gamma \sqsubseteq (\alpha \sqcap \gamma \cdot \beta^{\sharp}) \cdot \beta.
```

```
Lemma dedekind2 {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ B \ C} {gamma : Rel \ A \ C}: ((alpha \cdot beta) \ gamma) ((alpha \ (gamma \cdot beta \#)) \cdot beta). Proof. apply (@inc\_trans \_ \_ \_  (@dedekind \_ \_ \_ )).
```

### CHAPTER 7. LIBRARY DEDEKIND

```
apply comp\_inc\_compat\_ab\_ab'. apply cap\_l. Qed.
```

```
Lemma 169 (relation_rel_inv_rel) Let \alpha : A \rightarrow B. Then
```

$$\alpha \sqsubseteq \alpha \cdot \alpha^{\sharp} \cdot \alpha$$
.

```
Lemma relation\_rel\_inv\_rel {A \ B : eqType} {alpha : Rel \ A \ B}: alpha ((alpha \cdot alpha \#) \cdot alpha).

Proof.

move : (@dedekind1 = -alpha \ (Id \ B) \ alpha) \Rightarrow H.

rewrite comp\_id\_r \ cap\_idem \ in \ H.

apply (@inc\_trans = - - - H).

rewrite comp\_assoc.

apply comp\_inc\_compat\_ab\_ab'.

apply cap\_r.

Qed.
```

## 7.2 Dedekind formula と全関係

```
Lemma 170 (dedekind_universal1) Let \alpha : B \rightarrow C. Then
```

$$\nabla_{AC} \cdot \alpha^{\sharp} \cdot \alpha = \nabla_{AB} \cdot \alpha.$$

```
Lemma dedekind\_universal1 {A \ B \ C : eqType} {alpha : Rel \ B \ C}: ( A \ C \cdot alpha \ \#) • alpha = A \ B \cdot alpha.

Proof.

apply inc\_antisym.

apply comp\_inc\_compat\_ab\_a'b.

apply inc\_alpha\_universal.

apply (@inc\_trans\_\_\_\_(A \ B \cdot ((alpha \cdot alpha \ \#) \cdot alpha))).

apply comp\_inc\_compat\_ab\_ab'.

apply relation\_rel\_inv\_rel.

rewrite -comp\_assoc -comp\_assoc.

apply comp\_inc\_compat\_ab\_a'b.

apply comp\_inc\_compat\_ab\_a'b.

apply comp\_inc\_compat\_ab\_a'b.

apply inc\_alpha\_universal.

Qed.
```

Lemma 171 (dedekind\_universal2a, dedekind\_universal2b,

```
dedekind_universal2c) Let \alpha : A \rightarrow B and \beta : C \rightarrow B. Then
                    \nabla_{IC} \cdot \beta \sqsubseteq \nabla_{IA} \cdot \alpha \Leftrightarrow \nabla_{CC} \cdot \beta \sqsubseteq \nabla_{CA} \cdot \alpha \Leftrightarrow \beta \sqsubseteq \beta \cdot \alpha^{\sharp} \cdot \alpha.
Lemma dedekind\_universal2a {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ C \ B}:
 (i \ C \cdot beta) \ (i \ A \cdot alpha) \rightarrow (C \ C \cdot beta) \ (C \ A \cdot alpha).
Proof.
move \Rightarrow H.
rewrite -unit_universal -(@lemma_for_tarski2 C A).
rewrite comp_assoc comp_assoc.
apply (comp\_inc\_compat\_ab\_ab', H).
Qed.
Lemma dedekind_universal2b {A B C : eqType} {alpha : Rel A B} {beta : Rel C B}:
 (CC \cdot beta) \quad (CA \cdot alpha) \rightarrow beta \quad ((beta \cdot alpha \#) \cdot alpha).
Proof.
move \Rightarrow H.
apply (@inc\_trans \_ \_ \_ (beta)).
apply inc\_cap.
split.
apply inc\_reft.
apply comp\_inc\_compat\_b\_ab.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ (beta ( CA \cdot alpha))).
apply (cap\_inc\_compat\_l\ H).
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_r.
Qed.
Lemma dedekind\_universal2c {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ C \ B}:
           ((beta \cdot alpha \#) \cdot alpha) \rightarrow (i \ C \cdot beta) \ (i \ A \cdot alpha).
 beta
Proof.
move \Rightarrow H.
apply (@inc\_trans \_ \_ \_ ( i C \cdot ((beta \cdot alpha \#) \cdot alpha))).
apply (comp\_inc\_compat\_ab\_ab', H).
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
```

#### CHAPTER 7. LIBRARY DEDEKIND

 $\beta: A \rightarrow C$ . Then

```
\beta \cdot \nabla_{CI} \sqsubseteq \alpha \cdot \nabla_{BI} \Leftrightarrow \beta \cdot \nabla_{CC} \sqsubseteq \alpha \cdot \nabla_{BC} \Leftrightarrow \beta \sqsubseteq \alpha \cdot \alpha^{\sharp} \cdot \beta.
Lemma dedekind_universal3a {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
                        (alpha \cdot B i) \leftrightarrow (beta \cdot C C) \quad (alpha \cdot C C)
 (beta •
               C(i)
Proof.
split; move \Rightarrow H.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_universal inv_universal.
apply dedekind_universal2a.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_universal inv_universal.
apply dedekind_universal2c.
apply dedekind_universal2b.
apply inv_inc_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
Qed.
Lemma dedekind\_universal3b {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ A \ C}:
 (beta •
               C(i) (alpha \cdot B(i) \leftrightarrow beta) ((alpha \cdot alpha \#) \cdot beta).
Proof.
split; move \Rightarrow H.
apply inv_inc_invol.
rewrite comp_inv comp_inv -comp_assoc.
apply dedekind_universal2b.
apply dedekind_universal2a.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_invol inv_invol inv_universal inv_universal.
apply H.
apply inv\_inc\_invol.
rewrite comp_inv comp_inv inv_universal inv_universal.
apply dedekind_universal2c.
rewrite -comp_inv -comp_inv -comp_assoc.
apply inc_{-}inv.
apply H.
Qed.
```

Lemma 172 (dedekind\_universal3a, dedekind\_universal3b) Let  $\alpha : A \rightarrow B$  and

```
Lemma universal_total \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: alpha \bullet B \ i = A \ i \leftrightarrow total\_r \ alpha.

Proof.

move : (@dedekind\_universal3b\_\_\_\_ alpha \ (Id \ A)) \Rightarrow H.

rewrite comp\_id\_l \ comp\_id\_r \ in \ H.

rewrite /total\_r.

rewrite -H.

split; move \Rightarrow H0.

rewrite H0.

apply inc\_refl.

apply inc\_antisym.

apply inc\_antisym.

apply inc\_alpha\_universal.
```

Lemma 173 (universal\_total) Let  $\alpha : A \rightarrow B$ . Then

## 7.3 Dedekind formula と恒等関係

apply H0.

Qed.

```
Lemma 174 (dedekind_id1) Let \alpha : A \rightarrow A. Then
                                        \alpha \sqsubseteq id_A \Rightarrow \alpha^{\sharp} = \alpha.
Lemma dedekind\_id1 \{A: eqType\} \{alpha: Rel\ A\ A\}: alpha Id\ A \rightarrow alpha \# = alpha.
Proof.
move \Rightarrow H.
assert (alpha #
                       alpha).
move: (@dedekind1 \_ \_ \_ (alpha \#) (Id A) (Id A)) \Rightarrow H0.
rewrite comp\_id\_r comp\_id\_r inv\_invol in H0.
replace (alpha #
                        Id\ A) with (alpha\ \#) in H0.
                  alpha) with alpha in H0.
replace (Id A
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot alpha)).
apply H0.
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_inc_move.
rewrite inv_{-}id.
apply H.
rewrite cap\_comm.
apply inc\_def1.
```

### CHAPTER 7. LIBRARY DEDEKIND

```
apply H.
apply inc\_def1.
rewrite -inv\_inc\_move.
rewrite inv_id.
apply H.
apply inc\_antisym.
apply H0.
apply inv\_inc\_move.
apply H0.
Qed.
  Lemma 175 (dedekind_id2) Let \alpha : A \rightarrow A. Then
                                           \alpha \sqsubseteq id_A \Rightarrow \alpha \cdot \alpha = \alpha.
Lemma dedekind\_id2 \{A : eqType\} \{alpha : Rel A A\}:
            Id A \rightarrow alpha \cdot alpha = alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply (comp\_inc\_compat\_ab\_a\ H).
move: (dedekind\_id1 \ H) \Rightarrow H0.
apply (@inc_trans _ _ _ ((alpha • Id A)
                                                        Id\ A)).
rewrite comp_{-}id_{-}r.
apply inc\_cap.
split.
apply inc\_reft.
apply H.
apply (@inc_trans _ _ _ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite H0 comp_{-}id_{-}r.
apply cap_{-}r.
Qed.
  Lemma 176 (dedekind_id3) Let \alpha, \beta : A \rightarrow A. Then
                                  \alpha \sqsubseteq id_A \wedge \beta \sqsubseteq id_A \Rightarrow \alpha \cdot \beta = \alpha \sqcap \beta.
Lemma dedekind\_id3 {A: eqType} {alpha beta: Rel A A}:
            Id \ A \rightarrow \mathtt{beta} \quad Id \ A \rightarrow alpha \ \bullet \ \mathtt{beta} = alpha
 alpha
                                                                             beta.
Proof.
move \Rightarrow H H0.
apply inc\_antisym.
```

```
apply inc_-cap.
split.
apply (comp\_inc\_compat\_ab\_a\ H0).
apply (comp\_inc\_compat\_ab\_b\ H).
replace (alpha
                     beta) with ((alpha)
                                               beta) • (alpha
                                                                    beta)).
apply comp\_inc\_compat.
apply cap_{-}l.
apply cap_{-}r.
apply dedekind_id2.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply cap_{-}l.
Qed.
  Lemma 177 (dedekind_id4) Let \alpha, \beta : A \rightarrow A. Then
                     \alpha \sqsubseteq id_A \land \beta \sqsubseteq id_A \Rightarrow (\alpha \rhd \beta) \sqcap id_A = (\alpha \Rightarrow \beta) \sqcap id_A.
Lemma dedekind\_id4 {A : eqType} {alpha beta : Rel A A}:
           Id A \rightarrow beta \qquad Id A \rightarrow (alpha)
 alpha
                                                            Id A = (alpha \gg beta)
                                                                                             Id\ A.
                                                  beta)
Proof.
move \Rightarrow H H0.
apply inc\_lower.
move \Rightarrow qamma.
rewrite inc\_cap inc\_cap.
split; elim \Rightarrow H1 H2.
split.
rewrite inc\_rpc\ cap\_comm.
rewrite -(@dedekind_id3 _ _ _ H H2).
rewrite -(@dedekind_id1 \_ _ H).
apply inc\_residual.
apply H1.
apply H2.
split.
rewrite inc_residual (@dedekind_id1 _ _ H) (@dedekind_id3 _ _ _ H H2).
rewrite cap\_comm - inc\_rpc.
apply H1.
apply H2.
Qed.
```

# Chapter 8

# Library Rationality

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.
```

## 8.1 有理性から導かれる系

```
Lemma 178 (rationality_corollary1) Let u: A \to A and u \sqsubseteq id_A. Then, \exists R, \exists j: R \rightarrowtail A, u = j^{\sharp} \cdot j.
```

```
Lemma rationality\_corollary1 {A: eqType} {u: Rel\ A\ A}:
       Id A \to \exists (R : eqType)(j : Rel R A), injection_r j \land u = j \# \cdot j.
Proof.
move: (rationality \_ \_ u).
elim \Rightarrow R.
elim \Rightarrow f.
elim \Rightarrow g.
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2 H3.
\exists R.
\exists f.
assert (g = f).
apply (function\_inc\ H0\ H).
apply (@inc\_trans \_ \_ \_ ((f \cdot f \#) \cdot g)).
apply comp\_inc\_compat\_b\_ab.
apply H.
rewrite comp_assoc -H1.
```

### CHAPTER 8. LIBRARY RATIONALITY

Qed.

```
apply (comp\_inc\_compat\_ab\_a\ H3).
rewrite H4 in H1.
rewrite H_4 cap_idem in H_2.
split.
split.
apply H.
rewrite /univalent_r.
rewrite inv_{-}invol\ H2.
apply inc\_reft.
apply H1.
Qed.
  Lemma 179 (rationality_corollary2) Let f: A \to B be a function. Then,
                                \exists e: A \rightarrow R, \exists m: R \rightarrow B, f = e \cdot m.
Lemma rationality\_corollary2 {A B : eqType} {f : Rel A B}:
 function\_r \ f \rightarrow \exists \ (R : eqType)(e : Rel \ A \ R)(m : Rel \ R \ B), \ surjection\_r \ e \land injection\_r
m.
Proof.
elim \Rightarrow H H0.
move: (@rationality\_corollary1 \_ (f \# • f) H0).
elim \Rightarrow R.
elim \Rightarrow m.
elim \Rightarrow H1 H2.
\exists R.
\exists (f \cdot m \#).
\exists m.
split.
apply (injection_unique2 H1 (conj H H0) H2).
apply H1.
```

# Chapter 9

# Library Conjugate

```
Require Import Basic\_Notations.
Require Import Basic\_Lemmas.
Require Import Relation\_Properties.
Require Import Functions\_Mappings.
Require Import Dedekind.
```

### 9.1 共役性の定義

条件 P を満たす関係  $\alpha:A\to B$  と条件 Q を満たす関係  $\beta:A'\to B'$  が変換  $\alpha=\phi(\beta),\beta=\psi(\alpha)$  によって、1 対 1 (全射的) に対応することを、図式

$$\frac{\alpha:A \to B\ \{P\}}{\beta:A' \to B'\ \{Q\}} \ \frac{\alpha=\phi(\beta)}{\beta=\psi(\alpha)}$$

によって表す. また, Coq では以下のように表すことにする.

```
Definition conjugate
```

```
(A \ B \ C \ D : eqType) \ (P : Rel \ A \ B \to \mathsf{Prop}) \ (Q : Rel \ C \ D \to \mathsf{Prop})

(phi : Rel \ C \ D \to Rel \ A \ B) \ (psi : Rel \ A \ B \to Rel \ C \ D) :=

(\forall \ alpha : Rel \ A \ B, \ P \ alpha \to Q \ (psi \ alpha) \land phi \ (psi \ alpha) = alpha)

\land \ (\forall \ \mathsf{beta} : Rel \ C \ D, \ Q \ \mathsf{beta} \to P \ (phi \ \mathsf{beta}) \land psi \ (phi \ \mathsf{beta}) = \mathsf{beta}).
```

さらに、上の図式において条件 P または Q が不要な場合には、以下の  ${\tt True\_r}$  を代入する.

Definition  $True_r \{A \ B : eqType\} := fun_r : Rel \ A \ B \Rightarrow True.$ 

### 9.2 共役の例

Lemma 180 (inv\_conjugate) Inverse relation (\*) makes conjugate. That is,

$$\frac{\alpha: A \to B}{\beta: B \to A} \frac{\alpha = \beta^{\sharp}}{\beta = \alpha^{\sharp}}.$$

```
Lemma inv\_conjugate \{A \ B : eqType\}: \\ conjugate \ A \ B \ B \ A \ True\_r \ True\_r \ (@inverse \_ \_) \ (@inverse \_ \_).

Proof.

split.

move \Rightarrow alpha \ H.

split.

by [].

apply inv\_invol.

move \Rightarrow beta H.

split.

by [].

apply inv\_invol.

Qed.
```

**Lemma 181 (injection\_conjugate)** Let  $j: C \rightarrow B$  be an injection. Then,

$$\frac{f:A\to B\ \{f^{\sharp}\cdot f\sqsubseteq j^{\sharp}\cdot j\}}{h:A\to C}\ \frac{f=h\cdot j}{h=f\cdot j^{\sharp}}$$

```
Lemma injection\_conjugate \{A \ B \ C : eqType\} \{j : Rel \ C \ B\}:
 injection_r j \rightarrow
 conjugate A \ B \ A \ C \ (\mathbf{fun} \ f : Rel \ A \ B \Rightarrow ((f \# \bullet f) \ (j \# \bullet j)) \land function\_r \ f)
 (\mathbf{fun}\ h: Rel\ A\ C \Rightarrow function\_r\ h)\ (\mathbf{fun}\ h: Rel\ A\ C \Rightarrow h\ \boldsymbol{\cdot}\ j)\ (\mathbf{fun}\ f: Rel\ A\ B \Rightarrow f\ \boldsymbol{\cdot}
j \#).
Proof.
elim.
elim \Rightarrow H H0 H1.
split.
move \Rightarrow alpha.
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
assert (function_r (alpha \cdot j \#)).
split.
apply (@inc\_trans \_ \_ \_ \_ H3).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ j).
```

```
apply (@inc\_trans \_ \_ \_ (alpha \cdot ((alpha \# \cdot alpha) \cdot alpha \#))).
rewrite comp_assoc -comp_assoc.
apply (comp\_inc\_compat\_a\_ab\ H3).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_a'b H2).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_ab\_b.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply H\theta.
split.
apply H5.
apply function_inc.
apply function\_comp.
apply H5.
split.
apply H.
apply H0.
split.
apply H3.
apply H_4.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
move \Rightarrow beta.
elim \Rightarrow H2 H3.
assert (function_r (beta \cdot j)).
split.
apply (@inc\_trans \_ \_ \_ \_ H2).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ j).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
split.
split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ _ beta).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H3).
```

```
apply H_4.
rewrite comp\_assoc.
replace (j \cdot j \#) with (Id \ C).
apply comp_{-}id_{-}r.
apply inc\_antisym.
apply H.
rewrite /univalent_r in H1.
rewrite inv_{-}invol in H1.
apply H1.
Qed.
  Lemma 182 (injection_conjugate_corollary1, injection_conjugate_corollary2)
  Let j: C \rightarrow B be an injection and f: A \rightarrow B be a function. Then,
              f^{\sharp} \cdot f \sqsubseteq j^{\sharp} \cdot j \Leftrightarrow (\exists! h : A \to C, f = h \cdot j) \Leftrightarrow (\exists h' : A \to C, f \sqsubseteq h' \cdot j).
Lemma injection\_conjugate\_corollary1 \{A B C : eqType\} \{f : Rel A B\} \{j : Rel C B\}:
 injection_r j \rightarrow function_r f \rightarrow
 ((f \# \cdot f) \ (j \# \cdot j) \leftrightarrow \exists ! \ h : Rel \ A \ C, function\_r \ h \land f = h \cdot j).
Proof.
move \Rightarrow H H0.
move: (@injection\_conjugate\ A\_\_\_\ H).
elim \Rightarrow H1 H2.
split; move \Rightarrow H3.
\exists (f \cdot j \#).
split.
move: (H1 f (conj H3 H0)).
elim \Rightarrow H4 H5.
split.
apply H_4.
by [rewrite H5].
move \Rightarrow h.
elim \Rightarrow H4 H5.
rewrite H5 comp\_assoc.
replace (j \cdot j \#) with (Id \ C).
apply comp_{-}id_{-}r.
rewrite /injection_r/function_r/univalent_r in H.
rewrite inv\_invol in H.
apply inc\_antisym.
apply H.
apply H.
elim H3 \Rightarrow h.
elim.
```

```
elim \Rightarrow H4 \ H5 \ H6.
rewrite H5 comp_inv comp_assoc -(@comp_assoc _ _ _ _ h).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_b.
apply H_4.
Qed.
Lemma injection\_conjugate\_corollary2 \{A \ B \ C : eqType\} \{f : Rel \ A \ B\} \{j : Rel \ C \ B\}:
 injection\_r \ j \rightarrow function\_r \ f \rightarrow
 ((f \# \cdot f) \quad (j \# \cdot j) \leftrightarrow \exists h' : Rel \land C, f \quad (h' \cdot j)).
Proof.
move \Rightarrow H H0.
split; move \Rightarrow H1.
apply (injection_conjugate_corollary1 H H0) in H1.
elim H1 \Rightarrow h.
elim.
elim \Rightarrow H2 \ H3 \ H4.
\exists h.
rewrite H3.
apply inc\_reft.
elim H1 \Rightarrow h' H2.
replace (f \# \cdot f) with (f \# \cdot (f (h' \cdot j))).
apply (@inc\_trans \_ \_ \_ ((f \# \cdot f) \cdot (j \# \cdot j))).
rewrite comp\_assoc\ cap\_comm\ -(@comp\_assoc\ \_\ \_\ \_\ f).
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_r.
apply comp\_inc\_compat\_ab\_b.
apply H0.
apply f_equal.
apply inc_-def1 in H2.
by [rewrite -H2].
Qed.
```

Lemma 183 (surjection\_conjugate) Let  $e: A \rightarrow C$  be a surjection. Then,

$$\frac{f:A\to B\ \{e\cdot e^\sharp\sqsubseteq f\cdot f^\sharp\}}{h:C\to B}\ \frac{f=e\cdot h}{h=e^\sharp\cdot f}$$

```
Lemma surjection_conjugate {A B C : eqType} {e : Rel A C}: surjection_r e \rightarrow conjugate A B C B (fun f : Rel A B \Rightarrow ((e • e #) (f • f #)) \land function_r f)
```

```
(\operatorname{fun} h : Rel \ C \ B \Rightarrow function\_r \ h) \ (\operatorname{fun} h : Rel \ C \ B \Rightarrow e \ {}^{\bullet} \ h) \ (\operatorname{fun} f : Rel \ A \ B \Rightarrow e \ \#)
• f).
Proof.
elim.
elim \Rightarrow H H0 H1.
split.
move \Rightarrow alpha.
elim \Rightarrow H2.
elim \Rightarrow H3 H4.
assert (function_r (e \# \bullet alpha)).
split.
apply @inc\_trans \_ \_ \_ \_ H1).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H3).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H_4).
rewrite comp_inv inv_invol comp_assoc -(@comp_assoc _ _ _ e).
apply (@inc\_trans \_ \_ \_ (alpha # \cdot ((alpha \cdot alpha #) \cdot alpha))).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_a'b H2).
rewrite comp_assoc -comp_assoc.
apply (comp\_inc\_compat\_ab\_a\ H_4).
split.
apply H5.
apply Logic.eq_sym.
apply function_inc.
split.
apply H3.
apply H_4.
apply function_comp.
split.
apply H.
apply H0.
apply H5.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H.
move \Rightarrow beta.
elim \Rightarrow H2 H3.
assert (function_r (e \cdot beta)).
split.
apply @inc\_trans \_ \_ \_ \_ H).
```

```
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H2).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H3).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ e).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H0).
split.
split.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_b\_ab\ H2).
apply H_4.
rewrite -comp_-assoc.
replace (e \# \cdot e) with (Id \ C).
apply comp_{-}id_{-}l.
apply inc\_antisym.
rewrite /total_r in H1.
rewrite inv_-invol in H1.
apply H1.
apply H0.
Qed.
  Lemma 184 (surjection_conjugate_corollary) Let e: A \rightarrow C be a surjection and
  f: A \to B be a function. Then,
                            e \cdot e^{\sharp} \sqsubseteq f \cdot f^{\sharp} \Leftrightarrow (\exists! h : C \to B, f = e \cdot h).
Lemma surjection\_conjugate\_corollary \{A B C : eqType\} \{f : Rel A B\} \{e : Rel A C\}:
 surjection\_r \ e \rightarrow function\_r \ f \rightarrow
               (f \cdot f \#) \leftrightarrow \exists ! \ h : Rel \ C \ B, function\_r \ h \land f = e \cdot h).
 ((e \cdot e \#)
Proof.
move \Rightarrow H H0.
move: (@surjection\_conjugate \_ B \_ \_ H).
elim \Rightarrow H1 H2.
split; move \Rightarrow H3.
\exists (e \# \cdot f).
split.
move: (H1 f (conj H3 H0)).
elim \Rightarrow H4 H5.
split.
apply H_4.
by [rewrite H5].
```

```
move \Rightarrow h.
elim \Rightarrow H4 H5.
rewrite H5 -comp\_assoc.
replace (e \# \cdot e) with (Id \ C).
apply comp_{-}id_{-}l.
rewrite /surjection_r/function_r/total_r in H.
rewrite inv_invol in H.
apply inc\_antisym.
apply H.
apply H.
elim H3 \Rightarrow h.
elim.
elim \Rightarrow H4 \ H5 \ H6.
rewrite H5 comp_inv comp_assoc -(@comp_assoc _ _ _ h).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply H_4.
Qed.
```

**Lemma 185 (subid\_conjugate)** Subidentity  $u \sqsubseteq id_A$  corresponds  $\rho: I \rightarrow A$ . That is,

$$\frac{\rho: I \to A}{u: A \to A \ \{u \sqsubseteq id_A\}} \ \frac{\rho = \nabla_{IA} \cdot u}{u = id_A \sqcap \nabla_{AI} \cdot \rho}.$$

```
Lemma subid\_conjugate \{A : eqType\}:
 conjugate i A A A True_r (fun u : Rel A A \Rightarrow u Id A)
 (fun u : Rel \ A \ A \Rightarrow i \ A \cdot u) (fun rho : Rel \ i \ A \Rightarrow Id \ A ( A \ i \cdot rho)).
Proof.
split.
move \Rightarrow alpha H.
split.
apply cap_{-}l.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ( i A \cdot ( A i \cdot alpha))).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_r.
rewrite -comp_-assoc.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
rewrite -(@inv\_universal\ i\ A).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (dedekind1)).
rewrite comp_id_r cap_comm cap_universal.
```

```
apply inc_refl.
move \Rightarrow beta H.
split.
by [].
apply inc\_antisym.
rewrite cap_comm -comp_assoc lemma_for_tarski2.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite comp_id_l cap_comm cap_universal.
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_inc_move inv_id.
apply H.
apply inc_-cap.
split.
apply H.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_b\_ab.
rewrite lemma\_for\_tarski2.
apply inc\_alpha\_universal.
Qed.
```

Lemma 186 (subid\_conjugate\_corollary1) Let  $u, v : A \rightarrow A$  and  $u, v \sqsubseteq id_A$ . Then,

```
\nabla_{IA} \cdot u = \nabla_{IA} \cdot v \Rightarrow u = v.
```

```
Lemma subid\_conjugate\_corollary1 \{A : eqType\} \{u \ v : Rel \ A \ A\}:
                                   i A \cdot u = i A \cdot v \rightarrow u = v.
       Id A \rightarrow v
                       Id A \rightarrow
 u
Proof.
move \Rightarrow H H0 H1.
move: (@subid\_conjugate\ A).
elim \Rightarrow H2 H3.
move: (H3 \ u \ H).
elim \Rightarrow H4 H5.
rewrite -H5.
move: (H3 \ v \ H0).
elim \Rightarrow H6 H7.
rewrite -H?.
apply f_equal.
apply f_equal.
apply H1.
Qed.
```

Lemma 187 (subid\_conjugate\_corollary2) Let  $\rho, \rho' : I \to A$ . Then,

$$id_A \sqcap \nabla_{AI} \cdot \rho = id_A \sqcap \nabla_{AI} \cdot \rho' \Rightarrow \rho = \rho'.$$

```
Lemma subid\_conjugate\_corollary2\ \{A: eqType\}\ \{rho\ rho': Rel\ i\ A\}:\ Id\ A\quad (\quad A\ i\quad rho)=Id\ A\quad (\quad A\ i\quad rho')\rightarrow rho=rho'. Proof.

move \Rightarrow H.

move: (@subid\_conjugate\ A).

elim \Rightarrow H0\ H1.

move: (H0\ rho\ I).

elim \Rightarrow H2\ H3.

rewrite-H3.

move: (H0\ rho'\ I).

elim \Rightarrow H4\ H5.

rewrite-H5.

apply f_equal.

apply H.

Qed.
```

## Chapter 10

# Library Domain

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.

Require Import Dedekind.

Require Import Logic.FunctionalExtensionality.
```

## 10.1 定義域の定義

関係  $\alpha: A \rightarrow B$  に対して、その定義域 (関係)  $\lfloor \alpha \rfloor: A \rightarrow A$  は、

$$\lfloor \alpha \rfloor = \alpha \cdot \alpha^{\sharp} \sqcap id_A$$

で表される. また、Coq では以下のように表すことにする.

**Definition** domain  $\{A \ B : eqType\}$   $(alpha : Rel \ A \ B) := (alpha \cdot alpha \#)$   $Id \ A.$ 

## 10.2 定義域の性質

### 10.2.1 基本的な性質

**Lemma 188 (domain\_another\_def)** Let  $\alpha : A \rightarrow B$ . Then,

$$|\alpha| = \alpha \cdot \nabla_{BA} \cap id_A.$$

Lemma  $domain\_another\_def$  { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }:  $domain \ alpha = (alpha \cdot B \ A) \quad Id \ A.$ Proof.

```
apply inc\_antisym.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply inc\_cap.
split.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm comp_id_r cap_universal.
apply inc\_reft.
apply cap_r.
Qed.
  Lemma 189 (domain_inv) Let \alpha : A \rightarrow B. Then,
                                            |\alpha|^{\sharp} = |\alpha|.
Lemma domain\_inv \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 (domain \ alpha) \# = domain \ alpha.
Proof.
apply dedekind_id1.
apply cap_{-}r.
Qed.
  Lemma 190 (domain_comp_alpha1, domain_comp_alpha2) Let \alpha : A \rightarrow B.
  Then,
                                   |\alpha| \cdot \alpha = \alpha \wedge \alpha^{\sharp} \cdot |\alpha| = \alpha^{\sharp}.
Lemma domain\_comp\_alpha1 {A B : eqType} {alpha : Rel A B}:
 (domain \ alpha) \cdot alpha = alpha.
Proof.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_r.
rewrite / domain.
rewrite cap\_comm.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (dedekind2)).
rewrite comp\_id\_l cap\_idem.
apply inc\_reft.
Qed.
```

Lemma  $domain\_comp\_alpha2$  {A B : eqType} {alpha : Rel A B}:

 $alpha \# \bullet (domain \ alpha) = alpha \#.$ 

```
Proof.
rewrite -domain_inv -comp_inv.
apply f_equal.
apply domain_comp_alpha1.
Qed.
```

```
Lemma 191 (domain_inc_compat) Let \alpha, \alpha' : A \rightarrow B. Then,
```

$$\alpha \sqsubseteq \alpha' \Rightarrow |\alpha| \sqsubseteq |\alpha'|.$$

```
Lemma domain\_inc\_compat {A \ B : eqType} {alpha \ alpha' : Rel \ A \ B}: alpha \ alpha' \rightarrow domain \ alpha \ domain \ alpha'.

Proof.

move \Rightarrow H.

apply cap\_inc\_compat\_r.

apply comp\_inc\_compat.

apply H.

apply (@inc\_inv \_ \_ \_ \_ H).

Qed.
```

### Lemma 192 (domain\_total) Let $\alpha : A \rightarrow B$ . Then,

" $\alpha$  is total"  $\Leftrightarrow \lfloor \alpha \rfloor = id_A$ .

```
Lemma domain\_total \{A \ B : eqType\} \{alpha : Rel \ A \ B\}: total\_r \ alpha \leftrightarrow domain \ alpha = Id \ A.
```

#### Proof.

```
split; move \Rightarrow H.

rewrite /domain.

rewrite cap\_comm.

apply Logic.eq\_sym.

apply inc\_def1.

apply H.

apply inc\_def1.

rewrite /domain in H.

by [rewrite cap\_comm H].

Qed.
```

Lemma 193 (domain\_inc\_id) Let  $u : A \rightarrow A$ . Then,

 $u \sqsubseteq id_A \Leftrightarrow |u| = u$ .

Lemma  $domain\_inc\_id$   $\{A: eqType\}$   $\{u: Rel\ A\ A\}: u \ Id\ A \leftrightarrow domain\ u=u.$ 

```
Proof. split; move \Rightarrow H. rewrite /domain. rewrite (dedekind\_id1\ H)\ (dedekind\_id2\ H). apply inc\_def1 in H. by [rewrite -H]. rewrite -H. apply cap\_r. Qed.
```

### 10.2.2 合成と定義域

```
Lemma 194 (comp_domain1, comp_domain2) Let \alpha : A \rightarrow B and \beta : B \rightarrow C.
  Then,
                                  |\alpha \cdot \beta| = |\alpha \cdot |\beta| |\sqsubseteq |\alpha|.
Lemma comp\_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 domain (alpha • beta)
                              domain alpha.
Proof.
rewrite / domain.
rewrite comp_{-}inv.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot (beta + alpha \#)))
                                                                        alpha \#))
                                                                                        Id\ A)).
replace (((alpha \cdot beta) \cdot (beta \# \cdot alpha \#)) Id A) with ((((alpha \cdot beta) \cdot
(\mathtt{beta} \ \# \ \bullet \ alpha \ \#)) \qquad Id \ A)
                                    Id\ A).
apply cap\_inc\_compat\_r.
rewrite comp_{-}assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite comp_{-}id_{-}r.
apply inc\_reft.
by [rewrite cap_assoc cap_idem].
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}r.
Qed.
Lemma comp\_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 domain (alpha \cdot beta) = domain (alpha \cdot domain beta).
Proof.
apply inc\_antisym.
replace (domain (alpha • beta)) with (domain ((alpha • domain beta) • beta)).
apply comp\_domain1.
by [rewrite comp_assoc domain_comp_alpha1].
apply (@inc\_trans \_ \_ \_ (domain (alpha \cdot (beta \cdot beta \#)))).
```

```
CHAPTER 10. LIBRARY DOMAIN
apply domain\_inc\_compat.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
rewrite -comp\_assoc.
apply comp_domain1.
Qed.
  Lemma 195 (comp_domain3) Let \alpha : A \rightarrow B be a relation and \beta : B \rightarrow C be a total
  relation. Then,
                                           |\alpha \cdot \beta| = |\alpha|.
Lemma comp\_domain3 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total\_r \text{ beta} \rightarrow domain (alpha \cdot \text{beta}) = domain alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply comp\_domain1.
rewrite / domain.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ beta).
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
Qed.
  Lemma 196 (comp_domain4) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                  |\alpha^{\sharp}| \sqsubseteq |\beta| \Rightarrow |\alpha \cdot \beta| = |\alpha|.
Lemma comp\_domain4 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 domain (alpha \#)
                         domain \ \mathsf{beta} \to domain \ (alpha \cdot \mathsf{beta}) = domain \ alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply comp_domain1.
rewrite / domain.
rewrite -(@domain_comp_alpha1 _ _ (alpha #)) comp_inv comp_assoc -(@comp_assoc _ _
_ _ beta).
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
```

apply  $(@inc\_trans \_ \_ \_ \_ H)$ .

apply  $cap_{-}l$ .

Qed.

```
Lemma 197 (comp_domain5) Let \alpha : A \rightarrow B be a univalent relation and \beta : B \rightarrow C.
  Then,
                                   |\alpha^{\sharp}| \sqsubset |\beta| \Leftrightarrow |\alpha \cdot \beta| = |\alpha|.
Lemma comp_domain5 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow
 (domain (alpha \#)
                           domain beta \leftrightarrow domain (alpha \cdot beta) = domain alpha).
Proof.
move \Rightarrow H.
split; move \Rightarrow H0.
apply (comp\_domain \not\downarrow H0).
rewrite /domain.
rewrite inv_invol.
apply cap\_inc\_compat\_r.
replace (alpha \# \cdot alpha) with (alpha \# \cdot (domain (alpha \cdot beta) \cdot alpha)).
rewrite /domain.
rewrite comp_{-}inv.
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot (((alpha \cdot beta) \cdot (beta \# \cdot alpha \#)) \cdot alpha))).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp_assoc comp_assoc -comp_assoc -(@comp_assoc _ _ _ beta).
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_b H)).
apply (comp\_inc\_compat\_ab\_a\ H).
by [rewrite H0 domain_comp_alpha1].
Qed.
  Lemma 198 (comp_domain6) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                       \alpha \cdot |\beta| \sqsubseteq |\alpha \cdot \beta| \cdot \alpha.
Lemma comp_domain6 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 (alpha • domain beta) (domain (alpha • beta) • alpha).
Proof.
apply (@inc\_trans \_ \_ \_ \_ (@comp\_cap\_distr\_l \_ \_ \_ \_ )).
rewrite cap\_comm.
replace (alpha \cdot Id B) with (Id A \cdot alpha).
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite cap_comm -comp_assoc comp_assoc -comp_inv.
apply inc\_reft.
by [rewrite comp\_id\_l comp\_id\_r].
Qed.
```

```
Lemma 199 (comp_domain7) Let \alpha : A \rightarrow B be a univalent relation and \beta : B \rightarrow C.
  Then,
                                          \alpha \cdot |\beta| = |\alpha \cdot \beta| \cdot \alpha.
Lemma comp_domain7 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 univalent_r \ alpha \rightarrow alpha \cdot domain \ \mathsf{beta} = domain \ (alpha \cdot \mathsf{beta}) \cdot alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply comp\_domain6.
apply (@inc\_trans \_ \_ \_ \_ \_ (@comp\_cap\_distr\_r \_ \_ \_ \_ \_)).
rewrite comp_id_l comp_inv comp_assoc comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
apply (fun H' \Rightarrow cap\_inc\_compat \ H' \ H).
rewrite comp_assoc -comp_assoc.
apply (comp\_inc\_compat\_ab\_a\ H).
Qed.
  Lemma 200 (comp_domain8) Let u: A \rightarrow A, \alpha: A \rightarrow B and u \sqsubseteq id_A. Then,
                                            |u \cdot \alpha| = u \cdot |\alpha|.
Lemma comp\_domain8 \{A \ B : eqType\} \{u : Rel \ A \ A\} \{alpha : Rel \ A \ B\}:
       Id A \rightarrow domain (u \cdot alpha) = u \cdot domain alpha.
```

```
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite -(@cap\_idem \_ \_ (domain (u \cdot alpha))).
rewrite (dedekind_id3 H).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_domain1)).
apply domain\_inc\_id in H.
rewrite H.
apply inc_refl.
apply domain_inc_compat.
apply (comp\_inc\_compat\_ab\_b\ H).
apply cap_{-}r.
apply (@inc\_trans \_ \_ \_ \_ (comp\_domain6)).
apply (comp\_inc\_compat\_ab\_a\ H).
Qed.
```

### 10.2.3 その他の性質

```
Lemma 201 (cap_domain) Let \alpha, \alpha' : A \rightarrow B. Then,
                                      |\alpha \cap \alpha'| = \alpha \cdot \alpha'^{\sharp} \cap id_A.
Lemma cap\_domain \{A \ B : eqType\} \{alpha \ alpha' : Rel \ A \ B\}:
 domain (alpha
                      alpha') = (alpha \cdot alpha' \#)
Proof.
apply inc\_antisym.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat.
apply cap_{-}l.
apply inc_{-}inv.
apply cap_{-}r.
rewrite -(@cap_idem _ _ (Id A)) -cap_assoc.
apply cap\_inc\_compat\_r.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite inv_invol comp_id_l comp_id_r -inv_cap_distr (@cap_comm _ _ alpha').
apply inc\_reft.
Qed.
  Lemma 202 (cupP_domain_distr, cup_domain_distr) Let f: (C \rightarrow D) \rightarrow (A \rightarrow D)
  B) and P: predicate. Then,
                                    |\sqcup_{P(\alpha)} f(\alpha)| = \sqcup_{P(\alpha)} |f(\alpha)|.
Lemma cupP\_domain\_distr {A B C D : eqType} {f : Rel\ C\ D \rightarrow Rel\ A\ B} {P : Rel\ C\ D
\rightarrow Prop:
 domain ( _{\{P\}} f) = _{\{P\}} (fun \ alpha : Rel \ C \ D \Rightarrow domain \ (f \ alpha)).
Proof.
rewrite / domain.
rewrite inv_cupP_distr comp_cupP_distr_l cap_cupP_distr_r.
apply cupP_-eq.
move \Rightarrow alpha H.
rewrite -cap\_domain - cap\_domain.
apply f_equal.
rewrite cap\_idem.
apply inc\_antisym.
apply cap_r.
apply inc_-cap.
split.
```

```
move: alpha H.
apply inc\_cupP.
apply inc\_reft.
apply inc\_reft.
Qed.
Lemma cup\_domain\_distr \{A \ B : eqType\} \{alpha \ alpha' : Rel \ A \ B\}:
 domain (alpha
                     alpha') = domain \ alpha \ domain \ alpha'.
Proof.
rewrite cup\_to\_cupP (@cup\_to\_cupP _ _ _ _ id).
apply cupP\_domain\_distr.
Qed.
  Lemma 203 (domain_universal1) Let \alpha : A \rightarrow B. Then,
                                      |\alpha| \cdot \nabla_{AC} = \alpha \cdot \nabla_{BC}.
Lemma domain\_universal1 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\}:
 domain alpha \cdot A C = alpha \cdot B C.
Proof.
apply inc\_antisym.
apply @inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot A C)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ ((domain alpha \cdot alpha) \cdot B C)).
rewrite domain_comp_alpha1.
apply inc_refl.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
  Lemma 204 (domain_universal2) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then,
                                     \alpha \cdot |\beta| = \alpha \sqcap \nabla_{AC} \cdot \beta^{\sharp}.
Lemma domain\_universal2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 alpha \cdot domain beta = alpha \quad (A C \cdot beta \#).
Proof.
apply inc\_antisym.
apply inc\_cap.
```

domain alpha

```
split.
apply comp\_inc\_compat\_ab\_a.
apply cap_{-}r.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite -comp_-assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
rewrite -inv_universal -comp_inv -domain_universal1.
rewrite comp_inv inv_universal domain_inv cap_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite cap_comm cap_universal domain_inv.
apply comp\_inc\_compat\_ab\_a.
apply cap_r.
Qed.
  Lemma 205 (domain_lemma1) Let \alpha, \beta : A \rightarrow B and \beta is univalent. Then,
                                    \alpha \sqsubseteq \beta \land |\alpha| = |\beta| \Rightarrow \alpha = \beta.
Lemma domain\_lemma1 \{A B : eqType\} \{alpha beta : Rel A B\}:
                                  beta \rightarrow domain \ alpha = domain \ beta \rightarrow alpha = beta.
 univalent_r beta \rightarrow alpha
Proof.
move \Rightarrow H H0 H1.
apply inc\_antisym.
apply H0.
rewrite -(@domain_comp_alpha1 _ _ beta) -H1.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc_inv_{-} - H0).
Qed.
  Lemma 206 (domain_lemma2a, domain_lemma2b) Let \alpha : A \rightarrow B and \beta : A \rightarrow B
  C. Then,
                        |\alpha| \subseteq |\beta| \Leftrightarrow \alpha \cdot \nabla_{BB} \subseteq \beta \cdot \nabla_{CB} \Leftrightarrow \alpha \subseteq \beta \cdot \beta^{\sharp} \cdot \alpha.
Lemma domain\_lemma2a \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
```

(beta •


 $domain beta \leftrightarrow (alpha \cdot B B)$ 

```
Proof.
split; move \Rightarrow H.
rewrite -(@domain_comp_alpha1 _ _ alpha) comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_a'b H)).
apply (@inc\_trans \_ \_ \_ \_ (comp\_inc\_compat\_ab\_a'b (cap\_l))).
rewrite comp_{-}assoc.
apply comp_inc_compat_ab_ab'.
apply inc\_alpha\_universal.
apply (@inc\_trans \_ \_ \_ (domain ((beta • beta #) • alpha))).
apply domain_inc_compat.
apply (@inc_trans _ _ _ (alpha
                                    (beta • C(B))).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cap\_inc\_compat\_l H)).
                   (alpha \cdot B B)) with ((alpha \cdot Id B) \quad (alpha \cdot
                                                                                B(B)).
replace (alpha
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (comp\_cap\_distr\_l)).
rewrite cap_universal comp_id_r.
apply inc\_reft.
by [rewrite comp_{-}id_{-}r].
rewrite cap\_comm\ comp\_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite cap_comm cap_universal.
apply inc_refl.
rewrite comp_assoc.
apply comp_domain1.
Qed.
Lemma domain\_lemma2b {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta : Rel \ A \ C}:
                   domain \ beta \leftrightarrow alpha \quad ((beta \cdot beta \#) \cdot alpha).
 domain alpha
Proof.
split; move \Rightarrow H.
apply domain\_lemma2a in H.
apply (@inc\_trans \_ \_ \_ (alpha (beta \cdot CB))).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cap\_inc\_compat\_l H)).
                   (alpha \cdot B B)) with ((alpha \cdot Id B))
                                                                 (alpha \cdot B B).
replace (alpha
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (comp\_cap\_distr\_l)).
rewrite cap_universal comp_id_r.
apply inc_refl.
by [rewrite comp_{-}id_{-}r].
rewrite cap_comm comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
rewrite cap_comm cap_universal.
apply inc_refl.
apply domain\_inc\_compat in H.
apply @inc\_trans \_ \_ \_ \_ H).
```

```
rewrite comp_assoc.
apply comp_domain1.
Qed.
```

### Lemma 207 (domain\_corollary1) In below figure,


"\alpha and \beta are total" \land \alpha^\pm \cdot \beta \subseteq \rho^\pm \cdot \tau \infty \subseteq \alpha^\pm \cdot \ta \subseteq \subseteq \alpha \cdot \rho^\pm \subseteq \alpha \cdot \rho^\pm \subseteq \alpha \cdot \tau^\pm is total".



```
Lemma domain\_corollary1 \{X \ Y \ Z \ T : eqType\}
 \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ X \ Z\} \ \{rho : Rel \ T \ Y\} \ \{tau : Rel \ T \ Z\}:
 total\_r \ alpha \rightarrow total\_r \ \mathsf{beta} \rightarrow (alpha \ \# \ \bullet \ \mathsf{beta}) \qquad (rho \ \# \ \bullet \ tau) \rightarrow
 total\_r ((alpha \cdot rho \#) (beta \cdot tau \#)).
Proof.
move \Rightarrow H H0 H1.
move: (comp\_inc\_compat\ H\ H0) \Rightarrow H2.
rewrite comp\_id\_l -comp\_assoc (@comp\_assoc _ _ _ alpha) in H2.
rewrite /total_r.
replace (Id\ X) with (((alpha \cdot (rho \# \cdot tau)) \cdot beta \#)
                                                                         Id\ X).
rewrite -comp_assoc comp_assoc.
apply (@inc_trans _ _ _ _ (@dedekind _ _ _ _ _)).
rewrite comp_id_l comp_id_r comp_inv comp_inv inv_invol inv_invol.
rewrite inv_cap_distr comp_inv comp_inv inv_invol inv_invol (@cap_comm _ _ (tau •
beta #)).
apply inc\_reft.
apply Logic.eq_sym.
rewrite cap\_comm.
apply inc\_def1.
apply (@inc\_trans \_ \_ \_ \_ H2).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp\_inc\_compat\_ab\_ab' H1).
Qed.
```

### Lemma 208 (domain\_corollary2) In below figure,

" $\alpha$  and  $\beta$  are univalent"  $\wedge \rho \cdot \rho^{\sharp} \sqcap \tau \cdot \tau^{\sharp} = id_T \Rightarrow "\alpha \cdot \rho^{\sharp} \sqcap \beta \cdot \tau^{\sharp}$  is univalent".



```
Lemma domain\_corollary2 \{X \ Y \ Z \ T : eqType\}
 \{alpha: Rel\ X\ Y\}\ \{beta: Rel\ X\ Z\}\ \{rho: Rel\ T\ Y\}\ \{tau: Rel\ T\ Z\}:
 univalent_r \ alpha \rightarrow univalent_r \ \mathsf{beta} \rightarrow (rho \ \cdot \ rho \ \#) \ (tau \ \cdot \ tau \ \#) = Id \ T \rightarrow
 univalent_r ((alpha \cdot rho \#) (beta \cdot tau \#)).
Proof.
move \Rightarrow H H0 H1.
rewrite /univalent_r.
rewrite -H1 inv_cap_distr.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_inv inv_invol -comp_assoc (@comp_assoc _ _ _ rho).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp\_inc\_compat\_ab\_a\ H).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
rewrite comp_inv inv_invol -comp_assoc (@comp_assoc _ _ _ tau).
apply comp\_inc\_compat\_ab\_a'b.
apply (comp\_inc\_compat\_ab\_a\ H0).
Qed.
```

### 10.2.4 矩形関係

$$\alpha: A \rightarrow B$$
  $\not\!\! m$ 

$$\alpha \cdot \nabla_{BA} \cdot \alpha \sqsubseteq \alpha$$

を満たすとき,  $\alpha$  は 矩形関係 (rectangular relation) であると言われる.

```
Definition rectangular \{A \ B : eqType\} (alpha : Rel A B):= ((alpha \cdot B \ A) \cdot alpha) alpha.
```

**Lemma 209 (rectangular\_inv)** Let  $\alpha : A \to B$  is a rectangular relation, then  $\alpha^{\sharp}$  is also a rectangular relation.

```
Lemma rectangular_inv \{A B : eqType\} \{alpha : Rel A B\}:
    rectangular\ alpha \rightarrow rectangular\ (alpha\ \#).
Proof.
move \Rightarrow H.
apply inv\_inc\_move.
rewrite comp_inv comp_inv inv_invol inv_universal -comp_assoc.
apply H.
Qed.
      Lemma 210 (rectangular_capP, rectangular_cap) Let f(\alpha) is always a rectangu-
       lar relation and P: predicate, then \sqcap_{P(\beta)} f(\beta) is also a rectangular relation.
Lemma rectangular_capP \{A \ B \ C \ D : eqType\} \{f : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C \ D \rightarrow Rel \ A \ B\} \{P : Rel \ C
   (\forall alpha : Rel \ C \ D, P \ alpha \rightarrow rectangular \ (f \ alpha)) \rightarrow rectangular \ ( \ _{\{P\}} \ f).
Proof.
move \Rightarrow H.
rewrite / rectangular.
apply (@inc_trans _ _ _ ( _{-}\{P\}\ (\mathbf{fun}\ alpha: Rel\ C\ D \Rightarrow (f\ alpha \cdot B\ A) \cdot f\ alpha))).
apply (@inc\_trans \_ \_ \_ \_ (comp\_capP\_distr\_l)).
apply inc\_capP.
move \Rightarrow beta H\theta.
\texttt{apply} \ (@inc\_trans \ \_ \ \_ \ \_ \ ((( \ \ \_\{P\} \ f) \ \bullet \ \ B \ A) \ \bullet \ f \ \texttt{beta})).
move : beta H0.
apply inc\_capP.
apply inc\_reft.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a'b.
move: H\theta.
apply inc\_capP.
apply inc\_reft.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (H beta H\theta)).
move: beta H0.
apply inc\_capP.
apply inc\_reft.
Qed.
Lemma rectangular_cap {A B : eqType} {alpha beta : Rel A B}:
```

```
rectangular\ alpha 
ightarrow rectangular\ beta 
ightarrow rectangular\ (alpha
Proof.
move \Rightarrow H H0.
rewrite (@cap\_to\_capP\_\_\_\_\_id).
apply rectangular_capP.
move \Rightarrow qamma.
case \Rightarrow H1; rewrite H1.
apply H.
apply H\theta.
Qed.
  Lemma 211 (rectangular_comp) Let \alpha : A \rightarrow B, \beta : B \rightarrow C and \alpha or \beta is a
  rectangular relation, then \alpha \cdot \beta is also a rectangular relation.
Lemma rectangular\_comp {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 rectangular\ alpha\ \lor\ rectangular\ beta \to rectangular\ (alpha\ •\ beta).
Proof.
rewrite / rectangular.
case; move \Rightarrow H.
rewrite - comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_-assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
rewrite comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
rewrite -comp\_assoc -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
Qed.
  Lemma 212 (rectangular_unit) Let \alpha : A \rightarrow B. Then,
                     "\alpha is rectangular" \Leftrightarrow \exists \mu : I \to A, \exists \rho : I \to B, \alpha = \rho^{\sharp} \cdot \mu.
Lemma rectangular\_unit \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
 rectangular \ alpha \leftrightarrow \exists \ (mu : Rel \ i \ A)(rho : Rel \ i \ B), \ alpha = mu \ \# \cdot rho.
Proof.
```

```
split; move \Rightarrow H.
\exists (i B \cdot alpha \#).
\exists (i A \cdot alpha).
\verb"rewrite" comp\_inv inv\_invol inv\_universal.
rewrite -comp_assoc (@comp_assoc _ _ _ alpha) lemma_for_tarski2.
apply inc\_antisym.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply H.
\texttt{elim}\ H \Rightarrow mu.
elim \Rightarrow rho H0.
rewrite H0.
rewrite / rectangular.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
rewrite \ unit\_identity\_is\_universal.
apply inc\_alpha\_universal.
Qed.
```

# Chapter 11

# Library Residual

```
Require Import Basic\_Notations.
Require Import Basic\_Lemmas.
Require Import Relation\_Properties.
Require Import Functions\_Mappings.
Require Import Dedekind.
Require Import Domain.
Require Import Logic\_FunctionalExtensionality.
```

# 11.1 剰余合成関係の性質

# 11.1.1 基本的な性質

```
Lemma 213 (double_residual) Let \alpha: A \to B, \beta: B \to C and \gamma: C \to D. Then \alpha \rhd (\beta \rhd \gamma) = (\alpha \cdot \beta) \rhd \gamma.
```

```
Lemma double\_residual \{A \ B \ C \ D : eqType\} \ \{alpha : Rel \ A \ B\} \ \{beta : Rel \ B \ C\} \ \{gamma : Rel \ C \ D\}: alpha \ (beta \ gamma) = (alpha \cdot beta) \ gamma.

Proof.

apply inc\_lower.

move \Rightarrow delta.

split; move \Rightarrow H.

apply inc\_residual.

rewrite comp\_inv \ comp\_assoc.

rewrite -inc\_residual \ -inc\_residual.

apply H.

rewrite inc\_residual \ inc\_residual.

rewrite -comp\_assoc \ -comp\_inv.
```

Qed.

```
apply inc\_residual.
apply H.
Qed.
  Lemma 214 (residual_to_complement) Let \alpha : A \to B and \beta : B \to C. Then
                                     \alpha \triangleright \beta = (\alpha \cdot \beta^{-})^{-}.
Lemma residual_to_complement {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 alpha
          beta = (alpha \cdot beta \hat{)} \hat{.}
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
rewrite bool_lemma2 complement_invol cap_comm.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
replace (beta \hat{} (alpha # • gamma)) with ( B C).
rewrite comp\_empty\_r.
apply inc\_reft.
apply Logic.eq_sym.
rewrite cap\_comm.
apply bool_lemma2.
apply inc\_residual.
apply H.
apply inc\_empty\_alpha.
apply inc\_residual.
apply bool_lemma2.
apply inc\_antisym.
apply (@inc_trans _ _ _ _ (dedekind1)).
rewrite inv_-invol.
                     (alpha \cdot beta \hat{}) with (AC).
replace (gamma
rewrite comp\_empty\_r.
apply inc\_reft.
apply Logic.eq_sym.
rewrite -(@complement_invol _ _ (alpha • beta ^)).
apply bool_lemma2.
apply H.
apply inc\_empty\_alpha.
```

**Lemma 215 (inv\_residual\_inc)** Let  $\alpha : A \rightarrow B$  and  $\beta : B \rightarrow C$ . Then

$$\alpha^{\sharp} \cdot (\alpha \rhd \beta) \sqsubseteq \beta.$$

Lemma  $inv\_residual\_inc$  {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}: alpha # • (alpha beta) beta.

#### Proof.

apply  $inc\_residual$ .

apply  $inc_refl.$ 

Qed.

**Lemma 216 (inc\_residual\_inv)** Let  $\alpha : A \rightarrow B$  and  $\gamma : A \rightarrow C$ . Then

$$\gamma \sqsubseteq \alpha \rhd \alpha^{\sharp} \cdot \gamma.$$

Lemma  $inc\_residual\_inv$  {A B C : eqType} {alpha : Rel A B} {gamma : Rel A C}: gamma (alpha (alpha # • gamma)).

#### Proof.

apply  $inc\_residual$ .

apply  $inc_refl.$ 

Qed.

Lemma 217 (id\_inc\_residual) Let  $\alpha : A \rightarrow B$ . Then

$$id_A \sqsubseteq \alpha \rhd \alpha^{\sharp}$$
.

Lemma  $id\_inc\_residual$  { $A \ B : eqType$ } { $alpha : Rel \ A \ B$ }:  $Id \ A \ (alpha \ alpha \ \#)$ . Proof.

apply *inc\_residual*.

rewrite  $comp_{-}id_{-}r$ .

apply  $inc\_reft$ .

Qed.

## Lemma 218 (residual\_universal) Let $\alpha : A \rightarrow B$ . Then

$$\alpha \triangleright \nabla_{BC} = \nabla_{AC}$$
.

Lemma  $residual\_universal$  {A B C : eqType} {alpha : Rel A B}: alpha B C = A C. Proof.

apply  $inc\_antisym$ .

apply  $inc\_alpha\_universal$ .

apply  $inc\_residual$ .

apply  $inc\_alpha\_universal$ .

Qed.

## 11.1.2 単調性と分配法則

```
Lemma 219 (residual_inc_compat) Let \alpha, \alpha' : A \to B and \beta, \beta' : B \to C. Then \alpha' \sqsubseteq \alpha \land \beta \sqsubseteq \beta' \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha' \rhd \beta'.
```

```
Lemma residual\_inc\_compat
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ B\} \{beta \ beta' : Rel \ B \ C\}:
             alpha \rightarrow \mathtt{beta} \quad beta' \rightarrow (alpha \quad \mathtt{beta})
                                                                     (alpha')
 alpha'
                                                                                 beta').
Proof.
move \Rightarrow H H0.
apply inc\_residual.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H0).
move: (@inc\_refl\_\_(alpha)
                                      beta)) \Rightarrow H1.
apply inc\_residual in H1.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H1).
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply H.
Qed.
```

Lemma 220 (residual\_inc\_compat\_l) Let  $\alpha : A \to B$  and  $\beta, \beta' : B \to C$ . Then  $\beta \sqsubseteq \beta' \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha \rhd \beta'$ .

```
Lemma residual\_inc\_compat\_l {A \ B \ C : eqType} {alpha : Rel \ A \ B} {beta beta' : Rel \ B \ C}: beta beta' \to (alpha \ beta) (alpha \ beta').

Proof.

move \Rightarrow H.

apply (@residual\_inc\_compat \_ \_ \_ \_ \_ \_ (@inc\_refl \_ \_ \_) H).

Qed.
```

Lemma 221 (residual\_inc\_compat\_r) Let  $\alpha, \alpha' : A \to B$  and  $\beta : B \to C$ . Then  $\alpha' \sqsubseteq \alpha \Rightarrow \alpha \rhd \beta \sqsubseteq \alpha' \rhd \beta$ .

```
Lemma residual\_inc\_compat\_r {A \ B \ C : eqType} {alpha \ alpha' : Rel \ A \ B} {beta : Rel \ B \ C}: alpha' \ alpha \rightarrow (alpha \ beta) \ (alpha' \ beta).

Proof.
```

```
move \Rightarrow H.
apply (@residual\_inc\_compat \_ \_ \_ \_ \_ H (@inc\_refl \_ \_ \_)).
Qed.
  Lemma 222 (residual_capP_distr_l, residual_cap_distr_l) Let \alpha : A \rightarrow B, f :
  (D \rightarrow E) \rightarrow (B \rightarrow C) and P: predicate. Then
                               \alpha \rhd (\sqcap_{P(\beta)} f(\beta)) = \sqcap_{P(\beta)} (\alpha \rhd f(\beta)).
\{alpha: Rel\ A\ B\}\ \{f: Rel\ D\ E \rightarrow Rel\ B\ C\}\ \{P: Rel\ D\ E \rightarrow {\tt Prop}\}:
           ( -\{P\} f) = -\{P\} \text{ (fun beta : } Rel \ D \ E \Rightarrow alpha
 alpha
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow beta H\theta.
apply inc\_residual.
move: beta H0.
apply inc\_capP.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inc\_capP.
move \Rightarrow beta H0.
apply inc\_residual.
move : beta H0.
apply inc\_capP.
apply H.
Qed.
Lemma residual\_cap\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta \ gamma : Rel \ B \ C\}:
                    gamma) = (alpha  beta)
           (beta
                                                         (alpha
 alpha
                                                                    qamma).
Proof.
rewrite cap\_to\_capP (@cap\_to\_capP\_\_\_\_\_id).
apply residual\_capP\_distr\_l.
Qed.
```

 $(A \rightarrow B), \beta: B \rightarrow C \text{ and } P: \text{ predicate. Then}$ 

```
(\sqcup_{P(\alpha)} f(\alpha)) \rhd \beta = \sqcap_{P(\alpha)} (f(\alpha) \rhd \beta).
Lemma residual\_cupP\_distr\_r {A \ B \ C \ D \ E : eqType}
 \{ beta : Rel \ B \ C \} \{ f : Rel \ D \ E \rightarrow Rel \ A \ B \} \{ P : Rel \ D \ E \rightarrow Prop \} :
                beta = _{\{P\}} (fun \ alpha : Rel \ D \ E \Rightarrow f \ alpha
Proof.
apply inc\_lower.
move \Rightarrow gamma.
split; move \Rightarrow H.
apply inc\_capP.
move \Rightarrow alpha H0.
apply inc\_residual.
move: alpha\ H0.
apply inc\_cupP.
rewrite -comp_cupP_distr_r -inv_cupP_distr.
apply inc\_residual.
apply H.
apply inc\_residual.
rewrite inv\_cupP\_distr\_comp\_cupP\_distr\_r.
apply inc\_cupP.
move \Rightarrow alpha H0.
apply inc\_residual.
move: alpha\ H0.
apply inc\_capP.
apply H.
Qed.
Lemma residual\_cup\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ beta : Rel \ A \ B\} \{gamma : Rel \ B \ C\}:
            beta) qamma = (alpha qamma)
                                                           (beta
Proof.
rewrite (@cup\_to\_cupP\_\_\_\_\_id) (@cap\_to\_capP\_\_\_\_\_\_(fun x \Rightarrow x)
apply residual\_cupP\_distr\_r.
Qed.
```

Lemma 223 (residual\_cupP\_distr\_r, residual\_cup\_distr\_r) Let  $f:(D \to E) \to$ 

## 11.1.3 剰余合成と関数

```
Lemma 224 (total_residual) Let \alpha : A \to B be a total relation and \beta : B \to C. Then
                                            \alpha \rhd \beta \sqsubseteq \alpha \cdot \beta.
Lemma total_residual {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
 total\_r \ alpha \rightarrow (alpha
                              beta) (alpha \cdot beta).
Proof.
move \Rightarrow H.
apply (@inc_trans _ _ _ ((alpha • alpha #) • (alpha
                                                                 beta))).
apply (comp\_inc\_compat\_b\_ab\ H).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inv_residual_inc.
Qed.
  Lemma 225 (univalent_residual) Let \alpha : A \to B be a univalent relation and \beta :
  B \rightarrow C. Then
                                            \alpha \cdot \beta \sqsubseteq \alpha \rhd \beta.
Lemma univalent\_residual \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}:
 univalent_r \ alpha \rightarrow (alpha \cdot beta) \quad (alpha
                                                          beta).
Proof.
move \Rightarrow H.
apply (@inc_trans _ _ _ (@inc_residual_inv _ _ alpha _)).
apply residual_inc_compat_l.
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
  Lemma 226 (function_residual1) Let \alpha : A \to B be a function and \beta : B \to C.
  Then
                                            \alpha \triangleright \beta = \alpha \cdot \beta.
Lemma function\_residual1 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\}:
 function_r \ alpha \rightarrow alpha
                                beta = alpha • beta.
Proof.
elim \Rightarrow H H0.
apply inc\_antisym.
apply (total\_residual\ H).
apply (univalent_residual H0).
Qed.
```

```
Lemma 227 (residual_id) Let \alpha : A \to B. Then
                                              id_A \rhd \alpha = \alpha.
Lemma residual\_id {A B : eqType} {alpha : Rel A B}:
           alpha = alpha.
 Id A
Proof.
move: (@function\_residual1 \_ \_ \_ (Id A) alpha (@id\_function A)) \Rightarrow H.
rewrite comp_{-}id_{-}l in H.
apply H.
Qed.
  Lemma 228 (universal_residual) Let \alpha : A \to B. Then
                                             \nabla_{AA} \triangleright \alpha \sqsubseteq \alpha.
Lemma universal\_residual \{A \ B : eqType\} \{alpha : Rel \ A \ B\}:
              alpha
                         alpha.
     A A
Proof.
apply (@inc_trans _ _ _ (Id A
                                         alpha)).
apply residual_inc_compat_r.
apply inc\_alpha\_universal.
rewrite residual_id.
apply inc_refl.
Qed.
  Lemma 229 (function_residual2) Let \alpha: A \to B be a function, \beta: B \to C and
  \gamma: C \rightarrow D. Then
                                       \alpha \cdot (\beta \triangleright \gamma) = (\alpha \cdot \beta) \triangleright \gamma.
Lemma function\_residual2
 \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ C \ D\}:
 function\_r \ alpha \rightarrow alpha \cdot (beta \ gamma) = (alpha \cdot beta) \ gamma.
Proof.
move \Rightarrow H.
rewrite -(@function_residual1 _ _ _ _ H).
apply double_residual.
Qed.
```

**Lemma 230 (function\_residual3)** Let  $\alpha:A \to B,\ \beta:B \to C$  be relations and  $\gamma:D\to C$  be a function. Then

$$(\alpha \rhd \beta) \cdot \gamma^{\sharp} = \alpha \rhd (\beta \cdot \gamma^{\sharp}).$$

```
Lemma function_residual3
     \{A \ B \ C \ D : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ D \ C\}: \{gamma : Rel \ D \ C\}
    function\_r\ gamma \rightarrow (alpha \ beta) \cdot gamma \# = alpha \ (beta \cdot gamma \#).
Proof.
move \Rightarrow H.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H0.
apply inc_residual.
rewrite -(@function\_move2\_\_\_\_\_H).
rewrite comp\_assoc.
apply inc\_residual.
rewrite (@function_move2 _ _ _ _ H).
apply H0.
rewrite -(@function\_move2\_\_\_\_\_H).
apply inc\_residual.
rewrite -comp_-assoc.
rewrite (@function\_move2\_\_\_\_\_H).
apply inc\_residual.
apply H0.
Qed.
```

**Lemma 231 (function\_residual4)** Let  $\alpha:A\rightarrow B,\ \gamma:C\rightarrow D$  be relations and  $\beta:B\rightarrow C$  be a function. Then

$$\alpha \cdot \beta \rhd \gamma = \alpha \rhd \beta \cdot \gamma.$$

```
Lemma function_residual4  \{A \ B \ C \ D : eqType\} \ \{alpha : Rel \ A \ B\} \ \{beta : Rel \ B \ C\} \ \{gamma : Rel \ C \ D\}: function\_r \ beta \rightarrow (alpha \cdot beta) \quad gamma = alpha \quad (beta \cdot gamma).  Proof.  move \Rightarrow H.  rewrite -double\_residual.  by [rewrite (function\_residual1 \ H)]. Qed.
```

# 11.2 Galois 同値とその系

```
Lemma 232 (galois) Let \alpha: A \rightarrow B, \beta: B \rightarrow C and \gamma: A \rightarrow C. Then
                                         \gamma \sqsubseteq \alpha \rhd \beta \Leftrightarrow \alpha \sqsubseteq \gamma \rhd \beta^{\sharp}.
Lemma galois \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ A \ C\}:
 qamma
               (alpha
                            beta) \leftrightarrow alpha
                                                    (qamma)
                                                                   beta \#).
Proof.
split; move \Rightarrow H.
apply inc\_residual.
apply inv\_inc\_move.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inv_inc_invol.
rewrite comp_{-}inv \ inv_{-}invol.
apply inc\_residual.
apply H.
Qed.
  Lemma 233 (galois_corollary1) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then
                                              \alpha \sqsubset (\alpha \rhd \beta) \rhd \beta^{\sharp}.
Lemma galois_corollary1 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
             ((alpha
                          beta) beta \#).
 alpha
Proof.
rewrite -galois.
apply inc\_reft.
Qed.
  Lemma 234 (galois_corollary2) Let \alpha : A \rightarrow B and \beta : B \rightarrow C. Then
                                        ((\alpha \rhd \beta) \rhd \beta^{\sharp}) \rhd \beta = \alpha \rhd \beta.
Lemma galois_corollary2 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}:
                         \mathtt{beta} \ \#) \qquad \mathtt{beta} = \mathit{alpha}
 ((alpha
               beta)
Proof.
apply inc\_antisym.
apply residual_inc_compat_r.
```

```
apply galois\_corollary1.

move: (@galois\_corollary1\_\_\_(alpha beta) (beta #)) \Rightarrow H.

rewrite inv\_invol in H.

apply H.


Qed.
```

Lemma 235 (galois\_corollary3) Let  $\alpha : A \to B$  and  $\beta : B \to C$ . Then  $\alpha = (\alpha \rhd \beta) \rhd \beta^{\sharp} \Leftrightarrow \exists \gamma : A \to C, \alpha = \gamma \rhd \beta^{\sharp}.$ 

```
Lemma galois\_corollary3 {A B C : eqType} {alpha : Rel A B} {beta : Rel B C}: alpha = (alpha \quad beta) \quad beta \# \leftrightarrow (\exists \ gamma : Rel A C, alpha = gamma \quad beta \#) Proof. split; move \Rightarrow H. \exists \ (alpha \quad beta). apply H. elim H \Rightarrow gamma \ H0. rewrite H0. move : (@galois\_corollary2 \ \_ \ \_ \ gamma \ (beta \#)) \Rightarrow H1. rewrite inv\_invol in H1. by [rewrite H1]. Qed.
```

# 11.3 その他の性質

この節では、特記が無い限り、記号は以下の図式に従って割り振られるものとする.



Lemma 236 (residual\_property1)

$$(\alpha \rhd \beta) \cdot \gamma \sqsubseteq \alpha \rhd \beta \cdot \gamma.$$

```
Lemma residual\_property1 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}: ((alpha \ beta) \cdot gamma) \ (alpha \ (beta \cdot gamma)).
Proof.
```

## Lemma 237 (residual\_property2)

$$(\alpha \rhd \beta) \cdot (\beta^{\sharp} \rhd \eta) \sqsubseteq \alpha \rhd \eta.$$

```
Lemma residual\_property2 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{eta : Rel \ Y \ V\}: ((alpha beta) \cdot (beta \# eta)) (alpha eta).

Proof.

apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).

apply residual\_inc\_compat\_l.

move : (@inv\_residual\_inc \_ \_ (beta \#) eta).

by [rewrite \ inv\_invol].

Qed.
```

## Lemma 238 (residual\_property3)

Qed.

$$\alpha \rhd \beta \sqsubseteq \alpha \cdot \eta \rhd \eta^{\sharp} \cdot \beta.$$

```
Lemma residual\_property3 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{eta : Rel \ Y \ V\}: (alpha \ beta) \ ((alpha \cdot eta) \ (eta \# \cdot beta)).

Proof.

apply (@inc\_trans \_ \_ \_ \_ \_ (@inc\_residual\_inv \_ \_ \_ (alpha \cdot eta) \ (alpha \ beta))).

apply residual\_inc\_compat\_l.

rewrite comp\_inv \ comp\_assoc.

apply comp\_inc\_compat\_ab\_ab.

apply inv\_residual\_inc.
```

## Lemma 239 (residual\_property4a, residual\_property4b)

$$(\alpha \rhd \beta) \cdot \gamma \sqsubseteq (\alpha \rhd \beta \cdot \gamma) \sqcap \nabla_{XZ} \cdot \gamma \sqsubseteq (\alpha \rhd \beta \cdot \gamma) \cdot \gamma^{\sharp} \cdot \gamma.$$

```
Lemma residual\_property4a {W X Y Z : eqType} {alpha : Rel X Y} {beta : Rel Y Z} {gamma : Rel Z W}:
```

```
beta) • gamma)
                                                 (beta • qamma))
                                                                             X Z \cdot qamma)).
 ((alpha
                                     ((alpha
Proof.
rewrite -(@cap_universal _ _ (alpha
                                            beta)).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_r.
apply residual_property1.
Qed.
Lemma residual_property4b
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}:
             (beta \cdot gamma)) \quad (XZ \cdot gamma)) \quad ((alpha
                                                                             (beta \cdot gamma)) \cdot
(gamma \# \bullet gamma)).
Proof.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite cap_comm cap_universal comp_assoc.
apply inc\_reft.
Qed.
  Lemma 240 (residual_property5) Let \tau be a univalent relation. Then,
                             (\alpha \rhd \beta) \cdot \tau^{\sharp} = (\alpha \rhd \beta \cdot \tau^{\sharp}) \sqcap \nabla_{XZ} \cdot \tau^{\sharp}.
Lemma residual\_property5
 \{V \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{tau : Rel \ V \ Z\}:
 univalent_r tau \rightarrow
           beta) • tau \# = (alpha \quad (beta • tau \#)) \quad (XZ • tau \#).
 (alpha
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite -(@cap_universal _ _ (alpha
apply (@inc\_trans \_ \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat\_r.
apply residual_property1.
rewrite cap\_comm.
apply (@inc_trans _ _ _ (dedekind2)).
{\tt rewrite}\ cap\_comm\ cap\_universal\ inv\_invol.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply residual_inc_compat_l.
rewrite comp_{-}assoc.
apply (comp\_inc\_compat\_ab\_a\ H).
Qed.
```

## Lemma 241 (residual\_property6)

$$\alpha \rhd (\gamma^{\sharp} \rhd \beta^{\sharp})^{\sharp} = (\gamma^{\sharp} \rhd (\alpha \rhd \beta)^{\sharp})^{\sharp}.$$

```
Lemma residual_property6
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\} \{gamma : Rel \ Z \ W\}:
          (gamma \ \# \ beta \ \#) \ \# = (gamma \ \#)
 alpha
                                                       (alpha
                                                                   beta) #) #.
Proof.
apply inc\_lower.
move \Rightarrow delta.
split; move \Rightarrow H.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_move.
apply inc\_residual.
rewrite -comp_inv comp_assoc.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_invol.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply H.
apply inc\_residual.
apply inv\_inc\_move.
apply inc\_residual.
apply inv\_inc\_move.
rewrite comp_inv inv_invol inv_invol comp_assoc.
apply inc\_residual.
apply inv\_inc\_invol.
rewrite comp_{-}inv.
apply inc\_residual.
apply inv\_inc\_move.
apply H.
Qed.
```

# Lemma 242 (residual\_property7a, residual\_property7b)

$$\alpha \rhd (\beta \Rightarrow \beta') \sqsubseteq (\alpha \cdot \beta \Rightarrow \alpha \cdot \beta') \sqsubseteq \alpha \rhd (\beta \Rightarrow \alpha^{\sharp} \cdot \alpha \cdot \beta').$$

Lemma  $residual\_property7a$  { $X \ Y \ Z : eqType$ } { $alpha : Rel \ X \ Y$ } {beta  $beta' : Rel \ Y \ Z$ }: (alpha (beta » beta')) (( $alpha \cdot beta'$ )) » ( $alpha \cdot beta'$ )).

Proof. move  $\Rightarrow H$ .

```
apply inc_-rpc.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp_inc_compat_ab_ab'.
rewrite cap\_comm.
apply inc_-rpc.
apply inv\_residual\_inc.
Lemma residual\_property7b {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta beta' : Rel \ Y \ Z}:
 ((alpha \cdot beta) * (alpha \cdot beta')) (alpha \cdot (beta * (alpha # \cdot (alpha \cdot beta')))).
Proof.
rewrite inc_residual inc_rpc.
apply @inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite inv_invol -inc_rpc.
apply inc_refl.
Qed.
  Lemma 243 (residual_property8) Let \alpha be a univalent relation. Then,
                                  \alpha \rhd (\beta \Rightarrow \beta') = (\alpha \cdot \beta \Rightarrow \alpha \cdot \beta').
Lemma residual\_property8 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta \ beta' : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \quad (beta \gg beta') = (alpha \cdot beta) \gg (alpha \cdot beta').
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply residual_property7a.
apply (@inc_trans _ _ _ residual_property7b).
apply residual_inc_compat_l.
apply rpc\_inc\_compat\_l.
rewrite - comp_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
  Lemma 244 (residual_property9) Let \alpha be a univalent relation. Then,
                                     \alpha \rhd \beta = (\alpha \cdot \nabla_{YZ} \Rightarrow \alpha \cdot \beta).
Lemma residual\_property9 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \quad beta = (alpha \cdot Y Z) \otimes (alpha \cdot beta).
```

```
by [rewrite -(residual\_property8\ H) rpc\_universal\_alpha]. Qed.
```

Lemma 245 (residual\_property10) Let  $\alpha$  be a univalent relation. Then,

$$\alpha \cdot \beta = \lfloor \alpha \rfloor \cdot (\alpha \rhd \beta).$$

```
Lemma residual\_property10 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
 univalent_r \ alpha \rightarrow alpha \cdot beta = domain \ alpha \cdot (alpha
Proof.
move \Rightarrow H.
apply inc\_antisym.
replace (alpha • beta) with (domain alpha • (alpha • beta)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite inc\_residual -comp\_assoc.
apply (comp\_inc\_compat\_ab\_b\ H).
by [rewrite -comp_assoc domain_comp_alpha1].
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot (alpha beta))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inv\_residual\_inc.
Qed.
```

## Lemma 246 (residual\_property11)

```
(\alpha \cdot \beta \Rightarrow \delta) \sqsubseteq \alpha \rhd (\beta \Rightarrow \alpha^{\sharp} \cdot \delta).
```

```
Lemma residual\_property11 \{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{delta : Rel \ X \ Z\} : ((alpha \cdot beta) \otimes delta) \ (alpha \ (beta \otimes (alpha \# \cdot delta))).

Proof.

apply inc\_residual.

apply inc\_rpc.

apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).

rewrite inv\_invol.

apply inc\_rpc.

apply inc\_rpc.

apply inc\_rpc.

apply inc\_rpc.

apply inc\_rpc.
```

```
Lemma 247 (residual_property12a, residual_property12b) Let u \sqsubseteq id_X. Then,
```

```
u \rhd \alpha = u \cdot \nabla_{XY} \Rightarrow \alpha = u \rhd u \cdot \alpha.
```

```
Lemma residual\_property12a {X Y : eqType} {u : Rel X X} {alpha : Rel X Y}:
      Id X \to u
                    alpha = (u \cdot X Y) * alpha.
Proof.
move \Rightarrow H.
apply inc\_antisym.
assert (univalent_r u).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' H).
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_{-}id.
apply (@inc_inv_{-} - H).
rewrite (residual_property9 H0).
apply rpc\_inc\_compat\_l.
apply (comp\_inc\_compat\_ab\_b\ H).
apply (@inc_trans _ _ _ residual_property11).
apply residual_inc_compat_l.
rewrite rpc_universal_alpha.
apply comp\_inc\_compat\_ab\_b.
rewrite -inv_{-}id.
apply (@inc_inv_{-} - H).
Qed.
Lemma residual\_property12b {X Y : eqType} {u : Rel X X} {alpha : Rel X Y}:
      Id X \rightarrow u
                    alpha = u \quad (u \cdot alpha).
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite (residual_property12a H).
apply (@inc_trans _ _ _ residual_property11).
apply residual_inc_compat_l.
rewrite rpc\_universal\_alpha.
apply comp\_inc\_compat\_ab\_a'b.
rewrite (dedekind_id1 H).
apply inc_reft.
apply residual\_inc\_compat\_l.
apply (comp\_inc\_compat\_ab\_b\ H).
Qed.
```

## Lemma 248 (residual\_property13)

```
(\alpha \cdot \nabla_{YZ} \sqcap \delta) \rhd \gamma = (\alpha \cdot \nabla_{YW} \Rightarrow (\delta \rhd \gamma)).
```

```
Lemma residual_property13
 \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{gamma : Rel \ Z \ W\} \{delta : Rel \ X \ Z\}:
                                   qamma = (alpha \cdot Y W) \gg (delta)
 ((alpha \cdot Y Z)  delta)
Proof.
apply inc\_antisym.
rewrite inc\_rpc inc\_residual.
remember (((alpha \cdot Y Z) \quad delta) \quad gamma) \text{ as } sigma1.
apply (@inc\_trans \_ \_ \_ (((alpha \cdot Y Z) \quad delta) \# \cdot sigma1)).
                                                                            (alpha \cdot
apply (@inc\_trans \_ \_ \_ (((alpha \cdot Y Z) \quad delta) \# \cdot (sigma1))
                                                                                            Y
W)))).
assert ((delta \# \cdot (sigma1 \quad (alpha \cdot Y W))) (delta \# \cdot sigma1)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
apply inc_-def1 in H.
rewrite H.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite (@inv_cap_distr _ _ _ delta) cap_comm.
apply cap\_inc\_compat\_r.
rewrite inv\_cap\_distr.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply @inc\_trans \_ \_ \_ \_ (cap\_r).
rewrite comp\_inv \ comp\_inv \ -comp\_assoc \ (@inv\_universal \ Y \ Z).
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_ab'.
apply cap_{-}l.
rewrite Hegsigma1.
apply inc\_residual.
apply inc\_reft.
rewrite inc_residual.
remember \ ((alpha \ \cdot \ Y \ W) \ * \ (delta \ gamma)) \ as \ sigma2.
apply @inc\_trans \_ \_ \_ (delta \# \cdot ((alpha \cdot Y W)))
                                                                 sigma2))).
apply (@inc\_trans\_\_\_(((alpha \cdot YZ) \quad delta) \# \cdot ((alpha \cdot YW) \quad sigma2))).
assert ((((alpha \cdot YZ) delta) # sigma2) (delta # sigma2)).
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply cap_r.
```

Proof.

```
apply inc\_def1 in H.
rewrite H.
apply @inc\_trans \_ \_ \_ \_ (dedekind1).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm inv_invol.
apply cap\_inc\_compat\_r.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot Y Z) \cdot (delta \# \cdot sigma2))).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply inc_{-}inv.
apply cap_r.
rewrite Hegsigma2.
rewrite -inc_residual cap_comm -inc_rpc.
apply inc\_reft.
Qed.
  Lemma 249 (residual_property14) Let \nabla_{XX} \cdot \alpha \sqsubseteq \alpha. Then,
                                      \nabla_{XX} \cdot (\alpha \rhd \beta) \sqsubset \alpha \rhd \beta.
Lemma residual\_property14 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
 (XX \cdot alpha) \quad alpha \rightarrow (XX \cdot (alpha))
                                                            beta))
                                                                          (alpha
                                                                                     beta).
Proof.
move \Rightarrow H.
apply (@inc\_trans \_ \_ \_ ( X X \cdot ( X X (alpha beta)))).
apply comp\_inc\_compat\_ab\_ab'.
rewrite double_residual.
apply (residual\_inc\_compat\_r\ H).
rewrite -inv_universal -inc_residual inv_universal.
apply inc\_reft.
Qed.
  Lemma 250 (residual_property15) Let \beta \cdot \nabla_{ZZ} \subseteq \beta. Then,
                                      (\alpha \rhd \beta) \cdot \nabla_{ZZ} \sqsubset \alpha \rhd \beta.
Lemma residual\_property15 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
            ZZ) beta \rightarrow ((alpha \text{ beta}) \cdot ZZ) (alpha)
```

```
\begin{array}{l} \operatorname{move} \Rightarrow H. \\ \operatorname{apply} \left( @inc\_trans \_ \_ \_ \_ \left( residual\_property1 \right) \right). \\ \operatorname{apply} \left( residual\_inc\_compat\_l \ H \right). \\ \\ \operatorname{Qed}. \\ \\ \hline \\ \operatorname{Lemma 251} \left( \operatorname{residual\_property16} \right) \end{array}
```

 $id_X \sqsubseteq \alpha \rhd \alpha^{\sharp} \land (\alpha \rhd \alpha^{\sharp}) \cdot (\alpha \rhd \alpha^{\sharp}) \sqsubseteq \alpha \rhd \alpha^{\sharp}.$ 

```
Lemma residual_property16 \{X \mid Y : eqType\} \{alpha : Rel \mid X \mid Y\}:

Id X (alpha alpha \#) \bullet (alpha alpha \#)) (alpha alpha \#).

Proof.

split.

rewrite inc\_residual comp\_id\_r.

apply inc\_refl.

move : (@residual\_property2 _ _ _ _ alpha (alpha \#) (alpha \#)) \Rightarrow H.

rewrite inv\_invol in H.

apply H.

Qed.
```

```
Lemma 252 (residual_property17) Let P(y) := "y : I \to Y \text{ is a function"}. Then, \sqcup_{P(y)} y^{\sharp} \cdot y = id_Y \Rightarrow \alpha \rhd \beta = \sqcap_{P(y)} (\alpha \cdot y^{\sharp} \cdot \nabla_{IZ} \Rightarrow \alpha \cdot y^{\sharp} \cdot y \cdot \beta).
```

```
Lemma residual\_property17 \{X \ Y \ Z : eqType\}
 \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{P : Rel \ i \ Y \rightarrow Prop\}:
 P = (\mathbf{fun} \ y : Rel \ i \ Y \Rightarrow function_r \ y) \rightarrow
    \{P\} (fun y: Rel \ i \ Y \Rightarrow y \# \cdot y) = Id \ Y \rightarrow
 alpha beta = _{-}\{P\} (fun y: Rel i Y \Rightarrow
  ((alpha \cdot y \#) \cdot i Z) \gg ((alpha \cdot y \#) \cdot (y \cdot beta))).
Proof.
move \Rightarrow H H0.
replace (alpha
                      beta) with ((alpha • Id Y)
rewrite -H0 comp_cupP_distr_l residual_cupP_distr_r.
apply capP_{-}eq.
move \Rightarrow y H1.
rewrite H in H1.
rewrite -comp_assoc (function_residual4 H1).
apply residual_property9.
rewrite /univalent_r.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
```

by [rewrite  $comp_{-}id_{-}r$ ]. Qed.

# 11.4 順序の関係と左剰余合成

## 11.4.1 max, sup, min, inf

 $\xi: X \to X$  を集合 X における順序と見なしたときの、関係  $\rho: V \to X$  の 最大値  $(\max)$ 、上限  $(\sup)$ 、最小値  $(\min)$ 、下限  $(\inf)$  はそれぞれ、以下のように定義される.

- $max(\rho, \xi) := \rho \sqcap (\rho \rhd \xi)$
- $sup(\rho, \xi) := (\rho \triangleright \xi) \sqcap ((\rho \triangleright \xi) \triangleright \xi^{\sharp})$
- $min(\rho, \xi) := \rho \sqcap (\rho \rhd \xi^{\sharp}) (= max(\rho, \xi^{\sharp}))$
- $inf(\rho, \xi) := (\rho \triangleright \xi^{\sharp}) \sqcap ((\rho \triangleright \xi^{\sharp}) \triangleright \xi) (= sup(\rho, \xi^{\sharp}))$

```
Definition max \{ V \mid X : eqType \} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
             (rho
 := rho
                      xi).
Definition sup \{V \mid X : eqType\} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
                      ((rho
 := (rho)
                                xi)
                                     xi \#).
Definition min \{V \mid X : eqType\} (rho : Rel \mid V \mid X) (xi : Rel \mid X \mid X)
 := rho
             (rho
                      xi \#).
Definition inf \{V \mid X : eqType\} (rho: Rel V X) (xi: Rel X X)
 := (rho
              xi \#
                       ((rho
                                  xi \#
                                              xi).
```

```
Lemma 253 (max_inc_sup) Let \rho: V \to X and \xi: X \to X. Then, max(\rho, \xi) \sqsubseteq sup(\rho, \xi).
```

```
Lemma max\_inc\_sup {V X : eqType} {rho : Rel V X} {xi : Rel X X}: max \ rho \ xi \quad sup \ rho \ xi.
```

### Proof.

rewrite /max/sup.
rewrite  $cap\_comm$ .
apply  $cap\_inc\_compat\_l$ .
apply  $galois\_corollary1$ .
Qed.

```
Lemma 254 (min_inc_inf) Let \rho: V \to X and \xi: X \to X. Then,
                                      min(\rho, \xi) \sqsubseteq inf(\rho, \xi).
Lemma min\_inc\_inf {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 min rho xi
                 inf rho xi.
Proof.
rewrite /min/inf.
rewrite cap\_comm.
apply cap\_inc\_compat\_l.
move: (@galois\_corollary1\_\_\_rho(xi\#)) \Rightarrow H.
rewrite inv\_invol in H.
apply H.
Qed.
  Lemma 255 (inf_to_sup) Let \rho: V \to X and \xi: X \to X. Then,
                                   inf(\rho,\xi) = sup(\rho \triangleright \xi^{\sharp}, \xi).
Lemma inf\_to\_sup \{V \ X : eqType\} \{rho : Rel \ V \ X\} \{xi : Rel \ X \ X\}:
 inf \ rho \ xi = sup \ (rho
                             xi \#) xi.
Proof.
rewrite /sup/inf.
rewrite cap\_comm.
move: (@galois\_corollary2 \_ \_ \_ rho (xi \#)) \Rightarrow H.
rewrite inv\_invol in H.
by [rewrite H].
Qed.
  Lemma 256 (sup_to_inf) Let \rho: V \to X and \xi: X \to X. Then,
                                    sup(\rho, \xi) = inf(\rho \triangleright \xi, \xi).
Lemma sup\_to\_inf {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 sup \ rho \ xi = inf \ (rho
                             xi) xi.
Proof.
rewrite /sup/inf.
rewrite cap\_comm.
by [rewrite qalois_corollary2].
Qed.
```

```
Lemma 257 (residual_inc_sup1, residual_inc_sup2) Let \rho: V \to X and \xi: X \to X
  X. Then,
                                 sup(\rho, \xi) \sqsubseteq \rho \rhd \xi \sqsubseteq sup(\rho, \xi) \rhd \xi.
Lemma residual\_inc\_sup1 { V X : eqType} { rho : Rel V X} { xi : Rel X X}:
                 (rho
 sup rho xi
                          xi).
Proof.
apply cap_{-}l.
Qed.
Lemma residual\_inc\_sup2 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                  ((sup \ rho \ xi)
 (rho
          xi
                                     xi).
Proof.
rewrite qalois.
apply cap_{-}r.
Qed.
  Lemma 258 (max_inc_xi_cap) Let \rho: V \to X and \xi: X \to X. Then,
                                 (max(\rho,\xi))^{\sharp} \cdot max(\rho,\xi) \sqsubseteq \xi \sqcap \xi^{\sharp}.
Lemma max\_inc\_xi\_cap {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (max \ rho \ xi \ \# \ \bullet \ max \ rho \ xi) (xi)
                                               xi \#).
Proof.
rewrite /max.
rewrite inv\_cap\_distr.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat.
apply inc\_residual.
apply cap_{-}r.
apply inv\_inc\_move.
rewrite comp_inv inv_invol.
apply inc\_residual.
apply residual\_inc\_compat\_r.
apply cap_{-}l.
Qed.
  Lemma 259 (sup_inc_xi_cap) Let \rho: V \to X and \xi: X \to X. Then,
                                  (sup(\rho,\xi))^{\sharp} \cdot sup(\rho,\xi) \sqsubseteq \xi \sqcap \xi^{\sharp}.
Lemma sup\_inc\_xi\_cap {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (sup \ rho \ xi \ \# \ \bullet \ sup \ rho \ xi) (xi)
                                            xi \#).
```

```
Proof.
move: (@max\_inc\_xi\_cap\_\_(rho
                                           xi) (xi \#).
rewrite /max/sup.
by [rewrite inv_invol (@cap_comm _ _ xi)].
Qed.
  Lemma 260 (transitive_sup1) Let \rho: V \to X, \xi: X \to X and \xi \cdot \xi \sqsubseteq \xi. Then,
                                  sup(\rho, \xi) \cdot (\xi \sqcap \xi^{\sharp}) = sup(\rho, \xi).
Lemma transitive\_sup1 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                xi \rightarrow sup \ rho \ xi \cdot (xi \quad xi \#) = sup \ rho \ xi.
 (xi \cdot xi)
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite /sup.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply (residual\_inc\_compat\_l\ H).
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
apply (@inc\_trans \_ \_ \_ \_ (residual\_property1)).
apply residual_inc_compat_l.
rewrite -comp_inv inv_inc_move inv_invol.
apply H.
apply (@inc_trans _ _ _ _ (relation_rel_inv_rel)).
rewrite comp\_assoc.
apply (comp\_inc\_compat\_ab\_ab' sup\_inc\_xi\_cap).
Qed.
  Lemma 261 (transitive_sup2) Let \rho: V \to X, \xi: X \to X and \xi \cdot \xi \sqsubseteq \xi. Then,
                                sup(\rho, \xi) \cdot \xi = |sup(\rho, \xi)| \cdot (\rho \triangleright \xi).
Lemma transitive\_sup2 {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
                xi \rightarrow sup \ rho \ xi \cdot xi = domain (sup \ rho \ xi) \cdot (rho
 (xi \cdot xi)
Proof.
move \Rightarrow H.
apply inc\_antisym.
replace (sup rho xi • xi) with (domain (sup rho xi) • (sup rho xi • xi)).
```

apply  $comp\_inc\_compat\_ab\_ab$ '.

```
apply @inc\_trans \_ \_ \_ ((rho
                                    xi) \cdot xi).
apply (comp\_inc\_compat\_ab\_a'b cap\_l).
apply (@inc_trans _ _ _ _ (residual_property1) (residual_inc_compat_l H)).
by [rewrite -comp_assoc domain_comp_alpha1].
apply (@inc_trans _ _ _ (domain (sup rho xi) • (sup rho xi)
                                                                     xi))).
apply comp_inc_compat_ab_ab'.
apply galois.
apply cap_r.
rewrite / domain.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp\_assoc.
apply comp_inc_compat_ab_ab'.
apply inc\_residual.
apply inc_refl.
Qed.
  Lemma 262 (domain_sup_inc) Let \rho: V \to X and \xi: X \to X. Then,
                                |sup(\rho,\xi)| \cdot \rho \sqsubseteq sup(\rho,\xi) \cdot \xi^{\sharp}.
Lemma domain\_sup\_inc {V X : eqType} {rho : Rel V X} {xi : Rel X X}:
 (domain (sup rho xi) \cdot rho) (sup rho xi \cdot xi \#).
Proof.
apply (@inc\_trans \_ \_ \_ (domain (sup rho xi) \cdot (sup rho xi xi \#))).
apply comp\_inc\_compat\_ab\_ab'.
rewrite -galois.
apply cap_{-}l.
rewrite / domain.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_residual.
apply inc\_reft.
Qed.
  Lemma 263 (sup_function) Let \rho: V \to X, \xi: X \to X be relations and f: W \to V
  be a function. Then,
                                  f \cdot sup(\rho, \xi) = sup(f \cdot \rho, \xi).
```

```
function_r f \to f • sup rho xi = \sup (f \cdot rho) xi.
Proof.
move \Rightarrow H.
rewrite /sup.
rewrite (function_cap_distr_l H).
by [rewrite (function_residual2 H) (function_residual2 H) (function_residual2 H)].
Qed.
  Lemma 264 (max_univalent) Let \rho: V \to X, \xi: X \to X be relations and \varphi: W \to X
  V be a univalent relation. Then,
                                  \varphi \cdot max(\rho, \xi) = max(\varphi \cdot \rho, \xi).
Lemma max\_univalent \{ V \ W \ X : eqType \}
 \{rho: Rel\ V\ X\}\ \{xi: Rel\ X\ X\}\ \{phi: Rel\ W\ V\}:
 univalent_r \ phi \rightarrow phi \cdot max \ rho \ xi = max \ (phi \cdot rho) \ xi.
Proof.
move \Rightarrow H.
rewrite /max.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat\_l.
apply (@inc\_trans \_ \_ \_ \_ (univalent\_residual H)).
rewrite double_residual.
apply inc\_reft.
apply (@inc_trans _ _ _ _ (dedekind1)).
```

# 11.4.2 左剰余合成

apply  $cap\_inc\_compat\_l$ .

apply  $inc\_reft$ .

Qed.

apply  $comp\_inc\_compat\_ab\_ab$ '.

rewrite -inc\_residual double\_residual.

```
関係 \alpha: X \to Y, \beta: Y \to Z に対し、左剰余合成を \alpha \triangleleft \beta := (\beta^{\sharp} \triangleright \alpha^{\sharp})^{\sharp} で定義する.
```

```
Definition leftres \{X \ Y \ Z : eqType\} (alpha : Rel X Y) (beta : Rel Y Z) := (beta \# alpha \#) \#.
```

Lemma 265 (inc\_leftres) Let  $\alpha: X \to Y, \ \beta: Y \to Z \ and \ \delta: X \to Z$ . Then,  $\delta \sqsubseteq \alpha \lhd \beta \Leftrightarrow \delta \cdot \beta^{\sharp} \sqsubseteq \alpha.$ 

Lemma  $inc\_leftres$  { $X \ Y \ Z : eqType$ }

 $\{alpha: Rel\ X\ Y\}\ \{beta: Rel\ Y\ Z\}\ \{delta: Rel\ X\ Z\}:$ 

 $delta \qquad leftres \ alpha \ beta \leftrightarrow (delta \ \cdot \ beta \ \#) \qquad alpha$ 

### Proof.

rewrite / leftres.

by [rewrite  $inv\_inc\_move\ inc\_residual\ -comp\_inv\ inv\_inc\_move\ inv\_invol$ ]. Qed.

**Lemma 266 (residual\_leftres\_assoc)** Let  $\alpha: X \to Y$ ,  $\beta: Y \to Z$  and  $\gamma: Z \to W$ . Then,

$$(\alpha \rhd \beta) \lhd \gamma = \alpha \rhd (\beta \lhd \gamma).$$

 $\{alpha: Rel\ X\ Y\}\ \{beta: Rel\ Y\ Z\}\ \{gamma: Rel\ Z\ W\}:$ 

leftres (alpha beta) gamma = alpha leftres beta gamma.

### Proof.

apply  $inc\_lower$ .

 $move \Rightarrow delta$ .

by [rewrite inc\_leftres inc\_residual -comp\_assoc -inc\_leftres -inc\_residual]. Qed.

# Chapter 12

# Library Schroder

```
Require Import Basic\_Notations.

Require Import Basic\_Lemmas.

Require Import Relation\_Properties.

Require Import Functions\_Mappings.


Require Import Dedekind.

Require Import Residual.

Require Import Logic\_FunctionalExtensionality.
```

# 12.1 Schröder 圏の性質

この節では、特記が無い限り、記号は以下の図式に従って割り振られるものとする.



Lemma 267 (schroder\_equivalence1, schroder\_equivalence2)

$$\alpha \cdot \beta \sqsubseteq \delta \Leftrightarrow \alpha^{\sharp} \cdot \delta^{-} \sqsubseteq \beta^{-} \Leftrightarrow \delta^{-} \cdot \beta^{\sharp} \sqsubseteq \alpha^{-}.$$

```
Lemma schroder\_equivalence1 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z} {delta : Rel \ X \ Z}: (alpha \cdot beta) delta \leftrightarrow (alpha \# \cdot delta ^) beta ^{\circ}. Proof. split; move \Rightarrow H. rewrite bool\_lemma2 \ complement\_invol.
```

## CHAPTER 12. LIBRARY SCHRODER

```
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply bool\_lemma2 in H.
rewrite cap_comm inv_invol H comp_empty_r.
apply inc_refl.
apply inc\_empty\_alpha.
rewrite bool_lemma2.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply bool\_lemma2 in H.
rewrite cap_comm -(@complement_invol _ _ beta) H comp_empty_r.
apply inc_refl.
apply inc\_empty\_alpha.
Qed.
Lemma schroder_equivalence2
 \{X \ Y \ Z : eqType\} \ \{alpha : Rel \ X \ Y\} \ \{beta : Rel \ Y \ Z\} \ \{delta : Rel \ X \ Z\}:
 (alpha \cdot beta) \quad delta \leftrightarrow (delta \cdot beta \#)
Proof.
split; move \Rightarrow H.
rewrite bool_lemma2 complement_invol.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply bool\_lemma2 in H.
rewrite cap_comm inv_invol H comp_empty_l.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite bool\_lemma2.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply bool\_lemma2 in H.
rewrite cap_comm -(@complement_invol _ _ alpha) H comp_empty_l.
apply inc\_reft.
apply inc\_empty\_alpha.
Qed.
```

```
Lemma 268 (function_inv_complement) Let \alpha and \tau be functions. Then,
```

$$(\alpha \cdot \beta \cdot \tau^{\sharp})^{-} = \alpha \cdot \beta^{-} \cdot \tau^{\sharp}.$$

```
CHAPTER 12. LIBRARY SCHRODER
 ((alpha \cdot beta) \cdot tau \#) = (alpha \cdot beta) \cdot tau \#.
Proof.
move \Rightarrow H H0.
apply inc\_antisym.
rewrite bool_lemma1 complement_invol.
apply inc\_antisym.
rewrite -comp_cup_distr_r -comp_cup_distr_l complement_classic.
apply (@inc\_trans \_ \_ \_ (((alpha \cdot alpha \#) \cdot
                                                     X V) • (tau \cdot tau \#)).
apply (@inc\_trans \_ \_ \_ ((alpha \cdot alpha \#) \cdot X V)).
apply comp\_inc\_compat\_b\_ab.
apply H.
apply comp\_inc\_compat\_a\_ab.
apply H0.
rewrite -comp_assoc (@comp_assoc _ _ _ alpha) (@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
rewrite bool_lemma2 complement_invol.
apply inc\_antisym.
rewrite - (function\_cap\_distr\ H\ H0)\ cap\_comm\ cap\_complement\_empty\ comp\_empty\_r\ comp\_empty\_l.
apply inc_refl.
apply inc\_empty\_alpha.
Qed.
  Lemma 269 (schroder_univalent1) Let \alpha be a univalent relation and \beta \sqsubseteq \beta'. Then,
                                \alpha \cdot (\beta' \sqcap \beta^-) = \alpha \cdot \beta' \sqcap (\alpha \cdot \beta)^-.
Lemma schroder\_univalent1
```

### CHAPTER 12. LIBRARY SCHRODER

```
apply comp\_cap\_distr\_l.
replace (X Z) with ((alpha \cdot beta)
                                                (alpha • beta ^)).
apply cap\_inc\_compat\_l.
apply cap_r.
apply inc\_antisym.
move: (@univalent\_residual \_ \_ \_ \_ beta H) \Rightarrow H1.
rewrite -inc_{-}rpc.
rewrite residual_to_complement in H1.
apply H1.
apply inc\_empty\_alpha.
apply inc\_def2 in H0.
rewrite -comp_cup_distr_l cup_cap_distr_l.
rewrite -H0 complement_classic cap_universal.
rewrite cup\_cap\_distr\_l -comp\_cup\_distr\_l.
by rewrite -H0 complement_classic cap_universal.
Qed.
  Lemma 270 (schroder_univalent2) Let \alpha be a univalent relation. Then,
                                  \alpha \cdot \beta^- = \alpha \cdot \nabla_{YZ} \sqcap (\alpha \cdot \beta)^-.
```

```
Lemma schroder\_univalent2 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow alpha \cdot beta \hat{\ } = (alpha \cdot Y Z) \quad (alpha \cdot beta) \hat{\ }.
Proof.
move \Rightarrow H.
move: (@schroder_univalent1 _ _ alpha beta ( Y Z) H (@inc_alpha_universal _ _ ))
\Rightarrow H0.
rewrite cap_comm cap_universal in H0.
apply H0.
Qed.
```

Lemma 271 (schroder\_univalent3) Let  $\alpha$  be a univalent relation. Then,

$$(\alpha \cdot \beta)^- = (\alpha \cdot \nabla_{YZ})^- \sqcup \alpha \cdot \beta^-.$$

```
Lemma schroder\_univalent3 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}:
 univalent_r \ alpha \rightarrow (alpha \cdot beta) \hat{} = (alpha \cdot Y Z) \hat{} (alpha \cdot beta \hat{}).
Proof.
move \Rightarrow H.
rewrite (schroder_univalent2 H).
rewrite cup_cap_distr_l cup_comm complement_classic cap_comm cap_universal.
apply inc\_def2.
apply rpc\_inc\_compat\_r.
```

```
apply comp_inc_compat_ab_ab'.
apply inc_alpha_universal.
Qed.
```

Lemma 272 (schroder\_univalent4) Let  $\alpha$  be a univalent relation. Then,

$$\alpha \rhd \beta = (\alpha \cdot \nabla_{YZ})^- \sqcup \alpha \cdot \beta.$$

```
Lemma schroder\_univalent4 {X \ Y \ Z : eqType} {alpha : Rel \ X \ Y} {beta : Rel \ Y \ Z}: univalent\_r \ alpha \rightarrow alpha beta = (alpha \cdot Y \ Z) \cdot (alpha \cdot beta).

Proof.

move \Rightarrow H.

rewrite (residual\_property9 \ H).

apply Logic.eq\_sym.

apply cup\_to\_rpc.

Qed.
```

Lemma 273 (schroder\_universal) Let  $\nabla_{XZ} \cdot \nabla_{ZW} = \nabla_{XW}$ . Then,

$$(\alpha \cdot \nabla_{YZ})^{-} \cdot \nabla_{ZW} = (\alpha \cdot \nabla_{YW})^{-}.$$

```
Lemma schroder\_universal \{W \ X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\}:
                          X W \rightarrow
 (XZ \cdot ZW) =
 (alpha \cdot YZ) \hat{} \cdot ZW = (alpha \cdot YW) \hat{}.
Proof.
move \Rightarrow H.
apply (@cap\_cup\_unique\_\_(alpha \cdot Y W)).
rewrite cap_complement_empty cap_comm.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply (@inc\_trans\_\_\_ (((alpha \cdot YZ) \hat{Y} Z) \hat{Y} (alpha \cdot YZ)) \cdot ZW)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap\_inc\_compat\_l.
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
rewrite cap_comm cap_complement_empty comp_empty_l.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite complement_classic.
apply inc\_antisym.
apply inc\_alpha\_universal.
rewrite -H -(@complement\_classic _ _ (alpha • Y Z)) comp\_cup\_distr\_r.
```

```
apply cup_inc_compat_r.
rewrite comp_assoc.
apply comp_inc_compat_ab_ab'.
apply inc_alpha_universal.
Qed.
```

## Lemma 274 (residual\_inv)

$$(\alpha \rhd \beta)^{\sharp} = \beta^{-\sharp} \rhd \alpha^{-\sharp}.$$

Lemma  $residual\_inv$  { $X \ Y \ Z : eqType$ } { $alpha : Rel \ X \ Y$ } {beta :  $Rel \ Y \ Z$ }: ( $alpha \ beta$ ) # ( $alpha \ )$  #.

Proof.

rewrite  $residual\_to\_complement \ residual\_to\_complement$ .

by [rewrite  $-inv\_complement \ complement\_invol \ inv\_complement \ comp\_inv$ ].

Qed.

Lemma 275 (residual\_cupP\_distr\_l, residual\_cup\_distr\_l) Let  $\alpha$  be a univalent relation,  $f: (V \to W) \to (Y \to Z)$  and  $\exists \beta, P(\beta)$ . Then,

$$\alpha \rhd (\sqcup_{P(\beta)} f(\beta)) = \sqcup_{P(\beta)} (\alpha \rhd f(\beta)).$$

```
Lemma residual\_cupP\_distr\_l { V W X Y Z : eqType}
 \{alpha: Rel\ X\ Y\}\ \{f: Rel\ V\ W \rightarrow Rel\ Y\ Z\}\ \{P: Rel\ V\ W \rightarrow Prop\}:
 univalent_r \ alpha \rightarrow (\exists \ beta' : Rel \ V \ W, P \ beta') \rightarrow
          ( _{P} f) = _{P} (fun beta : Rel V W \Rightarrow alpha f beta).
 alpha
Proof.
move \Rightarrow H.
elim \Rightarrow beta' H0.
rewrite (schroder_univalent4 H) comp_cupP_distr_l.
replace ( _{-}\{P\} (fun beta : Rel\ V\ W \Rightarrow alpha
                                                         f beta)) with ( _{-}\{P\} (fun beta:
Rel\ V\ W \Rightarrow (alpha\ \cdot\ Y\ Z)\ \hat{}\ (alpha\ \cdot\ f\ beta)).
apply (@cap_cup_unique _ _ (alpha •
                                            (Y Z)).
rewrite cap_cup_distr_l cap_cupP_distr_l cap_complement_empty cup_comm cup_empty.
rewrite cap\_cupP\_distr\_l.
apply cupP_{-}eq.
move \Rightarrow gamma H1.
by [rewrite cap_cup_distr_l cap_complement_empty cup_comm cup_empty].
rewrite -cup_assoc complement_classic cup_comm cup_universal.
rewrite -(@complement_invol _ _ (alpha •
apply bool_lemma1.
rewrite complement_invol.
apply (@inc\_trans \_ \_ \_ ((alpha \cdot Y Z) ^ (alpha \cdot f beta'))).
```

```
apply cup_{-}l.
move: beta' H0.
apply inc\_cupP.
apply inc\_reft.
apply cupP_-eq.
move \Rightarrow qamma\ H1.
by [rewrite (schroder_univalent4 H)].
Qed.
Lemma residual\_cup\_distr\_l
 \{X \ Y \ Z : eqType\} \{alpha : Rel \ X \ Y\} \{beta \ beta' : Rel \ Y \ Z\}:
 univalent_r \ alpha \rightarrow
 alpha
            (beta
                        beta') = (alpha)
                                                beta)
                                                            (alpha
                                                                         beta').
Proof.
move \Rightarrow H.
rewrite cup\_to\_cupP (@cup\_to\_cupP\_\_\_\_\_id).
apply (residual\_cupP\_distr\_l\ H).
\exists beta.
by [left].
Qed.
  Lemma 276 (residual_capP_distr_r, residual_cap_distr_r) Let f:(Y \to Z) \to \mathbb{R}
  (I \rightarrow X) and \exists \alpha, P(\alpha). Then,
                                  (\sqcap_{P(\alpha)} f(\alpha)^{\sharp}) \rhd \rho = \sqcup_{P(\alpha)} (f(\alpha)^{\sharp} \rhd \rho).
Lemma residual\_capP\_distr\_r
 \{X \ Y \ Z : eqType\} \ \{rho : Rel \ i \ X\} \ \{f : Rel \ Y \ Z \rightarrow Rel \ i \ X\} \ \{P : Rel \ Y \ Z \rightarrow Prop\}:
 (\exists alpha' : Rel Y Z, P alpha') \rightarrow
 ( _{P} (fun \ alpha : Rel \ Y \ Z \Rightarrow f \ alpha \#)) \quad rho = _{P} (fun \ alpha : Rel \ Y \ Z \Rightarrow f \ alpha \#))
f alpha \#
                 rho).
Proof.
elim \Rightarrow alpha' H.
rewrite residual_to_complement.
rewrite -(@complement_invol _ _ ( _{-}\{P\}\ (fun\ alpha: Rel\ Y\ Z \Rightarrow f\ alpha\ \#
                                                                                                         rho))).
apply f_equal.
rewrite de_{-}morgan3.
replace (fun alpha: Rel Y Z \Rightarrow (f alpha # rho) \hat{}) with (fun alpha: Rel Y Z \Rightarrow f
alpha \# \cdot rho \hat{}).
apply inc\_antisym.
apply comp\_capP\_distr\_r.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
apply (@inc\_trans \_ \_ \_ ((( \_{P} (fun alpha : Rel Y Z \Rightarrow f alpha \# \cdot rho ^))) \cdot (fun alpha : Rel Y Z \Rightarrow f alpha \# \cdot rho ^))
```

## CHAPTER 12. LIBRARY SCHRODER

```
alpha' \# \cdot rho \hat{)} \# \cdot (f \ alpha' \# \cdot rho \hat{)})).
apply comp_inc_compat.
apply comp\_inc\_compat\_ab\_ab'.
move: alpha'H.
apply inc\_capP.
rewrite inv\_capP\_distr.
apply inc\_reft.
move: alpha'H.
apply inc\_capP.
apply inc\_reft.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (comp\_capP\_distr\_r)).
apply inc\_capP.
move \Rightarrow beta H0.
\texttt{apply} \ (@inc\_trans \_ \_ \_ \ ((f \ beta \ \# \ \cdot \ rho \ \hat{\ }) \ \cdot \ ((f \ alpha' \ \# \ \cdot \ rho \ \hat{\ }) \ \# \ \cdot \ f \ alpha' \ \#))).
move: beta H0.
apply inc\_capP.
apply inc\_reft.
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
apply functional_extensionality.
move \Rightarrow x.
by [rewrite residual_to_complement complement_invol].
Qed.
```

# Chapter 13

# Library Sum\_Product

```
Require Import Basic\_Notations.
Require Import Basic\_Lemmas.
Require Import Relation\_Properties.
Require Import Functions\_Mappings.
Require Import Dedekind.
Require Import Conjugate.
Require Import Domain.
Require Import Logic\_IndefiniteDescription.
```

# 13.1 関係の直和

# 13.1.1 入射対,関係直和の定義

入射対の存在公理 (Axiom 23) で入射対が存在することまでは仮定済みなので、実際に入射対  $j:A \rightarrow A+B, k:B \rightarrow A+B$  を定義する関数を定義する.

```
Definition sum_r (A \ B : eqType):
 \{x : (Rel \ A \ (sum_eqType \ A \ B)) \times (Rel \ B \ (sum_eqType \ A \ B)) \mid \\ (fst \ x) \cdot (fst \ x) \ \# = Id \ A \wedge (snd \ x) \cdot (snd \ x) \ \# = Id \ B \wedge \\ (fst \ x) \cdot (snd \ x) \ \# = A \ B \wedge \\ ((fst \ x) \ \# \cdot (fst \ x)) \quad ((snd \ x) \ \# \cdot (snd \ x)) = Id \ (sum_eqType \ A \ B)\}. 
 apply \ constructive\_indefinite\_description. 
 elim \ (@pair_of\_inclusions \ A \ B) \Rightarrow j. 
 elim \Rightarrow k \ H. 
 \exists \ (j,k). 
 simpl. 
 apply \ H. 
 Defined. 
 Defined. 
 Definition \ inl_r \ (A \ B : eqType) := fst \ (sval \ (sum_r \ A \ B)).
```

```
Definition inr_r (A B : eqType) := snd (sval (sum_r A B)).
```

またこの定義による入射対が、入射対としての性質  $(Axiom\ 23)+\alpha$  を満たしていることも事前に証明しておく.

```
Lemma inl\_id {A B : eqType}: inl\_r A B • inl\_r A B \# = Id A.
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inr_id \{A B : eqType\}: inr_r A B \cdot inr_r A B \# = Id B.
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inl\_inr\_empty \{A B : eqType\}: inl\_r A B \cdot inr\_r A B \# = 1
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inr\_inl\_empty {A B : eqType}: inr\_r A B • inl\_r A B # =
Proof.
apply inv_invol2.
rewrite comp_inv inv_invol inv_empty.
apply inl_inr_empty.
Qed.
Lemma inl\_inr\_cup\_id \{A \ B : eqType\}:
 (inl\_r \ A \ B \ \# \ \cdot \ inl\_r \ A \ B) (inr\_r \ A \ B \ \# \ \cdot \ inr\_r \ A \ B) = Id \ (sum\_eqType \ A \ B).
Proof.
apply (proj2\_sig\ (sum\_r\ A\ B)).
Lemma inl\_function \{A \ B : eqType\}: function\_r (inl\_r \ A \ B).
Proof.
move: (proj2\_sig\ (sum\_r\ A\ B)).
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2.
split.
rewrite /total_{-}r.
rewrite H.
apply inc\_reft.
rewrite /univalent_r.
rewrite -H2.
apply cup_l.
Qed.
```

```
Lemma inr\_function \{A \ B : eqType\}: function\_r (inr\_r \ A \ B).

Proof.

move: (proj2\_sig\ (sum\_r\ A\ B)).
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1\ H2.
split.
rewrite /total\_r.
rewrite H0.
apply inc\_refl.
rewrite -H2.
apply cup\_r.
Qed.
```

さらに  $\alpha:A\to C$  と  $\beta:B\to C$  の関係直和  $\alpha\perp\beta:A+B\to C$  を,  $\alpha\perp\beta:=j^{\sharp}\cdot\alpha\sqcup k^{\sharp}\cdot\beta$  で定義する.

```
Definition Rel\_sum \{A \ B \ C : eqType\} \ (alpha : Rel \ A \ C) \ (beta : Rel \ B \ C) := (inl\_r \ A \ B \ \# \ \bullet \ alpha) \ (inr\_r \ A \ B \ \# \ \bullet \ beta).
```

#### 13.1.2 関係直和の性質

```
Lemma 277 (sum_inc_compat) Let \alpha, \alpha' : A \to C and \beta, \beta' : B \to C. Then, \alpha \sqsubseteq \alpha' \land \beta \sqsubseteq \beta' \Rightarrow \alpha \bot \beta \sqsubseteq \alpha' \bot \beta'.
```

```
Lemma sum\_inc\_compat \{A \ B \ C : eqType\} \ \{alpha \ alpha' : Rel \ A \ C\} \ \{beta \ beta' : Rel \ B \ C\}: \ alpha \ alpha' \rightarrow beta \ beta' \rightarrow Rel\_sum \ alpha \ beta \ Rel\_sum \ alpha' \ beta'.

Proof.

move \Rightarrow H \ H0.

apply cup\_inc\_compat.

apply (comp\_inc\_compat\_ab\_ab' \ H).

apply (comp\_inc\_compat\_ab\_ab' \ H0).

Qed.
```

```
Lemma 278 (sum_inc_compat_l) Let \alpha : A \to C and \beta, \beta' : B \to C. Then, \beta \sqsubset \beta' \Rightarrow \alpha \bot \beta \sqsubset \alpha \bot \beta'.
```

Lemma  $sum\_inc\_compat\_l$ 

```
\{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta \ beta' : Rel \ B \ C\}:
           beta' \rightarrow Rel\_sum \ alpha \ beta
                                                Rel_sum alpha beta'.
 beta
Proof.
move \Rightarrow H.
apply (sum\_inc\_compat (@inc\_refl \_ \_ alpha) H).
Qed.
  Lemma 279 (sum_inc_compat_r) Let \alpha, \alpha' : A \rightarrow C and \beta : B \rightarrow C. Then,
                                        \alpha \sqsubseteq \alpha' \Rightarrow \alpha \bot \beta \sqsubseteq \alpha' \bot \beta.
Lemma sum\_inc\_compat\_r
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ C\} \{beta : Rel B \ C\}:
            alpha' \rightarrow Rel\_sum \ alpha \ \texttt{beta}
                                                    Rel_sum alpha' beta.
Proof.
move \Rightarrow H.
apply (sum_inc_compat H (@inc_refl _ _ beta)).
Qed.
  Lemma 280 (total_sum) Let \alpha: A \rightarrow C and \beta: B \rightarrow C are total relations, then
  \alpha \perp \beta is also a total relation.
Lemma total\_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 total\_r \ alpha \rightarrow total\_r \ beta \rightarrow total\_r \ (Rel\_sum \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /total_r/Rel_sum.
\texttt{rewrite-} inl\_inr\_cup\_id\ inv\_cup\_distr\ comp\_cup\_distr\_l\ comp\_cup\_distr\_r\ comp\_cup\_distr\_r.
rewrite comp_inv comp_inv inv_invol inv_invol.
apply cup\_inc\_compat.
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cup\_l)).
rewrite comp_assoc -(@comp_assoc _ _ _ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
apply (fun H' \Rightarrow @inc\_trans \_ \_ \_ \_ H' (cup\_r)).
rewrite comp_assoc -(@comp_assoc _ _ _ beta).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H0).
Qed.
```


**Lemma 281 (univalent\_sum)** Let  $\alpha : A \rightarrow C$  and  $\beta : B \rightarrow C$  are univalent relations, then  $\alpha \perp \beta$  is also a univalent relation.

```
Lemma univalent\_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ \overline{B \ C}\}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (Rel\_sum \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /univalent_r/Rel_sum.
rewrite inv_cup_distr comp_cup_distr_l comp_cup_distr_r comp_cup_distr_r.
rewrite comp_inv comp_inv inv_invol inv_invol.
rewrite comp_assoc -(@comp_assoc _ _ _ (inl_r A B)) inl_id comp_id_l.
rewrite comp_assoc -(@comp_assoc _ _ _ (inr_r A B)) inr_inl_empty comp_empty_l
comp\_empty\_r cup\_empty.
rewrite-cup_assoc comp_assoc -(@comp_assoc _ _ _ (inl_r A B)) inl_inr_empty comp_empty_l
comp\_empty\_r \ cup\_empty.
rewrite comp_assoc -(@comp_assoc _ _ _ (inr_r A B)) inr_id comp_id_l.
apply inc\_cup.
split.
apply H.
apply H\theta.
Qed.
  Lemma 282 (function_sum) Let \alpha: A \to C and \beta: B \to C are functions, then \alpha \perp \beta
  is also a function.
Lemma function_sum \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 function\_r \ alpha \rightarrow function\_r \ beta \rightarrow function\_r \ (Rel\_sum \ alpha \ beta).
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (total_sum H H1).
apply (univalent\_sum\ H0\ H2).
Qed.
  Lemma 283 (sum_conjugate) Let \alpha: A \rightarrow C, \beta: B \rightarrow C and \gamma: A+B \rightarrow C be
  relations, j:A\to A+B and k:B\to A+B be inclusions. Then,
                                j \cdot \gamma = \alpha \wedge k \cdot \gamma = \beta \Leftrightarrow \gamma = \alpha \perp \beta.
Lemma sum\_conjugate
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta : Rel \ B \ C\} \{gamma : Rel \ (sum\_eqType \ A \ B)\}
C}:
 inl\_r \ A \ B \cdot gamma = alpha \land inr\_r \ A \ B \cdot gamma = beta \leftrightarrow
 gamma = Rel\_sum \ alpha \ beta.
Proof.
```

```
 \begin{split} & \text{split; move} \Rightarrow H. \\ & \text{elim } H \Rightarrow H0 \ H1. \\ & \text{rewrite -} (@comp\_id\_l\_\_\_gamma). \\ & \text{rewrite -} inl\_inr\_cup\_id \ comp\_cup\_distr\_r \ comp\_assoc \ comp\_assoc. \\ & \text{by [rewrite } H0 \ H1]. \\ & \text{split.} \\ & \text{rewrite } H \ comp\_cup\_distr\_l - comp\_assoc - comp\_assoc. \\ & \text{rewrite } inl\_id \ inl\_inr\_empty \ comp\_id\_l \ comp\_empty\_l. \\ & \text{by [rewrite } cup\_empty]. \\ & \text{rewrite } H \ comp\_cup\_distr\_l - comp\_assoc - comp\_assoc. \\ & \text{rewrite } inr\_id \ inr\_inl\_empty \ comp\_id\_l \ comp\_empty\_l. \\ & \text{by [rewrite } cup\_comm \ cup\_empty]. \\ & \text{Qed.} \\ \end{split}
```

#### Lemma 284 (sum\_comp) In below figure,

$$(\alpha \perp \beta)^{\sharp} \cdot (\gamma \perp \delta) = \alpha^{\sharp} \cdot \gamma \sqcup \beta^{\sharp} \cdot \delta.$$



```
Lemma sum\_comp {W \ X \ Y \ Z : eqType} {alpha : Rel \ Y \ X} {beta : Rel \ Z \ X} {gamma : Rel \ Y \ W} {delta : Rel \ Z \ W}: (Rel\_sum \ alpha \ beta) # • Rel\_sum \ gamma \ delta = (alpha \ \# \ \bullet \ gamma) (beta # • delta).
```

Proof.

rewrite  $/Rel_sum$ .

 $\textbf{rewrite} \ inv\_cup\_distr\_comp\_cup\_distr\_l \ comp\_cup\_distr\_r \ comp\_cup\_distr\_r.$ 

rewrite comp\_inv comp\_inv inv\_invol inv\_invol.

apply  $f_{-}equal2$ .

rewrite  $comp\_assoc$  -(@ $comp\_assoc$  \_ \_ \_ ( $inl\_r$  Y Z))  $inl\_id$   $comp\_id\_l$ .

by [rewrite  $comp\_assoc$  -(@ $comp\_assoc$  \_ \_ \_ ( $inr\_r$  Y Z))  $inr\_inl\_empty$   $comp\_empty\_l$   $comp\_empty\_r$   $cup\_empty$ ].

rewrite  $comp\_assoc$  -(@ $comp\_assoc$  \_ \_ \_ ( $inl\_r$  Y Z))  $inl\_inr\_empty$   $comp\_empty\_l$   $comp\_empty\_r$   $cup\_comm$   $cup\_empty$ .

by [rewrite  $comp\_assoc$  -(@ $comp\_assoc$  \_ \_ \_ \_ ( $inr\_r$  Y Z))  $inr\_id$   $comp\_id\_l$ ]. Qed.

#### 13.1.3 分配法則

```
Lemma 285 (sum_cap_distr_l) Let \alpha : A \to C and \beta, \beta' : B \to C. Then,
                                    \alpha \perp (\beta \sqcap \beta') \sqsubseteq (\alpha \perp \beta) \sqcap (\alpha \perp \beta').
Lemma sum\_cap\_distr\_l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta \ beta' : Rel \ B \ C\}:
 Rel_sum alpha (beta beta') (Rel_sum alpha beta Rel_sum alpha beta').
Proof.
rewrite -cup\_cap\_distr\_l.
apply cup\_inc\_compat\_l.
apply comp\_cap\_distr\_l.
Qed.
  Lemma 286 (sum_cap_distr_r) Let \alpha, \alpha' : A \to C and \beta : B \to C. Then,
                                    (\alpha \sqcap \alpha') \bot \beta \sqsubseteq (\alpha \bot \beta) \sqcap (\alpha' \bot \beta).
Lemma sum\_cap\_distr\_r
 \{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ C\} \{beta : Rel \ B \ C\}:
 Rel_sum (alpha
                        alpha') beta
                                          (Rel_sum alpha beta Rel_sum alpha' beta).
Proof.
rewrite -cup\_cap\_distr\_r.
apply cup\_inc\_compat\_r.
apply comp\_cap\_distr\_l.
Qed.
  Lemma 287 (sum_cup_distr_l) Let \alpha : A \rightarrow C and \beta, \beta' : B \rightarrow C. Then,
                                    \alpha \bot (\beta \sqcup \beta') = (\alpha \bot \beta) \sqcup (\alpha \bot \beta').
Lemma sum_-cup_-distr_-l
 \{A \ B \ C : eqType\} \{alpha : Rel \ A \ C\} \{beta \ beta' : Rel \ B \ C\}:
 Rel\_sum\ alpha\ (beta) = Rel\_sum\ alpha\ beta Rel\_sum\ alpha\ beta.
rewrite -cup_assoc (@cup_comm _ _ (Rel_sum alpha beta)) -cup_assoc.
by [rewrite cup_idem cup_assoc -comp_cup_distr_l].
Qed.
```

```
Lemma 288 (sum_cup_distr_r) Let \alpha, \alpha' : A \to C and \beta : B \to C. Then, (\alpha \sqcup \alpha') \bot \beta = (\alpha \bot \beta) \sqcup (\alpha' \bot \beta).
```

Lemma  $sum\_cup\_distr\_r$  { $A \ B \ C : eqType$ } { $alpha \ alpha' : Rel \ A \ C$ } {beta :  $Rel \ B \ C$ }:  $Rel\_sum \ (alpha \ alpha')$  beta = ( $Rel\_sum \ alpha$  beta  $Rel\_sum \ alpha'$  beta). Proof. rewrite  $cup\_assoc \ (@cup\_comm\_\_ \ (inr\_r \ A \ B \ \# \ \ beta)) \ cup\_assoc.$  by [rewrite  $cup\_idem \ -cup\_assoc \ -comp\_cup\_distr\_l$ ]. Qed.

**Lemma 289 (comp\_sum\_distr\_r)** Let  $\alpha:A\rightarrow C,\ \beta:B\rightarrow C$  and  $\gamma:C\rightarrow D.$  Then,

$$(\alpha \perp \beta) \cdot \gamma = \alpha \cdot \gamma \perp \beta \cdot \gamma.$$

```
Lemma comp\_sum\_distr\_r {A B C D : eqType} {alpha : Rel A C} {beta : Rel B C} {gamma : Rel C D}: (Rel\_sum alpha beta) • gamma = Rel\_sum (alpha • gamma) (beta • gamma). Proof. by [rewrite comp\_cup\_distr\_r comp\_assoc comp\_assoc]. Qed.
```

#### 13.2 関係の直積

#### 13.2.1 射影対,関係直積の定義

射影対の存在公理  $(Axiom\ 24)$  で射影対が存在することまでは仮定済みなので、実際に射影対  $p:A\times B\to A, k:A\times B\to B$  を定義する関数を定義する.

```
Definition prod_r (A B : eqType): \{x : (Rel (prod_eqType A B) A) \times (Rel (prod_eqType A B) B) \mid (fst x) \# \cdot (snd x) = A B \land ((fst x) \cdot (fst x) \#) ((snd x) \cdot (snd x) \#) = Id (prod_eqType A B) \land univalent_r (fst x) \land univalent_r (snd x)\}. apply constructive\_indefinite\_description. elim (@pair\_of\_projections A B) \Rightarrow p. elim \Rightarrow q H. \exists (p,q). simpl. apply H.
```

も事前に証明しておく.

# Defined. Definition $fst_r$ (A B : eqType):= fst (sval ( $prod_r A B$ )). Definition $snd_r$ (A B : eqType):= snd (sval ( $prod_r A B$ )). またこの定義による射影対が、射影対としての性質 ( $Axiom\ 24$ ) + $\alpha$ を満たしていること

Lemma  $fst\_snd\_universal \{A B : eqType\}: fst\_r A B \# \bullet snd\_r A B =$ A B. Proof. apply  $(proj2\_sig\ (prod\_r\ A\ B))$ . Qed. Lemma  $snd\_fst\_universal\ \{A\ B: eqType\}:\ snd\_r\ A\ B\ \#\ \bullet\ fst\_r\ A\ B=$ B A. Proof. apply  $inv_invol2$ . rewrite comp\_inv inv\_invol inv\_universal. apply  $fst\_snd\_universal$ . Qed. Lemma  $fst\_snd\_cap\_id \{A B : eqType\}$ :  $(fst_r \ A \ B \cdot fst_r \ A \ B \#) \quad (snd_r \ A \ B \cdot snd_r \ A \ B \#) = Id (prod_eqType \ A \ B).$ Proof. apply  $(proj2\_sig\ (prod\_r\ A\ B))$ . Qed. Lemma  $fst\_function \{A \ B : eqType\}$ :  $function\_r (fst\_r \ A \ B)$ . Proof. move:  $(proj2\_sig\ (prod\_r\ A\ B))$ .  $elim \Rightarrow H$ .  $elim \Rightarrow H0 \ H1.$ split. rewrite  $/total_{-}r$ . rewrite -H0. apply  $cap_{-}l$ . apply H1. Lemma  $snd\_function \{A B : eqType\}$ :  $function\_r (snd\_r A B)$ . Proof. move:  $(proj2\_sig\ (prod\_r\ A\ B))$ .  $elim \Rightarrow H$ .  $elim \Rightarrow H0 \ H1.$ split. rewrite  $/total_r$ . rewrite -H0. apply  $cap_{-}r$ .

apply H1.

Qed.

さらに  $\alpha:A \to B$  と  $\beta:A \to C$  の関係直積  $\alpha \top \beta:A \to B \times C$  を,  $\alpha \top \beta:=\alpha \cdot p^\sharp \sqcap \beta \cdot q^\sharp$  で定義する.

```
Definition Rel\_prod \{A \ B \ C : eqType\} \ (alpha : Rel \ A \ B) \ (beta : Rel \ A \ C) := (alpha \cdot fst\_r \ B \ C \ \#) \ (beta \cdot snd\_r \ B \ C \ \#).
```

#### 13.2.2 関係直積の性質

Lemma 290 (prod\_inc\_compat) Let  $\alpha, \alpha' : A \to B$  and  $\beta, \beta' : A \to C$ . Then,  $\alpha \sqsubset \alpha' \land \beta \sqsubset \beta' \Rightarrow \alpha \top \beta \sqsubset \alpha' \top \beta'.$ 

Lemma  $prod\_inc\_compat$ 

```
\{A\ B\ C: eqType\}\ \{alpha\ alpha': Rel\ A\ B\}\ \{beta\ beta': Rel\ A\ C\}: alpha\ alpha' 	o beta\ beta' 	o Rel\_prod\ alpha\ beta\ Rel\_prod\ alpha'\ beta'.

Proof.
```

move  $\Rightarrow H H0$ .

apply  $cap\_inc\_compat$ .

apply  $(comp\_inc\_compat\_ab\_a'b\ H)$ .

apply  $(comp_inc_compat_ab_a'b H0)$ .

Qed.

**Lemma 291 (prod\_inc\_compat\_l)** Let  $\alpha : A \rightarrow B$  and  $\beta, \beta' : A \rightarrow C$ . Then,

$$\beta \sqsubseteq \beta' \Rightarrow \alpha \top \beta \sqsubseteq \alpha \top \beta'.$$

Lemma  $prod\_inc\_compat\_l$ 

```
\{A\ B\ C: eqType\}\ \{alpha: Rel\ A\ B\}\ \{beta beta': Rel\ A\ C\}: beta beta' \rightarrow Rel_prod alpha beta Rel_prod alpha beta'. Proof.
```

move  $\Rightarrow H$ .

 ${\tt apply} \ (prod\_inc\_compat \ (@inc\_refl \ \_ \ alpha) \ H).$ 

Qed.

**Lemma 292** (prod\_inc\_compat\_r) Let  $\alpha, \alpha' : A \rightarrow B$  and  $\beta : A \rightarrow C$ . Then,

$$\alpha \sqsubseteq \alpha' \Rightarrow \alpha \top \beta \sqsubseteq \alpha' \top \beta.$$

Lemma  $prod\_inc\_compat\_r$ 

```
\{A \ B \ C : eqType\} \{alpha \ alpha' : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
```

```
alpha' \rightarrow Rel\_prod\ alpha\ beta
                                                   Rel_prod alpha' beta.
 alpha
Proof.
move \Rightarrow H.
apply (prod_inc_compat H (@inc_refl _ _ beta)).
Qed.
  Lemma 293 (total_prod) Let \alpha: A \rightarrow B and \beta: A \rightarrow C are total relations, then
  \alpha \top \beta is also a total relation.
Lemma total\_prod \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 total\_r \ alpha \rightarrow total\_r \ beta \rightarrow total\_r \ (Rel\_prod \ alpha \ beta).
Proof.
move \Rightarrow H H0.
{\tt rewrite} \ domain\_total \ cap\_domain \ cap\_comm.
apply Logic.eq_sym.
apply inc\_def1.
apply @inc\_trans \_ \_ \_ \_ H).
rewrite comp_inv inv_invol comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply (@inc\_trans \_ \_ \_ (alpha \# \cdot (beta \cdot beta \#))).
apply (comp\_inc\_compat\_a\_ab\ H0).
rewrite -comp_assoc -comp_assoc fst_snd_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
  Lemma 294 (univalent_prod) Let \alpha : A \rightarrow B and \beta : A \rightarrow C are univalent relations,
  then \alpha \top \beta is also a univalent relation.
Lemma univalent\_prod \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 univalent_r \ alpha \rightarrow univalent_r \ beta \rightarrow univalent_r \ (Rel_prod \ alpha \ beta).
Proof.
move \Rightarrow H H0.
rewrite /univalent_r/Rel_prod.
rewrite inv_cap_distr comp_inv inv_invol comp_inv inv_invol.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
rewrite -fst\_snd\_cap\_id.
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp\_assoc - (@comp\_assoc \_ \_ \_ \_ alpha).
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_ab\_b\ H).
```

rewrite (@comp\_assoc \_ \_ \_ \_ (

apply  $comp\_inc\_compat$ .

```
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_r)).
rewrite comp\_assoc - (@comp\_assoc \_ \_ \_ \_ \_ beta).
apply comp_inc_compat_ab_ab'.
apply (comp\_inc\_compat\_ab\_b\ H0).
Qed.
  Lemma 295 (function_prod) Let \alpha: A \to B and \beta: A \to C are functions, then
  \alpha \top \beta is also a function.
Lemma function_prod {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 function\_r \ alpha \rightarrow function\_r \ \mathsf{beta} \rightarrow function\_r \ (Rel\_prod \ alpha \ \mathsf{beta}).
Proof.
elim \Rightarrow H H0.
elim \Rightarrow H1 H2.
split.
apply (total_prod H H1).
apply (univalent_prod H0 H2).
Qed.
  Lemma 296 (prod_fst_surjection) Let p: B \times C \rightarrow B be a projection. Then,
                           "p is a surjection" \Leftrightarrow \forall D, \nabla_{BD} = \nabla_{BC} \cdot \nabla_{CD}.
Lemma prod\_fst\_surjection \{B \ C : eqType\}:
 surjection\_r (fst\_r \ B \ C) \leftrightarrow \forall \ D : eqType, \quad B \ D = \quad B \ C \cdot \quad C \ D.
Proof.
split; move \Rightarrow H.
move \Rightarrow D.
elim H \Rightarrow H0 \ H1.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((fst\_r \ B \ C \# \cdot (fst\_r \ B \ C \#) \#) \cdot B \ D)).
apply (comp\_inc\_compat\_b\_ab\ H1).
rewrite inv_invol.
apply (@inc\_trans \_ \_ \_ (((fst\_r \ B \ C \# \cdot snd\_r \ B \ C) \cdot (snd\_r \ B \ C \# \cdot fst\_r \ B \ C))
   B D)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc - (@comp\_assoc \_ \_ \_ \_ (snd\_r B C)).
apply comp_inc_compat_ab_ab'.
apply comp\_inc\_compat\_b\_ab.
apply snd_-function.
```

BD)).

```
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
split.
apply fst_-function.
rewrite /total_{-}r.
rewrite - (@cap_universal _ _ (Id B)) (H B) - (@fst_snd_universal B C) cap_comm comp_assoc.
apply (@inc\_trans\_\_\_\_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite comp_{-}id_{-}r.
apply cap_r.
Qed.
  Lemma 297 (prod_snd_surjection) Let q: B \times C \to C be a projection. Then,
                          "q is a surjection" \Leftrightarrow \forall D, \nabla_{CD} = \nabla_{CB} \cdot \nabla_{BD}.
Lemma prod\_snd\_surjection \{B \ C : eqType\}:
 surjection\_r (snd\_r \ B \ C) \leftrightarrow \forall \ D : eqType, \qquad C \ D = C \ B \bullet
                                                                               B D.
Proof.
split; move \Rightarrow H.
move \Rightarrow D.
elim H \Rightarrow H0 \ H1.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((snd\_r \ B \ C \# \cdot (snd\_r \ B \ C \#) \#) \cdot C \ D)).
apply (comp\_inc\_compat\_b\_ab\ H1).
rewrite inv_invol.
apply (@inc\_trans \_ \_ \_ (((snd\_r \ B \ C \ \# \ \cdot \ fst\_r \ B \ C) \ \cdot \ (fst\_r \ B \ C \ \# \ \cdot \ snd\_r \ B \ C))
   (CD)).
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp\_assoc -(@comp\_assoc _ _ _ _ (fst\_r B C)).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_b\_ab.
apply fst\_function.
rewrite (@comp\_assoc\_\_\_\_\_(CD)).
apply comp\_inc\_compat.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
apply inc\_alpha\_universal.
split.
apply snd\_function.
rewrite /total_{-}r.
```

 $rewrite - (@cap\_universal\_\_(Id\ C))(H\ C) - (@snd\_fst\_universal\ B\ C)\ cap\_comm\ comp\_assoc.$ 

```
apply (@inc\_trans \_ \_ \_ \_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite comp_{-}id_{-}r.
apply cap_{-}r.
Qed.
  Lemma 298 (prod_fst_domain1) Let p: B \times C \rightarrow B be a projection, \alpha: A \rightarrow B
  and \beta: A \rightarrow C. Then,
                                       (\alpha \top \beta) \cdot p = |\beta| \cdot \alpha.
Lemma prod_fst_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • fst\_r\ B\ C=domain\ beta • alpha.
Proof.
rewrite (@comp_inv_inv A A) domain_inv.
rewrite domain_universal2 inv_cap_distr comp_inv inv_invol inv_invol inv_universal.
rewrite -snd_-fst_-universal.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
rewrite comp\_assoc comp\_assoc.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_a.
apply fst\_function.
rewrite cap_comm -comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite cap\_comm.
apply inc\_reft.
Qed.
  Lemma 299 (prod_fst_domain2) Let p: B \times C \rightarrow B be a projection, \alpha: A \rightarrow B
  and \beta: A \rightarrow C. Then,
                                  (\alpha \top \beta) \cdot p = \alpha \Leftrightarrow |\alpha| \sqsubseteq |\beta|.
Lemma prod_fst_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) \cdot fst\_r\ B\ C = alpha \leftrightarrow domain\ alpha
                                                                          domain beta.
Proof.
rewrite prod_fst_domain1.
split; move \Rightarrow H.
apply domain_lemma2b.
assert ((domain beta • alpha)
                                        ((beta \cdot beta \#) \cdot alpha)).
apply comp\_inc\_compat\_ab\_a'b.
apply cap_{-}l.
rewrite H in H0.
apply H0.
```

```
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_{-}r.
apply (@inc_trans _ _ _ (domain alpha • alpha)).
rewrite domain_comp_alpha1.
apply inc_refl.
apply (comp\_inc\_compat\_ab\_a'b\ H).
Qed.
  Lemma 300 (prod_snd_domain1) Let q: B \times C \to C be a projection, \alpha: A \to B
  and \beta: A \rightarrow C. Then,
                                      (\alpha \top \beta) \cdot q = |\alpha| \cdot \beta.
Lemma prod_snd_domain1 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • snd\_r\ B\ C=domain\ alpha • beta.
Proof.
rewrite (@comp_inv_inv A A) domain_inv.
rewrite domain_universal2 inv_cap_distr comp_inv inv_invol inv_invol inv_universal.
rewrite -fst\_snd\_universal.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
rewrite comp_assoc comp_assoc cap_comm.
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_ab\_a.
apply snd_-function.
rewrite cap\_comm -comp\_assoc.
apply dedekind2.
Qed.
  Lemma 301 (prod_snd_domain2) Let q: B \times C \to C be a projection, \alpha: A \to B
  and \beta: A \rightarrow C. Then,
                                 (\alpha \top \beta) \cdot q = \beta \Leftrightarrow |\beta| \sqsubset |\alpha|.
Lemma prod\_snd\_domain2 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 (Rel\_prod\ alpha\ beta) • snd\_r\ B\ C = beta \leftrightarrow domain\ beta domain alpha.
Proof.
rewrite prod_snd_domain1.
split; move \Rightarrow H.
apply domain_lemma2b.
                                      ((alpha \cdot alpha \#) \cdot beta)).
assert ((domain alpha • beta)
apply comp\_inc\_compat\_ab\_a'b.
apply cap_l.
rewrite H in H0.
```

```
apply H0.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_b.
apply cap_{-}r.
apply (@inc_trans _ _ (domain beta • beta)).
rewrite domain_comp_alpha1.
apply inc\_reft.
apply (comp\_inc\_compat\_ab\_a'b\ H).
Qed.
  Lemma 302 (prod_to_cap) Let \alpha : A \rightarrow B and \beta : A \rightarrow C. Then,
                                        |\alpha \top \beta| = |\alpha| \sqcap |\beta|.
Lemma prod\_to\_cap \{A \ B \ C : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ A \ C\}:
 domain (Rel\_prod alpha beta) = domain alpha
                                                           domain beta.
Proof.
replace (domain (Rel_prod alpha beta)) with (domain (Rel_prod alpha beta • snd_r B
C)).
rewrite prod_snd_domain1 comp_domain8.
apply dedekind_id3.
apply cap_{-}r.
apply cap_r.
apply cap_{-}r.
apply comp\_domain3.
apply snd_-function.
Qed.
  Lemma 303 (prod_conjugate1) Let \alpha: A \to B and \beta: A \to C be functions, p:
  B \times C \to B and q: B \times C \to C be projections. Then,
                                  (\alpha \top \beta) \cdot p = \alpha \wedge (\alpha \top \beta) \cdot q = \beta.
Lemma prod\_conjugate1 {A B C : eqType} {alpha : Rel A B} {beta : Rel A C}:
 function\_r \ alpha \rightarrow function\_r \ \texttt{beta} \rightarrow
 Rel\_prod\ alpha\ beta\ \cdot\ fst\_r\ B\ C=alpha\ \wedge\ Rel\_prod\ alpha\ beta\ \cdot\ snd\_r\ B\ C=beta.
Proof.
move \Rightarrow H H0.
split.
rewrite prod_fst_domain1.
elim H0 \Rightarrow H1 \ H2.
apply inc\_def1 in H1.
rewrite / domain.
```

```
by [rewrite cap\_comm -H1 comp\_id\_l]. rewrite prod\_snd\_domain1. elim H \Rightarrow H1 H2. apply inc\_def1 in H1. rewrite /domain. by [rewrite cap\_comm -H1 comp\_id\_l]. Qed.
```

**Lemma 304 (prod\_conjugate2)** Let  $\gamma: A \to B \times C$  be a function,  $p: B \times C \to B$  and  $q: B \times C \to C$  be projections. Then,

$$(\gamma \cdot p) \top (\gamma \cdot q) = \gamma.$$

Lemma  $prod\_conjugate2$  { $A \ B \ C : eqType$ } { $gamma : Rel \ A \ (prod\_eqType \ B \ C)$ }:  $function\_r \ gamma \rightarrow Rel\_prod \ (gamma \cdot fst\_r \ B \ C) \ (gamma \cdot snd\_r \ B \ C) = gamma.$  Proof. move  $\Rightarrow H$ . rewrite  $/Rel\_prod$ . rewrite  $/Rel\_prod$ . rewrite  $comp\_assoc \ comp\_assoc \ -(function\_cap\_distr\_l \ H)$ . by [rewrite  $fst\_snd\_cap\_id \ comp\_id\_r$ ]. Qed.

**Lemma 305 (diagonal\_conjugate)** Let  $p: B \times C \rightarrow B$  and  $q: B \times C \rightarrow C$  be projections. Then,

$$\frac{\alpha:A \to B}{u \sqsubseteq id_{A \times B}} \ \frac{\alpha = p^{\sharp} \cdot u \cdot q}{u = |p \cdot \alpha \sqcap q|}.$$

```
Lemma diagonal\_conjugate \{A B : eqType\} \{alpha : Rel A B\}:
 conjugate A B (prod_eqType A B) (prod_eqType A B)
 True\_r (fun \ u \Rightarrow u \quad Id (prod\_eqType \ A \ B))
 (\mathbf{fun}\ u \Rightarrow (fst_r\ A\ B\ \#\ \cdot\ u)\ \cdot\ snd_r\ A\ B)
 (\mathbf{fun} \ alpha \Rightarrow domain \ ((fst_r \ A \ B \cdot alpha) \quad snd_r \ A \ B)).
Proof.
split.
move \Rightarrow alpha0 H.
split.
apply cap_{-}r.
rewrite cap\_domain.
apply inc\_antisym.
apply (@inc\_trans\_\_\_((fst\_r\ A\ B\ \#\ \cdot\ ((fst\_r\ A\ B\ \bullet\ alpha0)\ \bullet\ snd\_r\ A\ B\ \#))\ \bullet\ snd\_r
A B)).
apply comp_inc_compat_ab_a'b.
apply comp\_inc\_compat\_ab\_ab'.
```

```
apply cap_{-}l.
rewrite comp_assoc comp_assoc -(@comp_assoc _ _ _ _ (fst_r A B #)).
apply (@inc\_trans \_ \_ \_ ((fst\_r \ A \ B \ \# \cdot fst\_r \ A \ B) \cdot alpha0)).
apply comp\_inc\_compat\_ab\_a.
apply snd_-function.
apply comp\_inc\_compat\_ab\_b.
apply fst_-function.
                                       ((fst_r \ A \ B \ \# \ \cdot \ Id \ (prod_eqType \ A \ B)) \ \cdot \ snd_r \ A
apply (@inc_trans _ _ _ (alpha0
B))).
rewrite comp_id_r fst_snd_universal cap_universal.
apply inc\_reft.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc_trans _ _ _ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm inv_invol comp_assoc.
apply inc\_reft.
move \Rightarrow u H.
split.
by ||.
replace ((fst_r \ A \ B \cdot ((fst_r \ A \ B \# \cdot u) \cdot snd_r \ A \ B)) \quad snd_r \ A \ B) with (u \cdot snd_r \ A \ B)
A B).
apply domain\_inc\_id in H.
move: (@snd\_function \ A \ B) \Rightarrow H0.
elim H0 \Rightarrow H1 \ H2.
by [rewrite (comp\_domain3 \ H1) \ H].
rewrite comp_assoc -comp_assoc.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ ((u \cdot snd\_r A B) \quad snd\_r A B)).
apply inc\_cap.
split.
apply inc\_reft.
apply (comp\_inc\_compat\_ab\_b\ H).
apply cap\_inc\_compat\_r.
apply comp\_inc\_compat\_b\_ab.
apply fst\_function.
apply (@inc_trans _ _ _ (dedekind2)).
apply comp\_inc\_compat\_ab\_b.
rewrite -fst\_snd\_cap\_id.
apply cap\_inc\_compat\_l.
apply comp\_inc\_compat\_ab\_ab'.
```


```
apply inc\_inv. apply (comp\_inc\_compat\_ab\_b\ H). Qed.
```

#### 13.2.3 鋭敏性

#### この節の補題は以下の 1 つのみだが、証明が異様に長いため単独の節を設ける.

Lemma 306 (sharpness) In below figure,

$$\alpha \cdot \gamma^{\sharp} \sqcap \beta \cdot \delta^{\sharp} = (\alpha \cdot p^{\sharp} \sqcap \beta \cdot q^{\sharp}) \cdot (p \cdot \gamma^{\sharp} \sqcap q \cdot \delta^{\sharp}).$$



```
Lemma sharpness \{ W X Y Z : eqType \}
 \{alpha: Rel\ X\ Y\}\ \{beta: Rel\ X\ Z\}\ \{gamma: Rel\ W\ Y\}\ \{delta: Rel\ W\ Z\}:
 (alpha \cdot gamma \#) \quad (beta \cdot delta \#) =
 ((alpha \cdot fst_r \ Y \ Z \ \#) \quad (beta \cdot snd_r \ Y \ Z \ \#))
  • ((fst_r \ Y \ Z \ \bullet \ qamma \ \#) \ (snd_r \ Y \ Z \ \bullet \ delta \ \#)).
Proof.
apply inc\_antisym.
move: (rationality \_ \_ alpha) \Rightarrow H.
move: (rationality \_ \_ beta) \Rightarrow H0.
move: (rationality \_ \_ (gamma \#)) \Rightarrow H1.
move: (rationality \_ \_ (delta \#)) \Rightarrow H2.
elim H \Rightarrow R.
elim \Rightarrow f\theta.
elim \Rightarrow g\theta H3.
elim H\theta \Rightarrow R\theta.
elim \Rightarrow f1.
elim \Rightarrow g1 H_4.
elim H1 \Rightarrow R1.
elim \Rightarrow h\theta.
elim \Rightarrow k0 H5.
elim H2 \Rightarrow R2.
elim \Rightarrow h1.
elim \Rightarrow k1 H6.
```

```
\overline{\text{move}}: (rationality \_ \_ (g0 \cdot h0 \#)) \Rightarrow H7.
move: (rationality \_ \_ (g1 \cdot h1 \#)) \Rightarrow H8.
move: (rationality \_ \_ ((alpha \cdot gamma \#))
                                                         (beta \cdot delta \#)) \Rightarrow H9.
elim H7 \Rightarrow R3.
elim \Rightarrow s\theta.
elim \Rightarrow t0 \ H10.
elim H8 \Rightarrow R4.
elim \Rightarrow s1.
elim \Rightarrow t1 \ H11.
elim H9 \Rightarrow R5.
elim \Rightarrow x.
elim \Rightarrow z H12.
assert (alpha \cdot gamma \# = (f0 \# \cdot (s0 \# \cdot t0)) \cdot k0).
replace alpha with (f0 \# \cdot g0).
replace (gamma \#) with (h0 \# \cdot k0).
rewrite -comp\_assoc (@comp\_assoc\_\_\_\_ (f0 \#)).
apply f_{-}equal2.
apply f_equal.
apply H10.
by [].
apply Logic.eq\_sym.
apply H5.
apply Logic.eq\_sym.
apply H3.
assert (beta • delta \# = (f1 \# \bullet (s1 \# \bullet t1)) \bullet k1).
replace beta with (f1 \# \cdot q1).
replace (delta \#) with (h1 \# \cdot k1).
rewrite -comp\_assoc (@comp\_assoc\_\_\_\_ (f1 \#)).
apply f_{-}equal2.
apply f_equal.
apply H11.
by [].
apply Logic.eq\_sym.
apply H6.
apply Logic.eq_sym.
apply H_4.
assert (t\theta \cdot h\theta = s\theta \cdot g\theta).
apply function_inc.
apply function_comp.
apply H10.
apply H5.
apply function_comp.
```

```
apply H10.
apply H3.
apply (@inc\_trans\_\_\_(s\theta \cdot ((s\theta \# \cdot t\theta) \cdot h\theta))).
rewrite comp_assoc -comp_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H10.
apply comp_inc_compat_ab_ab'.
replace (s\theta \# \cdot t\theta) with (g\theta \cdot h\theta \#).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H5.
apply H10.
assert (t1 \cdot h1 = s1 \cdot g1).
apply function_inc.
apply function\_comp.
apply H11.
apply H6.
apply function_comp.
apply H11.
apply H_4.
apply (@inc\_trans \_ \_ \_ (s1 \cdot ((s1 \# \cdot t1) \cdot h1))).
rewrite comp_assoc -comp_assoc.
apply comp\_inc\_compat\_b\_ab.
apply H11.
apply comp\_inc\_compat\_ab\_ab'.
replace (s1 \# \cdot t1) with (q1 \cdot h1 \#).
rewrite comp_{-}assoc.
apply comp\_inc\_compat\_ab\_a.
apply H6.
apply H11.
remember ((x \cdot (s0 \cdot f0) \#) (z \cdot (t0 \cdot k0) \#)) as m0.
remember ((x \cdot (s1 \cdot f1) \#) (z \cdot (t1 \cdot k1) \#)) as m1.
assert (total_r \ m\theta).
rewrite Heqm0.
apply domain_corollary1.
apply H12.
apply H12.
replace (x \# \cdot z) with ((alpha \cdot gamma \#) \pmod{beta} \cdot delta \#)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
rewrite comp_inv H13 -comp_assoc comp_assoc.
apply inc\_reft.
apply H12.
```

```
assert (total_r m1).
rewrite Heqm1.
apply domain_corollary1.
apply H12.
apply H12.
replace (x \# \cdot z) with ((alpha \cdot gamma \#) (beta \cdot delta \#)).
apply @inc\_trans \_ \_ \_ \_ (cap\_r).
rewrite comp_inv H14 -comp_assoc comp_assoc.
apply inc\_reft.
apply H12.
remember (m\theta \cdot (s\theta \cdot g\theta)) as n\theta.
remember (m1 \cdot (s1 \cdot g1)) as n1.
assert (total_r \ n\theta).
rewrite Hegn0.
apply (total_comp H17).
apply total_comp.
apply H10.
apply H3.
assert (total_r \ n1).
rewrite Hegn1.
apply (total_comp H18).
apply total_comp.
apply H11.
apply H_4.
assert (total_r ((n0 \cdot fst_r Y Z \#) (n1 \cdot snd_r Y Z \#))).
apply (domain_corollary1 H19 H20).
rewrite fst\_snd\_universal.
apply inc\_alpha\_universal.
assert ((x \# \cdot n\theta))
                          alpha).
replace alpha with (f0 \# \cdot g0).
rewrite Heqn0 Heqm0.
apply (@inc\_trans \_ \_ \_ (((x \# \cdot x) \cdot f0 \#) \cdot ((s0 \# \cdot s0) \cdot q0))).
rewrite comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
\verb"rewrite" -comp\_assoc -comp\_assoc -comp\_assoc -comp\_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc -comp_inv.
apply cap_{-}l.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_b.
apply H12.
```

```
apply comp\_inc\_compat\_ab\_b.
apply H10.
apply Logic.eq_sym.
apply H3.
assert ((x \# \cdot n1)  beta).
replace beta with (f1 \# \bullet q1).
rewrite Heqn1 Heqm1.
apply (@inc\_trans \_ \_ \_ (((x \# \cdot x) \cdot f1 \#) \cdot ((s1 \# \cdot s1) \cdot g1))).
rewrite comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply comp_inc_compat_ab_a'b.
rewrite comp_assoc -comp_inv.
apply cap_{-}l.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_b.
apply H12.
apply comp\_inc\_compat\_ab\_b.
apply H11.
apply Logic.eq_sym.
apply H_4.
assert ((n0 \# \cdot z) \quad gamma \#).
replace (gamma \#) with (h0 \# \cdot k0).
rewrite Heqn0 Heqm0 -H15 comp_inv comp_inv inv_cap_distr.
apply (@inc\_trans \_ \_ \_ ((h0 \# \cdot (t0 \# \cdot t0)) \cdot (k0 \cdot (z \# \cdot z)))).
rewrite -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp_assoc (@comp_inv _ _ z) inv_invol.
apply cap_r.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a.
apply H10.
apply comp\_inc\_compat\_ab\_a.
apply H12.
apply Logic.eq_sym.
apply H5.
assert ((n1 \# \cdot z) \text{ delta } \#).
replace (delta \#) with (h1 \# \cdot k1).
```

```
rewrite Hegn1 Hegm1 -H16 comp_inv comp_inv inv_cap_distr.
apply (@inc\_trans \_ \_ \_ ((h1 \# \cdot (t1 \# \cdot t1)) \cdot (k1 \cdot (z \# \cdot z)))).
rewrite -comp_assoc -comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
rewrite comp_assoc comp_assoc comp_assoc.
apply comp\_inc\_compat\_ab\_ab'.
apply comp_inc_compat_ab_ab'.
rewrite -comp_assoc (@comp_inv _ _ z) inv_invol.
apply cap_r.
apply comp\_inc\_compat.
apply comp\_inc\_compat\_ab\_a.
apply H11.
apply comp\_inc\_compat\_ab\_a.
apply H12.
apply Logic.eq\_sym.
apply H6.
replace ((alpha \cdot gamma \#) (beta \cdot delta \#)) with (x \# \cdot z).
apply (@inc\_trans\_\_\_((x \# \cdot (((n0 \cdot fst\_r Y Z \#) (n1 \cdot snd\_r Y Z \#)) \cdot (((n0 \cdot fst\_r Y Z \#)))))
• fst_r Y Z \#) (n1 \cdot snd_r Y Z \#))) \#)) • z)).
apply comp_inc_compat_ab_a'b.
apply (comp\_inc\_compat\_a\_ab\ H21).
rewrite -comp_assoc comp_assoc.
apply comp\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
rewrite -comp\_assoc.
apply (comp\_inc\_compat\_ab\_a'b H22).
rewrite - comp_assoc.
apply (comp\_inc\_compat\_ab\_a'b H23).
rewrite inv_cap_distr comp_inv comp_inv inv_invol inv_invol.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply cap\_inc\_compat.
rewrite comp_{-}assoc.
apply (comp\_inc\_compat\_ab\_ab', H24).
rewrite comp\_assoc.
apply (comp\_inc\_compat\_ab\_ab', H25).
apply Logic.eq_sym.
apply H12.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_l)).
apply cap\_inc\_compat.
apply (@inc\_trans \_ \_ \_ \_ (comp\_cap\_distr\_r)).
apply (@inc\_trans \_ \_ \_ \_ (cap\_l)).
```

```
rewrite -comp_assoc (@comp_assoc _ _ _ alpha).

apply comp_inc_compat_ab_a'b.

apply fst_function.

apply (@inc_trans _ _ _ (comp_cap_distr_r)).

apply (@inc_trans _ _ _ (cap_r)).

rewrite -comp_assoc (@comp_assoc _ _ _ beta).

apply comp_inc_compat_ab_a'b.

apply comp_inc_compat_ab_a.

apply snd_function.

Qed.
```

#### 13.2.4 分配法則

```
Lemma 307 (prod_cap_distr_l) Let \alpha: A \to B and \beta, \beta': A \to C. Then, \alpha \top (\beta \sqcap \beta') = (\alpha \top \beta) \sqcap (\alpha \top \beta').
Lemma prod_cap_distr_l { A B C : eaType} { calpha : Rel_A B } { heta beta' : Rel_A C } :
```

Lemma  $prod\_cap\_distr\_l$  { $A \ B \ C : eqType$ } { $alpha : Rel \ A \ B$ } {beta  $beta' : Rel \ A \ C$ }:  $Rel\_prod \ alpha$  (beta  $beta' ) = Rel\_prod \ alpha$  beta  $Rel\_prod \ alpha \ beta'$ .

Proof.

rewrite  $/Rel\_prod$ .

rewrite  $-cap\_assoc$  (@ $cap\_comm \ \_ \ \_ \ (alpha \ \cdot \ fst\_r \ B \ C \ \#)$ )  $-cap\_assoc \ cap\_idem \ cap\_assoc$ .

apply f\_equal.

apply  $function\_cap\_distr\_r$ .

apply  $snd\_function$ .

Qed.

```
Lemma 308 (prod_cap_distr_r) Let \alpha, \alpha' : A \to B and \beta : A \to C. Then, (\alpha \sqcap \alpha') \top \beta = (\alpha \top \beta) \sqcap (\alpha' \top \beta).
```

```
Lemma prod\_cap\_distr\_r {A \ B \ C : eqType} {alpha \ alpha' : Rel \ A \ B} {beta : Rel \ A \ C}: Rel\_prod \ (alpha \ alpha') beta = Rel\_prod \ alpha beta Rel\_prod \ alpha' beta. Proof. rewrite /Rel\_prod. rewrite /Rel\_prod. rewrite cap\_assoc (@cap\_comm\_\_ (beta • snd\_r B C #)) cap\_assoc cap\_idem -cap\_assoc. apply (@f\_equal \_\_ (fun x \Rightarrow @cap\_\_\_x (beta • snd\_r B C #))). apply function\_cap\_distr\_r. apply fst\_function. Qed.
```

**Lemma 309** (prod\_cup\_distr\_l) Let  $\alpha : A \rightarrow B$  and  $\beta, \beta' : A \rightarrow C$ . Then,

$$\alpha \top (\beta \sqcup \beta') = (\alpha \top \beta) \sqcup (\alpha \top \beta').$$

Proof.

by [rewrite  $-cap\_cup\_distr\_l$   $-comp\_cup\_distr\_r$ ]. Qed.

**Lemma 310** (prod\_cup\_distr\_r) Let  $\alpha, \alpha' : A \to B$  and  $\beta : A \to C$ . Then,

$$(\alpha \sqcup \alpha') \top \beta = (\alpha \top \beta) \sqcup (\alpha' \top \beta).$$

Lemma  $prod\_cup\_distr\_r$  { $A \ B \ C : eqType$ } { $alpha \ alpha' : Rel \ A \ B$ } {beta :  $Rel \ A \ C$ }:  $Rel\_prod \ (alpha \ alpha')$  beta =  $Rel\_prod \ alpha$  beta  $Rel\_prod \ alpha'$  beta.

Proof.

by [rewrite  $-cap\_cup\_distr\_r$   $-comp\_cup\_distr\_r$ ]. Qed.

**Lemma 311 (comp\_prod\_distr\_l)** Let  $\alpha:A\rightarrow B,\ \beta:B\rightarrow C$  and  $\gamma:B\rightarrow D.$  Then,

$$\alpha \cdot (\beta \top \gamma) \sqsubseteq \alpha \cdot \beta \top \alpha \cdot \gamma.$$

Lemma comp\_prod\_distr\_l

 $\{A\ B\ C\ D: eqType\}\ \{alpha: Rel\ A\ B\}\ \{beta: Rel\ B\ C\}\ \{gamma: Rel\ B\ D\}: alpha \cdot Rel\_prod\ beta\ gamma \quad Rel\_prod\ (alpha \cdot beta)\ (alpha \cdot gamma).$ 

Proof.

rewrite  $/Rel_prod$ .

rewrite  $comp\_assoc$   $comp\_assoc$ .

apply  $comp\_cap\_distr\_l$ .

Qed.

**Lemma 312 (function\_prod\_distr\_l)** Let  $\alpha : A \rightarrow B$  be a function,  $\beta : B \rightarrow C$  and  $\gamma : B \rightarrow D$ . Then,

$$\alpha \cdot (\beta \top \gamma) = \alpha \cdot \beta \top \alpha \cdot \gamma.$$

 $\{A\ B\ C\ D: eqType\}\ \{alpha: Rel\ A\ B\}\ \{$ beta:  $Rel\ B\ C\}\ \{gamma: Rel\ B\ D\}$ :  $function\_r\ alpha \rightarrow alpha \cdot Rel\_prod\$ beta  $gamma=Rel\_prod\ (alpha \cdot beta)\ (alpha \cdot gamma)$ .

Proof.

```
\begin{split} & \text{move} \Rightarrow H. \\ & \text{rewrite} \ / Rel\_prod. \\ & \text{rewrite} \ comp\_assoc \ comp\_assoc. \\ & \text{apply} \ (function\_cap\_distr\_l \ H). \\ & \text{Qed.} \end{split}
```

```
Lemma 313 (comp_prod_universal) Let \alpha: A \rightarrow B, \ \beta: B \rightarrow C \ and \ \gamma: D \rightarrow E. Then,
```

```
\alpha \cdot (\beta \top \nabla_{BD} \cdot \gamma) = \alpha \cdot \beta \top \nabla_{AD} \cdot \gamma.
```

```
Lemma comp\_prod\_universal
 \{A \ B \ C \ D \ E : eqType\} \{alpha : Rel \ A \ B\} \{beta : Rel \ B \ C\} \{gamma : Rel \ D \ E\}:
 alpha \cdot Rel\_prod \ \mathsf{beta} \ ( B \ D \cdot gamma) = Rel\_prod \ (alpha \cdot \mathsf{beta}) \ (
                                                                                A D \cdot qamma).
Proof.
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (comp\_prod\_distr\_l)).
apply prod_inc_compat_l.
rewrite - comp_assoc.
apply comp_inc_compat_ab_a'b.
apply inc\_alpha\_universal.
rewrite /Rel_prod.
rewrite comp_assoc.
apply (@inc\_trans\_\_\_\_ (dedekind1)).
apply comp\_inc\_compat\_ab\_ab'.
apply cap\_inc\_compat\_l.
rewrite comp_assoc comp_assoc -comp_assoc.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
```

**Lemma 314 (fst\_cap\_snd\_distr)** Let  $u, v : A \times B \rightarrow A \times B$  and  $u, v \sqsubseteq id_{A \times B}$ ,  $p : B \times C \rightarrow B$  and  $q : B \times C \rightarrow C$  be projections. Then,

$$p^{\sharp} \cdot (u \sqcap v) \cdot q = p^{\sharp} \cdot u \cdot q \sqcap p^{\sharp} \cdot v \cdot q.$$

```
Lemma fst\_cap\_snd\_distr \{A\ B: eqType\}\ \{u\ v: Rel\ (prod\_eqType\ A\ B)\ (prod\_eqType\ A\ B)\}: u\ Id\ (prod\_eqType\ A\ B) \rightarrow v\ Id\ (prod\_eqType\ A\ B) \rightarrow fst\_r\ A\ B\ \#\ \cdot\ (u\ v)\ \cdot\ snd\_r\ A\ B = ((fst\_r\ A\ B\ \#\ \cdot\ u)\ \cdot\ snd\_r\ A\ B)\ ((fst\_r\ A\ B\ \#\ \cdot\ v)\ \cdot\ snd\_r\ A\ B). Proof. move \Rightarrow H\ H0. apply inc\_antisym.
```

```
apply (fun H' \Rightarrow @inc\_trans\_\_\_\_H' (comp\_cap\_distr\_r)).
apply comp\_inc\_compat\_ab\_a'b.
apply comp\_cap\_distr\_l.
apply (@inc_trans _ _ _ (dedekind1)).
rewrite -(dedekind_id3 H H0) -(@comp_assoc _ _ _ u) (@comp_assoc _ _ _ (fst_r A
B \# \cdot u) v).
apply comp\_inc\_compat\_ab\_ab'.
rewrite cap_comm comp_assoc -comp_assoc.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
apply comp\_inc\_compat\_ab\_b.
rewrite comp_inv comp_inv inv_invol -fst_snd_cap_id.
apply cap\_inc\_compat.
rewrite comp_assoc (dedekind_id1 H).
apply (comp\_inc\_compat\_ab\_b\ H).
rewrite -comp_assoc (dedekind_id1 H0).
apply (comp_inc_compat_ab_a H0).
Qed.
```

## Chapter 14

### Library Point\_Axiom

```
Require Import Basic\_Notations.
Require Import Basic\_Lemmas.
Require Import Relation\_Properties.
Require Import Functions\_Mappings.
Require Import Dedekind.
Require Import Rationality.
```

#### 14.1 I-点

#### 14.1.1 I-点の定義

Dedekind 圏における域 X の I-点 x とは、関数  $x:I \to X$  のことであり、記号  $x \in X$  によって表される。 また関係  $\rho:I \to X$  と I-点  $x:I \to X$  に対して、記号  $x \in \rho$  で  $x \sqsubseteq \rho$  を表すものとする.

ちなみに I-点の定義  $x \in X$  は  $x \in \nabla_{IX}$  と言い換えることも可能である.

```
Definition point\_inc\ \{X: eqType\}\ (x\ rho: Rel\ i\ X):=function\_r\ x \wedge x rho.
Definition point\ \{X: eqType\}\ (x: Rel\ i\ X):=point\_inc\ x\ (\ i\ X).
```

#### 14.1.2 I-点の性質

```
Lemma 315 (point_property1) Let x, y \in X. Then, x = y \Leftrightarrow x \cdot y^{\sharp} = id_{I}.
```

```
Lemma point_property1 \{X : eqType\}\ \{x \ y : Rel \ i \ X\}: point \ x \to point \ y \to (x = y \leftrightarrow x \cdot y \# = Id \ i).
Proof.
```

#### CHAPTER 14. LIBRARY POINT\_AXIOM

```
move \Rightarrow H H0.
split; move \Rightarrow H1.
apply inc\_antisym.
rewrite unit_identity_is_universal.
apply inc_alpha_universal.
rewrite H1.
apply H0.
apply Logic.eq_sym.
apply function_inc.
apply H0.
apply H.
rewrite -(@comp\_id\_l\_\_y) -H1 comp\_assoc.
apply comp\_inc\_compat\_ab\_a.
apply H0.
Qed.
  Lemma 316 (point_property2a, point_property2b) Let \rho: I \to X be a total re-
  lation. Then,
                                      \rho \cdot \rho^{\sharp} = \rho \cdot \nabla_{XI} = id_I.
Lemma point\_property2a \{X : eqType\} \{rho : Rel \ i \ X\}:
 total\_r \ rho \rightarrow rho \cdot rho \# = Id \ i.
Proof.
move \Rightarrow H.
apply inc\_antisym.
rewrite unit\_identity\_is\_universal.
apply inc\_alpha\_universal.
apply H.
Qed.
Lemma point\_property2b \{X : eqType\} \{rho : Rel \ i \ X\}:
 total\_r \ rho \rightarrow rho \cdot rho \# = rho \cdot X \ i.
Proof.
move \Rightarrow H.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_ab'.
apply inc\_alpha\_universal.
rewrite (point_property2a H) unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
```

#### CHAPTER 14. LIBRARY POINT\_AXIOM

Lemma 317 (point\_property3) Let  $\rho: I \to X$ . Then,

```
\exists x \in \rho \Rightarrow "\rho \text{ is total"} \land \rho \neq \phi_{IX}.
Lemma point\_property3 \{X : eqType\} \{rho : Rel \ i \ X\}:
 (\exists x : Rel \ i \ X, point\_inc \ x \ rho) \rightarrow total\_r \ rho \land rho \neq
                                                                           i X.
Proof.
elim \Rightarrow x H.
assert (total_r rho).
elim H \Rightarrow H0 \ H1.
elim H0 \Rightarrow H2 \ H3.
apply @inc_trans_H = H2.
apply comp\_inc\_compat.
apply H1.
apply (@inc_inv_{-1} - H1).
split.
apply H0.
move \Rightarrow H1.
rewrite /total_r in H0.
rewrite H1 comp_empty_l in H0.
apply unit_identity_not_empty.
apply inc\_antisym.
apply H0.
apply inc\_empty\_alpha.
Qed.
  Lemma 318 (point_property4)
                                \exists x \in X \Rightarrow "\nabla_{IX} \text{ is total"} \land \nabla_{IX} \neq \phi_{IX}.
Lemma point\_property4 \{X : eqType\}:
 (\exists x : Rel \ i \ X, \ point \ x) \rightarrow total\_r \ (i \ X) \land (i \ X) \neq
                                                                             i X.
Proof.
move \Rightarrow H.
apply (@point\_property3 \_ ( i X) H).
Qed.
```

#### 14.2 I-点に関する諸公理

#### 14.2.1 点公理

```
この"点公理"を使えば、I-点に関する様々な定理や補題が導出できる.
```

Lemma 319 (point\_axiom) Let  $\rho: I \to X$ . Then,

```
\rho = \sqcup_{x \in \rho} x.
```

```
Lemma lemma\_for\_PA \{X : eqType\} \{rho : Rel \ i \ X\}:
                i(X) \rightarrow False) \rightarrow False) \rightarrow rho = i(X).
Proof.
move \Rightarrow H.
case (@unit_empty_or_universal (rho • rho \#)) \Rightarrow H0.
apply inc\_antisym.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
rewrite H0 \ comp\_empty\_l.
apply inc\_reft.
apply inc\_empty\_alpha.
apply False_ind.
apply H.
move \Rightarrow H1.
rewrite H1 comp_-empty_-l in H0.
apply (unit_empty_not_universal H0).
Qed.
Lemma point\_axiom \{X : eqType\} \{rho : Rel \ i \ X\}:
 rho =
            \{ fun \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho \} \ id.
Proof.
apply inc\_antisym.
apply bool_lemma2.
assert ((\exists x : Rel \ i \ X, point\_inc \ x \ (( \_{fun} \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho) \ id)
( _{\text{-}}\{\text{fun } x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho\} \ id) \hat{\ })) \rightarrow False).
move \Rightarrow H.
move: (point\_property3 \ H) \Rightarrow H0.
apply H0.
apply cap\_complement\_empty.
assert ((\exists x : Rel \ i \ X, point\_inc \ x \ (rho) \ ( \_\{fun \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho\} \ id)
\hat{})) \rightarrow False).
move \Rightarrow H0.
apply H.
elim H0 \Rightarrow x H1.
```

#### CHAPTER 14. LIBRARY POINT\_AXIOM

```
\exists x.
split.
apply H1.
apply inc\_cap.
split.
assert (point_inc \ x \ rho).
split.
apply H1.
elim H1 \Rightarrow H2 \ H3.
apply inc\_cap in H3.
apply H3.
clear H1.
move: x H2.
apply inc\_cupP.
apply inc_refl.
elim H1 \Rightarrow H2 \ H3.
apply inc\_cap in H3.
apply H3.
apply lemma\_for\_PA.
\mathtt{move} \Rightarrow \mathit{H1}.
apply H0.
apply weak_axiom_of_choice.
rewrite /total_r.
remember (rho
                    ( \{ fun \ x : Rel \ i \ X \Rightarrow point\_inc \ x \ rho \} \ id) \hat{)}  as rho'.
case (@unit\_empty\_or\_universal\ (rho' \cdot rho' \#)) \Rightarrow H2.
apply False\_ind.
apply H1.
apply inc\_antisym.
apply (@inc_trans _ _ _ _ (relation_rel_inv_rel)).
rewrite H2 comp\_empty\_l.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite H2.
apply inc_alpha_universal.
apply inc\_cupP.
move \Rightarrow beta H.
apply H.
Qed.
```

```
Lemma 320 (PA_corollary1)
```

 $\nabla_{IX} = \sqcup_{x \in X} x.$ 

#### CHAPTER 14. LIBRARY POINT\_AXIOM

```
Lemma PA\_corollary1 \{X:eqType\}: i\ X = \_\{\text{fun } x:Rel\ i\ X\Rightarrow point\ x\}\ id. Proof. apply point\_axiom. Qed.
```

#### Lemma 321 (PA\_corollary2)

$$id_X = \sqcup_{x \in X} x^{\sharp} \cdot x.$$

```
Lemma PA\_corollary2 \{X : eqType\}:
             \{ \text{fun } x : Rel \ i \ X \Rightarrow point \ x \} \ (\text{fun } x : Rel \ i \ X \Rightarrow x \# \cdot x \}.
 Id X =
Proof.
rewrite -(@cap_universal _ _ (Id X)) -lemma_for_tarski2 PA_corollary1.
rewrite comp\_cupP\_distr\_l cap\_cupP\_distr\_l.
apply cupP_{-}eq.
move \Rightarrow alpha H.
apply inc\_antisym.
rewrite cap\_comm.
apply (@inc\_trans \_ \_ \_ \_ (dedekind2)).
rewrite comp_id_l cap_comm cap_universal.
apply inc_refl.
apply inc\_cap.
split.
apply H.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
Qed.
```

#### Lemma 322 (PA\_corollary3) Let $\alpha, \beta: X \to Y$ . Then,

$$(\forall x \in X, x \cdot \alpha = x \cdot \beta) \Rightarrow \alpha = \beta.$$

```
Lemma PA\_corollary3 \{X \ Y : eqType\} \{alpha \ beta : Rel \ X \ Y\}: (\forall \ x : Rel \ i \ X, \ point \ x \to x \ \bullet \ alpha = x \ \bullet \ beta) \to alpha = beta. Proof. move \Rightarrow H. rewrite -(@comp\_id\_l\_\_\_alpha) -(@comp\_id\_l\_\_\_beta) PA\_corollary2. rewrite comp\_cupP\_distr\_r comp\_cupP\_distr\_r. apply cupP\_eq. move \Rightarrow gamma \ H0. by [rewrite comp\_assoc \ comp\_assoc \ (H \ gamma \ H0)]. Qed.
```

Lemma 323 (total\_axiom) Let  $\rho: I \to X$ . Then,

#### 14.2.2 その他の公理

```
\rho \neq \phi_{IX} \Rightarrow id_I = \rho \cdot \rho^{\sharp}.
Lemma total\_axiom \{X : eqType\} \{rho : Rel \ i \ X\}:
            i X \rightarrow Id i = rho \cdot rho \#.
 rho \neq
Proof.
move \Rightarrow H.
case (@unit_empty_or_universal (rho • rho \#)) \Rightarrow H0.
apply False_ind.
apply H.
apply inc\_antisym.
apply (@inc_trans _ _ _ (relation_rel_inv_rel)).
rewrite H0 comp_-empty_-l.
apply inc\_reft.
apply inc\_empty\_alpha.
by [rewrite H0 unit_identity_is_universal].
Qed.
  Lemma 324 (nonempty_axiom) Let \rho: I \to X. Then,
```

 $\rho \neq \phi_{IX} \Rightarrow \exists x \dot{\in} \rho.$ 

```
Lemma nonempty\_axiom \{X: eqType\} \{rho: Rel \ i \ X\}: rho \neq i \ X \rightarrow \exists \ x: Rel \ i \ X, \ point\_inc \ x \ rho.

Proof.

move: (@weak\_axiom\_of\_choice \_ rho) \Rightarrow H.

move \Rightarrow H0.

apply H.

rewrite /total\_r.

rewrite (total\_axiom \ H0).

apply inc\_reft.

Qed.
```

```
Lemma 325 (axiom_of_subobjects) Let \rho: I \to X. Then, \exists S, \exists j: S \to X, \rho = \nabla_{IS} \cdot j \wedge j \cdot j^{\sharp} = id_{S}.
```

```
Lemma axiom\_of\_subobjects \{X : eqType\} \{rho : Rel \ i \ X\}: \exists \ (S : eqType)(j : Rel \ S \ X), \ rho = i \ S \ \cdot \ j \land j \ \cdot \ j \ \# = Id \ S. Proof.
```

#### CHAPTER 14. LIBRARY POINT\_AXIOM

```
elim (@rationality \_ \_ rho) \Rightarrow R.
elim \Rightarrow f.
elim \Rightarrow q.
elim \Rightarrow H.
elim \Rightarrow H0.
elim \Rightarrow H1 H2.
\exists R.
\exists q.
split.
rewrite H1.
apply inc\_antisym.
apply comp\_inc\_compat\_ab\_a'b.
apply inc\_alpha\_universal.
apply comp\_inc\_compat\_ab\_a'b.
apply (@inc\_trans \_ \_ \_ ( i R \cdot (f \cdot f \#))).
apply comp\_inc\_compat\_a\_ab.
apply H.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
rewrite -H2 cap_comm inc_def1.
assert ((f \# \cdot g)
                         rho).
rewrite H1.
apply inc\_reft.
apply (function\_move1 \ H) in H3.
apply (@inc\_trans \_ \_ \_ ((f \cdot rho) \cdot (f \cdot rho) \#)).
apply comp\_inc\_compat.
apply H3.
apply (@inc_inv_- - H3).
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ rho).
apply comp\_inc\_compat\_ab\_ab'.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
Qed.
```

```
Lemma 326 (axiom_of_choice) Let \alpha: X \to Y be a total relation. Then, \exists f: X \to Y, f \sqsubseteq \alpha.
```

```
Lemma axiom\_of\_choice \{X \ Y : eqType\} \{alpha : Rel \ X \ Y\}: total\_r \ alpha \rightarrow \exists \ (f : Rel \ X \ Y), function\_r \ f \land f \ alpha.
```

```
Proof.
move \Rightarrow H.
case (@unit\_empty\_or\_universal\ (\ i\ X\ \cdot\ X\ i)) \Rightarrow H0.
\exists (X Y).
repeat split.
assert (Id X =
                      XX).
apply inc\_antisym.
apply (@inc\_trans \_ \_ \_ \_ (@inc\_alpha\_universal \_ \_ \_)).
apply (@inc\_trans\_\_\_(XX \cdot XX)).
apply comp\_inc\_compat\_a\_ab.
apply inc\_alpha\_universal.
rewrite -unit\_universal\ comp\_assoc\ -(@comp\_assoc\ \_\_\_\_\ (\ i\ X)).
rewrite H0 comp_empty_l comp_empty_r.
apply inc\_reft.
apply inc\_empty\_alpha.
rewrite /total_r.
rewrite H1 comp\_empty\_l.
apply inc\_reft.
rewrite /univalent_r.
rewrite comp\_empty\_r.
apply inc\_empty\_alpha.
apply inc\_empty\_alpha.
assert (\forall x : Rel \ i \ X, \ point \ x \rightarrow \exists \ g : Rel \ i \ Y, \ function\_r \ g \land g \qquad (x \cdot alpha)).
move \Rightarrow x H1.
apply weak_axiom_of_choice.
apply total\_comp.
apply H1.
apply H.
assert (\forall x : Rel \ i \ X, point \ x \rightarrow \exists \ q' : Rel \ X \ Y, univalent_r \ q' \land q'
                                                                                    alpha).
move \Rightarrow x H2.
elim (H1 \ x \ H2) \Rightarrow g \ H3.
\exists (x \# \cdot q).
split.
rewrite /univalent_r.
rewrite comp_inv inv_invol.
apply (@inc\_trans\_\_\_(g \# \bullet g)).
rewrite comp\_assoc.
apply comp\_inc\_compat\_ab\_ab'.
rewrite -comp\_assoc.
apply comp\_inc\_compat\_ab\_b.
rewrite unit_identity_is_universal.
apply inc\_alpha\_universal.
```

#### CHAPTER 14. LIBRARY POINT\_AXIOM

```
apply H3.
apply function_move1.
apply H2.
apply H3.
assert (\exists f' : Rel \ i \ X \rightarrow Rel \ X \ Y, function_r \ ( \_\{\mathbf{fun} \ x \Rightarrow point \ x\} \ f') \land ( \_\{\mathbf{fun} \ x \Rightarrow point \ x\} \ f') \land ( \_\{\mathbf{fun} \ x \Rightarrow point \ x\} \ f')
\Rightarrow point x \} f'
                        alpha).
move: (@weak\_axiom\_of\_choice\_(iX \cdot alpha)) \Rightarrow H1.
assert (total_r ( i X \cdot alpha)).
rewrite /total_{-}r.
rewrite comp_inv comp_assoc -(@comp_assoc _ _ _ alpha).
{\tt rewrite} \ unit\_identity\_is\_universal \ -H0 \ inv\_universal.
apply comp\_inc\_compat\_ab\_ab'.
apply (comp\_inc\_compat\_b\_ab\ H).
apply H1 in H2.
elim H2 \Rightarrow g H3.
Qed.
```

## Bibliography

[1] R. Affeldt and M. Hagiwara. Formalization of Shannon 's Theorems in SSReflect-Coq. In 3rd Conference on Interactive Theorem Proving, LNCS 7406, 233–249, 2012.