

Mathematical Foundations of Political Methodology

Robert Gulotty

University of Chicago

November 7, 2019

Day 3: Random Variables: Discrete

- Random Variables
- Expectation, Variance
- Discrete Probability Mass Function
- Cumulative Distribution Functions
- Joint Probability Mass Functions

Random Variables

Given a sample space Ω , and a probability law P :

Definition

*A random variable is a **function** that assigns real numbers (usually) to events in a sample space Ω .*

$$X(\omega) : \Omega \rightarrow \mathbb{R}$$

Random Variables: Example

Sample space from tossing three coins:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

Let $Y(\omega)$ be the number of heads in outcome $\omega \in \Omega$.

Random Variables: Example

Sample space from tossing three coins:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

Let $Y(\omega)$ be the number of heads in outcome $\omega \in \Omega$.

- $Y(HHH) = 3$

Random Variables: Example

Sample space from tossing three coins:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

Let $Y(\omega)$ be the number of heads in outcome $\omega \in \Omega$.

- $Y(HHH) = 3$
- $Y(HHT) = Y(HTH) = Y(THH) = 2$

Random Variables: Example

Sample space from tossing three coins:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

Let $Y(\omega)$ be the number of heads in outcome $\omega \in \Omega$.

- $Y(HHH) = 3$
- $Y(HHT) = Y(HTH) = Y(THH) = 2$
- $Y(HTT) = Y(THT) = Y(TTH) = 1$

Random Variables: Example

Sample space from tossing three coins:

$$\Omega = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$$

Let $Y(\omega)$ be the number of heads in outcome $\omega \in \Omega$.

- $Y(HHH) = 3$
- $Y(HHT) = Y(HTH) = Y(THH) = 2$
- $Y(HTT) = Y(THT) = Y(TTH) = 1$
- $Y(TTT) = 0$

Random Variables: Example II

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|c|} \hline \bullet & \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet & \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet \bullet & \bullet \bullet \\ \hline \end{array} \right\}$$

Let $Z(\omega)$ be the dots on the die face $\omega \in \Omega$.

Random Variables: Example II

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|c|} \hline \bullet & \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet & \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet & \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet & \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|c|} \hline \bullet \bullet \bullet & \bullet \bullet \\ \hline \end{array} \right\}$$

Let $Z(\omega)$ be the dots on the die face $\omega \in \Omega$.

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet & \\ \hline \end{array}\right)=1$$

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet & \bullet \\ \hline \end{array}\right)=2$$

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet & \bullet \bullet \\ \hline \end{array}\right)=3$$

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet \bullet & \bullet \\ \hline \end{array}\right)=4$$

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet \bullet & \bullet \bullet \\ \hline \end{array}\right)=5$$

$$\blacksquare Z\left(\begin{array}{|c|c|} \hline \bullet \bullet \bullet & \bullet \bullet \\ \hline \end{array}\right)=6$$

Random Variables: Example III

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array} \right\}$$

Let $X(\omega)$ be 3 to the power of the number of dots on the die face $\omega \in \Omega$ minus 1.

Random Variables: Example III

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \\ \hline \end{array} \right\}$$

Let $X(\omega)$ be 3 to the power of the number of dots on the die face $\omega \in \Omega$ minus 1.

$$\blacksquare X\left(\begin{array}{|c|} \hline \bullet \\ \hline \end{array}\right) = 3^{1-1}$$

Random Variables: Example III

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}, \begin{array}{|c|c|c|c|c|c|} \hline \bullet & \bullet & \bullet & \bullet & \bullet & \bullet \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array} \right\}$$

Let $X(\omega)$ be 3 to the power of the number of dots on the die face $\omega \in \Omega$ minus 1.

$$\blacksquare X(\begin{array}{|c|c|c|c|c|c|} \hline \bullet & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline & & & & & \\ \hline \end{array}) = 3^{1-1} = 3^0 = 1$$

Random Variables: Example III

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array} \right\}$$

Let $X(\omega)$ be 3 to the power of the number of dots on the die face $\omega \in \Omega$ minus 1.

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \\ \hline \end{array}) = 3^{1-1} = 3^0 = 1$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array}) = 3^{2-1} = 3$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \\ \hline \end{array}) = 3^{3-1} = 9$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{4-1} = 27$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{5-1} = 81$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{6-1} = 243$$

Random Variables: Example III

Sample space from the number of dots on a die roll.

$$\Omega = \left\{ \begin{array}{|c|} \hline \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}, \begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array} \right\}$$

Let $X(\omega)$ be 3 to the power of the number of dots on the die face $\omega \in \Omega$ minus 1.

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \\ \hline \end{array}) = 3^{1-1} = 3^0 = 1$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \\ \hline \end{array}) = 3^{2-1} = 3$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \\ \hline \end{array}) = 3^{3-1} = 9$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{4-1} = 27$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{5-1} = 81$$

$$\blacksquare X(\begin{array}{|c|} \hline \bullet \bullet \bullet \bullet \bullet \bullet \\ \hline \end{array}) = 3^{6-1} = 243$$

$$X(\omega) = 3^{\omega-1}$$

functions

- Random Variables are functions.
- In R, we see functions look like a word followed by ():
 - mean(), median(), head(), str()
- Our own functions are written using:

```
function(input){  
  do something  
  return(output)}
```


Properties of Random Variables

Random variables (for us) will always give a number.

- X, Y, Z : capital letters for a random variable.
- x, y, z : are the output of the function.

Kinds of Random Variables

Definition

A random variable may be

- 1 *discrete: if outcomes are countable.*
- 2 *continuous: If outcomes can be any value in some interval.*
- 3 *mixed: if outcomes are a combination of events and intervals.*

Connecting random variables to probability

- X assigns some numbers to events.

Connecting random variables to probability

- X assigns some numbers to events.
- Remember, probability assigns a likelihood to an event.

Connecting random variables to probability

- X assigns some numbers to events.
- Remember, probability assigns a likelihood to an event.
- What is the probability that $X(\omega)$ gives some number x ?

Connecting random variables to probability

- X assigns some numbers to events.
- Remember, probability assigns a likelihood to an event.
- What is the probability that $X(\omega)$ gives some number x ?
- $P(X = x)$

Connecting random variables to probability

- X assigns some numbers to events.
- Remember, probability assigns a likelihood to an event.
- What is the probability that $X(\omega)$ gives some number x ?
- $P(X = x) \equiv P(\{\omega \in \Omega : X(\omega) = x\})$
- This is another function (of a function)

Discrete Probability Function

Definition

If X is a discrete random variable, the probability of each outcome x is provided by a probability function:

$$p_X(x) = P(X = x) = P(\{\omega \in \Omega : X(\omega) = x\})$$

- this is also called the *probability mass function* (pmf)
- We often just write $p(x)$, some people use $f_X(x)$

Building a Discrete Probability Function: Example

- Sample space: $\Omega = \{\text{Pashto}, \text{Not Pashto}\}$
- $X=1$ if $\omega = \text{Pashto}$
- $X=0$ if $\omega = \text{Not Pashto}$
- $P(X=1)=.3$

Building a Discrete Probability Function: Example

- Sample space: $\Omega = \{\text{Pashto}, \text{Not Pashto}\}$
- $X=1$ if $\omega = \text{Pashto}$
- $X=0$ if $\omega = \text{Not Pashto}$
- $P(X=1)=.3$

Often we will generalize the probability of success with a parameter p .

Often we will generalize the probability of success with a parameter p . A pmf can take on any value of x , so we must consider all possibilities of x .

- $x = 1$ with probability p
- $x = 0$ with probability $1 - p$
- all other values of x occur with probability 0.

- If $x = 1$

$$P(X = x) = P(X = 1) = P(\text{Pashto}) = p = .3$$

- If $x = 1$

$$P(X = x) = P(X = 1) = P(\text{Pashto}) = p = .3$$

- If $x = 0$

$$P(X = x) = P(X = 0) = P(\text{Not Pashto}) = 1 - p = .7$$

- If $x = 1$

$$P(X = x) = P(X = 1) = P(\text{Pashto}) = p = .3$$

- If $x = 0$

$$P(X = x) = P(X = 0) = P(\text{Not Pashto}) = 1 - p = .7$$

- all other values of x .

$$P(X = x) = 0$$

- If $x = 1$

$$P(X = x) = P(X = 1) = P(\text{Pashto}) = p = .3$$

- If $x = 0$

$$P(X = x) = P(X = 0) = P(\text{Not Pashto}) = 1 - p = .7$$

- all other values of x .

$$P(X = x) = 0$$

Combining these results, we get the probability function.

$$f_X(x) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \\ 0 & \text{otherwise} \end{cases}$$

Condensed Math notation for Bernoulli PMF

$$f_X(x) = \begin{cases} p & \text{if } x = 1 \\ 1 - p & \text{if } x = 0 \\ 0 & \text{otherwise} \end{cases}$$

can be written shorthand as:

$$f_X(x) = p^x(1-p)^{(1-x)}, \quad x \in \{0, 1\}$$

Where the 0 is left implicit.

Dichotomous process

To summarize

- If X only has two values 0 and 1 and $p = P(X = 1)$ then X has the probability function:

$$f_X(x) = p^x(1 - p)^{(1-x)}, \quad x \in \{0, 1\}$$

Dichotomous process

To summarize

- If X only has two values 0 and 1 and $p = P(X = 1)$ then X has the probability function:

$$f_X(x) = p^x(1 - p)^{(1-x)}, \quad x \in \{0, 1\}$$

- This is called a Bernoulli distribution:

$$X \sim \text{Bernoulli}(p)$$

- p is a *parameter*

Bernoulli Random Variable

Suppose we flip a fair coin and $Y = 1$ if the outcome is Heads .

$$Y \sim \text{Bernoulli}(1/2)$$

$$p(1) = (1/2)^1(1 - 1/2)^{1-1} = 1/2$$

$$p(0) = (1/2)^0(1 - 1/2)^{1-0} = (1 - 1/2)$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose.

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose.

Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose.

Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war. Engaging in the war will cost the country c .

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

$$E[U(\text{war})] = (\text{Utility}|\text{win}) \times P(\text{win}) + (\text{Utility}|\text{lose}) \times P(\text{lose})$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

$$\begin{aligned} E[U(\text{war})] &= (\text{Utility}|\text{win}) \times P(\text{win}) + (\text{Utility}|\text{lose}) \times P(\text{lose}) \\ &= (B - c)P(Y = 1) + (-c)P(Y = 0) \end{aligned}$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

$$\begin{aligned} E[U(\text{war})] &= (\text{Utility}|\text{win}) \times P(\text{win}) + (\text{Utility}|\text{lose}) \times P(\text{lose}) \\ &= (B - c)P(Y = 1) + (-c)P(Y = 0) \\ &= B \times p(Y = 1) - c(P(Y = 1) + P(Y = 0)) \end{aligned}$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

$$\begin{aligned} E[U(\text{war})] &= (\text{Utility}|\text{win}) \times P(\text{win}) + (\text{Utility}|\text{lose}) \times P(\text{lose}) \\ &= (B - c)P(Y = 1) + (-c)P(Y = 0) \\ &= B \times p(Y = 1) - c(P(Y = 1) + P(Y = 0)) \\ &= B \times \pi - c \end{aligned}$$

Example: Winning a War

Suppose country 1 is engaged in a conflict and can either win or lose. Define $Y = 1$ if the country wins and $Y = 0$ otherwise.

Then,

$$Y \sim \text{Bernoulli}(\pi)$$

Suppose country 1 is deciding whether to fight a war.

Engaging in the war will cost the country c .

If they win, country 1 receives B .

What is 1's expected utility from fighting a war?

$$\begin{aligned} E[U(\text{war})] &= (\text{Utility}|\text{win}) \times P(\text{win}) + (\text{Utility}|\text{lose}) \times P(\text{lose}) \\ &= (B - c)P(Y = 1) + (-c)P(Y = 0) \\ &= B \times p(Y = 1) - c(P(Y = 1) + P(Y = 0)) \\ &= B \times \pi - c \end{aligned}$$

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement.**

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement.**

Consider a coin, flipped twice. Suppose $X_1(T) = 0$, $X_1(H) = 1$,
 $X_2(T) = 0$, $X_2(H) = 1$

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement**.

Consider a coin, flipped twice. Suppose $X_1(T) = 0$, $X_1(H) = 1$,
 $X_2(T) = 0$, $X_2(H) = 1$

$$P(X_1 = 1, X_2 = 1) = P(X_1 = 1 \cap X_2 = 1) = P(X_1 = 1)P(X_2 = 1)$$

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement.**

Consider a coin, flipped twice. Suppose $X_1(T) = 0$, $X_1(H) = 1$,
 $X_2(T) = 0$, $X_2(H) = 1$

$$P(X_1 = 1, X_2 = 1) = P(X_1 = 1 \cap X_2 = 1) = P(X_1 = 1)P(X_2 = 1)$$

$$p(x_1, x_2) = p^{x_1}(1 - p)^{1-x_1}p^{x_2}(1 - p)^{1-x_2}$$

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement**.

Consider a coin, flipped twice. Suppose $X_1(T) = 0$, $X_1(H) = 1$,
 $X_2(T) = 0$, $X_2(H) = 1$

$$P(X_1 = 1, X_2 = 1) = P(X_1 = 1 \cap X_2 = 1) = P(X_1 = 1)P(X_2 = 1)$$

$$p(x_1, x_2) = p^{x_1}(1 - p)^{1-x_1}p^{x_2}(1 - p)^{1-x_2}$$

$$p(x_1, x_2) = p^{x_1+x_2}(1 - p)^{2-x_1-x_2}$$

Example: Sequence of Bernoulli trials

We might be interested in the outcome of several independent trials **ie. with replacement**.

Consider a coin, flipped twice. Suppose $X_1(T) = 0$, $X_1(H) = 1$,
 $X_2(T) = 0$, $X_2(H) = 1$

$$P(X_1 = 1, X_2 = 1) = P(X_1 = 1 \cap X_2 = 1) = P(X_1 = 1)P(X_2 = 1)$$

$$p(x_1, x_2) = p^{x_1}(1-p)^{1-x_1}p^{x_2}(1-p)^{1-x_2}$$

$$p(x_1, x_2) = p^{x_1+x_2}(1-p)^{2-x_1-x_2}$$

Let $X_1, \dots, X_n$ be independent Bernoulli random variables with the same parameter p .

$$p(x_1, \dots, x_n) = p(x_1)p(x_2)\dots p(x_n) = p^{x_1+\dots+x_n}(1-p)^{n-x_1-\dots-x_n}$$

$$\text{for } x_i \in \{0, 1\} \text{ and } i = 1, \dots, n$$

Expectation

What can we **expect** from a trial?

Expectation

What can we **expect** from a trial?

Value of random variable for any outcome weighted by the probability of observing that outcome

Expectation

What can we **expect** from a trial?

Value of random variable for any outcome weighted by the probability of observing that outcome
probability weighted average

Expectation

What can we **expect** from a trial?

Value of random variable for any outcome weighted by the probability of observing that outcome
probability weighted average

Definition

Expected Value: define the expected value of a function X as,

$$E[X] = \sum_{x:p(x)>0} xp(x)$$

Expectation

What can we **expect** from a trial?

Value of random variable for any outcome weighted by the probability of observing that outcome
probability weighted average


Definition

Expected Value: define the expected value of a function X as,


$$E[X] = \sum_{x:p(x)>0} xp(x)$$

In words: for all values of x with $p(x)$ greater than zero, take the weighted average of the values


$$E(X) = \sum_{x:p(x)>0} x * p(x) = 0 * .75 + 2 * .25 = \frac{1}{2}$$



$$E(X) = \sum_{x:p(x)>0} x * p(x) = 0 * .6 + 2 * .2 + 6 * .2 = 1.6$$



$$E(X) = \sum_{x:p(x)>0} x * p(x) = 0 * .6 + 2 * .2 + 9 * .2 = 2.2$$



Expectation informally

The expectation is based on the principle that the value of a future gain should be directly proportional to the chance of getting it.

$$E[X + Y] = E[X] + E[Y]$$

$$E[a] = a$$

$$E[aX] = aE[X]$$

$$E[E[X]] = E[X]$$

$$E[XY] \neq E[X] \times E[Y]$$

Properties of the Expectation

$$E[a] = a$$

Properties of the Expectation

$$E[a] = a$$

Proof: Suppose Y is a random variable $= a$ with probability 1 and 0 otherwise:

$$E[Y] = \sum_{y:p(y)>0} yp(y) = a * 1$$

Properties of the Expectation

$$E[a] = a$$

Proof: Suppose Y is a random variable $= a$ with probability 1 and 0 otherwise:

$$E[Y] = \sum_{y:p(y)>0} yp(y) = a * 1$$

Also:

$$E[aX] = \sum_{x:p(x)>0} a \times p(x)$$

Properties of the Expectation

$$E[a] = a$$

Proof: Suppose Y is a random variable $= a$ with probability 1 and 0 otherwise:

$$E[Y] = \sum_{y:p(y)>0} yp(y) = a * 1$$

Also:

$$E[aX] = \sum_{x:p(x)>0} a x p(x)$$

$$E[aX] = a \sum_{x:p(x)>0} x p(x)$$

Properties of the Expectation

$$E[a] = a$$

Proof: Suppose Y is a random variable $= a$ with probability 1 and 0 otherwise:

$$E[Y] = \sum_{y:p(y)>0} yp(y) = a * 1$$

Also:

$$E[aX] = \sum_{x:p(x)>0} a x p(x)$$

$$E[aX] = a \sum_{x:p(x)>0} x p(x)$$

Indicator Variables and Probabilities

Indicator Variables and Probabilities

Proposition

Indicator Variables and Probabilities

Proposition

Suppose A is an event. Define random variable I such that $I = 1$ if an outcome in A occurs and $I = 0$ if an outcome in A^c occurs. Then,

Indicator Variables and Probabilities

Proposition

Suppose A is an event. Define random variable I such that $I = 1$ if an outcome in A occurs and $I = 0$ if an outcome in A^c occurs. Then,

$$E[I] = P(A)$$

Indicator Variables and Probabilities

Proposition

Suppose A is an event. Define random variable I such that $I = 1$ if an outcome in A occurs and $I = 0$ if an outcome in A^c occurs. Then,

$$E[I] = P(A)$$

Proof.

Indicator Variables and Probabilities

Proposition

Suppose A is an event. Define random variable I such that $I = 1$ if an outcome in A occurs and $I = 0$ if an outcome in A^c occurs. Then,

$$E[I] = P(A)$$

Proof.

$$E[I] = 1 \times P(A) + 0 \times P(A^c)$$

Indicator Variables and Probabilities

Proposition

Suppose A is an event. Define random variable I such that $I = 1$ if an outcome in A occurs and $I = 0$ if an outcome in A^c occurs. Then,

$$E[I] = P(A)$$

Proof.

$$\begin{aligned} E[I] &= 1 \times P(A) + 0 \times P(A^c) \\ &= P(A) \end{aligned}$$



Variance

Expected value is a measure of **central tendency**.

Variance

Expected value is a measure of **central tendency**.

What about spread?

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

$$E[(X - E[X])^2] = \sum_x (x - E[X])^2 p(x)$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned} E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\ &= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x)\end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x) \\&= \sum_x x^2 p(x) - \sum_x 2xE[X] p(x) + \sum_x E[X]^2 p(x)\end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[x])^2 = (x - E[x])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x) \\&= \sum_x x^2 p(x) - \sum_x 2xE[X]p(x) + \sum_x E[X]^2 p(x) \\&= \sum_x x^2 p(x) - 2E[X] \sum_x xp(x) + \sum_x E[X]^2 p(x)\end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[X])^2 = (x - E[X])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x) \\&= \sum_x x^2 p(x) - \sum_x 2xE[X]p(x) + \sum_x E[X]^2 p(x) \\&= \sum_x x^2 p(x) - 2E[X] \sum_x xp(x) + \sum_x E[X]^2 p(x) \\&= E[X^2] - 2E[X]^2 + E[X]^2\end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[X])^2 = (x - E[X])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x) \\&= \sum_x x^2 p(x) - \sum_x 2xE[X]p(x) + \sum_x E[X]^2 p(x) \\&= \sum_x x^2 p(x) - 2E[X] \sum_x xp(x) + \sum_x E[X]^2 p(x) \\&= E[X^2] - 2E[X]^2 + E[X]^2 \\&= E[X^2] - E[X]^2\end{aligned}$$

Variance

Expected value is a measure of **central tendency**.

What about spread? **Variance**

- For each value, we might measure distance from center
 - Euclidean distance, squared $d(x, E[X])^2 = (x - E[X])^2$
- Then, we might take weighted average of these distances,

$$\begin{aligned}E[(X - E[X])^2] &= \sum_x (x - E[X])^2 p(x) \\&= \sum_x (x^2 - E[X]x - xE[X] + E[X]^2) p(x) \\&= \sum_x (x^2 - 2E[X]x + E[X]^2) p(x) \\&= \sum_x x^2 p(x) - \sum_x 2xE[X]p(x) + \sum_x E[X]^2 p(x) \\&= \sum_x x^2 p(x) - 2E[X] \sum_x xp(x) + \sum_x E[X]^2 p(x) \\&= E[X^2] - 2E[X]^2 + E[X]^2 \\&= E[X^2] - E[X]^2 \\&= \text{Var}(X)\end{aligned}$$

Variance

Definition

The variance of a random variable X , $\text{var}(X)$, is

$$\begin{aligned}\text{var}(X) &= E[(X - E[X])^2] \\ &= E[X^2] - E[X]^2\end{aligned}$$

- We will define the standard deviation of X , $\text{sd}(X) = \sqrt{\text{var}(X)}$
- $\text{var}(X) \geq 0$.

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\text{Var}(Y) = E[(Y - E[Y])^2]$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\ &= E[((aX + b) - E[aX + b])]^2\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\ &= E[((aX + b) - E[aX + b])]^2 \\ &= E[((aX + b) - (aE[X] + b))]^2\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\ &= E[((aX + b) - E[aX + b])]^2] \\ &= E[((aX + b) - (aE[X] + b))]^2] \\ &= E[(aX - aE[X])^2]\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\ &= E[((aX + b) - E[aX + b])]^2] \\ &= E[((aX + b) - (aE[X] + b))]^2] \\ &= E[(aX - aE[X])^2] \\ &= E[(a^2X^2 - 2a^2XE[X] + a^2E[X]^2)]\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\&= E[((aX + b) - E[aX + b])]^2] \\&= E[((aX + b) - (aE[X] + b))]^2] \\&= E[(aX - aE[X])^2] \\&= E[(a^2X^2 - 2a^2XE[X] + a^2E[X]^2)] \\&= a^2E[X^2] - 2a^2E[X]^2 + a^2E[X]^2\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\&= E[((aX + b) - E[aX + b])]^2] \\&= E[((aX + b) - (aE[X] + b))]^2] \\&= E[(aX - aE[X])^2] \\&= E[(a^2X^2 - 2a^2XE[X] + a^2E[X]^2)] \\&= a^2E[X^2] - 2a^2E[X]^2 + a^2E[X]^2 \\&= a^2(E[X^2] - E[X]^2)\end{aligned}$$

Variance Corollary

Corollary

$$\text{Var}(aX + b) = a^2 \text{Var}(X)$$

Proof.

Define $Y = aX + b$. Now, we know that $\text{Var}(Y) = E[(Y - E[Y])^2]$.

$$\begin{aligned}\text{Var}(Y) &= E[(Y - E[Y])^2] \\&= E[((aX + b) - E[aX + b])]^2] \\&= E[((aX + b) - (aE[X] + b))]^2] \\&= E[(aX - aE[X])^2] \\&= E[(a^2X^2 - 2a^2XE[X] + a^2E[X]^2)] \\&= a^2E[X^2] - 2a^2E[X]^2 + a^2E[X]^2 \\&= a^2(E[X^2] - E[X]^2) \\&= a^2\text{Var}(X)\end{aligned}$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$E[Y] = 1 \times P(Y = 1) + 0 \times P(Y = 0)$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$\begin{aligned} E[Y] &= 1 \times P(Y = 1) + 0 \times P(Y = 0) \\ &= \pi + 0(1 - \pi) = \pi \end{aligned}$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$\begin{aligned}E[Y] &= 1 \times P(Y = 1) + 0 \times P(Y = 0) \\&= \pi + 0(1 - \pi) = \pi \\E[Y^2] &= 1^2 P(Y = 1) + 0^2 P(Y = 0)\end{aligned}$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$\begin{aligned}E[Y] &= 1 \times P(Y = 1) + 0 \times P(Y = 0) \\&= \pi + 0(1 - \pi) = \pi \\E[Y^2] &= 1^2 P(Y = 1) + 0^2 P(Y = 0) \\&= \pi\end{aligned}$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$E[Y] = 1 \times P(Y = 1) + 0 \times P(Y = 0)$$

$$= \pi + 0(1 - \pi) = \pi$$

$$E[Y^2] = 1^2 P(Y = 1) + 0^2 P(Y = 0)$$

$$= \pi$$

$$\text{var}(Y) = E[Y^2] - E[Y]^2$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$E[Y] = 1 \times P(Y = 1) + 0 \times P(Y = 0)$$

$$= \pi + 0(1 - \pi) = \pi$$

$$E[Y^2] = 1^2 P(Y = 1) + 0^2 P(Y = 0)$$

$$= \pi$$

$$\text{var}(Y) = E[Y^2] - E[Y]^2$$

$$\text{var}(Y) = \pi - \pi^2$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$\begin{aligned} E[Y] &= 1 \times P(Y = 1) + 0 \times P(Y = 0) \\ &= \pi + 0(1 - \pi) = \pi \end{aligned}$$

$$\begin{aligned} E[Y^2] &= 1^2 P(Y = 1) + 0^2 P(Y = 0) \\ &= \pi \end{aligned}$$

$$\text{var}(Y) = E[Y^2] - E[Y]^2$$

$$\begin{aligned} \text{var}(Y) &= \pi - \pi^2 \\ &= \pi(1 - \pi) \end{aligned}$$

Bernoulli Random Variable Moments

Suppose $Y \sim \text{Bernoulli}(\pi)$

$$\begin{aligned} E[Y] &= 1 \times P(Y = 1) + 0 \times P(Y = 0) \\ &= \pi + 0(1 - \pi) = \pi \end{aligned}$$

$$\begin{aligned} E[Y^2] &= 1^2 P(Y = 1) + 0^2 P(Y = 0) \\ &= \pi \end{aligned}$$

$$\text{var}(Y) = E[Y^2] - E[Y]^2$$

$$\begin{aligned} \text{var}(Y) &= \pi - \pi^2 \\ &= \pi(1 - \pi) \end{aligned}$$

$$E[Y] = \pi$$

$\text{var}(Y) = \pi(1 - \pi)$ What is the maximum variance?

Trials with More than Two Outcomes

Definition

Suppose we observe a trial, which might result in J outcomes.

And that $P(\text{outcome} = i) = \pi_i$

$\mathbf{Y} = (Y_1, Y_2, \dots, Y_J)$ where $Y_j = 1$ if outcome j occurred and 0 otherwise.

*Then $\mathbf{Y}$ follows a **multinomial** distribution, with*

$$p(\mathbf{y}) = \pi_1^{y_1} \pi_2^{y_2} \dots \pi_k^{y_k}$$

if $\sum_{i=1}^k y_i = 1$ and the pmf is 0 otherwise.

Equivalently, we'll write

$$\mathbf{Y} \sim \text{Multinomial}(1, \boldsymbol{\pi})$$

$$\mathbf{Y} \sim \text{Categorical}(\boldsymbol{\pi})$$

Multinomial Properties + Notes

Computer scientists: commonly call $\text{Multinomial}(1, \boldsymbol{\pi})$ **Discrete**($\boldsymbol{\pi}$).

$$\begin{aligned} E[X_i] &= N\pi_i \\ \text{var}(X_i) &= N\pi_i(1 - \pi_i) \end{aligned}$$

Counting the Number of Events

Often interested in counting number of events that occur:

- 1) Number of wars started
- 2) Number of speeches made
- 3) Number of bribes offered
- 4) Number of people waiting for license

Generally referred to as **event counts**

Stochastic processes: a course provide introduction to many processes
(**Queing Theory**)

Definition

Suppose X is a random variable that counts the number of successes in N independent and identically distributed Bernoulli trials. Then X is a *Binomial* random variable,

$$p(k) = \binom{N}{k} \pi^k (1 - \pi)^{1-k}$$

for $k \in \{0, 1, 2, \dots, N\}$ and $p(k) = 0$ otherwise.

Equivalently,

$$Y \sim \text{Binomial}(N, \pi)$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

$$E[Z] = E[Y_1 + Y_2 + Y_3 + \dots + Y_N]$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

$$\begin{aligned} E[Z] &= E[Y_1 + Y_2 + Y_3 + \dots + Y_N] \\ &= \sum_{i=1}^N E[Y_i] \end{aligned}$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

$$\begin{aligned} E[Z] &= E[Y_1 + Y_2 + Y_3 + \dots + Y_N] \\ &= \sum_{i=1}^N E[Y_i] \\ &= N\pi \end{aligned}$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

$$\begin{aligned} E[Z] &= E[Y_1 + Y_2 + Y_3 + \dots + Y_N] \\ &= \sum_{i=1}^N E[Y_i] \\ &= N\pi \\ \text{var}(Z) &= \sum_{i=1}^N \text{var}(Y_i) \end{aligned}$$

Binomial Random Variable Moments

$$Z = \sum_{i=1}^N Y_i \text{ where } Y_i \sim \text{Bernoulli}(\pi)$$

$$E[Z] = E[Y_1 + Y_2 + Y_3 + \dots + Y_N]$$

$$= \sum_{i=1}^N E[Y_i]$$

$$= N\pi$$

$$\text{var}(Z) = \sum_{i=1}^N \text{var}(Y_i)$$

$$= N\pi(1 - \pi)$$

$$E[Z] = N\pi$$

$$\text{var}(Z) = N\pi(1 - \pi)$$

Example: Counts

We might be interested in the *sum* of several independent trials:

Example: Counts

We might be interested in the *sum* of several independent trials:

- n draws from a population size N with replacement.
- If Y is number of the sample that have a binary characteristic (support Trump, show up heads, etc).
- and M is the number of the people in the population that have a binary characteristic.

Our random variable, Y , is equal to the sum of successes.

Example: Counts

We might be interested in the *sum* of several independent trials:

- n draws from a population size N with replacement.
- If Y is number of the sample that have a binary characteristic (support Trump, show up heads, etc).
- and M is the number of the people in the population that have a binary characteristic.

Our random variable, Y , is equal to the sum of successes. Let y be an integer between 0 and n ,

$$P(Y = y) = \binom{n}{y} p^y (1 - p)^{(n-y)}$$

where $p = \frac{M}{N}$

Binomial Random Variable

- A model to count the number of successes across N trials

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1)$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$\begin{aligned} P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1) \\ = P(Y_1 = 1)P(Y_2 = 0) \cdots P(Y_N = 1) \end{aligned}$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$\begin{aligned} P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1) \\ &= P(Y_1 = 1)P(Y_2 = 0) \cdots P(Y_N = 1) \\ &= \underbrace{P(Y_1 = 1)P(Y_3 = 1) \cdots P(Y_z = 1)}_M \times \underbrace{P(Y_2 = 0) \cdots P(Y_N = 0)}_{N-M} \end{aligned}$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$\begin{aligned} P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1) \\ &= P(Y_1 = 1)P(Y_2 = 0) \cdots P(Y_N = 1) \\ &= \underbrace{P(Y_1 = 1)P(Y_3 = 1) \cdots P(Y_z = 1)}_M \times \underbrace{P(Y_2 = 0) \cdots P(Y_N = 0)}_{N-M} \\ &= \underbrace{\pi \pi \cdots \pi}_M \times \underbrace{(1 - \pi)(1 - \pi) \cdots (1 - \pi)}_{N-M} \end{aligned}$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$\begin{aligned} P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1) \\ &= P(Y_1 = 1)P(Y_2 = 0) \cdots P(Y_N = 1) \\ &= \underbrace{P(Y_1 = 1)P(Y_3 = 1) \cdots P(Y_z = 1)}_M \times \underbrace{P(Y_2 = 0) \cdots P(Y_N = 0)}_{N-M} \\ &= \underbrace{\pi \pi \cdots \pi}_M \times \underbrace{(1 - \pi)(1 - \pi) \cdots (1 - \pi)}_{N-M} \\ &= \pi^M (1 - \pi)^{N-M} \end{aligned}$$

Binomial Random Variable

- A model to count the number of successes across N trials
 - Assume the Bernoulli trials are independent
 - Each Bernoulli trial i is

$$Y_i \sim \text{Bernoulli}(\pi)$$

Independent and identically distributed.

- Z = number of successful trials
- Derive probability mass function $P(Z = M) = p(M)$
- One way to obtain M successful trials:

$$\begin{aligned} P(Y_1 = 1, Y_2 = 0, Y_3 = 1, \dots, Y_N = 1) \\ &= P(Y_1 = 1)P(Y_2 = 0) \cdots P(Y_N = 1) \\ &= \underbrace{P(Y_1 = 1)P(Y_3 = 1) \cdots P(Y_z = 1)}_M \times \underbrace{P(Y_2 = 0) \cdots P(Y_N = 0)}_{N-M} \\ &= \underbrace{\pi \pi \cdots \pi}_M \times \underbrace{(1 - \pi)(1 - \pi) \cdots (1 - \pi)}_{N-M} \\ &= \pi^M (1 - \pi)^{N-M} \end{aligned}$$

Are we done?

Are we done? No

Are we done? No

- This is just one instance of M successes

Are we done? No

- This is just one instance of M successes
- How many total instances?

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials
 - We want to select M

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials
 - We want to select M
- $\binom{N}{M} = \frac{N!}{(N-M)!M!}$

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials
 - We want to select M
- $\binom{N}{M} = \frac{N!}{(N-M)!M!}$

Then,

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials
 - We want to select M
- $\binom{N}{M} = \frac{N!}{(N-M)!M!}$

Then,

$$P(Z = M) = p(M) = \binom{N}{M} \pi^M (1 - \pi)^{N-M}$$

Are we done? **No**

- This is just one instance of M successes
- How many total instances?
 - N total trials
 - We want to select M
- $\binom{N}{M} = \frac{N!}{(N-M)!M!}$

Then,

$$P(Z = M) = p(M) = \binom{N}{M} \pi^M (1 - \pi)^{N-M}$$

Origins of Binomial

One head on one Coin:

$$P(Y = 1) = p$$

One head on two coins:

Origins of Binomial

One head on one Coin:

$$P(Y = 1) = p$$

One head on two coins:

$$P(Y = 1) = p(1 - p) + (1 - p)p = 2 * p(1 - p)^1$$

One head on three coins:

Origins of Binomial

One head on one Coin:

$$P(Y = 1) = p$$

One head on two coins:

$$P(Y = 1) = p(1 - p) + (1 - p)p = 2 * p(1 - p)^1$$

One head on three coins:

$$P(Y = 1) = p(1 - p)(1 - p) + (1 - p)p(1 - p) + (1 - p)(1 - p)p = 3 * p(1 - p)^2$$

One head on four coins:

Origins of Binomial

One head on one Coin:

$$P(Y = 1) = p$$

One head on two coins:

$$P(Y = 1) = p(1 - p) + (1 - p)p = 2 * p(1 - p)^1$$

One head on three coins:

$$P(Y = 1) = p(1 - p)(1 - p) + (1 - p)p(1 - p) + (1 - p)(1 - p)p = 3 * p(1 - p)^2$$

One head on four coins:

$$\begin{aligned} P(Y = 1) &= p(1 - p)^3 + (1 - p)p(1 - p)^2 \\ &+ (1 - p)^2 p(1 - p) + (1 - p)^3 p = 4 * p(1 - p)^3 \end{aligned}$$

Origins of Binomial

One head on four coins

$$P(Y = 1) = 4 * p(1 - p)^3$$

Two heads on four coins

Origins of Binomial

One head on four coins

$$P(Y = 1) = 4 * p(1 - p)^3$$

Two heads on four coins

$$P(Y = 2) = 6 * p^2(1 - p)^2$$

Three heads on four coins

Origins of Binomial

One head on four coins

$$P(Y = 1) = 4 * p(1 - p)^3$$

Two heads on four coins

$$P(Y = 2) = 6 * p^2(1 - p)^2$$

Three heads on four coins

$$P(Y = 3) = 4 * p^3(1 - p)$$

Number of different realizations with y successes: $\binom{n}{y}$

$$P(Y = y) = \binom{n}{y} p^y (1 - p)^{(n-y)}$$

where $p = \frac{M}{N}$

Origins of Binomial

If $X_1, \dots, X_n$ are Bernoulli distributed with probability p , then

$$Y = \sum_{k=1}^n X_k \sim \text{Binomial}(n, p)$$

n and p are parameters of the binomial.

Assumptions of Binomial Distribution

- The number of trials is fixed,
- The probability of success is the same for each trial.
- The trials are independent

Example: Baseball!

- The World Series of Baseball ends when one team wins 4 games.
- Each game is a Bernoulli trial.

Example: Baseball!

- The World Series of Baseball ends when one team wins 4 games.
- Each game is a Bernoulli trial.
- Length of World Series $\in \{4, 5, 6, 7\}$.
- What is the probability that the World Series lasts 7 games?

Example: Baseball!

- The World Series of Baseball ends when one team wins 4 games.
- Each game is a Bernoulli trial.
- Length of World Series $\in \{4, 5, 6, 7\}$.
- What is the probability that the World Series lasts 7 games?
- $P(X=7)$?

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.
- Number of diseased villages in a district.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.
- Number of diseased villages in a district.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.
- Number of diseased villages in a district.

Let X be the number of times an event occurs in a unit of time or space.

- The probability that the event occurs in a given unit of time or space is the same for all the units.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.
- Number of diseased villages in a district.

Let X be the number of times an event occurs in a unit of time or space.

- The probability that the event occurs in a given unit of time or space is the same for all the units.
- The number of events that occur in one unit of time is independent of the number of events in other units.

Example: Counts in a time period

We might be interested in the sum of events occurring in a specific time or space.

- Number of wars in a year.
- Number of diseased villages in a district.

Let X be the number of times an event occurs in a unit of time or space.

- The probability that the event occurs in a given unit of time or space is the same for all the units.
- The number of events that occur in one unit of time is independent of the number of events in other units.

Then X is a Poisson random variable, with rate $= \lambda$ and pf:

$$p(x) = \frac{\lambda^x}{x!} e^{-\lambda} \quad x = 0, 1, 2, \dots$$

Example: Poisson Distribution

If, on average, two customers sign up every hour, what is the probability that 4 customers show up in a given hour?

Example: Poisson Distribution

If, on average, two customers sign up every hour, what is the probability that 4 customers show up in a given hour?

$$\lambda = 2$$


Example: Poisson Distribution

If, on average, two customers sign up every hour, what is the probability that 4 customers show up in a given hour?

$$\lambda = 2$$

$$P(X = 4) = \frac{2^4}{4!} e^{-2} = .09$$

PF



Functions of Random Variables

Definition

If $Y = g(X)$, and X is a random variable with probability mass function $p_X(x)$, Y is a random variable with pmf:

$$p_Y(y) = \sum_{\{x|g(x)=y\}} p_X(x)$$