

Mathematical Foundations of Political Methodology

Robert Gulotty

University of Chicago

October 22, 2019

- Vectors
- Matrix
- Matrix Algebra
- Matrix Inversion

- Vectors
- Matrix
- Matrix Algebra
- Matrix Inversion

Points and Vectors

Definition

A point $\mathbf{x} \in R^n$ is an ordered n -tuple, $(x_1, x_2, \dots, x_n)$. The vector $\mathbf{x} \in R^n$ is the arrow pointing from the origin $(0, 0, \dots, 0)$ to $\mathbf{x}$.

Points and Vectors

Definition

A point $\mathbf{x} \in R^n$ is an ordered n -tuple, $(x_1, x_2, \dots, x_n)$. The vector $\mathbf{x} \in R^n$ is the arrow pointing from the origin $(0, 0, \dots, 0)$ to $\mathbf{x}$.

Some Examples:

Points and Vectors

Definition


A point $\mathbf{x} \in R^n$ is an ordered n -tuple, $(x_1, x_2, \dots, x_n)$. The vector $\mathbf{x} \in R^n$ is the arrow pointing from the origin $(0, 0, \dots, 0)$ to $\mathbf{x}$.

Some Examples:

- (Latitude, Longitude, Elevation)
- (1, 2, 3)
- (0, 1, 0)


Points and Vectors

Now, let's see a vector in a 2 dimensional space.



Points and Vectors

Now, let's see a vector in a 3 dimensional space.



Arithmetic with Vectors

Definition

Suppose $\mathbf{u}$ and $\mathbf{v}$ are vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$,

$$\mathbf{u} = (u_1, u_2, u_3, \dots, u_n)$$

$$\mathbf{v} = (v_1, v_2, v_3, \dots, v_n)$$

We will say $\mathbf{u} = \mathbf{v}$ if $u_1 = v_1, u_2 = v_2, \dots, u_n = v_n$

Arithmetic with Vectors

Definition

Suppose $\mathbf{u}$ and $\mathbf{v}$ are vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$,

$$\mathbf{u} = (u_1, u_2, u_3, \dots, u_n)$$

$$\mathbf{v} = (v_1, v_2, v_3, \dots, v_n)$$

We will say $\mathbf{u} = \mathbf{v}$ if $u_1 = v_1, u_2 = v_2, \dots, u_n = v_n$

Define the sum of $\mathbf{u} + \mathbf{v}$ as

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, u_3 + v_3, \dots, u_n + v_n)$$

Arithmetic with Vectors

Definition

Suppose $\mathbf{u}$ and $\mathbf{v}$ are vectors $\mathbf{u}, \mathbf{v} \in R^n$,

$$\mathbf{u} = (u_1, u_2, u_3, \dots, u_n)$$

$$\mathbf{v} = (v_1, v_2, v_3, \dots, v_n)$$

We will say $\mathbf{u} = \mathbf{v}$ if $u_1 = v_1, u_2 = v_2, \dots, u_n = v_n$

Define the sum of $\mathbf{u} + \mathbf{v}$ as

$$\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, u_3 + v_3, \dots, u_n + v_n)$$

Suppose $k \in R$. We will call k a scalar.

Define $k\mathbf{u}$ as the scalar product

$$k\mathbf{u} = (ku_1, ku_2, \dots, ku_n)$$

Examples

Suppose:

$$\mathbf{u} = (1, 2, 3, 4, 5)$$

$$\mathbf{v} = (1, 1, 1, 1, 1)$$

$$k = 2$$

Examples

Suppose:

$$\mathbf{u} = (1, 2, 3, 4, 5)$$

$$\mathbf{v} = (1, 1, 1, 1, 1)$$

$$k = 2$$

Then,

$$\mathbf{u} + \mathbf{v} = (1 + 1, 2 + 1, 3 + 1, 4 + 1, 5 + 1) = (2, 3, 4, 5, 6)$$

Examples

Suppose:

$$\mathbf{u} = (1, 2, 3, 4, 5)$$

$$\mathbf{v} = (1, 1, 1, 1, 1)$$

$$k = 2$$

Then,

$$\mathbf{u} + \mathbf{v} = (1 + 1, 2 + 1, 3 + 1, 4 + 1, 5 + 1) = (2, 3, 4, 5, 6)$$

$$k\mathbf{u} = (2 \times 1, 2 \times 2, 2 \times 3, 2 \times 4, 2 \times 5) = (2, 4, 6, 8, 10)$$

Examples

Suppose:

$$\mathbf{u} = (1, 2, 3, 4, 5)$$

$$\mathbf{v} = (1, 1, 1, 1, 1)$$

$$k = 2$$

Then,

$$\mathbf{u} + \mathbf{v} = (1 + 1, 2 + 1, 3 + 1, 4 + 1, 5 + 1) = (2, 3, 4, 5, 6)$$

$$k\mathbf{u} = (2 \times 1, 2 \times 2, 2 \times 3, 2 \times 4, 2 \times 5) = (2, 4, 6, 8, 10)$$

$$k\mathbf{v} = (2 \times 1, 2 \times 1, 2 \times 1, 2 \times 1, 2 \times 1) = (2, 2, 2, 2, 2)$$

Vector Lengths

Mathematics calls measures of distance a norm, written $\| \cdot \|$.

Vector Lengths

Mathematics calls measures of distance a norm, written $\|\cdot\|$.

The length of vectors in R^n can be measured using the **Euclidean** norm.

Vector Lengths

Mathematics calls measures of distance a norm, written $\|\cdot\|$.

The length of vectors in R^n can be measured using the **Euclidean** norm.

Definition

On an n -dimensional Euclidean space R^n , the Euclidean norm of vector, $x = (x_1, x_2, \dots, x_n)$, is defined by the formula

$$\|x\|_2 := \sqrt{x_1^2 + \dots + x_n^2}.$$

Vector Lengths

Mathematics calls measures of distance a norm, written $\|\cdot\|$.

The length of vectors in R^n can be measured using the **Euclidean** norm.

Definition

On an n -dimensional Euclidean space R^n , the Euclidean norm of vector, $x = (x_1, x_2, \dots, x_n)$, is defined by the formula

$$\|x\|_2 := \sqrt{x_1^2 + \dots + x_n^2}.$$

An example. What's the Euclidean norm of $x = (1, 3, 5)$?

Vector Lengths

Mathematics calls measures of distance a norm, written $\|\cdot\|$.

The length of vectors in R^n can be measured using the **Euclidean** norm.

Definition

On an n -dimensional Euclidean space R^n , the Euclidean norm of vector, $x = (x_1, x_2, \dots, x_n)$, is defined by the formula

$$\|x\|_2 := \sqrt{x_1^2 + \dots + x_n^2}.$$

An example. What's the Euclidean norm of $x = (1, 3, 5)$?

Its Euclidean norm equals to:

$$\sqrt{1^2 + 3^2 + 5^2} = \sqrt{35}$$

Dot Product

$$x = (x_1, x_2, \dots, x_n)$$

$$y = (y_1, y_2, \dots, y_n)$$

$$x \cdot y = \sum_{i=1}^n x_i y_i = x_1 * y_1 + x_2 * y_2 + \dots + x_n * y_n$$

- Vectors
- **Matrix**
- Matrix Algebra
- Matrix Inversion

Matrices

Matrices collect vectors for convenience.

Matrices

Matrices collect vectors for convenience.

The first example is the following system of equations:

Matrices

Matrices collect vectors for convenience.

The first example is the following system of equations:

$$2x + 2y + 3z = 1$$

$$3x + 5y + 7z = 2$$

$$5x + 8y + 3z = 3$$

Matrices

Matrices collect vectors for convenience.

The first example is the following system of equations:

$$2x + 2y + 3z = 1$$

$$3x + 5y + 7z = 2$$

$$5x + 8y + 3z = 3$$

The coefficient of unknowns can be written compactly as matrices:

$$\begin{bmatrix} 2 & 2 & 3 \\ 3 & 5 & 7 \\ 5 & 8 & 3 \end{bmatrix}$$

Matrices

A second example of matrix is the data set.

	id	name	year	urbanpop	totalpop	urbrate	fpresence	westelbe	protestant	latitude	longitude
1	1	Rhineland	1700	83000	.	.	19	1	.25	50.35	7.6
2	1	Rhineland	1750	129000	1438817	8.9657	19	1	.25	50.35	7.6
3	1	Rhineland	1800	274000	1860719	14.72549	19	1	.25	50.35	7.6
4	1	Rhineland	1850	572278	2846708	20.10317	19	1	.25	50.35	7.6
5	1	Rhineland	1875	1629051	3804381	42.8204	19	1	.25	50.35	7.6
6	1	Rhineland	1880	1850627	4074000	45.42531	19	1	.25	50.35	7.6
7	1	Rhineland	1885	2103753	4344527	48.42306	19	1	.25	50.35	7.6
8	1	Rhineland	1895	2836140	5106002	55.54522	19	1	.25	50.35	7.6
9	1	Rhineland	1900	3506587	5759798	60.88038	19	1	.25	50.35	7.6
10	1	Rhineland	1905	4145633	6436337	64.40982	19	1	.25	50.35	7.6
11	1	Rhineland	1910	4806725	7121140	67.49937	19	1	.25	50.35	7.6
12	2	Mark	1700	5000	117963	4.238618	6	1	.6	51.68333	7.816667
13	2	Mark	1750	5000	149869	3.336254	6	1	.6	51.68333	7.816667
14	2	Mark	1800	36000	193096	18.64353	6	1	.6	51.68333	7.816667
15	2	Mark	1850	55628	362392	15.35011	6	1	.6	51.68333	7.816667
16	2	Mark	1875	305301	721466	42.31675	6	1	.6	51.68333	7.816667
17	2	Mark	1880	.	.	.	6	1	.6	51.68333	7.816667
18	2	Mark	1885	.	.	.	6	1	.6	51.68333	7.816667
19	2	Mark	1895	.	.	.	6	1	.6	51.68333	7.816667
20	2	Mark	1900	1134655	1514546	74.91717	6	1	.6	51.68333	7.816667
21	2	Mark	1905	.	.	.	6	1	.6	51.68333	7.816667
22	2	Mark	1910	.	.	.	6	1	.6	51.68333	7.816667
23	3	Westphalia without Mark	1700	7000	.	.	6	1	.4	51.96667	7.633333
24	3	Westphalia without Mark	1750	14000	528851	2.647246	6	1	.4	51.96667	7.633333
25	3	Westphalia without Mark	1800	55000	763110	7.20735	6	1	.4	51.96667	7.633333
26	3	Westphalia without Mark	1850	61000	1120946	5.441833	6	1	.4	51.96667	7.633333
27	3	Westphalia without Mark	1875	190503	1184231	16.08664	6	1	.4	51.96667	7.633333
28	3	Westphalia without Mark	1880	.	.	.	6	1	.4	51.96667	7.633333

Matrices

Definition

A Matrix is a rectangular array of numbers

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

*If $\mathbf{A}$ has m rows n columns (**R**oman **C**atholic; **R**um & **C**oke) we will say that $\mathbf{A}$ is an $m \times n$ matrix.*

Suppose $\mathbf{X}$ and $\mathbf{Y}$ are $m \times n$ matrices. Then $\mathbf{X} = \mathbf{Y}$ if $x_{ij} = y_{ij}$ for all i and j

Simple Examples

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & 1 \end{pmatrix}$$

If $\mathbf{I}$ is an $n \times n$ matrix we will call an identity matrix.

Simple Examples

$$\mathbf{X} = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 4 \end{pmatrix}$$

$\mathbf{X}$ is an 2×3 matrix

- Vectors
- Matrix
- **Matrix Algebra**
- Matrix Inversion

Matrix Algebra

Definition

Suppose $\mathbf{X}$ and $\mathbf{Y}$ are $m \times n$ matrices. Then define

$$\begin{aligned}\mathbf{X} + \mathbf{Y} &= \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{pmatrix} + \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1n} \\ y_{21} & y_{22} & \cdots & y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ y_{m1} & y_{m2} & \cdots & y_{mn} \end{pmatrix} \\ &= \begin{pmatrix} x_{11} + y_{11} & x_{12} + y_{12} & \cdots & x_{1n} + y_{1n} \\ x_{21} + y_{21} & x_{22} + y_{22} & \cdots & x_{2n} + y_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} + y_{m1} & x_{m2} + y_{m2} & \cdots & x_{mn} + y_{mn} \end{pmatrix}\end{aligned}$$

Matrix Algebra

Definition

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $k \in R$. Then,

$$k\mathbf{X} = \begin{pmatrix} kx_{11} & kx_{12} & \dots & kx_{1n} \\ kx_{21} & kx_{22} & \dots & kx_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ kx_{m1} & kx_{m2} & \dots & kx_{mn} \end{pmatrix}$$

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

$$\mathbf{X} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{mn} \end{pmatrix}$$

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

$$\mathbf{X} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{mn} \end{pmatrix}$$
$$\mathbf{X}' = \begin{pmatrix} x_{11} & x_{21} & \dots & x_{m1} \\ x_{12} & x_{22} & \dots & x_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ x_{1n} & x_{2n} & \dots & x_{mn} \end{pmatrix}$$

If $\mathbf{X}$ is an $m \times n$ then $\mathbf{X}'$ is $n \times m$.

If $\mathbf{X} = \mathbf{X}'$ then we say $\mathbf{X}$ is symmetric.

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

$$\mathbf{X} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{mn} \end{pmatrix}$$
$$\mathbf{X}' = \begin{pmatrix} x_{11} & x_{21} & \dots & x_{m1} \\ x_{12} & x_{22} & \dots & x_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ x_{1n} & x_{2n} & \dots & x_{mn} \end{pmatrix}$$

If $\mathbf{X}$ is an $m \times n$ then $\mathbf{X}'$ is $n \times m$.

If $\mathbf{X} = \mathbf{X}'$ then we say $\mathbf{X}$ is symmetric.

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

$$\mathbf{X} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{mn} \end{pmatrix}$$
$$\mathbf{X}' = \begin{pmatrix} x_{11} & x_{21} & \dots & x_{m1} \\ x_{12} & x_{22} & \dots & x_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ x_{1n} & x_{2n} & \dots & x_{mn} \end{pmatrix}$$

If $\mathbf{X}$ is an $m \times n$ then $\mathbf{X}'$ is $n \times m$.

If $\mathbf{X} = \mathbf{X}'$ then we say $\mathbf{X}$ is symmetric.

Matrix Transpose

We will use matrix transpose to flip the dimensionality of a matrix

$$\mathbf{X} = \begin{pmatrix} x_{11} & x_{12} & \dots & x_{1n} \\ x_{21} & x_{22} & \dots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \dots & x_{mn} \end{pmatrix}$$
$$\mathbf{X}' = \begin{pmatrix} x_{11} & x_{21} & \dots & x_{m1} \\ x_{12} & x_{22} & \dots & x_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ x_{1n} & x_{2n} & \dots & x_{mn} \end{pmatrix}$$

If $\mathbf{X}$ is an $m \times n$ then $\mathbf{X}'$ is $n \times m$.

If $\mathbf{X} = \mathbf{X}'$ then we say $\mathbf{X}$ is symmetric.

Matrix Transpose

Example 1: $\mathbf{X} = \begin{pmatrix} 4 & 1 & 2 \\ 1 & 2 & 3 \end{pmatrix}$ then $\mathbf{X}' = \begin{pmatrix} 4 & 1 \\ 1 & 2 \\ 2 & 3 \end{pmatrix}$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\mathbf{A} = \mathbf{XY}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned} \mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned}\mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix}\end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned}\mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix}\end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned} \mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix} \end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned}\mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix}\end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned} \mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix} \end{aligned}$$

Matrix Multiplication

Matrix multiplication is at the root of most econometrics, statistical modeling, but the procedure is not straightforward.

Suppose we have two matrices

$$\mathbf{X} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \mathbf{Y} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$$

We will create a new matrix $\mathbf{A}$ by matrix multiplication:

$$\begin{aligned} \mathbf{A} &= \mathbf{XY} \\ &= \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 1 \times 1 + 1 \times 3 & 1 \times 2 + 1 \times 4 \\ 1 \times 1 + 2 \times 3 & 1 \times 2 + 2 \times 4 \end{pmatrix} \\ &= \begin{pmatrix} 4 & 6 \\ 7 & 10 \end{pmatrix} \end{aligned}$$

Matrix Multiplication

Definition

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Then define the matrix $\mathbf{A}$ an $m \times k$ matrix that obtains from multiplying $\mathbf{X}$ and $\mathbf{Y}$ as,

$$\mathbf{A} = \mathbf{XY}$$

$$= \begin{pmatrix} x_{11} & x_{12} & \cdots & x_{1n} \\ x_{21} & x_{22} & \cdots & x_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & \cdots & x_{mn} \end{pmatrix} \begin{pmatrix} y_{11} & y_{12} & \cdots & y_{1k} \\ y_{21} & y_{22} & \cdots & y_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ y_{n1} & y_{n2} & \cdots & y_{nk} \end{pmatrix}$$

$$= \begin{pmatrix} x_{11}y_{11} + x_{12}y_{21} + \cdots + x_{1n}y_{n1} & \cdots & x_{11}y_{1k} + x_{12}y_{2k} + \cdots + x_{1n}y_{nk} \\ \vdots & \ddots & \vdots \\ x_{m1}y_{11} + x_{m2}y_{21} + \cdots + x_{mn}y_{n1} & \cdots & x_{m1}y_{1k} + x_{m2}y_{2k} + \cdots + x_{mn}y_{nk} \end{pmatrix}$$

Row Vectors

Definition

Suppose $\mathbf{X}$ is an $m \times n$ matrix. We write the row vectors of $\mathbf{X} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_m \end{pmatrix}$

$$\mathbf{X} = \begin{pmatrix} 4 & 1 & 2 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\mathbf{X} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{pmatrix}$$

where

$$\mathbf{x}_1 = (4 \ 1 \ 2)$$

$$\mathbf{x}_2 = (1 \ 2 \ 3)$$

Column Vectors

Definition

Suppose $\mathbf{X}$ is an $m \times n$ matrix. We write the column vectors of $\mathbf{X} = (\mathbf{x}_1 \ \mathbf{x}_2 \ \dots \ \mathbf{x}_m)$

$$\mathbf{X} = \begin{pmatrix} 4 & 1 & 2 \\ 1 & 2 & 3 \end{pmatrix}$$

$$\mathbf{X} = (\mathbf{x}_1 \ \mathbf{x}_2 \ \mathbf{x}_3)$$

where

$$\mathbf{x}_1 = \begin{pmatrix} 4 \\ 1 \end{pmatrix}, \ \mathbf{x}_2 = \begin{pmatrix} 1 \\ 2 \end{pmatrix}, \ \mathbf{x}_3 = \begin{pmatrix} 2 \\ 3 \end{pmatrix}$$

Definition

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Write the row

vectors of $\mathbf{X} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_m \end{pmatrix}$ and $\mathbf{Y}$ as column vector $\mathbf{Y} = (\mathbf{y}_1 \ \mathbf{y}_2 \ \dots \ \mathbf{y}_k)$.

Then the $m \times k$ matrix $\mathbf{A} = \mathbf{X}\mathbf{Y}$ can be written as

$$\mathbf{A} = \begin{pmatrix} \mathbf{x}_1 \cdot \mathbf{y}_1 & \mathbf{x}_1 \cdot \mathbf{y}_2 & \dots & \mathbf{x}_1 \cdot \mathbf{y}_k \\ \mathbf{x}_2 \cdot \mathbf{y}_1 & \mathbf{x}_2 \cdot \mathbf{y}_2 & \dots & \mathbf{x}_2 \cdot \mathbf{y}_k \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{x}_m \cdot \mathbf{y}_1 & \mathbf{x}_m \cdot \mathbf{y}_2 & \dots & \mathbf{x}_m \cdot \mathbf{y}_k \end{pmatrix}$$

Matrix Multiplication

Let's work on an example together!

$$\mathbf{X} = \begin{pmatrix} 1 & 4 & 5 \\ 10 & 2 & 3 \end{pmatrix}$$

$$\mathbf{Y} = \begin{pmatrix} 2 & 3 \\ 1 & 5 \\ 3 & 5 \end{pmatrix} \text{ What is } \mathbf{XY}?$$

Matrix Multiplication

Let's work on an example together!

$$\mathbf{X} = \begin{pmatrix} 1 & 4 & 5 \\ 10 & 2 & 3 \end{pmatrix}$$

$$\mathbf{Y} = \begin{pmatrix} 2 & 3 \\ 1 & 5 \\ 3 & 5 \end{pmatrix} \text{ What is } \mathbf{XY}?$$

Not all matrices can be multiplied.

Matrix $\mathbf{AB}$ exists only if the number of columns in $\mathbf{A}$ = number of rows in $\mathbf{B}$. If $\mathbf{AB}$ exists we will say the matrices are conformable

Matrix Multiplication with a Vector

Suppose $\mathbf{X} = \begin{pmatrix} 2 & 3 & 4 & 5 \\ 1 & 5 & 1 & 2 \\ 3 & 5 & 3 & 4 \end{pmatrix}$ a 3×4 matrix and that $\mathbf{v} = \begin{pmatrix} 3 \\ 3 \\ 4 \\ 10 \end{pmatrix}$ a 4×1

matrix (or a column vector) what is

$\mathbf{X}\mathbf{v}$?

What is $\mathbf{X}'\mathbf{v}$?

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

- $\mathbf{XY} \neq \mathbf{YX}$ in general

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

- $\mathbf{XY} \neq \mathbf{YX}$ in general
- $\mathbf{XI} = \mathbf{X}$

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

- $\mathbf{XY} \neq \mathbf{YX}$ in general
- $\mathbf{XI} = \mathbf{X}$
- $(\mathbf{X}')' = \mathbf{X}$

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

- $\mathbf{XY} \neq \mathbf{YX}$ in general
- $\mathbf{XI} = \mathbf{X}$
- $(\mathbf{X}')' = \mathbf{X}$
- $(\mathbf{XY})' = \mathbf{Y}'\mathbf{X}'$

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix}$ as the identity matrix and that $k \in R$.

- $\mathbf{XY} \neq \mathbf{YX}$ in general
- $\mathbf{XI} = \mathbf{X}$
- $(\mathbf{X}')' = \mathbf{X}$
- $(\mathbf{XY})' = \mathbf{Y}'\mathbf{X}'$
- $(k\mathbf{X})' = k\mathbf{X}'$

Algebraic Properties

Suppose $\mathbf{X}$ is an $m \times n$ matrix and $\mathbf{Y}$ is an $n \times k$ matrix. Suppose that

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 1 \end{pmatrix} \text{ as the identity matrix and that } k \in R.$$

- $\mathbf{XY} \neq \mathbf{YX}$ in general
- $\mathbf{XI} = \mathbf{X}$
- $(\mathbf{X}')' = \mathbf{X}$
- $(\mathbf{XY})' = \mathbf{Y}'\mathbf{X}'$
- $(k\mathbf{X})' = k\mathbf{X}'$
- $(\mathbf{X} + \mathbf{Y})' = \mathbf{X}' + \mathbf{Y}'$

- Vectors
- Matrix
- Matrix Algebra
- Matrix Inversion

Matrix Inversion: Motivations

Big topic: suppose $\mathbf{X}$ is an $n \times n$ matrix. We want to find the matrix $\mathbf{X}^{-1}$ such that

$$\mathbf{X}^{-1}\mathbf{X} = \mathbf{X}\mathbf{X}^{-1} = \mathbf{I}$$

where $\mathbf{I}$ is the $n \times n$ identity matrix.

Matrix Inversion: Motivations

Big topic: suppose $\mathbf{X}$ is an $n \times n$ matrix. We want to find the matrix $\mathbf{X}^{-1}$ such that

$$\mathbf{X}^{-1}\mathbf{X} = \mathbf{X}\mathbf{X}^{-1} = \mathbf{I}$$

where $\mathbf{I}$ is the $n \times n$ identity matrix.

Why do we care about matrix inversion?

Matrix Inversion: Motivations

Big topic: suppose $\mathbf{X}$ is an $n \times n$ matrix. We want to find the matrix $\mathbf{X}^{-1}$ such that

$$\mathbf{X}^{-1}\mathbf{X} = \mathbf{X}\mathbf{X}^{-1} = \mathbf{I}$$

where $\mathbf{I}$ is the $n \times n$ identity matrix.

Why do we care about matrix inversion?

It will help us to determine whether a system of equations is solvable.

Matrix Inversion: Motivations

Consider the following equations:

$$\begin{aligned}x_1 + x_2 + x_3 &= 0 \\ 0x_1 + 5x_2 + 0x_3 &= 5 \\ 0x_1 + 0x_2 + 3x_3 &= 6\end{aligned}$$

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\mathbf{x} = (x_1, x_2, x_3)$$

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\mathbf{x} = (x_1, x_2, x_3)$$

$$\mathbf{b} = (0, 5, 6)$$

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\mathbf{x} = (x_1, x_2, x_3)$$

$$\mathbf{b} = (0, 5, 6)$$

The system of equations are now,

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\mathbf{x} = (x_1, x_2, x_3)$$

$$\mathbf{b} = (0, 5, 6)$$

The system of equations are now,

$$\mathbf{Ax} = \mathbf{b}$$

Matrix Inversion: Motivations

Consider the following equations:

$$x_1 + x_2 + x_3 = 0$$

$$0x_1 + 5x_2 + 0x_3 = 5$$

$$0x_1 + 0x_2 + 3x_3 = 6$$

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 5 & 0 \\ 0 & 0 & 3 \end{pmatrix}$$

$$\mathbf{x} = (x_1, x_2, x_3)$$

$$\mathbf{b} = (0, 5, 6)$$

The system of equations are now,

$$\mathbf{Ax} = \mathbf{b}$$

$\mathbf{A}^{-1}$ exists if and only if $\mathbf{Ax} = \mathbf{b}$ has only one solution.

Matrix Inversion, Definition

Definition

Suppose $\mathbf{X}$ is an $n \times n$ matrix. We will call $\mathbf{X}^{-1}$ the inverse of $\mathbf{X}$ if

$$\mathbf{X}^{-1}\mathbf{X} = \mathbf{X}\mathbf{X}^{-1} = \mathbf{I}$$

If $\mathbf{X}^{-1}$ exists then $\mathbf{X}$ is invertible. If $\mathbf{X}^{-1}$ does not exist, then we will say $\mathbf{X}$ is singular.

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Proof.

By way of contradiction, suppose not.

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Proof.

By way of contradiction, suppose not. Then there are at least two matrices $\mathbf{A}$ and $\mathbf{C}$ such that $\mathbf{AX} = \mathbf{I}$ and $\mathbf{CX} = \mathbf{I}$

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Proof.

By way of contradiction, suppose not. Then there are at least two matrices $\mathbf{A}$ and $\mathbf{C}$ such that $\mathbf{AX} = \mathbf{I}$ and $\mathbf{CX} = \mathbf{I}$

This implies that,

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Proof.

By way of contradiction, suppose not. Then there are at least two matrices $\mathbf{A}$ and $\mathbf{C}$ such that $\mathbf{AX} = \mathbf{I}$ and $\mathbf{CX} = \mathbf{I}$

This implies that,

$$\mathbf{AXC} = (\mathbf{AX})\mathbf{C}$$

Matrix Inversion

Now, let's discuss some properties of matrix inversion.

Theorem

The inverse of matrix $\mathbf{X}$, $\mathbf{X}^{-1}$, is unique

Proof.

By way of contradiction, suppose not. Then there are at least two matrices $\mathbf{A}$ and $\mathbf{C}$ such that $\mathbf{AX} = \mathbf{I}$ and $\mathbf{CX} = \mathbf{I}$

This implies that,

$$\begin{aligned}\mathbf{AXC} &= (\mathbf{AX})\mathbf{C} \\ &= \mathbf{IC} \\ &= \mathbf{C}\end{aligned}$$

Matrix Inversion

But it also implies that

Matrix Inversion

But it also implies that

$$\mathbf{A}\mathbf{X}\mathbf{C} = \mathbf{A}(\mathbf{X}\mathbf{C})$$

Matrix Inversion

But it also implies that

$$\begin{aligned} \mathbf{AXC} &= \mathbf{A(XC)} \\ &= \mathbf{A(I)} \end{aligned}$$

Matrix Inversion

But it also implies that

$$\begin{aligned}\mathbf{A}\mathbf{X}\mathbf{C} &= \mathbf{A}(\mathbf{X}\mathbf{C}) \\ &= \mathbf{A}(\mathbf{I}) \\ &= \mathbf{A}\end{aligned}$$

Matrix Inversion

But it also implies that

$$\begin{aligned}\mathbf{A}\mathbf{X}\mathbf{C} &= \mathbf{A}(\mathbf{X}\mathbf{C}) \\ &= \mathbf{A}(\mathbf{I}) \\ &= \mathbf{A}\end{aligned}$$

So $\mathbf{C} = \mathbf{A}\mathbf{X}\mathbf{C} = \mathbf{A}$ or $\mathbf{C} = \mathbf{A}$

Matrix Inversion

But it also implies that

$$\begin{aligned}\mathbf{AXC} &= \mathbf{A(XC)} \\ &= \mathbf{A(I)} \\ &= \mathbf{A}\end{aligned}$$

So $\mathbf{C} = \mathbf{AXC} = \mathbf{A}$ or $\mathbf{C} = \mathbf{A}$ but this contradicts our assumption that there are two unique inverses.

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Proof.

We need to show that $(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = (\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{I}$.

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Proof.

We need to show that $(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = (\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{I}$.

$$(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = \mathbf{B}^{-1}(\mathbf{A}^{-1}\mathbf{A})\mathbf{B}$$

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Proof.

We need to show that $(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = (\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{I}$.

$$\begin{aligned}(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) &= \mathbf{B}^{-1}(\mathbf{A}^{-1}\mathbf{A})\mathbf{B} \\ &= \mathbf{B}^{-1}\mathbf{I}\mathbf{B}\end{aligned}$$

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Proof.

We need to show that $(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = (\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{I}$.

$$\begin{aligned}(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) &= \mathbf{B}^{-1}(\mathbf{A}^{-1}\mathbf{A})\mathbf{B} \\ &= \mathbf{B}^{-1}\mathbf{I}\mathbf{B} \\ &= \mathbf{B}^{-1}\mathbf{B}\end{aligned}$$

Matrix Inversion

Theorem

Suppose $\mathbf{A}$ has inverse $\mathbf{A}^{-1}$ and $\mathbf{B}$ has inverse $\mathbf{B}^{-1}$. Then,

$$(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$$

Proof.

We need to show that $(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) = (\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) = \mathbf{I}$.

$$\begin{aligned}(\mathbf{B}^{-1}\mathbf{A}^{-1})(\mathbf{AB}) &= \mathbf{B}^{-1}(\mathbf{A}^{-1}\mathbf{A})\mathbf{B} \\&= \mathbf{B}^{-1}\mathbf{I}\mathbf{B} \\&= \mathbf{B}^{-1}\mathbf{B} \\&= \mathbf{I}\end{aligned}$$

Matrix Inversion

$$(AB)(B^{-1}A^{-1}) = A(BB^{-1})A^{-1}$$

Matrix Inversion

$$\begin{aligned}(\mathbf{A}\mathbf{B})(\mathbf{B}^{-1}\mathbf{A}^{-1}) &= \mathbf{A}(\mathbf{B}\mathbf{B}^{-1})\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{I}\mathbf{A}^{-1}\end{aligned}$$

Matrix Inversion

$$\begin{aligned}(\mathbf{A}\mathbf{B})(\mathbf{B}^{-1}\mathbf{A}^{-1}) &= \mathbf{A}(\mathbf{B}\mathbf{B}^{-1})\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{I}\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{A}^{-1}\end{aligned}$$

Matrix Inversion

$$\begin{aligned}(\mathbf{A}\mathbf{B})(\mathbf{B}^{-1}\mathbf{A}^{-1}) &= \mathbf{A}(\mathbf{B}\mathbf{B}^{-1})\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{I}\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{A}^{-1} \\ &= \mathbf{I}\end{aligned}$$

Matrix Inversion

$$\begin{aligned}(\mathbf{AB})(\mathbf{B}^{-1}\mathbf{A}^{-1}) &= \mathbf{A}(\mathbf{BB}^{-1})\mathbf{A}^{-1} \\ &= \mathbf{A}\mathbf{I}\mathbf{A}^{-1} \\ &= \mathbf{AA}^{-1} \\ &= \mathbf{I}\end{aligned}$$

So $\mathbf{AB}$ is invertible and $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$.

Inclass Exercise: Challenge Inversion Proofs

- Show that $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- Show that $(k\mathbf{A})^{-1} = \frac{1}{k}\mathbf{A}^{-1}$

Matrix Inversion

After we define matrix inversion and discuss its properties, a natural question to ask is that when do matrix inversion exist?

Matrix Inversion

After we define matrix inversion and discuss its properties, a natural question to ask is that when do matrix inversion exist?

The answer relates to a concept, linear independence.

Matrix Inversion

After we define matrix inversion and discuss its properties, a natural question to ask is that when do matrix inversion exist?

The answer relates to a concept, linear independence.

Intuitively, linear independence means that there is no repeated information in the matrix.

Matrix Inversion: Existence

Definition

Suppose we have a set of vectors $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r\}$

And consider the system of equations

$$k_1 \mathbf{v}_1 + k_2 \mathbf{v}_2 + \dots + k_r \mathbf{v}_r = \mathbf{0}$$

If the only solution is $k_1 = 0, k_2 = 0, k_3 = 0, \dots, k_r = 0$ then we say that the set is linearly independent. If there are other solutions, then the set is linearly dependent.

Matrix Inversion: Existence

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$

Can we write this as a combination of other vectors?

Matrix Inversion: Existence

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$

Can we write this as a combination of other vectors? no!

Matrix Inversion: Existence

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$

Can we write this as a combination of other vectors? no!

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$, $\mathbf{v}_4 = (1, 2, 3)$.

Can we write any of these vector as a combination of the remaining two vectors?

Matrix Inversion: Existence

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$

Can we write this as a combination of other vectors? no!

Consider $\mathbf{v}_1 = (1, 0, 0)$, $\mathbf{v}_2 = (0, 1, 0)$, $\mathbf{v}_3 = (0, 0, 1)$, $\mathbf{v}_4 = (1, 2, 3)$.

Can we write any of these vector as a combination of the remaining two vectors?

$$\mathbf{v}_4 = \mathbf{v}_1 + 2\mathbf{v}_2 + 3\mathbf{v}_3$$

Matrix Inversion: Existence

Theorem

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_K \in \mathbb{R}^n$. If $K > n$ then the set is linearly dependent

Matrix Inversion: Existence

Theorem

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_K \in \mathbb{R}^n$. If $K > n$ then the set is linearly dependent

- $\mathbf{v}_1 = (v_{11}, v_{21}, \dots, v_{n1})$

Matrix Inversion: Existence

Theorem

Suppose $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_K \in R^n$. If $K > n$ then the set is linearly dependent

- $\mathbf{v}_1 = (v_{11}, v_{21}, \dots, v_{n1})$
- Says that if there are more vectors in the set than elements in each vector, one must be linearly dependent

Matrix Inversion: Existence

The following theorem tells that

Theorem

Suppose $\mathbf{X}$ is an $n \times n$ matrix. Recall we can write this matrix as $\begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_n \end{pmatrix}$.

Then $\mathbf{X}$ has an inverse if and only if $S = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ is linearly independent

Matrix Inversion: Existence

The following theorem tells that

Theorem

Suppose $\mathbf{X}$ is an $n \times n$ matrix. Recall we can write this matrix as $\begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \\ \vdots \\ \mathbf{x}_n \end{pmatrix}$.

Then $\mathbf{X}$ has an inverse if and only if $S = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n\}$ is linearly independent

If this is true, we say $\mathbf{X}$ has full rank.

Matrix Inversion: Rank and Span

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Matrix Inversion: Rank and Span

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Columns 1 and 2 are linearly independent, but their sum equals Column 3; the rank of $\mathbf{A} = 2$

Matrix Inversion: Rank and Span

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Columns 1 and 2 are linearly independent, but their sum equals Column 3; the rank of $\mathbf{A} = 2$

These three vectors span $\mathbb{R}^2$ but not $\mathbb{R}^3$: Show that for any $(x_1, x_2) \in \mathbb{R}^2$, we can find a $\lambda_1, \lambda_2, \lambda_3$ such that

$$\lambda_1(1, 0, 1) + \lambda_2(0, 1, 1) + \lambda_3(1, 1, 2) = (x_1, x_2, x_1 + x_2)$$

Matrix Inversion: Rank and Span

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Columns 1 and 2 are linearly independent, but their sum equals Column 3; the rank of $\mathbf{A} = 2$

These three vectors span $\mathbb{R}^2$ but not $\mathbb{R}^3$: Show that for any $(x_1, x_2) \in \mathbb{R}^2$, we can find a $\lambda_1, \lambda_2, \lambda_3$ such that

$$\lambda_1(1, 0, 1) + \lambda_2(0, 1, 1) + \lambda_3(1, 1, 2) = (x_1, x_2, x_1 + x_2)$$

They span $\mathbb{R}^2$ and the space of all vectors in $\mathbb{R}^3$ of form $(x_1, x_2, x_1 + x_2)$

Matrix Inversion: Rank and Span

$$\mathbf{A} = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 1 & 1 & 2 \end{bmatrix}$$

Columns 1 and 2 are linearly independent, but their sum equals Column 3; the rank of $\mathbf{A} = 2$

These three vectors span $\mathbb{R}^2$ but not $\mathbb{R}^3$: Show that for any $(x_1, x_2) \in \mathbb{R}^2$, we can find a $\lambda_1, \lambda_2, \lambda_3$ such that

$$\lambda_1(1, 0, 1) + \lambda_2(0, 1, 1) + \lambda_3(1, 1, 2) = (x_1, x_2, x_1 + x_2)$$

They span $\mathbb{R}^2$ and the space of all vectors in $\mathbb{R}^3$ of form $(x_1, x_2, x_1 + x_2)$

Show that for any $(x_1, x_2, x_3) \in \mathbb{R}^3$, we *can't* find a $\lambda_1, \lambda_2, \lambda_3$ such that

$$\lambda_1(1, 0, 1) + \lambda_2(0, 1, 1) + \lambda_3(1, 1, 2) = (x_1, x_2, x_3)$$

Matrix Inversion: Rank and Span

Finding the rank of a matrix **A**

Matrix Inversion: Rank and Span

Finding the rank of a matrix **A**

- Reduce **A** to an echelon matrix **E** using Gaussian elimination

Matrix Inversion: Rank and Span

Finding the rank of a matrix $\mathbf{A}$

- Reduce $\mathbf{A}$ to an echelon matrix $\mathbf{E}$ using Gaussian elimination
- The number of non-zero rows of $\mathbf{E}$ is the rank of $\mathbf{A}$!

Matrix Inversion: Rank and Span

Finding the rank of a matrix $\mathbf{A}$

- Reduce $\mathbf{A}$ to an echelon matrix $\mathbf{E}$ using Gaussian elimination
- The number of non-zero rows of $\mathbf{E}$ is the rank of $\mathbf{A}$!

Properties of ranks

Matrix Inversion: Rank and Span

Finding the rank of a matrix **A**

- Reduce **A** to an echelon matrix **E** using Gaussian elimination
- The number of non-zero rows of **E** is the rank of **A**!

Properties of ranks

- The rank of an $n \times m$ matrix **A** can't be greater than either n or m :
 $\text{Rank}(\mathbf{A}) \leq \min(n, m)$

Matrix Inversion: Rank and Span

Finding the rank of a matrix $\mathbf{A}$

- Reduce $\mathbf{A}$ to an echelon matrix $\mathbf{E}$ using Gaussian elimination
- The number of non-zero rows of $\mathbf{E}$ is the rank of $\mathbf{A}$!

Properties of ranks

- The rank of an $n \times m$ matrix $\mathbf{A}$ can't be greater than either n or m :
 $\text{Rank}(\mathbf{A}) \leq \min(n, m)$
- A matrix has *full rank* if its rank is the biggest it could be:
 $\text{Rank}(\mathbf{A}) = \min(n, m)$

Matrix Inversion: Rank and Span

Finding the rank of a matrix $\mathbf{A}$

- Reduce $\mathbf{A}$ to an echelon matrix $\mathbf{E}$ using Gaussian elimination
- The number of non-zero rows of $\mathbf{E}$ is the rank of $\mathbf{A}$!

Properties of ranks

- The rank of an $n \times m$ matrix $\mathbf{A}$ can't be greater than either n or m :
 $\text{Rank}(\mathbf{A}) \leq \min(n, m)$
- A matrix has *full rank* if its rank is the biggest it could be:
 $\text{Rank}(\mathbf{A}) = \min(n, m)$
- A square matrix is *invertible* if and only if it has full rank

Matrix Inversion: Gauss-Jordan elimination

How to invert a matrix

Matrix Inversion: Gauss-Jordan elimination

How to invert a matrix

Take

$$[\mathbf{A} \mathbf{I}]$$

and use Gauss-Jordan elimination to convert it into

$$[\mathbf{I} \mathbf{A}^{-1}]$$

Matrix Inversion: Gauss-Jordan elimination

How to invert a matrix

Take

$$[\mathbf{A} \mathbf{I}]$$

and use Gauss-Jordan elimination to convert it into

$$[\mathbf{I} \mathbf{A}^{-1}]$$

Gauss-Jordan elimination is similar to Gaussian elimination, but turns matrix into a diagonal matrix, instead of just an echelon matrix

Matrix Inversion: Gauss-Jordan elimination

Let's invert

$$\mathbf{A} = \begin{bmatrix} 2 & 1 & 2 \\ 3 & 1 & 1 \\ 3 & 1 & 2 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

Let's invert

$$\mathbf{A} = \begin{bmatrix} 2 & 1 & 2 \\ 3 & 1 & 1 \\ 3 & 1 & 2 \end{bmatrix}$$

$$\text{Step 1: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 3 & 1 & 1 & 0 & 1 & 0 \\ 3 & 1 & 2 & 0 & 0 & 1 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

Let's invert

$$\mathbf{A} = \begin{bmatrix} 2 & 1 & 2 \\ 3 & 1 & 1 \\ 3 & 1 & 2 \end{bmatrix}$$

$$\text{Step 1: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 3 & 1 & 1 & 0 & 1 & 0 \\ 3 & 1 & 2 & 0 & 0 & 1 \end{bmatrix}$$

$$\text{Step 2: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & -\frac{1}{2} & -1 & -\frac{3}{2} & 0 & 1 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

$$\text{Step 2: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & -\frac{1}{2} & -1 & -\frac{3}{2} & 0 & 1 \end{bmatrix}$$

Since we are creating a diagonal we want to get rid of the “1” in (1, 2)

Matrix Inversion: Gauss-Jordan elimination

$$\text{Step 2: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & -\frac{1}{2} & -1 & -\frac{3}{2} & 0 & 1 \end{bmatrix}$$

Since we are creating a diagonal we want to get rid of the “1” in (1, 2)

$$\text{Step 3: } \begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

$$\text{Step 2: } \begin{bmatrix} 2 & 1 & 2 & 1 & 0 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & -\frac{1}{2} & -1 & -\frac{3}{2} & 0 & 1 \end{bmatrix}$$

Since we are creating a diagonal we want to get rid of the “1” in (1, 2)

$$\text{Step 3: } \begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

This added $2 \times \text{Row 2}$ to Row 1 and subtracted Row 2 from Row 3

Matrix Inversion: Gauss-Jordan elimination

Step 3:
$$\begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

Step 3:
$$\begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

We get a diagonal matrix by adding $2 \times$ Row 3 to Rows 1 & 2

Matrix Inversion: Gauss-Jordan elimination

$$\text{Step 3: } \begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

We get a diagonal matrix by adding $2 \times \text{Row 3}$ to Rows 1 & 2

$$\text{Step 4: } \begin{bmatrix} 2 & 0 & 0 & -2 & 0 & 2 \\ 0 & -\frac{1}{2} & 0 & -\frac{3}{2} & -1 & 2 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

Matrix Inversion: Gauss-Jordan elimination

$$\text{Step 3: } \begin{bmatrix} 2 & 0 & -2 & -2 & 2 & 0 \\ 0 & -\frac{1}{2} & -2 & -\frac{3}{2} & 1 & 0 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

We get a diagonal matrix by adding $2 \times \text{Row 3}$ to Rows 1 & 2

$$\text{Step 4: } \begin{bmatrix} 2 & 0 & 0 & -2 & 0 & 2 \\ 0 & -\frac{1}{2} & 0 & -\frac{3}{2} & -1 & 2 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix}$$

We finish by normalizing

$$\text{Step 5: } \begin{bmatrix} 1 & 0 & 0 & -1 & 0 & 1 \\ 0 & 1 & 0 & 3 & 2 & -4 \\ 0 & 0 & 1 & 0 & -1 & 1 \end{bmatrix} \text{ and } \mathbf{A}^{-1} = \begin{bmatrix} -1 & 0 & 1 \\ 3 & 2 & -4 \\ 0 & -1 & 1 \end{bmatrix}$$

Matrix Inversion: An Application

Suppose we have following system of equations:

$$2x + 1y + 2z = 1$$

$$3x + 1y + 1z = 3$$

$$3x + 1y + 2z = 5$$

Say, $\mathbf{x} = (x, y, z)'$, $\mathbf{b} = (1, 3, 5)'$.

Say, the coefficient matrix for unknowns is A , the inverse of A is computed by the previous example.

Then, we can solve the unknowns by:

$$\mathbf{x} = A^{-1}\mathbf{b} = (4, -11, 2)'$$