

Fiber bundles

Luka Horjak (luka1.horjak@gmail.com)

November 5, 2024

Contents

Introduction	3
1 Manifolds	4
1.1 Smooth manifolds	4
1.2 Structure sheaf, tangent bundle, and differentials	5
1.3 Coverings and group actions	7
2 Fiber bundles	8
2.1 Basic definitions	8
2.2 Vector bundles and principal bundles	11
2.3 Sections	14
2.4 Pull-backs, subbundles and quotient bundles	16
2.5 Cartier divisors	18
2.6 Kernel, image, and exact sequences	19
2.7 Line bundles and the Picard group	21
Index	22

Introduction

These are my lecture notes on the course Fiber bundles in the year 2023/24. The lecturer that year was doc. dr. Riccardo Ugolini.

The notes are not perfect. I did not write down most of the examples that help with understanding the course material. I also did not formally prove every theorem and may have labeled some as trivial or only wrote down the main ideas.

I have most likely made some mistakes when writing these notes – feel free to correct them.

1 Manifolds

1.1 Smooth manifolds

Definition 1.1.1. A space M is *paracompact* if for every open covering $\{U_j \mid j \in I\}$ there exists a locally finite refinement. That is, every point $p \in M$ has a neighbourhood that intersects only finitely many sets of the refinement.

Remark 1.1.1.1. Instead of second-countable, we require manifolds to be paracompact.

Definition 1.1.2. The *Grassmanian* is the space

$$G_{k,n}(\mathbb{C}) = \{V \leq \mathbb{C}^n \mid \dim V = k\}.$$

Proposition 1.1.3. The Grassmanian is a complex manifold of dimension $k(n - k)$.

Proof. Let $M_{k,n}(\mathbb{C})$ be the set of $k \times n$ matrices of maximal rank. This is an open subspace of $\mathbb{C}^{k \times n}$, therefore it is a manifold of dimension $k \cdot n$. We can then define a map $\pi: M_{k,n}(\mathbb{C}) \rightarrow G_{k,n}(\mathbb{C})$ mapping each matrix to the span of its rows. Note that $\pi(A) = \pi(B)$ if and only if there exists some $g \in \text{GL}_k(\mathbb{C})$ with $gA = B$. That is,

$$G_{k,n}(\mathbb{C}) = M_{k,n}(\mathbb{C}) / \text{GL}_k(\mathbb{C}).$$

For $A \in M_{k,n}(\mathbb{C})$, denote by A_1, \dots, A_ℓ its $k \times k$ -minors. As $\text{rank } A = k$, there exists an index j such that $\det A_j \neq 0$. Let

$$U_j = \{\pi(A) \mid \det A_j \neq 0\}.$$

By the above observation, this is an open covering of $G_{k,n}(\mathbb{C})$. Define $\varphi_j: U_j \rightarrow \mathbb{C}^{k(n-k)}$ as follows – choose a matrix A such that $A_j \neq 0$. Then define

$$\varphi_j(\pi(A)) = A_j^{-1} \cdot B_j,$$

where B_j is A without the columns of A_j . This is clearly a homomorphism. It is easy to see that the transition maps are holomorphic. \square

Definition 1.1.4. A smooth manifold is *orientable* if there exists an atlas such that the jacobians of its transition maps have positive determinants.

Proposition 1.1.5. Let M be a complex manifold of complex dimension n . Then M is also a smooth orientable manifold of real dimension $2n$.

Proof. We can clearly see that M is a smooth manifold of real dimension $2n$. Let $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ be a complex atlas for M . For $\alpha, \beta \in I$, take $F = \varphi_\alpha \circ \varphi_\beta^{-1}$ and write $F(z) = u(x, y) + iv(x, y)$, where $z = x + iy$ for $x, y \in \mathbb{R}^n$. Then

$$dF = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{bmatrix} = \begin{bmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ -\frac{\partial u}{\partial y} & \frac{\partial u}{\partial x} \end{bmatrix},$$

therefore

$$\det(dF) = \left(\det \frac{\partial u}{\partial x} \right)^2 + \left(\det \frac{\partial u}{\partial y} \right)^2 > 0. \quad \square$$

1.2 Structure sheaf, tangent bundle, and differentials

Definition 1.2.1. Let M be a smooth manifold and $p \in M$. Consider¹

$$G_p = \{(U, f) \mid p \in U \wedge f \in \mathcal{C}^\infty(U) \wedge U \text{ is open}\}.$$

We define an equivalence relation on G_p by $(U, f) \sim (V, g)$ if there exists an open set $W \subseteq U \cap V$ containing p such that $f|_W = g|_W$. The set G_p/\sim is the space of *germs* of functions at p , denoted by $\mathcal{C}_{M,p}^\infty$. The equivalence class of (U, f) is denoted by f_p .

Remark 1.2.1.1. The sets $\mathcal{C}_{M,p}^\infty$ and $\mathcal{O}_{M,p}$ are commutative rings with a neutral element. The latter is also an integral domain.

Remark 1.2.1.2. The map $\mathcal{C}_{M,p}^\infty \rightarrow \mathbb{R}$, given by $f_p \mapsto f(p)$, is well defined and the set $\mathcal{M}_{M,p} = \{f_p \mid f(p) = 0\}$ is a maximal ideal.

Definition 1.2.2. Let $U \subseteq M$ be an open set. Define

$$\mathcal{C}_M^\infty(U) = \{f: U \rightarrow \mathbb{R} \mid f \text{ is smooth}\}.$$

The operator \mathcal{C}_M^∞ is called the *structure sheaf* of M .

Definition 1.2.3. A *derivation* v on $\mathcal{C}_{M,p}^\infty$ is a linear operator $v: \mathcal{C}_{M,p}^\infty \rightarrow \mathbb{R}$ such that

$$v(f_p g_p) = f(p)v(g_p) + g(p)v(f_p)$$

for all $f_p, g_p \in \mathcal{C}_{M,p}^\infty$. The set of all derivations on $\mathcal{C}_{M,p}^\infty$ is denoted by $T_p M$.

Remark 1.2.3.1. Let (U, φ) be a local chart around p such that $\varphi(p) = 0$. Then $\varphi^*: \mathcal{C}_{\mathbb{R}^n,0}^\infty \rightarrow \mathcal{C}_{M,p}^\infty$, given by $\varphi^*(f) = f \circ \varphi$, is a ring isomorphism.

Remark 1.2.3.2. The map $d\varphi_p: T_p M \rightarrow T_0 \mathbb{R}^n$, given by

$$d\varphi_p(v)(f) = v(f \circ \varphi),$$

is an isomorphism of vector spaces.

Definition 1.2.4. We define

$$\frac{\partial}{\partial x_j^\alpha}(p)(f) = \left. \frac{\partial (f \circ \varphi_\alpha^{-1})}{\partial x_j} \right|_0$$

for $f \in \mathcal{C}_{M,p}^\infty$.

Lemma 1.2.5. The set

$$\left\{ \frac{\partial}{\partial x_j}(p) \mid j \leq n \right\}$$

is a basis of $T_p M$.

Proof. We can assume $M = \mathbb{R}^n$ and $p = 0$. Take $f \in \mathcal{C}_{\mathbb{R}^n,0}^\infty$ and $v \in T_0 \mathbb{R}^n$. Then

$$v(f) = v \left(f(0) + \sum_{j=1}^n c_j x_j + O(|x|^2) \right) = \sum_{j=1}^n c_j v(x_j),$$

¹ Replace \mathcal{C}^∞ with \mathcal{O} for complex manifolds.

since $v\left(O\left(|x|^2\right)\right) = 0$. Hence

$$v = \sum_{j=1}^n v(x_j) \frac{\partial}{\partial x_j}(p)$$

as $c_j = \frac{\partial f}{\partial x_j}(0)$. It follows that the above set is in fact a generator of $T_p M$. As they are clearly linearly independent, they indeed form a basis. \square

Definition 1.2.6. Let M and N be manifolds and $f: M \rightarrow N$ be a smooth map. For $p \in M$, define $df_p: T_p M \rightarrow T_{f(p)} N$ via

$$df_p(v)(h) = v(h \circ f)$$

for $v \in T_p M$ and $h \in \mathcal{C}_{N,f(p)}^\infty$.

Proposition 1.2.7. Let $p \in M$ and $f: M \rightarrow N$ be a smooth map. Let (U, φ) be a local chart around p and (V, ψ) be a local chart around $f(p)$. Then the matrix of the linear map df_p with respect to the standard bases is given by the jacobian matrix of $\psi \circ f \circ \varphi^{-1}$ in $\varphi(p)$.

Definition 1.2.8. Let M be a smooth manifold of dimension n . A topological subspace $N \subseteq M$ is a *regular submanifold* of codimension k if for every point $p \in N$ there exists a local chart (U, φ) near $p \in M$ such that

$$\varphi(N \cap U) = \{x \in \varphi(U) \subseteq \mathbb{R}^n \mid \forall j \leq k: x_j = 0\}.$$

Proposition 1.2.9. If N is a regular submanifold, then for every point $p \in N$ there exists a neighbourhood U of p and a smooth map $F: U \rightarrow \mathbb{R}^k$ such that $N \cap U = \{q \in U \mid f(q) = 0\}$ and dF_q is surjective for all $q \in N \cap U$.

Proof. Take F to be the projection on the first k coordinates. \square

Theorem 1.2.10 (Rank). Let M_1 and M_2 be smooth manifolds of dimensions n and m . Let $F: M_1 \rightarrow M_2$ be a smooth map. Assume that $\text{rank}(df_p) = k \in \mathbb{N}$ for all $p \in M_1$. Then for all $a \in F(M_1)$ we have that $F^{-1}(a)$ is a regular submanifold of dimension k in M_1 .

Proposition 1.2.11. Let $(a_0, \dots, a_n) \in \mathbb{C}^{n+1} \setminus \{0\}$ and define

$$H = \left\{ [z_0 : \dots : z_n] \in \mathbb{CP}^n \mid \sum_{j=0}^n a_j z_j = 0 \right\}.$$

Then H is a submanifold.

1.3 Coverings and group actions

Proposition 1.3.1. Let M be a smooth manifold with covering (\widetilde{M}, π) . Then \widetilde{M} has a unique smooth structure such that π is smooth.

Definition 1.3.2. A group G is a *Lie group* if it is a topological space admitting a smooth structure such that the operation $G \times G \rightarrow G$, given by $(g, h) \mapsto gh^{-1}$, is smooth.

Definition 1.3.3. For a smooth manifold M denote

$$\text{Aut}(M) = \{f: M \rightarrow M \mid f \text{ is a diffeomorphism}\}.$$

A group G *acts* on M via $\varphi: G \rightarrow \text{Aut}(M)$ if φ is a homomorphism. The action is *faithful* or *effective* if φ is injective.

Definition 1.3.4. Given an action $\varphi: G \rightarrow \text{Aut}(M)$, we define the *orbit* of $x \in M$ as

$$Gx = \{\varphi(g)(x) \mid g \in G\}.$$

Define an equivalence relation on M as $x \sim y \iff Gx = Gy$. Denote $M/G = M/\sim$.

Remark 1.3.4.1. For a smooth manifold M , $M/\text{Aut}(M)$ is a point.

Definition 1.3.5. A *complex torus* is the quotient \mathbb{C}^n/G , where

$$G = \left\{ z \mapsto z + \sum_{j=1}^{2n} \lambda_j w_j \mid \forall j: \lambda_j \in \mathbb{Z} \right\}$$

for \mathbb{R} -linearly independent vectors w_j .

2 Fiber bundles

2.1 Basic definitions

Definition 2.1.1. Let E and M be smooth manifolds and $\pi: E \rightarrow M$ be a smooth map. The triple (E, M, π) is a *submersion* if $d\pi_p: T_p E \rightarrow T_{\pi(p)} M$ is surjective for all $p \in E$.

Definition 2.1.2. A *fiber bundle* is a quadruple (E, M, F, π) , where E, M and F are smooth manifolds and $\pi: E \rightarrow M$ is a surjective submersion such that every point $x \in M$ has a neighbourhood $U \subseteq M$ and a diffeomorphism $\varphi: \pi^{-1}(U) \rightarrow U \times F$ with $\pi = \pi_1 \circ \varphi$, where $\pi_1: U \times F \rightarrow U$ is the projection.

$$\begin{array}{ccc} \pi^{-1}(U) & \xrightarrow{\varphi} & U \times F \\ & \searrow \pi & \downarrow \pi_1 \\ & & U \end{array}$$

Definition 2.1.3. Let G be a Lie group. A fiber bundle (E, M, F, π) is a *bundle with structure group G* if the following conditions hold:

- i) The group G acts effectively on F .
- ii) There exists an open covering $\{U_\alpha \mid \alpha \in I\}$ of M such that for all $\alpha \in I$ there exists diffeomorphism $\varphi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times F$, such that if $U_\alpha \cap U_\beta \neq \emptyset$ there exists a smooth map $g_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G$ satisfying

$$\varphi_\alpha \circ \varphi_\beta^{-1}(x, f) = (x, g_{\alpha,\beta}(x)f)$$

for all $x \in U_\alpha \cap U_\beta$ and $f \in F$.

The set $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ is called the *trivializing atlas*, while $\{g_{\alpha,\beta} \mid \alpha, \beta \in I\}$ are the *(local) transition functions*.

Remark 2.1.3.1. Transition functions satisfy $g_{\alpha,\alpha}(x) = \text{id}_F$ for all $x \in U_\alpha$. Furthermore, if $x \in U_\alpha \cap U_\beta$, then

$$g_{\beta,\alpha}(x) \cdot g_{\alpha,\beta}(x) = \text{id}_F.$$

Finally, if $x \in U_\alpha \cap U_\beta \cap U_\gamma$, then

$$g_{\alpha,\beta}(x) \cdot g_{\beta,\gamma}(x) \cdot g_{\gamma,\alpha}(x) = \text{id}_F.$$

These are called *cocycle conditions*.

Definition 2.1.4. Let M and M' be manifolds with fiber bundles $\pi_E: E \rightarrow M$ and $\pi_{E'}: E' \rightarrow M$. A *morphism of bundles* is a pair (f, φ) of maps $f: M \rightarrow M'$ and $\varphi: E \rightarrow E'$ with $\pi_{E'} \circ \varphi = f \circ \pi_E$.

$$\begin{array}{ccc} E & \xrightarrow{\varphi} & E' \\ \pi_E \downarrow & & \downarrow \pi_{E'} \\ M & \xrightarrow{f} & M' \end{array}$$

If f and φ are diffeomorphisms, the bundles are *equivalent*.

Definition 2.1.5. Let $U \subseteq M$ be an open set. The bundle E is *trivial over U* if the bundle $E|_U = \pi^{-1}(U)$ is equivalent to $U \times F$.

Proposition 2.1.6. Let M be a manifold, E a bundle over M with fiber F , and G its structure group. Let $\{(U_\alpha, \varphi_\alpha^E) \mid \alpha \in I\}$ be a trivializing atlas for E with transition functions $\{g_{\alpha,\beta} \mid \alpha, \beta \in I\}$. Let E' be another bundle over the manifold M with trivializing atlas $\{(U_\alpha, \varphi_\alpha^{E'}) \mid \alpha \in I\}$, fiber F' , structure group G' and transition functions $\{h_{\alpha,\beta} \mid \alpha, \beta \in I\}$. If a smooth map $\psi: E \rightarrow E'$ induces an equivalence (id, ψ) of bundles, then for all $\alpha \in I$, posing

$$\psi_\alpha = \varphi_\alpha^{E'} \circ \psi \circ (\varphi_\alpha^E)^{-1} = (\psi'_\alpha, \psi''_\alpha): U_\alpha \times F \rightarrow U_\alpha \times F',$$

we have $\psi'_\alpha = \text{id}$, $\psi_\alpha(x, \cdot): \{x\} \times F \rightarrow \{x\} \times F'$ is a diffeomorphism for all x , and if $U_\alpha \cap U_\beta \neq \emptyset$, then

$$\psi''_\beta(x, t) = h_{\beta,\alpha}(x) \cdot \psi''_\alpha(x, g_{\alpha,\beta}(x)t).$$

Vice-versa, suppose that there exists a family $\{\psi_\alpha = (\text{id}, \psi''_\alpha) \mid \alpha \in I\}$ of smooth maps $\psi_\alpha: U_\alpha \times F \rightarrow U_\alpha \times F'$ such that, for every $x \in U_\alpha$, $\psi''_\alpha(x, \cdot): \{x\} \times F \rightarrow \{x\} \times F'$ is a diffeomorphism satisfying the above equation. Then there exists a bundle equivalence (id, ψ) such that $\psi_\alpha = \varphi_\alpha^{E'} \circ \psi \circ (\varphi_\alpha^E)^{-1}$ for all $\alpha \in I$.

Proof. Let $(x, t) \in U_\alpha \times F$ and consider ψ_α . We see that

$$\begin{aligned} \psi_\beta &= \varphi_\beta^{E'} \circ \psi \circ (\varphi_\beta^E)^{-1} \\ &= \varphi_\beta^{E'} \circ (\varphi_\alpha^{E'})^{-1} \circ \varphi_\alpha^{E'} \circ \psi \circ (\varphi_\alpha^E)^{-1} \circ \varphi_\alpha^E \circ (\varphi_\beta^E)^{-1} \\ &= (\text{id}, h_{\beta,\alpha}) \circ \psi_\alpha \circ (\text{id}, g_{\alpha,\beta}), \end{aligned}$$

as required.

Now define $\psi: E \rightarrow E'$ as

$$\psi(e) = (\varphi_\alpha^{E'})^{-1} \circ \psi_\alpha \circ \varphi_\alpha^E(e)$$

for $e \in \pi^{-1}(U_\alpha)$. It can be verified that ψ is well defined and is an equivalence of fibrations. \square

Theorem 2.1.7. Let M and F be manifolds and G a Lie group acting effectively on F . Let $\{U_\alpha \mid \alpha \in I\}$ be an open cover of M with maps $g_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow G$ whenever $U_\alpha \cap U_\beta \neq \emptyset$ satisfying the cocycle condition. Then there exists a unique² bundle E with base M , fiber F , structure group G and transition functions $\{g_{\alpha,\beta} \mid \alpha, \beta \in I\}$.

Proof. Define

$$E = \bigsqcup_{\alpha \in I} U_\alpha \times F \Big/ \sim,$$

where $(x, f) \sim (y, f')$ if $x = y$ and there exist $\alpha, \beta \in I$ such that $x \in U_\alpha \cap U_\beta$ and $f = g_{\alpha,\beta}(x)f'$. By the cocycle condition, this is in fact an equivalence relation. We define $\pi([x, f]) = x$, which is well defined and continuous. The local trivializations

² Up to equivalence.

$\varphi_\alpha: \pi^{-1}(U_\alpha) \rightarrow U_\alpha \times F$ are given by $[\tilde{x}, \tilde{f}] \mapsto (x, f)$, where (x, f) is the unique representative of $[\tilde{x}, \tilde{f}]$ in $U_\alpha \times F$. We see that φ_α is bijective, its inverse being the quotient map ρ . It follows that it is a homeomorphism.

Up to refining $\{U_\alpha \mid \alpha \in I\}$, we can assume that (U_α, ψ_α) are local charts for M . Let $\{(W_j, \theta_j) \mid j \in J\}$ be another atlas for F . Then $\{[U_\alpha \times W_j] \mid \alpha \in I \wedge j \in J\}$ is an open cover of E . Define $\tilde{\varphi}_{\alpha,j}: \psi_\alpha(U_\alpha) \times \theta_j(W_j)$ as $[x, f] \mapsto (\psi_\alpha(x), \theta_j(f))$. We need to check that transition maps are smooth. Let $(p, t) \in \psi_\beta(U_\alpha \cap U_\beta) \times \theta_k(W_j \cap W_k)$. Then,

$$\begin{aligned} \tilde{\varphi}_{\alpha,j} \circ (\tilde{\varphi}_{\beta,k})^{-1}(p, t) &= \tilde{\varphi}_{\alpha,j} \left([\psi_\beta^{-1}(p), \theta_k^{-1}(t)] \right) \\ &= \tilde{\varphi}_{\alpha,j} \left([\psi_\beta^{-1}(p), g_{\alpha,\beta}(\psi_\beta^{-1}(p)) \cdot \theta_k^{-1}(t)] \right) \\ &= (\psi_\alpha \circ \psi_\beta^{-1}(p), \theta_j(g_{\alpha,\beta}(\psi_\beta^{-1}(p)) \theta_k^{-1}(t))), \end{aligned}$$

which is smooth, hence E is a smooth manifold. We can check that E has the given transition functions. \square

2.2 Vector bundles and principal bundles

Definition 2.2.1. A bundle (E, M, π) with fiber \mathbb{R}^k and structure group $\mathrm{GL}_k(\mathbb{R})$ is a *vector bundle* of rank k , if there exists a trivializing atlas $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ for E such that for every point $x \in U_\alpha$, the map $\varphi_\alpha|_{E_x} : E_x \rightarrow \{x\} \times \mathbb{R}^k$ is a vector space isomorphism.

Remark 2.2.1.1. Let M be a manifold and $f: \mathbb{R} \rightarrow \mathbb{R}$ a non-linear diffeomorphism. Define $E = M \times \mathbb{R}$ with cover $\{M\}$ that trivializes E with the map $\varphi: E \rightarrow M \times \mathbb{R}$, $\varphi(x, v) \mapsto (x, f(v))$. This is a bundle with fiber \mathbb{R} and structure group $\mathrm{GL}_1(\mathbb{R})$, but not a vector bundle.

Definition 2.2.2. Let M and M' be manifolds with vector bundles E and E' . A bundle morphism (f, ρ) is a *vector bundle morphism* if the map $\varphi_x = \varphi|_{E_x} : E_x \rightarrow E'_{f(x)}$ is a linear map for all $x \in M$.

Lemma 2.2.3. Let M be a manifold and E a bundle over M with fiber \mathbb{R}^k and structure group $\mathrm{GL}_k(\mathbb{R})$. Then there exists a vector space bundle E' on M that is equivalent to E as a bundle.

Definition 2.2.4. A Lie group G *acts on the right* on a manifold F if $R: G \rightarrow \mathrm{Diff}(F)$ is such that $R(e) = \mathrm{id}$, $R(g^{-1}) = R(g)^{-1}$ and $R(gh) = R(h)R(g)$.

Remark 2.2.4.1. If L is an action, then $R(g) = L(g^{-1})$ is a right action.

Example 2.2.4.2. Let E be a bundle with fiber $F \cong G$ and structure group G . Then G acts on F as follows: Let $\theta: F \rightarrow G$ be a diffeomorphism. Define

$$R_g(f) = \theta^{-1}(\theta(f)g)$$

for $g \in G$, $f \in F$. Then $R_g = R(g) \in \mathrm{Diff}(F)$. We write $R_g(f) = fg$ when the choice for θ is clear.

Definition 2.2.5. Let G be a Lie group. A bundle (P, M, π) with fiber G and structure group G is a *principal bundle* if there exists a trivializing atlas $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ for P such that $\varphi_\alpha|_{P_x} : P_x \rightarrow \{x\} \times G$ is G -equivariant for every $x \in U_\alpha$. That is, if $\varphi_\alpha(v) = (x, \varphi''_\alpha(x, v))$ for $v \in P_x$, then $\varphi''_\alpha(x, vg) = \varphi''_\alpha(x, v)g$ for all $g \in G$.

Remark 2.2.5.1. This action on the right commutes with the action we have from the fact that P is a fiber bundle with structure group G .

Definition 2.2.6. Let M and M' be manifolds with principal bundles P and P' with groups G and G' . Let $\rho: G \rightarrow G'$ be a Lie group morphism. A bundle morphism (f, φ) is a *principal bundle ρ -morphism* if $\varphi_x(pg) = \varphi_x(p)\rho(g)$ holds for all $x \in M$, $p \in P_x$ and $g \in G$.

Lemma 2.2.7. Let M be a manifold with bundle P with fiber G and structure group G . Then there exists a principal bundle P' on M which is equivalent to P as a bundle.

Definition 2.2.8. Let G be a Lie group with Lie subgroup H and let $f: H \hookrightarrow G$ be an immersion. Let P and P' be principal bundles on a manifold M with group G and H respectively. Then P' is a *reduction* of P if there exists a ρ -morphism of principal bundles (id, h) , where with $h: P' \rightarrow P$ is injective.

Proposition 2.2.9. Let P be a principal bundle over M with group G . Let $H \leq G$ be a Lie subgroup. Then we can reduce G to H if and only if there exists a trivializing atlas of P with transition functions in H .

Proof. Suppose that $\{h_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow H \mid \alpha, \beta \in I\}$ are transition functions. They define a principal bundle P' on M with structure group H . Recall that

$$P' = \bigsqcup_{\alpha \in I} U_\alpha \times H \Big/ \sim \quad \text{and} \quad P = \bigsqcup_{\alpha \in I} U_\alpha \times G \Big/ \sim.$$

Note that there is a well defined map $U_\alpha \times H \hookrightarrow U_\alpha \times G$ given by $(x, h) \mapsto (x, h)$.

Let $[x, g] = [x, g_{\alpha,\beta}(x)g] \in P'$ for $g \in H$. The above map hence induces a well defined map on the quotient, therefore we get an injective morphism of principal bundles.

Now assume that $h: P' \rightarrow P$ is a reduction. Let $\{U_\alpha \mid \alpha \in I\}$ be a trivializing atlas for P and P' , and denote by $\varphi'_\alpha: P'|_{U_\alpha} \rightarrow U_\alpha \times H$ the local trivialization for P' . We see that

$$\varphi'_\alpha(p') = (x, \tilde{\varphi}'_\alpha(x, p')) = (x, \tilde{\varphi}'_\alpha(x, e)p').$$

Let $p \in P_x$ for $x \in U_\alpha$. Then there exists some $g \in G$ and $p' \in P'_x$ such that $p = h(p')g$ and if $p = h(p'_1)g_1$, then

$$gg_1^{-1} = h(p')^{-1}h(p'_1) = h((p')^{-1}p'_1) \in H,$$

hence $p' = p'_1(g_1g^{-1})$. We obtain that

$$\tilde{\varphi}'_\alpha(x, p')g = \tilde{\varphi}'_\alpha(x, p'_1g_1g^{-1}) = \tilde{\varphi}'_\alpha(x, p'_1)g_1.$$

We can now define $\psi_\alpha: P|_{U_\alpha} \rightarrow U_\alpha \times G$ by $p \mapsto (x, \tilde{\varphi}'_\alpha(x, p')g)$, where $p = h(p')$.

Consider the transition functions $\varphi'_{\alpha,\beta}$ for P' . Observe that

$$\varphi'_{\alpha,\beta}(x) = \tilde{\varphi}'_\alpha(x, e) \left(\tilde{\varphi}'_\beta(x, e) \right)^{-1}$$

For $x \in U_\alpha \cap U_\beta$ and $t \in G$ we can now write

$$\psi_\alpha \circ \psi_\beta^{-1}(x, t) = \psi_\alpha(p)$$

for $p = h(p')g \in P_x$, where $p' \in P'_x$ and $g \in G$. This is now further equal to

$$\begin{aligned} \psi_\alpha(p) &= (x, \tilde{\varphi}'_\alpha(x, p')g) \\ &= \left(x, \tilde{\varphi}'_\alpha(x, p') \left(\tilde{\varphi}'_\beta(x, p') \right)^{-1} t \right) \\ &= \left(x, \tilde{\varphi}'_\alpha(x, e)p' (p')^{-1} \tilde{\varphi}'_\beta(x, e)^{-1} t \right) \\ &= \left(x, \varphi'_{\alpha,\beta}(x)t \right). \end{aligned}$$

□

Example 2.2.9.1. Let E be a vector bundle over a manifold M . Then its transition maps $g_{\alpha,\beta}: U_\alpha \cap U_\beta \rightarrow \text{GL}_n(\mathbb{R})$ give rise to a principal bundle $P(E)$ with fiber $\text{GL}_n(\mathbb{R})$. Vice-versa is also true – principal $\text{GL}_n(\mathbb{R})$ bundles have unique associated vector bundles.

Definition 2.2.10. A *complex vector bundle of complex rank k* is a vector bundle over a manifold M with fiber \mathbb{C}^k and structure group $\text{GL}_k(\mathbb{C})$. If M is a complex manifold and $\pi: E \rightarrow M$ is holomorphic, then E is a *holomorphic bundle*.

Proposition 2.2.11. Let L and L' be holomorphic fiber bundles of rank 1 in a complex manifold M . Denote by $\{g_{\alpha,\beta} \mid \alpha, \beta \in I\}$ and $\{g'_{\alpha,\beta} \mid \alpha, \beta \in I\}$ their respective transition functions. A holomorphic vector bundle isomorphism $f: L \rightarrow L'$ exists if and only if there exist holomorphic functions $f_\alpha: U_\alpha \rightarrow \mathbb{C}^*$ such that

$$\frac{f_\beta}{f_\alpha} \Big|_{U_\alpha \cap U_\beta} = \frac{g_{\alpha,\beta}}{g'_{\alpha,\beta}}$$

whenever $U_\alpha \cap U_\beta \neq \emptyset$.

Definition 2.2.12. The *tangent bundle* of a manifold M is defined as $TM = \bigsqcup_{p \in M} T_p M$.

Proposition 2.2.13. The tangent bundle is a vector bundle of dimension $n = \dim M$. If $\varphi_{\alpha,\beta}$ are the transition maps of the manifold, then $g_{\alpha,\beta}(p)$ is the matrix of $(d\varphi_{\alpha,\beta})_p$ in the standard basis of \mathbb{R}^n .

Definition 2.2.14. Let E be a vector bundle of rank r on a manifold M . For $p \in M$ let $F(E)_p$ denote the ordered bases of E_p . Then $F(E) = \bigsqcup_{p \in M} F(E)_p$ with the trivial projection is the *frame bundle*.

Proposition 2.2.15. The frame bundle is a principal bundle with fiber $GL_r(\mathbb{R})$. It is equivalent to $P(E)$ as a principal bundle.

Definition 2.2.16. When $E = TM$, we write $FM = F(TM)$.

Example 2.2.16.1. The map $S^{2n+1} \rightarrow \mathbb{CP}^n$, given by $(z_0, \dots, z_n) \mapsto [z_0 : \dots : z_n]$,³ is a principal bundle with fiber S^1 .

Example 2.2.16.2 (Homogeneous spaces). Let G be a Lie group and $H \subseteq G$ be a Lie subgroup. Then $G \rightarrow G/H$ is a principal bundle with fiber H .

³ The *Hopf fibration*.

2.3 Sections

Definition 2.3.1. Let M be a manifold with fiber bundle $\pi: E \rightarrow M$. Let $U \subseteq M$ be an open set. A *section* of E over U is a smooth map $s: U \rightarrow E$ such that $\pi \circ s = \text{id}_U$. The set of sections of E over U is denoted by $\mathcal{C}^\infty(U, E)$. Sections with $U = M$ are called *global sections*.

Proposition 2.3.2. Let E be a fiber bundle over M with fiber F and structure group G . Let $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ be its trivializing atlas with transition maps $\{g_{\alpha, \beta} \mid \alpha, \beta \in I\}$. Let $U \subseteq M$ be an open set. If $s \in \mathcal{C}^\infty(U, E)$, denoting $s_\alpha = \pi_F \circ \varphi_\alpha \circ s$ we have

$$s_\alpha = g_{\alpha, \beta} s_\beta$$

for $U_\alpha \cap U_\beta \neq \emptyset$.

Vice-versa, if $\{s_\alpha \mid \alpha \in I\}$ are smooth functions satisfying the above equation, there exists a unique $s \in \mathcal{C}^\infty(U, E)$ with $s_\alpha = \pi_F \circ \varphi_\alpha \circ s$.

Proof. Note that $s(x) = \varphi_\alpha^{-1}(x, s_\alpha(x))$, therefore

$$(x, s_\alpha(x)) = \varphi_\alpha(s(x)) = \varphi_\alpha(\varphi_\beta^{-1}(x, s_\beta(x))) = (x, g_{\alpha, \beta}(x) s_\beta(x)).$$

If this equation holds, we can define $s(x) = \varphi_\alpha^{-1}(x, s_\alpha(x))$. The same computation as above shows that this is in fact a well-defined section. \square

Definition 2.3.3. The maps s_α are called the *local data* of s .

Definition 2.3.4. Let E be a vector bundle of rank k over M and $U \subseteq M$ an open subset. Then E is trivial over U if and only if there exist $s_1, \dots, s_k \in \mathcal{C}^\infty(U, E)$ such that $\{s_j(x) \mid j \leq k\} \subseteq E_x$ is a basis for every $x \in U$.

Proof. The proof is obvious and need not be mentioned. \square

Proposition 2.3.5. A principal bundle P over M admits a global section if and only if P is equivalent to $M \times G$ as a principal bundle.

Proof. Trivial bundles clearly have sections. Now suppose that $s: M \rightarrow P$ is a section. Then for every $g \in P_x$ there exists a unique $h \in G$ with $s(x)h = g$. We can now define $\Phi: P \rightarrow M \times G$ as $g \mapsto (x, h)$. \square

Corollary 2.3.5.1. A principal bundle $P \rightarrow M$ is equivalent to $M \times G$ if and only if its structure group can be reduced to $\{e\}$.

Proof. If a reduction exists, there exists an embedding $\Phi: M \times \{e\} \hookrightarrow P$. But then $x \mapsto \Phi(x, e)$ is a global section. \square

Definition 2.3.6. The *alternator* $A \in \text{End}(V^{\otimes r})$ is defined as

$$A(v_1 \otimes \cdots \otimes v_r) = \frac{1}{r!} \sum_{\sigma \in S_r} \text{sgn}(\sigma) v_{\sigma(1)} \otimes \cdots \otimes v_{\sigma(r)}.$$

Remark 2.3.6.1. It holds that $A^2 = A$.

Definition 2.3.7. The *external product* is defined as

$$\bigwedge^r V = A(V^{\otimes r}).$$

We write $v_1 \wedge \cdots \wedge v_r = A(v_1 \otimes \cdots \otimes v_r)$.

Definition 2.3.8. A *differential form* is a section of T^*M .

2.4 Pull-backs, subbundles and quotient bundles

Definition 2.4.1. Let M and N be manifolds, $f: M \rightarrow N$ a smooth map and $\pi: E \rightarrow N$ a bundle with fiber F and structure group G . The *pullback* f^*E with fiber F and structure group G over M is the bundle

$$f^*E = \{(m, e) \in M \times E \mid f(m) = \pi(e)\}.$$

Proposition 2.4.2. If $\{g_{\alpha,\beta} \mid \alpha, \beta \in I\}$ are the transition functions for E , then the functions $\{g_{\alpha,\beta} \circ f \mid \alpha, \beta \in I\}$ are the transition functions of f^*E .

Proof. Let $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ be the local trivialization for E . Define the map

$$\psi_\alpha: (\pi')^{-1}(f^{-1}(U_\alpha)) \rightarrow f^{-1}(U_\alpha) \times F$$

by $\psi_\alpha(m, e) = (m, \pi_F \circ \varphi_\alpha(e))$. These are trivializations for f^*E . Note that

$$\pi_F \circ \varphi_\beta(e) = \pi_F(g_{\beta,\alpha}(\pi(e)) \cdot \varphi_\alpha(e)) = \pi_F(g_{\beta,\alpha}(f(m))) \cdot \pi_F(\varphi_\alpha(e)),$$

as required. \square

Proposition 2.4.3. Let $f: M \rightarrow N$ be a smooth map. Let $\pi: E \rightarrow N$ and $\rho: E' \rightarrow M$ be fiber bundles with fibers F and structure groups G . Suppose that $(f, g): E' \rightarrow E$ is a fiber bundle equivalence. Then $E' \cong f^*E$.

Proof. Let $\Phi: E' \rightarrow E$ be defined as $\tilde{e} \mapsto (\rho(\tilde{e}), g(\tilde{e}))$.

$$\begin{array}{ccc} E' & \xrightarrow{g} & E \\ \rho \downarrow & & \downarrow \pi \\ M & \xrightarrow{f} & N \end{array}$$

\square

Definition 2.4.4. Let E and F' be vector bundles over M . If there exists a vector bundle morphism $i: F' \rightarrow E$ which is injective on the fibers, we call $F = i(F')$ a *subbundle* of E .

Proposition 2.4.5. Let M be a manifold with a vector bundle $\pi: E \rightarrow M$ of rank k and let $F = i(F')$ be a subbundle of rank $\ell \leq k$. Then i is an embedding and $F \subseteq E$ is a submanifold. Moreover, $\pi|_F: F \rightarrow M$ is a vector bundle and there exists a trivializing atlas $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ of E such that $\{(U_\alpha, \pi_\ell \circ \varphi_\alpha) \mid \alpha \in I\}$ is a trivializing atlas for F , where π_ℓ is the projection on the first ℓ coordinates.

Example 2.4.5.1. Let $S \subseteq M$ be a submanifold and $i: S \hookrightarrow M$ the inclusion map. Then $di: TS \rightarrow TM$ is a fiber-wise injection. In particular, $TS \subseteq TM|_S$ is a subbundle over S .

Definition 2.4.6. Let M be a manifold with a vector bundle $E \rightarrow M$ of rank k . Let $F \subseteq E$ be a subbundle of rank ℓ . Define a relation $e \sim e' \iff e, e' \in E_x \wedge e - e' \in F_x$. We denote the set of equivalence classes by E/F .

Proposition 2.4.7. The set E/F is a vector bundle such that $\rho: E \rightarrow E/F$, given by $e \mapsto [e]$, is a morphism, and $(E/F)_x = E_x/F_x$ for all $x \in M$. Furthermore, if $Q \rightarrow M$ is a vector bundle with fibers E_x/F_x and there exists a morphism $\rho': E \rightarrow Q$ with $v \mapsto [v]$, then $Q \cong E/F$.

Proof. Let $\pi': E/F \rightarrow M$ be given by $[e] \mapsto \pi(e)$, which is clearly well-defined. Let $\{(U_\alpha, \varphi_\alpha) \mid \alpha \in I\}$ be a trivializing atlas, adapted to F . Define $P': U_\alpha \times \mathbb{R}^k \rightarrow U_\alpha \times \mathbb{R}^{k-\ell}$ as the projection onto the last $k - \ell$ coordinates. Note that $e \sim e'$ if and only if we have $P'(\varphi_\alpha(e)) = P'(\varphi_\alpha(e'))$.

Define a map $\psi_\alpha: E/F|_{U_\alpha} \rightarrow U_\alpha \times \mathbb{R}^{k-\ell}$ by $\psi_\alpha([e]) = (\pi'(e), P' \circ \varphi_\alpha(e))$. This is clearly well defined and bijective. The quotient topology on E/F is the same as the one induced by $\{\psi_\alpha \mid \alpha \in I\}$. The transition maps are given by the block-matrix structure of subbundles transition maps. \square

Definition 2.4.8. The bundle E/F is called the *quotient bundle*.

Definition 2.4.9. Let $S \subseteq M$ be a submanifold. The *normal bundle* to S in M is the quotient bundle $NS = TM|_S / TS$.

2.5 Cartier divisors

Definition 2.5.1. Let M be a complex manifold. An *effective Cartier divisor* D on M is given by an open cover $\{U_\alpha \mid \alpha \in I\}$ and holomorphic functions $f_\alpha: U \rightarrow \mathbb{C}$, not identically zero, such that the quotients $f_{\alpha,\beta} = \frac{f_\alpha}{f_\beta}$ are holomorphic functions on $U_\alpha \cap U_\beta$ without zeros.

Remark 2.5.1.1. The functions $\{f_{\alpha,\beta} \mid \alpha, \beta \in I\}$ satisfy the cocycle conditions, hence they induce a line bundle $\mathcal{O}(D)$. Vice-versa, given a line bundle and a non-zero global holomorphic section, its local data define a divisor.

2.6 Kernel, image, and exact sequences

Proposition 2.6.1. Let $\varphi: E \rightarrow F$ be a morphism of vector bundles over a manifold M . Then $\ker \varphi$ and $\operatorname{im} \varphi$ are vector subbundles if and only if $\operatorname{rank} \varphi_x$ is constant for $x \in M$.

Proof. Let $k = \operatorname{rank} \varphi_x$, $m = \operatorname{rank} E$ and $l = \operatorname{rank} F$. Let $U \subseteq M$ be a trivializing open set for E and F .

Locally, the map $\varphi: U \times \mathbb{R}^m \rightarrow U \times \mathbb{R}^l$ is given by $\varphi(x, a) = (x, A(x)a)$ for a map $A: U \rightarrow \mathbb{R}^{l \times m}$. Let $x_0 \in U$ and assume that the first k rows of $A(x_0)$ are independent. Replace U by a smaller set such that the first k rows of $A(x)$ are independent for all $x \in U$. Denote the first k rows of $A(x)$ by $B(x)$. Then

$$\ker \varphi = \{(x, a) \in U \times \mathbb{R}^m \mid B(x)a = 0\}.$$

Note that the map $F: U \times \mathbb{R}^m \rightarrow \mathbb{R}^k$ has maximal rank, hence $F^{-1}(0) = \ker \varphi \cap E|_U$ is a submanifold.

We can also suppose that the $k \times k$ minor $B'(x)$ of $B(x)$, given by the first k columns, is non-singular – that is, $\det B'(x) \neq 0$. Let B'' be the remainder of the matrix. Then $B(x)a = 0$ if and only if $a' = -B'(x)^{-1}B''(x)a''$.

We found $(m - k)$ independent solutions depending smoothly on $x \in U$ and pointwise generating $\ker \varphi$. Let $v_j(x) \in E_x$ be the image of $a_j(x)$ under the trivialization. We can complete $\{v_j(x) \mid j \leq m - k\}$ to a basis of sections of $E|_U$. This basis trivializes $E|_U$ as

$$\sum_{j=1}^m b_j v_j(x) \mapsto (x, b_1, \dots, b_m),$$

with

$$\ker \varphi = \{(x, b_1, \dots, b_m) \in U \times \mathbb{R}^m \mid \forall j > m - k: b_j = 0\}$$

locally. Thus,

$$\sum_{j=1}^{m-k} b_j v_j(x) \mapsto (x, b_1, \dots, b_{m-k}),$$

are local trivializations for $\ker \varphi$.

Let now $Q = E/\ker \varphi$. Then $\varphi: E \rightarrow F$ induces a map $\rho: Q \rightarrow F$ which is injective on the fibers, hence $\rho(Q) = \varphi(E) = \operatorname{im} \varphi$ is a subbundle. \square

Definition 2.6.2. Let E, E' and E'' be vector bundles over a manifold M . Let $\alpha: E' \rightarrow E$ and $\beta: E \rightarrow E''$ be morphisms. If α is injective, β is surjective, and $\operatorname{im} \alpha = \ker \beta$, then

$$0 \longrightarrow E' \xrightarrow{\alpha} E \xrightarrow{\beta} E'' \longrightarrow 0$$

is a *short exact sequence*.

Remark 2.6.2.1. In a short exact sequence, $E'' \cong E/E'$.

Proposition 2.6.3. Suppose E, E' and E'' form a short exact sequence with morphisms α and β as above, and let F be another vector bundle over M . Then the sequence

$$0 \longrightarrow E' \otimes F \xrightarrow{\alpha \otimes \operatorname{id}} E \otimes F \xrightarrow{\beta \otimes \operatorname{id}} E'' \otimes F \longrightarrow 0$$

is also exact. Similarly, the sequences

$$0 \longrightarrow \operatorname{Hom}(F, E') \xrightarrow{\alpha \circ \cdot} \operatorname{Hom}(F, E) \xrightarrow{\beta \circ \cdot} \operatorname{Hom}(F, E'') \otimes F \longrightarrow 0$$

and

$$0 \longrightarrow \operatorname{Hom}(E', F) \xrightarrow{\cdot \circ \alpha} \operatorname{Hom}(E, F) \xrightarrow{\cdot \circ \beta} \operatorname{Hom}(E'', F) \otimes F \longrightarrow 0$$

are exact. Finally, if $f: N \rightarrow M$ is smooth, the sequence

$$0 \longrightarrow f^*(E') \longrightarrow f^*(E) \longrightarrow f^*(E'') \longrightarrow 0$$

is also exact.

Theorem 2.6.4. Suppose that E , E' and E'' form a short exact sequence as above. Then $\det E \cong \det E' \otimes \det E''$.

Proof. Take a trivialization adapted to $\alpha(E')$, that is

$$g_{\alpha, \beta}^E = \begin{bmatrix} g_{\alpha, \beta}^{E'} & k_{\alpha, \beta} \\ 0 & g_{\alpha, \beta}^{E''} \end{bmatrix}.$$

Then $\det g_{\alpha, \beta}^E = \det g_{\alpha, \beta}^{E'} \cdot \det g_{\alpha, \beta}^{E''}$. □

Definition 2.6.5. Let M be a manifold. Then $K_M = \det(T^*M)$ is the *canonical bundle* of M .

Remark 2.6.5.1. Let L be a line bundle. Since $L \otimes L^* \cong \operatorname{Hom}(L, L)$, it admits a global section $x \mapsto \operatorname{id}_L$. Hence $L \otimes L^*$ is trivial and $E \otimes L \cong F$ if and only if $E \cong F \otimes L^*$ for any two vector bundles E and F .

Theorem 2.6.6. Let $S \subseteq M$ be a complex submanifold of codimension 1. Then

$$K_S = (K_M \otimes \mathcal{O}([S]))|_S.$$

Proof. We have a short exact sequence

$$0 \longrightarrow TS \longrightarrow TM|_S \longrightarrow NS \longrightarrow 0$$

over S . Taking the dual and using the previous theorem, we find that

$$K_M|_S \cong K_S \otimes \mathcal{O}([S])^*|_S.$$

As $\mathcal{O}([S])$ is a line bundle, we are done by the above remark. □

2.7 Line bundles and the Picard group

For this section, let M denote a complex manifold of dimension n .

Definition 2.7.1. A holomorphic vector bundle of rank 1 on M is called a *line bundle*. The set of all line bundles (modulo equivalence) is the *Picard group* $\text{Pic}(M)$.

Proposition 2.7.2. The set $\text{Pic}(M)$ is an abelian group with $L^{-1} = L^*$ and neutral element $M \times \mathbb{C}$.

Proof. The proof is obvious and need not be mentioned. □

Definition 2.7.3. The *tautological bundle* on \mathbb{CP}^n is given by

$$\mathcal{O}(-1) = \left\{ ([p], v) \in \mathbb{CP}^n \times \mathbb{C}^{n+1} \mid v \in \mathbb{C} \times p \right\}.$$

Definition 2.7.4. Define $\mathcal{O}(1) = \mathcal{O}(-1)^*$ and $\mathcal{O}(k) = \mathcal{O}(1)^k$.

Proposition 2.7.5. Let $k \geq 0$. The space $\mathcal{O}(\mathbb{CP}^n, \mathcal{O}(k))$ is isomorphic to the space of homogeneous polynomials of degree k in $n+1$ variables. In particular, the bundles $\mathcal{O}(k)$ are pairwise distinct for $k \in \mathbb{Z}$.

Proof. Let $s \in \mathcal{O}(\mathbb{CP}^n, \mathcal{O}(k))$ and let $s_\alpha: U_\alpha \rightarrow \mathbb{C}$ be its local data. Observe that

$$s_\alpha([z]) = \left(\frac{z_\alpha}{z_\beta} \right)^{-k} s_\beta([z])$$

on $U_\alpha \cap U_\beta$. Thus $p(z) = z_\alpha^k s_\alpha$ is independent of α and holomorphic on \mathbb{C}^{n+1} . Furthermore, $p(\lambda z) = \lambda^k p(z)$, hence p is homogeneous of degree k . Since p is holomorphic on \mathbb{C}^{n+1} , we get $k \geq 0$. □

Index

A

alternator, [14](#)

C

canonical bundle, [20](#)

Cartier divisor, [18](#)

cocycle conditions, [8](#)

complex torus, [7](#)

D

derivation, [5](#)

differential form, [15](#)

E

equivalent bundles, [8](#)

external product, [15](#)

F

faithful action, [7](#)

fiber bundle, [8](#)

frame bundle, [13](#)

G

germs, [5](#)

Grassmanian, [4](#)

group action, [7](#), [11](#)

H

holomorphic bundle, [12](#)

Hopf fibration, [13](#)

L

Lie group, [7](#)

line bundle, [21](#)

local data, [14](#)

M

morphism of bundles, [8](#)

N

normal bundle, [17](#)

O

orbit, [7](#)

orientable, [4](#)

P

paracompact, [4](#)

Picard group, [21](#)

principal bundle, [11](#)

principal bundle morphism, [11](#)

pullback, [16](#)

Q

quotient bundle, [17](#)

R

rank theorem, [6](#)

reduction, [11](#)

regular submanifold, [6](#)

S

section, [14](#)

structure group bundle, [8](#)

structure sheaf, [5](#)

subbundle, [16](#)

submersion, [8](#)

T

tangent bundle, [13](#)

tautological bundle, [21](#)

transition functions, [8](#)

trivial bundle, [9](#)

trivializing atlas, [8](#)

V

vector bundle, [11](#), [12](#)

vector bundle morphism, [11](#)