

Algebraic Number Theory

 tuyaux.winak.be/index.php/Algebraic_Number_Theory

Algebraic Number Theory

Richting Wiskunde

Jaar MWIS

November 2017

1. Prove or give a counterexample:
 - An integrally closed domain is also factorial.
 - Assume A a Dedekind domain. Every nonzero element of A is contained in only finitely many ideals of A .
 - A \mathbb{Z} -submodule of \mathbb{Q} is also a fractional ideal of \mathbb{Z} .
2. Which of the following rings are Dedekind domains?
 $\mathbb{Z}[X], \mathbb{C}, \mathbb{Z}[\sqrt{5}], \mathbb{Z}[\sqrt{6}], \mathbb{R}[X^2, X^3]$
 $\mathbb{Z}[X], \mathbb{C}, \mathbb{Z}[5], \mathbb{Z}[6], \mathbb{R}[X^2, X^3]$
3. Consider the equation $x^2 + y^2 = 7z^2$. Show that it has no nontrivial integer solution.
4. Assume $\omega \in \mathbb{C}$ such that $\omega^5 = 1$ ($\omega \neq 1$) and let $K = \mathbb{Q}(\omega)$. What is the discriminant of K ? Furthermore, prove that $(1, \omega, \omega^2, \omega^3)$ is an integral basis of K .

Categorieën:

- Wiskunde
- MWIS