



When Independent Particle Methods Fail: The Effect of Correlations



Christopher Lane

Physics of Condensed Matter and Complex Systems, Theoretical Division
Los Alamos National Laboratory
Los Alamos, NM

LA-UR-26-21459



Los Alamos Computational Condensed Matter Summer School 2025
June 18, 2025




Approaches to the Many-Electron Problem

Many-body
Hamiltonian

$$\hat{H} = \sum_{\alpha\beta} \int d^3r \underbrace{\hat{\psi}_\alpha^\dagger(r) \left(\frac{-\hbar^2 \nabla^2}{2m} + v_{ext}^{\alpha\beta}(r) \right) \hat{\psi}_\beta(r)}_{\substack{\alpha\beta\gamma\delta \in \text{spin} \\ \text{Electronic} \\ \text{Kinetic Energy}}} + \sum_{\alpha\beta\gamma\delta} \iint d^3r d^3r' \underbrace{\hat{\psi}_\alpha^\dagger(r) \hat{\psi}_\beta^\dagger(r') v_{\alpha\beta}^{\delta\gamma}(r, r') \hat{\psi}_\gamma(r') \hat{\psi}_\delta(r)}_{\text{Interactions}}$$

Wavefunction based approaches


Key Quantity: $\Psi(\{r_j\})$



Observables: $\langle \Psi | \hat{O} | \Psi \rangle$

Reduced quantity based approaches

Key Quantity: Simpler physical quantity, e.g., $\rho(r)$



Observables: $F[\rho]$

Correlations



Independent
electron theory



Weak
Correlations



Strong
Correlations

Correlations

Quantities A and B
are *Correlated* if

$$\langle AB \rangle \neq \langle A \rangle \langle B \rangle$$

Measure of
Correlation

$$C_{AB} = \langle AB \rangle - \langle A \rangle \langle B \rangle$$

Spatial $\langle A(rt)B(r't) \rangle \neq \langle A(rt) \rangle \langle B(r't) \rangle$

Temporal $\langle A(rt)B(rt') \rangle \neq \langle A(rt) \rangle \langle B(rt') \rangle$



Strong
Correlations

How Do We Measure Correlations?...




...Hit it with a hammer and listen

$$\langle n(rt)n(r't') \rangle$$

$$\langle \mathbf{S}(rt) \cdot \mathbf{S}(r't') \rangle$$

$$\langle \mathbf{j}(rt) \cdot \mathbf{j}(r't') \rangle$$

$$\langle \{\psi(rt), \psi^\dagger(r't')\} \rangle$$




Pair Correlation
Function for
Electrons and Spin


Electrical and
Optical Properties

Removing (Adding)
Electrons


Signatures of Correlations: Ground State / Thermodynamic




Delocalized → Localized
Behavior




Experimental





VASP Relaxation






WIEN2k Relaxation



Signatures of Correlations: Ground State / Thermodynamic



Signatures of Correlations: Electron Addition/Removal






Renormalization

Kinks

Waterfalls

Signatures of Correlations: Electron Addition/Removal

Not all doom and gloom...




D.W. Latzke et al.
Phys Rev. B 91, 235202 (2015)


Y. Xia et al.
Nature Physics 5, 398–402 (2009)

Z.K. Liu et al. Nature Materials 15, 27–31 (2016)
Su-Yang Xu et al. Nature Physics 11, 748–754 (2015)

Signatures of Correlations: Particle-Hole/Collective Excitations




D. M. Roessler and W. C. Walker, J.
Opt. Soc. Am. 57:835–836, 1967.



J. A. Camargo-Martínez and R. Baquero
Phys. Rev. B 86, 195106 (2012)

Signatures of Correlations: Metal-Insulator Transitions




Metal-Insulator Transitions can arise from:

- Mott transition: (The most common transition)
Arising from intense electron-electron correlation.
- Peierls transition: Electron-phonon interactions can give rise to a transition.
- Anderson transition: When an insulator behavior in metals arises from distortions and lattice defects.

Paths forward?

Ab initio Methods



Low Energy Models

- Exact diagonalization method
 - **Exact**
 - **Limited to finite sizes**
- Quantum Monte Carlo method
 - Non-perturbative and thermodynamic limit
 - **Negative sign problem for fermions**
- Density matrix renormalization group (DMRG) theory
 - Capturing both quantum temporal and spatial fluctuations
 - **Limited to one dimensional (1D) or quasi-1D systems**