



# Disaster Medicine:

Using Modeling and Simulation to Determine Medical Requirements for Responding to Natural and Man-Made Disasters

The past decade is likely to be remembered for its disasters. The first 10 years of the 21st century saw a tragic series of devastating man-made and natural disasters: major terror attacks in the United States, England, India, and Spain; the wrath of Hurricanes Katrina and Rita; and the sweeping destruction of the 2004 Indian Ocean earthquake and tsunami. Natural or man-made, these disasters strained the capabilities of local medical facilities and medical-response units to treat the resulting injured and sick. During this same period, U.S. military forces fought wars in Iraq and Afghanistan, stressing the capabilities of military medical providers. In the latter case, however, the evolution of medical modeling and simulation (M&S) helped ease the guesswork previously used to determine what type of treatment facilities were required and how much equipment and consumable supplies they should carry. Unfortunately, medical M&S for civilian disaster response has not kept pace.

#### MARTIN HILL, BA, CHS III, EMT-T, MEMS

Division of Expeditionary Medical Modeling, Simulation & Analysis Naval Health Research Center San Diego, California martin.hill@med.navy.mil This article is approved by the following for continuing education credit: The American Board for Certification in Homeland Security, CHS® provides this continuing education opportunity for those individuals Certified in Homeland Security, who are required to obtain 30 Certification Maintenance Unit's (CMU) per 3-year recertification cycle.

### AFTER STUDYING THIS ARTICLE, PARTICIPANTS SHOULD BE BETTER ABLE TO DO THE FOLLOWING:

- 1. Explain the concept of medical modeling and simulation.
- 2. Articulate the need to create a centralized database of disasterrelated injuries and illnesses.
- 3. Demonstrate the potential advantages of using medical modeling and simulation in medical disaster planning.

**KEYWORDS:** modeling and simulation, disaster, planning, medical response

**TARGET AUDIENCE:** Disaster response planners justice, government, military, and counterterrorism arenas.

**PROGRAM LEVEL:** Intermediate

**DISCLOSURE:** The author has nothing to disclose.

 $\textbf{PREREQUISITES:} \\ None$ 



espite increased awareness of the need for improved medical response to disasters, little has been done to improve the way local communities, states, and the nation as a whole prepare for major mass-casualty events. Despite millions of dollars in federal homeland security funding following the terrorist attacks on September 11, 2001, there was no systematic review of what was needed for a medical response to a major

disaster. Much of the funding requirements emphasized responding to chemical, biological, radiological, or nuclear events, with little regard given to the most likely mass casualty incidents, such as conventional terrorist bombings and natural disasters. As a result, little progress toward improved disaster response readiness was achieved, and much of what was purchased with that funding was useless (Kimery, May 13, 2008).

As early as 2005, the year of Hurricane Katrina, federal emergency preparedness money began drying up, falling as much as 25% by 2009 (Closing the Seams, 2007; Ferenc, 2010). In 2010, a large-scale federal and local disaster exercise in Las Vegas was canceled due to its high cost, and additional exercises may be canceled as well (Hsu, 2010). At the same time, America's network of trauma centers and hospital emergency rooms (ERs)—the frontline

"receivers" in the event of a terrorist attack or natural disaster—are in crisis. In 2004, the National Foundation for Trauma Care predicted that 10%–20% of the nation's trauma centers would close within 3 years (U.S. Trauma Center Crisis, 2004). Between 2006 and 2008, five trauma centers did close, and another five downgraded their level of care. More than 12% of the nation's hospitals

"Relatively few models have addressed response to man-made disasters such as conventional warfare, radiologic and chemical accidents, acts of terrorism, and large-scale industrial accidents."

(Brandeau, McCoy, Hupert, Holty, & Bravata, 2009)

closed their ERs between 1994 and 2004; many hospitals closed down entirely. In addition, there is a shortage of health care providers, especially trauma and critical care specialists (Kimery, May 20, 2008, 2010; Wilper et al., 2008).



Unprepared for the "Predictable Surprise"

Dwindling funding and emergency medical resources are leaving the United States dangerously unprepared for another disaster with mass casualties. A report issued by the House Committee on Oversight and Government Reform maintains the country is unprepared to deal with the "predictable surprise," which the committee described as a terrorist bombing like the one that struck commuter trains in Madrid, Spain, on March 11, 2004. On March 25, 2008, the committee conducted a survey of 41 Level 1 trauma centers in seven major U.S. cities. None of the cities—New York, Los Angeles, Chicago, Houston, Denver, Minneapolis, Washington, District of Columbia—had the capacity to treat the more than 2,000 casualties created by the Madrid blasts (Hospital Emergency Surge Capacity, 2008).

Military medical providers, like their civilian counterparts, have been stretched thin by two wars and an ongoing shortage of medical personnel (Gahol, 2005; Schrader, 2004). Furthermore, space limitations on cargo ships and aircraft mean battlefield medical treatment facilities (MTFs) must maintain a relatively small logisti-

cal footprint while still providing advanced medical care to hundreds of injured and sick service men and women. Military medical planners frequently use medical M&S to meet both the restrictions and the requirements of their military medical mission. Because military operations today span a wide range of scenarios (e.g., large-scale warfare, counterterrorism and antipiracy operations, peacekeeping, humanitarian and disaster response), medical M&S efforts provide an all-hazards approach to planning.

Civilian medical disaster M&S tools do exist. A study described in the August 2009 issue of *Medical Decision Making* reviewed 55 medical models and M&S programs. For the most part, these were special-purpose models focusing on specific disease outbreaks or pandemics. Only a few addressed natural disasters, and even fewer focused on



man-made disasters. "Despite the frequency and potentially catastrophic health consequences of natural disasters [...] remarkably few published models have focused on the public health and medical responses to such events," the study reported. It also found that "Relatively few models have addressed response to man-made disasters such as conventional warfare, radiologic and chemical accidents, acts of terrorism, and large-scale industrial accidents" (Brandeau, McCoy, Hupert, Holty, & Bravata, 2009).

**Optimize Patient Care** 

Medical M&S can help medical disaster planners prepare their communities for a variety of natural and man-made disasters with relatively low cost. For instance, M&S can test a variety of configurations for a proposed disaster MTF to find the best composition of equipment and personnel before committing to the cost of buying matériel or hiring people. The same process can be used to develop strategies to task-organize that response unit for different disaster scenarios (earthquake, tornado, hurricane, flood). Patient

flow and treatment can be optimized by M&S analysis for a single hospital or an entire community's health care network.

None of this can be done, however, without the appropriate empirical data. Unfortunately, little of that is available. Much of what we think we know about disaster medicine is based on conjecture or anecdotal information. The unexpected nature of disasters is at least partly responsible for this; once in the midst of treating disaster victims, providers give little thought to data collection (Dominici, Levy, & Louis, 2005; Heide, 2006; Quick & Hogan, 2002).

A decade ago, military medical M&S suffered from this same lack of data. With the invasions of Afghanistan and Iraq, a concerted, joint-forces effort was launched by military medicine to cap-

ture injury, illness, and treatment data from the battlefield. For example, the Naval Health Research Center (NHRC) maintains the Navy-Marine Corps Combat Trauma Registry Expeditionary Medical Encounter Database (CTR EMED), which has collected data on wounds, diseases, treatments, and patient outcomes from Iraq and Afghanistan since 2004. The data collected by the

CTR EMED make up part of the Joint Theater Trauma Registry (JTTR), which also includes similar data collected by other military branches. A small army of medical coders, nurses, epidemiologists and statisticians, overseen by an institutional review board, process these data. The data, in turn, help develop relevant patient streams for M&S analysis.

A similar data collection effort is needed to make medical M&S

"Knowledge based on systematically collected data from field disaster research studies might help planners avoid common disaster management pitfalls, thereby improving disaster response planning."
(Heide, 2006)



for civilian responders possible. This certainly is not a new idea. In 2005, Fenig and Cone recommended the establishment of a national disaster-victim database (NDVD) capable of collecting real-time data on the injuries and illnesses seen during disasters. "A disaster-victim database will facilitate collection of comprehensive disaster data," they wrote. "However, the data that the



NDVD abstracts can only be as good as the data collected by personnel and individual facilities. Therefore, facilities should be encouraged to develop better standardization and collection methodologies" (Fenig & Cone, 2005). The following year, Heide also urged a comprehensive data collection effort during disasters: "Knowledge based on systematically collected data from field disaster research studies might help planners avoid common disaster management pitfalls, thereby improving disaster response planning" (Heide, 2006).

#### **Collecting Data**

The logical center of gravity for this effort would appear, at first, to be the National Trauma Data Bank (NTDB), which has standardized injury reporting among state trauma registries. However, unless a disaster-related trauma is identified as such by an appropriate *International Classification of Diseases, Ninth Revision* (ICD-9) E code, it would not be identifiable as different from any other trauma. An attempt to use E codes to identify disaster-related traumas in the NTDB database during 2007 found only 244 records (Fantus & Hammond, 2007). This was nearly two years after Hurricane Katrina injured 2,018 persons in New Orleans alone. Many patient encounters likely were not properly recorded due to heavy patient volume (Willams et al., 2005).

## A Short History of Navy Medical Modeling and Simulation

U.S. Marine Corps doctrine calls for maintaining a small logistical footprint to allow for a rapid response from the sea to emerging contingencies. As such, Marine medical treatment facilities (MTFs) have to be small, relatively lightweight, and quickly deployable. In the late 1990s, the Marine Corps turned to modeling and simulation (M&S) to analyze the size of its field MTFs. With the support of the Office of Naval Research, the Naval Health Research Center (NHRC) in San Diego, California, was tasked with developing an evidence-based modeling tool to perform that analysis.

The result was the Estimating Supplies Program (ESP), a deterministic, computer-based, patient-driven model of clinical events that analyzes logistical requirements for specific battlefield scenarios. From the start, ESP was able to reduce the size of Marine Corps medical inventories by an average of 30% while increasing their treatment capabilities (Galarneau, Konoske, Emens-Hesslink, & Pang, 1998; U.S. Patent No. 7,707,042, 2010).

ESP's underlying database of service-specific medical equipment and supplies, clinical tasks, treatment protocols, and models is housed in the Expeditionary Medical Knowledge Warehouse (EMedKW). Besides being used to trim existing Marine MTFs, NHRC's modeling process has been used to develop new battlefield treatment facilities. Among these is the Forward Resuscitative Surgical System, essentially a trauma center packed into three Humvees that was spotlighted by CNN's medical correspondent Dr. Sanjay Gupta during his coverage of the invasion of Iraq. Another battlefield development that was assisted by NHRC's modeling process is the Marine Corps' airborne casualty evacuation system, which has kept hundreds of wounded Marines and sailors alive as they were flown from the front line to the surgical hospital (Galarneau, Konoske, & Pang, 1999; Hill, Galarneau, Pang, & Konoske, 2004).

continue on page 29


Navy medical modeling and simulation relies on real-world data. Naval Health Research Center combined its data collection form with a patient care report form to collect information directly from frontline care providers for the U.S. Navy-Marine Corps Combat Trauma Registry Expeditionary Medical Encounter Database.

Entries circled in red on the CTR /EMEDS form provide NHRC analysts with data for use in statistical analysis of injury causation, and for developing patient treatment profiles for modeling.

Circled entries on Page 1 show the estimated time of injury and the time of arrival at the MTF, providing an approximation of evacuation transit time.

Just below that, the entries indicate the method of injury and whether it was from combat action (gunshot wound, landmine, etc.) or a nonbattle accident (fall, animal bite, electrocution, etc.). Statisticians analyze this data to develop wound and injury occurrence rates for use in casualty estimation software.

Entries below that indicate whether the casualty was wearing protective equipment, such as body armor and Kevlar helmet. This data is used to analyze effectiveness of issued personal protective equipment.



Even if all incidences of disaster-related trauma were recorded in the NTDB with the proper E codes, they would still tell only part of the story. The greater part of a medical disaster response is providing primary care for illnesses that may or may not be related to the disaster itself. The Centers for Disease Control and Prevention (CDC) estimates the number of persons seeking treatment for illnesses after Hurricane Katrina was 4,169, more than twice the number of those suffering traumatic injuries. An

additional 1,321 patients seen after the hurricane only required medication refills or follow-up care (Willams et al., 2005), yet even those patients placed a logistical burden on the responding treatment facilities.

Each type of patient seen by responders, from the response phase of a disaster through the recovery phase, must be considered when modeling medical requirements for such events. In other words, the data collection effort must be patient-centric,

	<u> </u>
OEF Camp	Dwyer, Afghanistan
Name (Last, First MI):	Patient I.D. / SSN:
Prior Medical History: Medications Administered:	
Marrhine /	se #2/Time Dose #3/Time #1 #2 #3 Antibiotic (Name):
C Passister C PM Versed	/ / #1  #2  #3  Other (Name):
Seizure Ulcer (G.I.) Demerol	
Subjective (Chief Complete) / History	
the promote since IED in her of mos outside	of vehicle 5 meters from IED was thrown 10 feet landed
	t has shappel injuries to LEFT side of face, photophob in burs dust cloud & crawling away from blast,
having that his corpsman told him he was r himself the let day but that has since resolve	speating Meds, methaquine, centrum
smoters trove LED was thrown lo feet	de mone
Objective / Exam: has some dizziness, HA, pho	otophobia Denies Fluid Fromears or nose
ENT - [Ms intact B + 3 injun; PERRL 4-> Head/Face - lesion to Left cheek from shraphal	ow and stright, country, cho a tal
eyes - symmetrical, perrin, - foreign body >	Darm: biceps, tendons intact, point TTF
EARS edema. @ hearing difficulty	of medial cordyle of Delbow, so swelling/
Nose - edema - pND -foreign body	I de la la con sent
threat odor,	ofradial heald, & discoloration of skin,
and the second s	1.1-
chest - Normal findings	speech (woth speed, fluency + content), Diff DX:
neuro- CN 2-121, 3/6 strength x4, 14	Dift Dx
ødysmetria x4, pronatordrift, nl	- tocal reuro de lei
	- baro trauma - extremity fx - experience inju
Laboratory Davidso	
Laboratory Results:	X-Ray Results: to globes
11 750	6 10 10 10 10 10 10 10 10 10 10 10 10 10
Assessment / Diagnosis: Primary - wild 18	Secondary - a brasion Omaxil (Tertiary - Delbow pain)
OR Start Time: OR Step Time: Plant / Procedure(s) / Treatment / Operative Notes / Nursing N	Vent Time On: S/p Shrapma Vent Time Off:
slp IED blast & improving m	
	10 000000000000000000000000000000000000
Off may return to full duty	based on al neuro exam + mental
status as well as interval Oir	uprovement in symptoms. Tylenof
@ If sxs worsen or become foce	al, return for re-eval
3) If sxs continue, may need	
17 sxs continue, may need	in agric evas.
	ury: Work/Training Respiratory Dental Ury, Other STDs Dental Misc./Admin./Follow-up
☐ Dermatologic ☐ Injury, Recreational/Sports ☐ O	phthalmologic
Disposition	sych_ Mental Disorders
Disposition Up Chit (Av Date/Time;   Up Chit (Av Date/Time;   Up Chit (Av Date/Time;   Down Chit (	Aviation) Admit to x day(s) N/A Urgent (Medical)
SIQ xday(s)	Other:   Routine   Urgent (Surgical)
Date/Time of Death:	□ N/A
ANATOMIC: Airway Head Neck Chest Abdomen	Pelvis
PHYSIOLOGIC:	
☐ Breathing ☐ CNS ☐ Hemorrhage ☐ Total Body Disruption	
Provider Name (Printed or Typed):	Provider Signature:
MC DSN	

The SOAP notes (Subject, Objective, Assessment/ diagnosis and Planned treatment) on Page 2 provide more detailed information on the injury or injuries the casualty presents with, and are translated by medical coders into ICD-9 codes and injury severity scores for further statistical analysis. This information is also used to develop treatment task profiles for modeling.

not logistics-centric. Counting the number of saline bags used in a disaster response shows only what was used, not why it was used. More important is being able to identify what conditions patients presented with, such as hemorrhagic shock or dehydration, along with their ages and co-morbidities, and in what numbers. In addition to patient presentations of disease and illness, a properly configured and maintained database allows the determination of not only how much IV fluid should be carried, but also what fluid

types (crystalloids vs. colloids), what size catheters, what type of IV administration sets, and even whether other fluid resuscitation methods, such as oral rehydration salts, should be considered.

The most likely sources for these kinds of data in the United States are the medical teams operated by the National Disaster Medical System (NDMS) and other medical disaster response units. During the response to Hurricane Katrina, the CDC launched a well-organized data collection effort focused on the

Figure	1
--------	---

	PC	Medical	Functional	Task	%	Equipment/
	Code	Capability	Area	Profile	Patients	Consumables
CD-9	Code  001  002  003  004  005  006  007  008    200   350	1A 1B 2		Profile  Assign Priority  Assess Patient Insert Endotrach Neurological As: Apply C-Collar Oxygen Set-up Oxygen-Admin Cardiac Monitor Cardiac Resus Set-Up Pulse Or Arterial Puncture Control Hemorrh Order Blood Gar	100 100 100 100 100 60 60 20 20 20 20 20 100 100 100	

This abbreviated example of the NHRC modeling process shows the clinical task profile for patient condition 5 (PC 005), a head injury, being treated in the pre-op section of a Level II field surgical unit. The middle column shows the tasks to be performed, followed by the percentage of patients with PC 5 who will receive the task. The last column shows the equipment and supplies need to complete the task, in this case endotracheal intubation. Not shown in this illustration are data points for quantities of each supply, the number of times a task will be performed in the first 24 hours, and the number in each additional 24-hour period.

NDMS and military medical units that responded to the disaster. Data collection was accomplished using a standardized Injury and Illness Report form to be completed by caregivers for each patient at the time care was given. This Injury and Illness Report form, however, was heavily weighted to the collection of disease type and occurrence. A revision of the form was made to collect more data on injury type and occurrence, but not on patient treatment (Wolkin & Noe, 2007). A few additional changes in this form, to include treatment given to patients as well as the anatomical location of injuries, would give greater visibility to the full range of patient presentations and aid in the statistical analysis of the patients' treatment.

NHRC's battlefield data collection efforts were maximized by combining the elements of its data collection form with the patient care report (PCR) providers are required to fill out. This combination of data form and PCR is a two-page, self-duplicating form that provides spaces for injury/disease type(s), mechanism of injury, anatomical locations, treatment record, and even type body armor worn, if any. The attending provider fills out the form only once, which saves time. When completed, one page stays with the patient's medical record and the second is sent to NHRC for processing (Galarneau et al., 2004). Analysis of this data by NHRC researchers has resulted in numerous epidemiological studies on combat injuries, such as trauma brain injury and post trauma stress disorder, and deployment diseases. Another advantage of such data collection and analysis is the development of "patient streams" used in modeling to emulate various combat scenarios or deployment environments.

The success of military medical M&S is directly related to the concerted effort the military health care community has made in collecting injury and illness data, in real time, on the battlefield.

#### **Modeling Process**

For several years, military medical modeling used 401 patient conditions (PCs) created by the Defense Medical Material Program Office (DMMPO) to represent typical combat and non-battle injuries and diseases encountered in the deployed field environment. These PCs were assigned statistical probabilities of occurrence, identified from patient encounter data in NHRC's CTR EMED and the JTTR mentioned earlier. PC probabilities vary according to deployed scenarios (i.e., large-scale conflicts, stability and support operations, counterinsurgency warfare, special operations) and geographic regions (desert, jungle, mountain, urban). Casual-ty-projection programs use these probabilities to develop patient streams used in the modeling process. These DMMPO PCs are now in the process of being replaced by ICD-9 codes.

Like many other military medical modeling programs, NHRC's modeling process used the DMMPO PCs (and now ICD-9 codes) in its modeling method. That process links clinical tasks to each PC for each level of care (i.e., corpsman or medic, forward aid station, surgical facility) as well as the functional areas within those levels



of care. The medical supplies and quantities required for completion of each clinical task are then identified and mapped to the appropriate task. The modeling process also identifies the number of patients expected to receive a clinical task, since not all patients will receive the same treatment. These clinical "task profiles" are built with input from the appropriate providers to ensure the model simulates how they work in a real-world field environment.

Figure 1 provides an abbreviated representation of the NHRC supply modeling process. In this example, PC 005, a head injury, is treated in the triage area of a battlefield surgical facility. The task profile shows the clinical tasks likely to be performed on this patient type in that functional area, and the percentage of those patients expected to receive each task. The Equipment/Supplies column identifies the items needed to intubate a patient. Not shown in this figure are additional data fields used to calculate supply quantities. These fields include the supply amounts needed to complete the task, how often the task will be repeated in the first 24 hours of treatment, how often the task will be repeated in each subsequent 24-hour period, the time each task requires for completion, and the average length of stay at that level of care. This model is the underlying engine that runs NHRC's M&S programs.

It is important that patient-driven models like NHRC's reflect how the various medical responders work in the field with the equipment and supplies they carry. Battlefield health care is divided into four levels of care, also known as roles. First is far-forward emergency care, which includes the combat medic or corpsman at the point of injury, and the forward aid station, staffed by a physician's assistant or general medical officer and

# A Short History of Navy Medical Modeling and Simulation continued from page 25

In addition to ESP, EMedKW's database provides the foundation for several other NHRC M&S programs. The ReSupply Validation Program (RSVP) uses the ESP modeling process to develop time-phased medical resupply strategies, including push packages. RSVP is also used by the Marine Corps to develop its medical contingency file, a list required by the Pentagon every six months of the medical supplies each military branch would need in the event of a major contingency, such as a large–scale war.

EMedKW also provides the underlying database for the Tactical Medical Logistics Planning Tool (TML+), a medical risk-assessment planning tool. The TML+ graphical user interface allows planners to create either a single treatment facility, such as a field hospital, or an entire network of MTFs and the transportation assets (aircraft, ambulances) needed to move patients between them.

Being stochastic rather deterministic, TML+ uses multiple iterations of varying patient streams to identify logjams caused by shortages in equipment, consumables or transportation assets, shortfalls in personnel, and even increases in patient mortality due to delays in treatment. Planners can perform "what if" experiments to study the effects of increasing or reducing personnel, adding or subtracting treatment beds, or mixing ground and air evacuation assets. (Olson, Bohannan, Peel, Jeschonek, & Leap, 2007).

TML+ has the ability to simulate routine field medical operations, such as sick call, or contingency medical operations for large-scale combat assaults, peacekeeping and stability operations, and unexpected mass casualty events. Originally conceived as a battlefield–planning tool, TML+ also has been used to provide course-of-action analysis, manpower studies, even the viability of using unmanned aerial vehicles for casualty evacuation.

Powering NHRC's modeling effort is the statistical analysis of combat injuries and diseases. Originally, this focused on data from previous wars such as World War II, Korea, Vietnam, and Operation Desert Storm. In 2004, NHRC launched the Navy-Marine Corps Combat Trauma Registry, a long-term data collection effort tracking patient care of sailors and Marines from as far forward as frontline battalion aid stations through the entire continuum of military medical care, including definitive and rehabilitative care back in the United States (Galarneau et al., 2004). Because the registry collects data on disease and non-battle injuries as well as combat casualties, it now called the CTR Expeditionary Medical Encounter Database.

continue on page 31

a few enlisted medics. Next is resuscitative care, which includes Advanced Trauma Life Support, and life- and limb-saving "damage-control" surgery, but very limited holding capability. This is followed by definitive care, large, complex field hospitals where damage-control surgeries are completed, and moderately wounded soldiers can recover in ward beds before returning to duty. The final battlefield role is en route care, which provides ground and air transportation of freshly wounded soldiers from the battlefield to an MTF, and postsurgical patients from the operating room to the next highest level of care. Military medical M&S programs must reproduce not only the treatment patients receive at each of these levels of care, but also how each individual service branch operates in those roles.

Similarly, disaster medical M&S programs need to account for the individual skill sets of civilian providers and civilian clinical levels of care. The clinical capabilities of paramedics in the field are far less than those of a physician in a hospital setting. The capabilities of a disaster field hospital will also be less than those found in a hospital ER, just as the ER capabilities will be less than those found in a trauma center. All of these factors must be taken into consideration when modeling the impact of disasters on medical care.

#### **Advantages of Modeling**

The accompanying sidebar explains in detail the extent of NHRC's military medical modeling efforts. Similar medical M&S capabilities could provide civilian planners with the capability to test hospital surge operations or hospital diversion strategies during disasters. For instance, when examining hospital surge capacity, planners typically count the number of staffed beds a hospital could open up to receive disaster victims. Planners, however, do not have good visibility on such things as critical equipment and supplies, patient movement, or provider availability (DeLia & Wood, 2008; Embrey, Clerman, Gentilman, Cecere, & Klenke, 2010). It is usually assumed, for example, that hospitals will have enough personnel to care for the patients in those extra beds. But there is evidence that this may not be the case. Many part-time hospital workers are employed by multiple hospitals or clinics, and those hospitals count on those workers to show up at their facility in the event of a disaster. Furthermore, a recent survey of hospital workers indicated up to a third would be unwilling to report for work during a severe pandemic influenza (Balicer et al., 2010). A civilian medical M&S program similar to NHRC's Tactical Medical Logistics Planning Tool (see sidebar) could allow planners to test hospital surge capacity assumptions with varying staffing levels.

Since the 9/11 attacks, several states have purchased large, deployable field hospitals for use during disasters. Often the design, equipping, and staffing of these hospitals was accomplished largely by guesswork. With the proper medical M&S capabilities, state planners could test a variety of virtual field hospital configurations to find the best fit for their needs before spending tax dollars on tentage and equipment. Supply blocks could be customized for each functional area in the field hospitals based on each state's vulnerability and hazard assessments (i.e., likelihood of hurricane, tornado, earthquake, flood). Time-phased push packages could be developed and made ready to resupply the field hospitals on a just-in-time

basis. Critical equipment and supply shortfalls can be identified, as well as personnel shortages. Logjams in patient flow can be identified and corrected before the field hospital ever deploys.

#### Conclusion

An article in a recent issue of *Military Medicine* suggested the military's expertise in medical contingency planning should be leveraged for community-based medical response planning (Embrey et al., 2010). As discussed here, military medical planners make great use of medical M&S to take the guesswork out of the process. M&S has helped those planners cope with the stresses caused by fighting two wars and a shortage of medical manpower. The success of military medical M&S is directly related to the concerted effort the military health care community has made in collecting injury and illness data, in real time, on the battlefield.

Civilian medical disaster planners could learn from military medicine's data collection and M&S efforts. To take full advantage of medical M&S, however, a rigorous long-term effort must be made to collect disaster response patient care data during the event. The collected data must include the type and number of injuries and illnesses seen by responders, including anatomical locations when appropriate and the treatment provided. Once collected, the data must be centralized so they can be coded and made available to researchers. As with military expeditionary medicine, empirically based M&S can help civilian medical disaster planners prepare for the infrequent but "predictable surprise."

#### **AUTHOR BIO**



MARTIN HILL, a research analyst with Naval Health Research Center in San Diego, California, specializes in military expeditionary medical capabilities. A 16-year veteran of the U.S. Coast Guard and U.S. Navy Reserve, he currently serves as a Medical Service Corps officer in the California State Military Reserve, where he trains Army National Guard combat medics. A National Registry Emergency Medical Technician, Hill also

served six years as a medical specialist and tactical medic with the San Diego County Sheriff's Search and Rescue Detail. In addition, he served on a federal Disaster Medical Assistance Team and with the San Diego Medical Reserve Corps, and is an American Red Cross instructor. Hill holds a Level III Certification in Homeland Security from the American Board for Certification in Homeland Security, and he is a qualified Military Emergency Management Specialist.

#### Acknowledgments

This work represents report 10-38, supported by the U.S. Department of Defense, under work unit number 60102. The views expressed in this article are those of the author and do not reflect the official policy or position of the Department of the Navy, the Department of Defense, or the U.S. Government. Approved for public release; distribution is unlimited. This research was conducted in compliance with all applicable federal regulations governing the protection of human subjects in research.

#### References

Balicer, R., Barnett, D., Thompson, C., Hsu, E., Catlett, C., Watson, C., et al. (2010). Characterizing hospital workers' willingness to report to duty in an influenza pandemic through threat- and efficacy-based assessment. *BMC Public Health*, 10(436). Retrieved July 27, 2010, from http://www.biomedcentral.com/1471-2458/10/436

Brandeau, M. L., McCoy, J. H., & Hupert, N., Holty, J-E., & Bravata, D. M. (2009). Recommendations for modeling disaster responses in public health and medicine: a position paper of the Society for Medical Decision Making. *Medical Decision Making*, 29(4), 438–460.

Closing the seams: Developing an integrated approach to health system disaster preparedness (2007). PricewaterhouseCoopers' Health Research Institute. Retrieved July 21, 2010, from http://www.pwc.com/us/en/healthcare/publications/closing-the-seams.jhtml

DeLia, D., & Wood, E. (2008). The dwindling supply of empty beds: Implications for hospital surge capacity. *Health Affairs*, 27(6), 1688–1694.

Dominici, F., Levy, J. I., & Louis, T. A. (2005). Methodological challenges and contributions in disaster epidemiology. *Epidemiologic Reviews*, 27, 9–12.

Embrey, E. P., Clerman, R., Gentilman, M. F., Cecere, F., & Klenke, W. (2010). Community-based medical disaster planning: A role for the Department of Defense and the military health system. *Military Medicine*, 175, 298–300.

Fantus, R. J., & Hammond, J. S. (2007). What a disaster [Column]. Bulletin of the American College of Surgeons, 92(6), 51–52.

Fenig, M., & Cone, D. C. (2005). Advancing disaster epidemiology and response: Developing a national disaster-victim database. *Prehospital Emergency Care*, 9(3), 457–467.

Ferenc, J. (2010). How disaster-ready are hospitals? *Materials Management in Health Care.* Retrieved July 22, 2010, from http://www.matmanmag.com/matmanmag\_app/jsp/articledisplay.jsp?dcrpath=MATMANMAG/Article/data/03MAR2010/1003MMH\_interview

Gahol, P. R. (2005). Attrition of Army Nurse Corps officers: Looking at factors that affect retention and recruitment of Army Nurse Corps officers. Unpublished master's thesis, Command and General Staff College, Fort Leavenworth, KS.

Galarneau, M. R., Hancock, W. C., Konoske, P., Melcer, T., Vickers, R. R., Walker, G. J., et al. (2004, September). *Development and preliminary findings of a combat trauma registry for the U.S. Navy-Marine Corps.* Paper presented at the Combat Casualty Care in Ground Based Tactical Situation: Trauma Technology and Emergency Medical Procedures, St. Pete Beach, FL.

Galarneau, M. R., Konoske, P. J., Emens-Hesslink, K. E., & Pang, G. (1998). Reducing the logistical footprint of forward resuscitative surgical units using a patient-driven model of clinical events (NHRC Report No. 98-1). San Diego, CA: Naval Health Research Center.

Galarneau, M. R., Konoske, P. J., & Pang, G. (1999). Projecting medical supply requirements for a highly mobile Foward Resuscitative Surgery System (NHRC Report No. 99-29). San Diego, CA: Naval Health Research Center.

Heide, E. A. d. (2006). The importance of evidence-based disaster planning. *Annals of Emergency Medicine*, 47(1), 34–49.

Hill, M., Galarneau, M., Pang, G., & Konoske, P. (2003). Marine Corps NBC warfare: Determining clinical supply requirements for treatment of battlefield casualties from chemical and biological warfare (NHRC Report No. 03-13). San Diego, CA: Naval Health Research Center.

Hill, M., Konoske, P., Pang, G., & Galarneau, M. (2006). Preparing for the worst: Using modeling to determine medical supply Requirements for terrorism response. *The Forensic Examiner*, 15(1), 24–31.

Hill, M. R., Galarneau, M., Pang, G., & Konoske, P. (2004). Marine Corps CASEVAC: Determining medical supply needs for long- and short-range airborne casualty evacuation. (NHRC Report No. 04-32). San Diego, CA: Naval Health Research Center.

Hospital emergency surge capacity: Not ready for the "predictable surprise" (2008). Washington, DC: U.S. Congress.

Hsu, S. S. (2010, April 2). National disaster exercises, called too costly and scripted, may be scaled back. *Washington Post*, p. A1.

Kimery, A. L. (2008, May 20). Crisis in mass casualty medical care – Part II. *Homeland Security Today*. McLean, VA: KMD Media, LLC.

Kimery, A. L. (2008, May 13). The crisis in trauma, ER care - Part I.

## A Short History of Navy Medical Modeling and Simulation continued from page 29

With two wars straining our military's personnel and funding, M&S has become an important tool in all fields of military planning in all branches of the service. Besides the Marine Corps, today EMedKW houses ESP medical models for both Navy surface-fleet and ground-combat assets, as well as the U.S. Air Force. Those models represent the entire battlefield continuum of care, from point of injury to definitive treatment in the USAF's Expeditionary Medical Support System field hospitals. Special Operations units are also represented, as are terrorism-response teams such as the Marine Corps' Chemical Biological Incident Response Force, which can response to terrorist attacks within the United States and abroad (Hill, Galarneau, Pang, & Konoske, 2003; Hill, Konoske, Pang, & Galarneau, 2006).

EMedKW is an all-hazards approach and models noncombat injuries, disease, field preventive medicine, and environmental injuries, as well as combat injuries. With an increased national focus on military response to disasters and humanitarian crises, NHRC researchers are currently studying patient presentations from such disasters as the 2004 tsunami, the 2005 Sumatra earthquake, and the devastating 2010 earthquake in Haiti. Analysis of those data will eventually lead to modeling tools to aid medical planners in preparing for HA/DR deployments.

Homeland Security Today. McLean, VA: KMD Media, LLC.

Kimery, A. L. (2010). Preparedness for WMD attack requires critical care. *Homeland Security Today*. McLean, VA: KMD Media, LLC.

Konoske, P., Galarneau, M., Pang, G., & Daly, T. (2010). *United States Patent No. 7,707,042*. Washington, DC: U.S. Patent and Trademark Office.

Olson, C., Bohannan, B., Peel, R., Jeschonek, R., & Leap, T. (2007). *Test and evaluation of the Tactical Medical Logistics Planning Tool (TML+)* (NHRC Report No. 07-4D). San Diego, CA: Naval Health Research Center.

Quick, G., & Hogan, D. E. (2002). Research in disaster medicine. In D. E. Hogan & J. L. Burstein (Eds.), *Disaster medicine* (pp. 395–403). Philadelphia, PA: Lippincott Williams & Wilkins.

Schrader, E. (2004). Army doctors scrambling, report says. *Los Angeles Times*. Retrieved July 22, 2010, from http://articles.latimes.com/2004/dec/09/world/fg-casualties9

U.S. trauma center crisis: Lost in the scramble for terror resources (2004). Washington, DC: National Foundation for Trauma Care.

Willams, W., Guarisco, J., Guillot, K., Wales, J., Revels, C., Barre, G., et al. (2005). Surveillance for illness and injury after Hurricane Katrina. *Morbidity and Mortality Weekly Report*, 54(40), 1018-1021. Retrieved July 26, 2010, from http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5440a4. htm#tab1

Wilper, A. P., Woolhandler, S., Lasser, K. E., McCormick, D., Cutrona, S. L., Bor, D. H., et al. (2008). Waits to see an emergency department physician: U.S. trends and predictors, 1997–2004. *Health Affairs*, 27(2), w84–w95.

Wolkin, A., & Noe, R. (2007, Dec. 18). *Disaster surveillance*. CDC Clinician Outreach and Community Activity. Atlanta, GA: Centers for Disease Control and Prevention.