

# BI2113 Ecology and Evolution

## Assignment 1

Aumshree P. Shah  
20231059

October 2024

**Q1)** For the population described by the transition matrix given below

$$\begin{pmatrix} 0.4 & 3 \\ 0.32 & 0.7 \end{pmatrix}$$

- a) What is the ratio of the total population sizes (i.e. sum of both stage classes in a generation) in two successive generations at equilibrium?
- b) At equilibrium what is the fraction of the juveniles in any given generation? This refers to the ratio of juveniles to the total population size.

### *Solution:*

a) Let's start with assuming that  $J$  is the fraction of juveniles at equilibrium and similarly  $A$  is the fraction of adults at equilibrium. At equilibrium, after any iteration, the fractions of the adults and juveniles will remain the same. This comes from Perron–Frobenius theorem, that every matrix with real positive numbers will have an eigenvalue and an eigenvector.

$$\begin{pmatrix} 0.4 & 3 \\ 0.32 & 0.7 \end{pmatrix} \begin{pmatrix} J \\ A \end{pmatrix} = r \begin{pmatrix} J \\ A \end{pmatrix}$$

Here  $r$  is the eigenvalue of the transition matrix which represents the ratio of the population size after each iteration (since the matrix multiplication is the same as multiplication by a scalar at equilibrium) and  $\begin{pmatrix} J \\ A \end{pmatrix}$  is the eigenvector of the matrix.

To find the eigenvalue and eigenvector we make use of properties of matrices, in which the eigenvalue of a  $2 \times 2$  matrix will be the roots of the quadratic equation  $A - \lambda I = 0$  where  $A$  is our given transition matrix and  $I$  is the identity matrix. From this we get:

$$\begin{aligned} & \begin{pmatrix} 0.4 - \lambda & 3 \\ 0.32 & 0.7 - \lambda \end{pmatrix} = 0 \\ \implies & (0.4 - \lambda)(0.7 - \lambda) - 3 \times 0.32 = 0 \\ \implies & 0.28 - 0.4\lambda - 0.7\lambda + \lambda^2 - 0.96 = 0 \\ \implies & \lambda^2 - 1.1\lambda - 0.68 = 0 \\ \implies & \lambda = \frac{-(-1.1) \pm \sqrt{1.1^2 + 4 \times 0.68}}{2} \quad \text{or} \quad \lambda \approx 1.541 \dots, -0.441 \dots \end{aligned}$$

### Final Eigenvalue

Since the other eigenvalue is negative (which does not biologically make sense), we choose the greater eigenvalue, which is  $r \approx 1.5412$ . Hence, the ratio of population sizes  $\left(\frac{T_{n+1}}{T_n}\right)$  of any two successive generations at equilibrium is approximately 1.54.

b) To find the ratio of juveniles, we first find the ratio of juveniles to adults to get  $\frac{J}{A}$ . We again make use of the properties of matrices in which the eigenvector of a  $2 \times 2$  matrix will solve  $(A - \lambda I)\vec{v} = 0$  where  $\vec{v}$  is the eigenvector. Hence we get:

$$\begin{pmatrix} 0.4 - \lambda & 3 \\ 0.32 & 0.7 - \lambda \end{pmatrix} \begin{pmatrix} J \\ A \end{pmatrix} = 0$$

From this, we get two linear equation as:

$$\begin{aligned} -(r - 0.4)J + 3A &= 0 \\ 0.32J + (r - 0.7)A &= 0 \end{aligned}$$

Where  $r$  is the greater eigenvalue found in part a). Solving any of the above equations we get the ratio of adults to juveniles as:

$$\frac{A}{J} = \frac{r - 0.4}{3}$$

But we need to find the ratio of juveniles to the total population, ie we need to find the ratio of  $\frac{J}{J+A}$ . we see that:

$$\begin{aligned} \frac{J}{J+A} &= \frac{\frac{J}{J}}{\frac{J+A}{J}} \\ &= \frac{1}{1 + \frac{A}{J}} \end{aligned}$$

Substituting the value of  $\frac{A}{J}$  in the above equation we get the fraction of the juveniles in any given generation as:  $\frac{1}{1 + \frac{r-0.4}{3}}$

#### Final Eigenvector

Hence we get the final ratio of juveniles to the total population  $\left(\frac{J}{J+A}\right)$ , by simplifying it to  $\approx 0.7244$

**Q2)** Assume a Lotka-Volterra competition scenario with the following parameter values:

Species 1 :  $r_1 = 1, K_1 = 200, \alpha_{12} = 0.8$ ; Species2 :  $r_2 = 0.5, K_2 = 300, \alpha_{21} = 2$

Given below are four starting points for the system. For each case, what will be the population sizes of Species 1 and Species 2 at equilibrium? For this your answer should consist of the filled table as well as the four isocline diagrams showing the trajectory of the system (not merely the vectors).


	Starting Populatoin Sizes		Equilibrium population sizes	
S. No.	Species 1	Species 2	Species 1	Species 2
1	70	175		
2	70	75		
3	30	120		
4	30	100		

Table 1


#### Solution

We first plot the isocline diagram with its vector field using python libraries where  $N_1$  is the population of species 1 and  $N_2$  is the population of species 2:

Here we make the length of the vector equal to get a proper visulization and represent the strength of those vectors by colours which have a linear change with logarithmic increase in magnitude.



Form this we see that the given competition scenario forms a feild with one unstable equilibrium point. Hence at the equilibrium the population of any one of the species will go extint and the other population will be at its maximum capacity. Now we plot the trajectory of the given four different types of starting population sizes: Here we see that in two of the cases, case-2 and case-3, where the



starting population sizes is  $[70, 75]$  and  $[30, 120]$  respectively, the species  $N_1$  goes to extint whereas in the case-2 and case-4, the  $N_1$  species two goes to extint.

Here we assume that the population follows a continious change which doesn't happen in real popula-tion but since the population number are large it doesn't affect our result that much. And from those results, we fill the table as follows:

Final population values

S. No.	Starting Population Sizes		Equilibrium population sizes	
	Species 1	Species 2	Species 1	Species 2
1	70	175	0	300
2	70	75	200	0
3	30	120	0	300
4	30	100	200	0

**Q3 a)** Suppose we have a hypothetical insect population with a cohort lifetable that looks as shown in ?? . Over time, a population of this insect will be increasing or decreasing in numbers?

Age (in days)	No. surviving	Avg. No. of eggs per capita
Egg	250	0
1	239	0
2	210	0
3	156	0.551
4	129	0.412
5	109	0.251
6	76	0.99
7	43	0.95
8	32	0.71
9	19	0.49
10	9	0.135
11	0	0

Table 2: Survival and Egg Production Data

**Solution:**

a) For this, We first find the fraction of surviving people by dividing the number of surviving people ( $l_x$ ) by 250, the number of eggs per capita ( $m_x$ ) is given, using this we calculate  $l_x m_x$  or the average number of offsprings by that particular age. Then we find the net per capita reproductive rate ( $\sum l_x m_x$ ), using the cohort lifetable as follows:

Age (days)	No. surviving	Fraction surviving ( $l_x$ )	No. eggs per capita ( $m_x$ )	$l_x m_x$
Egg	250	1.000	0	0
1	239	0.956	0	0
2	210	0.840	0	0
3	156	0.624	0.551	0.344
4	129	0.516	0.412	0.213
5	109	0.436	0.251	0.109
6	76	0.304	0.990	0.301
7	43	0.172	0.950	0.164
8	32	0.128	0.710	0.091
9	19	0.076	0.490	0.037
10	9	0.036	0.135	0.005
11	0	0.000	0	0
				$\sum l_x m_x = 1.264$

Table 3: Complete Survival and Egg Production Data

**Conclusion**

$\sum l_x m_x = 1.264$  means each approximately individual produces 1.26 offspring in his lifetime. Because the net per capita rate for the given cohort lifetable is greater than 1, the population over time will increase.

**Q3b)** Suppose there was a mutation in the insect population mentioned in Part a due to which this insect now lives two days longer with an enhanced egg-output during late life. However, the insect now starts reproducing one day later. The modified life-table is given to the right. From the perspective of long-term dynamics, is this a beneficial mutation or a harmful one?

Age (in days)	No. surviving	Avg. No. of eggs per capita
Egg	250	0
1	239	0
2	210	0
3	156	0
4	129	0.412
5	109	0.251
6	76	0.99
7	43	0.95
8	32	0.71
9	19	0.49
10	9	0.27
11	9	0.27
12	9	0.27
13	0	0

Table 4: Survival and Egg Production Data

Age (days)	No. surviving	Fraction surviving ( $l_x$ )	No. eggs per capita ( $m_x$ )	$l_x m_x$
Egg	250	1.000	0	0
1	239	0.956	0	0
2	210	0.840	0	0
3	156	0.624	0	0
4	129	0.516	0.412	0.213
5	109	0.436	0.251	0.109
6	76	0.304	0.990	0.301
7	43	0.172	0.950	0.164
8	32	0.128	0.710	0.091
9	19	0.076	0.490	0.037
10	9	0.036	0.270	0.010
11	9	0.036	0.270	0.010
12	9	0.036	0.270	0.010
13	0	0.000	0	0
				$\sum l_x m_x = 0.945$

Table 5: Survival and Egg Production Data

$\sum l_x m_x = 0.945$  means each approximately individual produces only 0.945 offspring in his lifetime. Because the net per capita rate for the given cohort lifetable in part b) is less than 1, the population over time will decrease.

### Conclusion

Hence from the perspective of long term dynamic the given mutation is a harmful one for the insect. As it not only decreases the rate of growth, but it actually makes it negative meaning after a finite amount of time the population will go extinct. The cause of the mutation giving lesser number of eggs per capita is due to the fact that the mutation skips the egg production by 1 day during which, without mutation the insect produces the most number of eggs compared to any age class, and in the latter age groups the population is already low which doesn't contribute much to the total eggs produced.

**Q4)** The government is planning to reintroduce 10 Namibian cheetahs to Madhya Pradesh's Gandhi Sagar Wildlife Sanctuary in January 2025. At the same time, they plan to introduce 30 Sangai deers from Manipur into the sanctuary as prey. No further introduction of either prey or the predator is planned. The government has decided to celebrate the introduction after a few years through a major celebration in the park inviting many world leaders. Park authorities have determined that the optimal viewing densities for the dignitaries would be 15 cheetahs and 18 sangais in the park. They have also determined that the cheetah-sangai system can be modelled as a Lotka-Volterra preypredator system

$$\frac{dN}{dt} = r_1N - CNP$$

$$\frac{dP}{dt} = -d_2P + gCNP$$

where  $N$  and  $P$  denote the prey and the predator population respectively, and the other constants have their usual meaning as per the Lotka-Volterra prey-predator model. As per data from the scientists, the yearly rates are  $r_1 = 1.05$ ,  $C = 0.1$ ,  $d_2 = 0.6$ ,  $g = 0.45$ .


Based on this information, in which year will the celebratory event happen for the first time? Assume that all the assumptions of the Lotka-Volterra continuous prey-predator model are true for this system, and one needs to have both the cheetah's and the sangai's numbers to be simultaneously as per specifications.

**Solution:**

We determine the year of the celebratory event by doing the following:

- First we plot the vector map of the Prey-Predator dynamics.
- Then we run the system for with the initial population of  $[30, 10]$ .
- We then see the time evolution of the system and determine the time period using it.

Plotting the vector field with time evolution we get: We run the population over the timescale of 30




years, we observe that the trajectory of the population oscillates constantly. This implies that the population never reaches the required number and hence the celebratory event never occurs.

But here we have assumed that the population which is given by the dynamics is continuous, which is **not the case in real life**. If we assume that the population at any given point is given by  $(a, b)$  then the population only has 4 ways in which it can change which are:  $(a+1, b)$ ,  $(a, b+1)$ ,  $(a-1, b)$ ,  $(a, b-1)$ . After the change in population it will have a different change in vector which will drive it to a different point.

Now how do we determine in which of the 4 directions it goes to? Well a simple way to determine that will be to see which direction the vector field favours the most. So we start at  $(30, 10)$  and see the time evolution of the system by iterating over steps rather than time.

Plotting the evolution of the system using the discrete step method mentioned above we get:



Here we see that the population does infact reach the desired value after **seventeen** iterations. But we arrive at a problem here. We need to determine the time taken to reach the required point but each iteration will have diffrenet time, if the vector aligns to x or y axis and has high magnitude, then the population will increase rapidly. Whereas if the population has a small magnitude and is aligned halfway between the axis then the time taken for that iteration would be larger. For this we complicate the code used which is given after remarks which makes use of the method mentioned above to get the time period as:

## Final Conclusion

REMARKS



this is 1

table refarence error

add marks

Let's start with visualizing the terms of the matix. Here  $a_{11} = 0.4$  represents the fraction of juveniles which remains juveniles after one iteration,  $a_{12} = 0.3$  represents the per capita child birth,  $a_{21} = 0.32$  represents the fraction of juveniles which become adult after an iteration and finally  $a_{22}$  represents the fraction of adults which survive each iteration.

check 1

box aj in q1b

check 2

q3 table shrink

assumptions?

non steps in q2

steps in q3

code for the polots in github too

red dot in q4b

refrecne code, link remarks in q4