

Title

A Senior Comprehensive Project

by

name

Allegheny College

Meadville, PA

October 20, 2016

Submitted to the Department of Mathematics in partial fulfillment of the
requirements for the degree of Bachelor of Science.

Project Advisor: Dr. Rachel J. Weir

Second Reader: [Enter name here.]

I hereby recognize and pledge to fulfill my responsibilities, as defined in the
Honor Code, and to maintain the integrity of both myself and the College
community as a whole.

Pledge:

name

Acknowledgements

I want to thank everybody for everything. Keep typing to see what happens with a new paragraph.

Start a new paragraph thanking people.

Abstract

This comp is about everything!

Contents

1	Introduction	1
2	Preliminaries	2
3	Chaos	3
4	L_p-spaces	4
5	Main results	5
	References	11

1 Introduction

This purpose of this comp is to show that I've learned something really wonderful.

As a bonus, when I'm done and have passed the oral exam, I will finally graduate!

2 Preliminaries

a

3 Chaos

The following is a well known criterion for chaos, known as the Eigenvalue Criterion. [2,3] provide proofs for the Criterion, and [9,10,12,14,15] provide examples using the Criterion.

Theorem 3.1. *Let $T : X \rightarrow X$ be an operator on a separable complex Banach space X . Consider the subspaces*

$$X_0 := \text{Span}\{x \in X : T(x) = \lambda x \text{ for some } \lambda \in \mathbb{C} \text{ with } |\lambda| < 1\},$$

$$Y_0 := \text{Span}\{x \in X : T(x) = \lambda x \text{ for some } \lambda \in \mathbb{C} \text{ with } |\lambda| > 1\},$$

$$Z_0 := \text{Span}\{x \in X : T(x) = e^{i\alpha} x \text{ for some } \alpha \in \mathbb{Q}\}.$$

If X_0, Y_0 , and Z_0 are all dense in X , then T is chaotic.

Since the set of eigenvalues $\sigma_p(B) = \mathbb{D}$ in our framework, this Criterion says that $\varphi(B)$ is chaotic on l^p if and only if $\varphi(\mathbb{D})$ intersects the unit circle.

Exercise 3.2. sldkfjlskdjldksjf

Theorem 3.3. as;ldkfjlasdkjfl

Proof. alsdkjflaskdjf

□

4 Lp-spaces

Exercise 4.1. sldkfjlskdfjldksjf

5 Main results

The following result gives a geometrical description of $\varphi(\mathbb{D})$ when the pole of φ lies outside the closed unit disc.

Lemma 5.1. *Let φ be a linear fractional transformation (LFT) with $c \neq 0$ and $|d| > |c|$. Then $\varphi(\mathbb{D})$ is the disc $P + r\mathbb{D}$ with center P and radius r given by*

$$P = \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2}, \quad r = \frac{|bc - ad|}{|d|^2 - |c|^2}.$$

Proof. Note that LFTs map circles and lines to circles and lines. Indeed, if f is a LFT and E is a circle or a line in \mathbb{C} , the image of E , $f(E)$, is mapped to a line if it passes through the pole. If E avoids the pole, $f(E)$ is a circle.

Observe that $\bar{\mathbb{D}} = \{z : |z| \leq 1\}$ (i.e. the closure of \mathbb{D}) is clearly a bounded and convex set. Because we imposed that $|d| > |c|$, we have that $|d/c| > 1$ and so the pole at $z = -d/c$ lies outside of $\bar{\mathbb{D}}$. Since LFTs are conformal at every point except at the pole, $\varphi(\bar{\mathbb{D}})$ must be bounded and convex. Furthermore, $\varphi(\bar{\mathbb{D}})$ is a circle whose boundary is $\varphi(\partial\mathbb{D})$, where $\partial\mathbb{D}$ denotes the boundary of \mathbb{D} .

Now, take three distinct points in the unit circle. We choose $z_1 = 1$, $z_2 = -1$, and $z_3 = i$ as they are three very simple points on the unit circle. Since φ is linear, it is also one-to-one. Thus, $A = f(z_1)$, $B = f(z_2)$, and $C = f(z_3)$ are three distinct points. Since z_1 , z_2 , and z_3 are on the unit circle which is equivalent to $\partial\mathbb{D}$, A , B , and C are in fact three distinct points in the circle $\varphi(\partial\mathbb{D})$. That is, circle circumscribed over A , B , and C coincides with $\varphi(\partial\mathbb{D})$.

To verify that $\varphi(\partial\mathbb{D})$ indeed has center P and radius r , we just need to show that

$$|A - P| = |B - P| = |C - P| = r.$$

For this, we use the equalities $|z|^2 = z\bar{z}$, and $|c + d| = |\bar{c} + \bar{d}|$.

So, we have

$$\begin{aligned}
|A - P| &= \left| \frac{a + b}{c + d} - \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(a + b)(|d|^2 - |c|^2) - (b\bar{d} - a\bar{c})(c + d)}{c + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{add\bar{d} - bc\bar{c} - bc\bar{d} + ad\bar{c}}{c + d} \right|, \text{ by using the first equality} \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(\bar{c} + \bar{d})(bc - ad)}{c + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|\bar{c} + \bar{d}||bc - ad|}{|c + d|} \right) \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|c + d||bc - ad|}{|c + d|} \right), \text{ by using the second equality} \\
&= \frac{bc - ad}{|d|^2 - |c|^2} \\
&= r.
\end{aligned}$$

Showing $|B - P| = r$ is analagous:

$$\begin{aligned}
|B - P| &= \left| \frac{-a + b}{-c + d} - \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(-a + b)(|d|^2 - |c|^2) - (b\bar{d} - a\bar{c})(-c + d)}{-c + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{-add\bar{d} - bc\bar{c} + bcd\bar{d} + ad\bar{c}}{-c + d} \right|, \text{ by using the first equality} \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(-\bar{c} + \bar{d})(bc - ad)}{-c + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|-\bar{c} + \bar{d}||bc - ad|}{| -c + d|} \right) \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|-c + d||bc - ad|}{| -c + d|} \right), \text{ by using the second equality} \\
&= \frac{bc - ad}{|d|^2 - |c|^2} \\
&= r.
\end{aligned}$$

Using a third equality, $|ci + d| = |\bar{c} + \bar{d}i|$, we have

$$\begin{aligned}
|C - P| &= \left| \frac{ai + b}{ci + d} - \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(ai + b)(|d|^2 - |c|^2) - (b\bar{d} - a\bar{c})(ci + d)}{ci + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{add\bar{i} - bc\bar{c} - bc\bar{d}i + ad\bar{c}}{ci + d} \right|, \text{ by using the first equality} \\
&= \frac{1}{|d|^2 - |c|^2} \left| \frac{(\bar{c} + \bar{d}i)(ad - bc)}{ci + d} \right| \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|\bar{c} + \bar{d}i||ad - bc|}{|ci + d|} \right) \\
&= \frac{1}{|d|^2 - |c|^2} \left(\frac{|ci + d||ad - bc|}{|ci + d|} \right), \text{ by using the third equality} \\
&= \frac{bc - ad}{|d|^2 - |c|^2} \\
&= r.
\end{aligned}$$

Hence the circle circumscribed over the points A , B , and C indeed has center P and radius r . Thus $\varphi(\partial\mathbb{D})$ has center P and radius r . Thus $\varphi(\mathbb{D})$ is the disc $P + r\mathbb{D}$. \square

In [10], DeLaubenfels and Emamirad showed that, for a non-constant polynomial $P(z)$, $P(B)$ (where B is the backwards shift operator) is chaotic on l^p , $1 \leq p \leq \infty$ whenever $P(\mathbb{D})$ intersects the unit disc. We provide a generalization of this result which can be applied to Linear Fractional Transformations.

Theorem 5.2. *Let φ be a LFT with $c \neq 0$ and $|d| > |c|$. The operator $\varphi(B)$*

is chaotic if and only if

$$||d|^2 - |c|^2 - |b\bar{d} - a\bar{c}| < |bc - ad|.$$

Proof. We showed in Lemma 2 that $\varphi(\mathbb{D}) = P + r\mathbb{D}$ with center P and radius r given where

$$P = \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2}, \quad r = \frac{|bc - ad|}{|d|^2 - |c|^2}.$$

Theorem 1, the Eigenvalue Criterion, showed that $\varphi(B)$ is chaotic on l^p if and only if $\varphi(\mathbb{D})$ intersects the unit circle. So, we have that $\varphi(B)$ is chaotic if and only if the disc $P + r\mathbb{D}$ intersects the unit circle.

In order for the disc to intersect the unit disc, we have two possibilities: the center of the disc P is contained within the unit circle, or P is outside the closed unit disc. If P is inside the unit disc, then $|P| + |r| > 1$; if P is outside the closed unit disc, then we must have $|P| - |r| < 1$.

These conditions lead to

$$-|r| < 1 - |P| < |r|.$$

After substituting in the values of P and r , we have

$$-\frac{|bc - ad|}{|d|^2 - |c|^2} < 1 - \left| \frac{b\bar{d} - a\bar{c}}{|d|^2 - |c|^2} \right| < \frac{|bc - ad|}{|d|^2 - |c|^2}.$$

Multiplying by $|d|^2 - |c|^2$ gives

$$-|bc - ad| < |d|^2 - |c|^2 - |b\bar{d} - a\bar{c}| < |bc - ad|.$$

So finally, we have

$$||d|^2 - |c|^2 - |b\bar{d} - a\bar{c}|| < |bc - ad|.$$

So, we have shown that if $\varphi(B)$ is chaotic, then $||d|^2 - |c|^2 - |b\bar{d} - a\bar{c}|| < |bc - ad|$. The other direction is completely analogous. It requires the exact same algebra, done in reverse, to show that if $||d|^2 - |c|^2 - |b\bar{d} - a\bar{c}|| < |bc - ad|$, then $P + r\mathbb{D}$ intersects the unit circle, and thus $\varphi(B)$ is chaotic. \square

References

- [1] Burton, David M., *Elementary Number Theory*, Second Ed., W.C. Brown Publishers, 1989.
- [2] Duren, P., Khavinson, D., Shapiro, H.S., and Sundberg, C., Contractive zero-divisors in Bergman spaces, *Pacific J. Math*, **157** (1993), 37-56.
- [3] Fraleigh, John B., *A First Course in Abstract Algebra*, Fifth Ed., Addison-Wesley, 1994.