

# Modern Bayesian Inference

## Theory and Practice

Will Handley  
wh260@cam.ac.uk

Astrophysics Group  
Cavendish Laboratory  
University of Cambridge

July 4, 2017

# Introduction

# Introduction

- ▶ Statistics  $\equiv$  Inference

# Introduction

- ▶ Statistics  $\equiv$  Inference
- ▶ How to extract information about scientific models from data.

# Introduction

- ▶ Statistics  $\equiv$  Inference
- ▶ How to extract information about scientific models from data.
- ▶ Most cosmologists work in a *Bayesian* framework of inference, although *Frequentist* methods are also sometimes used.

# Introduction

- ▶ Statistics  $\equiv$  Inference
- ▶ How to extract information about scientific models from data.
- ▶ Most cosmologists work in a *Bayesian* framework of inference, although *Frequentist* methods are also sometimes used.
- ▶ **Bayesians use Probability Distributions to quantify uncertainty.**

# Probability distributions

# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.



# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.
- ▶ Age of the universe (*Planck*):

# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.
- ▶ Age of the universe (*Planck*):

$13.73 \pm 0.12$  billion years old.

# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.
- ▶ Age of the universe (*Planck*):

$13.73 \pm 0.12$  billion years old.

- ▶ Masses of LIGO GW150914 binary merger:

$$m_1 = 36.4_{-4.9}^{+5.5} M_{\odot}, \quad m_2 = 30.9_{-4.4}^{+4.8} M_{\odot}$$

# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.
- ▶ Age of the universe (*Planck*):

13.73  $\pm$  0.12 billion years old.

- ▶ Masses of LIGO GW150914 binary merger:

$$m_1 = 36.4^{+5.5}_{-4.9} M_{\odot}, \quad m_2 = 30.9^{+4.8}_{-4.4} M_{\odot}$$

- ▶ These are called *credible intervals*, state that we are e.g. 90% confident of the value lying in this range.

# Probability distributions

- ▶ As scientists, we are used to seeing error bars on results.
- ▶ Age of the universe (*Planck*):


13.73  $\pm$  0.12 billion years old.

- ▶ Masses of LIGO GW150914 binary merger:

$$m_1 = 36.4^{+5.5}_{-4.9} M_{\odot}, \quad m_2 = 30.9^{+4.8}_{-4.4} M_{\odot}$$

- ▶ These are called *credible intervals*, state that we are e.g. 90% confident of the value lying in this range.
- ▶ More importantly, these are *summary statistics*.


# LIGO binary merger



$$m_1 = 36.4^{+5.5}_{-4.9} M_{\odot},$$

$$m_2 = 30.9^{+4.8}_{-4.4} M_{\odot}$$

# LIGO binary merger




$$m_1 = 36.4^{+5.5}_{-4.9} M_\odot,$$

$$m_2 = 30.9^{+4.8}_{-4.4} M_\odot$$

- Summary statistics summarise a full probability distribution.

# LIGO binary merger



$$m_1 = 36.4^{+5.5}_{-4.9} M_\odot,$$

$$m_2 = 30.9^{+4.8}_{-4.4} M_\odot$$

- Summary statistics summarise a full probability distribution.
- One goal of inference is to produce these probability distributions.




## Extended example of inference: LIGO

- ▶ We will introduce the key concepts by discussing an extended example of the inference process.


# Theory

Extended example of inference: LIGO



# The model $M$

Extended example of inference: LIGO



# The parameters $\Theta$ of the model $M$

Extended example of inference: LIGO

Theoretical signal depends on:

# The parameters $\Theta$ of the model $M$

Extended example of inference: LIGO

Theoretical signal depends on:

- ▶  $m_1, m_2$ : mass of binary

# The parameters $\Theta$ of the model $M$

Extended example of inference: LIGO

Theoretical signal depends on:

- ▶  $m_1, m_2$ : mass of binary
- ▶  $\theta, \phi$ : sky location

# The parameters $\Theta$ of the model $M$

Extended example of inference: LIGO

Theoretical signal depends on:

- ▶  $m_1, m_2$ : mass of binary
- ▶  $\theta, \phi$ : sky location
- ▶  $r$ : luminosity distance

# The parameters $\Theta$ of the model $M$

Extended example of inference: LIGO

Theoretical signal depends on:

- ▶  $m_1, m_2$ : mass of binary
- ▶  $\theta, \phi$ : sky location
- ▶  $r$ : luminosity distance
- ▶  $\Phi_c, t_c$ : phase and time of coalescence



# The parameters $\Theta$ of the model $M$


Extended example of inference: LIGO

Theoretical signal depends on:

- ▶  $m_1, m_2$ : mass of binary
- ▶  $\theta, \phi$ : sky location
- ▶  $r$ : luminosity distance
- ▶  $\Phi_c, t_c$ : phase and time of coalescence
- ▶  $i, \theta_{\text{sky}}$ : inclination and angle on sky (orbital parameters)


# The data $D$

Extended example of inference: LIGO




# The data $D$

Extended example of inference: LIGO




# The Likelihood: well matched

Extended example of inference: LIGO




# The Likelihood: coalescence off

Extended example of inference: LIGO




# The Likelihood: too large luminosity distance

Extended example of inference: LIGO




# The Likelihood: incorrect inclination

Extended example of inference: LIGO



# The Likelihood: 'Correct parameters'

Extended example of inference: LIGO





## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed
  - ▶  $h(t; \Theta)$ : Theoretical strain

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed
  - ▶  $h(t; \Theta)$ : Theoretical strain

$$P(h_i|\Theta, M) = \frac{1}{\sqrt{2\pi}\sigma_i} \exp\left(-\frac{[h_i - h(t_i; \Theta)]^2}{2\sigma_i^2}\right)$$

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed
  - ▶  $h(t; \Theta)$ : Theoretical strain

$$P(D|\Theta, M) = \prod_i \frac{1}{\sqrt{2\pi}\sigma_i} \exp\left(-\frac{[h_i - h(t_i; \Theta)]^2}{2\sigma_i^2}\right)$$

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed
  - ▶  $h(t; \Theta)$ : Theoretical strain

$$L(\Theta) = P(D|\Theta, M) = \prod_i \frac{1}{\sqrt{2\pi}\sigma_i} \exp\left(-\frac{[h_i - h(t_i; \Theta)]^2}{2\sigma_i^2}\right)$$

## The likelihood $L(\Theta)$

- ▶ We assume that observed signal is combination of theoretical signal and Gaussian noise  $O = S + N$ .
- ▶ Elementary probability tells us:
  - ▶  $(t_i, h_i \pm \sigma_i)$ : strain observed
  - ▶  $h(t; \Theta)$ : Theoretical strain

$$L(\Theta) = P(D|\Theta, M) = \prod_i \frac{1}{\sqrt{2\pi}\sigma_i} \exp\left(-\frac{[h_i - h(t_i; \Theta)]^2}{2\sigma_i^2}\right)$$

- ▶ We normally work with log-likelihoods, which turn  $\prod \rightarrow \sum$ .



# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood

# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood  $\equiv$  Probability of data, given model parameters:

# Bayes' Theorem

Extended example of inference: LIGO

- Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- ▶ What we want:
- ▶ *Posterior*

# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- ▶ What we want:
- ▶ *Posterior*  $\equiv$  Probability of model parameters, given data:

# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- ▶ What we want:
- ▶ *Posterior*  $\equiv$  Probability of model parameters, given data:

$$\mathcal{P}(\Theta) \equiv P(\Theta|D, M)$$

# Bayes' Theorem

Extended example of inference: LIGO

- ▶ Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- ▶ What we want:
- ▶ *Posterior*  $\equiv$  Probability of model parameters, given data:

$$\mathcal{P}(\Theta) \equiv P(\Theta|D, M)$$

- ▶ Use Bayes theorem:

# Bayes' Theorem

Extended example of inference: LIGO

- Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- What we want:
- *Posterior*  $\equiv$  Probability of model parameters, given data:

$$\mathcal{P}(\Theta) \equiv P(\Theta|D, M)$$

- Use Bayes theorem:

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$



# Bayes' Theorem

Extended example of inference: LIGO

- Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- What we want:
- *Posterior*  $\equiv$  Probability of model parameters, given data:

$$\mathcal{P}(\Theta) \equiv P(\Theta|D, M)$$

- Use Bayes theorem:

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$

$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

# Bayes' Theorem

Extended example of inference: LIGO

- Likelihood  $\equiv$  Probability of data, given model parameters:

$$L(\Theta) \equiv P(D|\Theta, M)$$

- What we want:
- *Posterior*  $\equiv$  Probability of model parameters, given data:

$$\mathcal{P}(\Theta) \equiv P(\Theta|D, M)$$

- Use Bayes theorem:

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$

$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

$$\text{Posterior} = \frac{\text{Likelihood} \times \text{Prior}}{\text{Evidence}}$$

# Prior $\pi(\Theta)$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$

$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

# Prior $\pi(\Theta)$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

► Prior:  $\pi(\Theta) \equiv P(\Theta|M)$ .

# Prior $\pi(\Theta)$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Prior:  $\pi(\Theta) \equiv P(\Theta|M)$ .
- ▶ Represents initial degree of knowledge about the system:

# Prior $\pi(\Theta)$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Prior:  $\pi(\Theta) \equiv P(\Theta|M)$ .
- ▶ Represents initial degree of knowledge about the system:
- ▶ e.g.  $\pi(\theta, \phi) d\theta d\phi = -\frac{1}{2} \sin \theta d\theta d\phi$

# Prior $\pi(\Theta)$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Prior:  $\pi(\Theta) \equiv P(\Theta|M)$ .
- ▶ Represents initial degree of knowledge about the system:
- ▶ e.g.  $\pi(\theta, \phi) d\theta d\phi = -\frac{1}{2} \sin \theta d\theta d\phi$
- ▶ Most Bayesian approaches are sensitive to this, and rightly so.

# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

► Evidence:  $Z \equiv P(D|M)$



# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

► Evidence:  $Z \equiv P(D|M) = \int P(D|\Theta, M)P(\Theta|M)d\Theta$ .

# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

► Evidence:  $Z \equiv P(D|M) = \int L(\Theta)\pi(\Theta)d\Theta$ .

# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Evidence:  $Z \equiv P(D|M) = \int L(\Theta)\pi(\Theta)d\Theta$ .
- ▶ Normalising constant.

# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Evidence:  $Z \equiv P(D|M) = \int L(\Theta)\pi(\Theta)d\Theta$ .
- ▶ Normalising constant.
- ▶ Difficult to compute.

# Evidence $Z$

Extended example of inference: LIGO

$$P(\Theta|D, M) = \frac{P(D|\Theta, M)P(\Theta|M)}{P(D|M)}$$
$$\mathcal{P}(\Theta) = \frac{L(\Theta)\pi(\Theta)}{Z}$$

- ▶ Evidence:  $Z \equiv P(D|M) = \int L(\Theta)\pi(\Theta)d\Theta$ .
- ▶ Normalising constant.
- ▶ Difficult to compute.
- ▶ Still extremely important.

# Posterior $\mathcal{P}$

Extended example of inference: LIGO

- ▶ Cannot plot the full posterior distribution:


$$\mathcal{P}(\Theta) \equiv P(m_1, m_2, \theta, \phi, r, \Phi_c, t_c, i, \theta_{\text{sky}} | D, M)$$

- ▶ Can plot 1D and 2D *marginalised* distributions e.g:

$$P(m_1, m_2 | D, M) = \int P(m_1, m_2, \theta, \phi, r, \Phi_c, t_c, i, \theta_{\text{sky}} | D, M) d\theta d\phi dr d\Phi_c dt_c di d\theta_{\text{sky}}$$


# Posterior $\mathcal{P}$

Extended example of inference: LIGO



# Posterior $\mathcal{P}$

Extended example of inference: LIGO




► May do this for each pair of parameters



# Posterior $\mathcal{P}$


Extended example of inference: LIGO



- ▶ May do this for each pair of parameters
- ▶ Generates a *triangle plot*


# Posterior $\mathcal{P}$

Extended example of inference: LIGO



# Posterior $\mathcal{P}$


## Extended example of inference: LIGO



► Does give insight

# Posterior $\mathcal{P}$

## Extended example of inference: LIGO



- Does give insight
- Not the full picture

# Evidences and model comparison

Extended example of inference: LIGO

# Evidences and model comparison

Extended example of inference: LIGO

- ▶ Up until now, we have discussed *Parameter estimation*:  
inferring what data tell us about parameters  $\Theta$  of a model  $M$ .

# Evidences and model comparison

Extended example of inference: LIGO

- ▶ Up until now, we have discussed *Parameter estimation*:  
inferring what data tell us about parameters  $\Theta$  of a model  $M$ .
- ▶ Scientifically speaking, this is only half the story.

# Evidences and model comparison

Extended example of inference: LIGO

- ▶ Up until now, we have discussed *Parameter estimation*:  
inferring what data tell us about parameters  $\Theta$  of a model  $M$ .
- ▶ Scientifically speaking, this is only half the story.
- ▶ In general, we will have several competing models that describe the data, and we want to know which is the “best”.



## Bayes' Theorem (again)

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

$$P(M_i|D) = \frac{Z_i \mu_i}{\sum_k Z_k \mu_k}$$

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

$$P(M_i|D) = \frac{Z_i \mu_i}{\sum_k Z_k \mu_k}$$

- Evidences allow us to determine what weighting we should give to each model in light of the data.

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

$$P(M_i|D) = \frac{Z_i \mu_i}{\sum_k Z_k \mu_k}$$

- ▶ Evidences allow us to determine what weighting we should give to each model in light of the data.
- ▶ e.g. EOBNR (Effective One Body) vs IMRPhenom (Inspiral-merger-ringdown).

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

$$P(M_i|D) = \frac{Z_i \mu_i}{\sum_k Z_k \mu_k}$$

- ▶ Evidences allow us to determine what weighting we should give to each model in light of the data.
- ▶ e.g. EOBNR (Effective One Body) vs IMRPhenom (Inspiral-merger-ringdown).
- ▶ Or testing modified theories of gravity.

## Bayes' Theorem (again)

What does data tell us about our model  $M_i$ ?

$$P(M_i|D) = \frac{P(D|M_i)P(M_i)}{P(D)}$$

$$P(M_i|D) = \frac{Z_i \mu_i}{\sum_k Z_k \mu_k}$$

- ▶ Evidences allow us to determine what weighting we should give to each model in light of the data.
- ▶ e.g. EOBNR (Effective One Body) vs IMRPhenom (Inspiral-merger-ringdown).
- ▶ Or testing modified theories of gravity.

### Model averaging:

- ▶ Multiple models with posterior on the same parameter:

$$P(y|M_i, D)$$

$$P(y|D) = \sum_i P(y|M_i, D)P(M_i|D)$$



# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell\}$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell\}$$

$$M = \Lambda\text{CDM}$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell\}$$

$$M = \Lambda\text{CDM}$$

$$\Theta = \Theta_{\Lambda\text{CDM}}$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell\}$$

$$M = \Lambda\text{CDM}$$

$$\Theta = \Theta_{\Lambda\text{CDM}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM}$$

$$\Theta = \Theta_{\Lambda\text{CDM}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$



# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}}, w)$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{CIB}, \xi^{tSZ-CIB}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}}, w, \Sigma m_\nu, m_{\nu, \text{sterile}}^{\text{eff}})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\} + \{\text{LSS}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{\text{CIB}}, \xi^{tSZ-\text{CIB}}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}}, w, \Sigma m_\nu, m_{\nu, \text{sterile}}^{\text{eff}})$$

# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\} + \{\text{LSS}\} + \{\text{"Big Data"}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{\text{CIB}}, \xi^{tSZ-\text{CIB}}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}}, w, \Sigma m_\nu, m_{\nu, \text{sterile}}^{\text{eff}})$$



# Parameter estimation

Another example.

$$\mathcal{L}(\Theta) = P(D|\Theta, M)$$

$$D = \{C_\ell^{(\text{Planck})}\} + \{\text{LSS}\} + \{\text{"Big Data"}\}$$

$$M = \Lambda\text{CDM} + \text{extensions}$$

$$\Theta = \Theta_{\Lambda\text{CDM}} + \Theta_{\text{Planck}} + \Theta_{\text{extensions}}$$

$$\Theta_{\Lambda\text{CDM}} = (\Omega_b h^2, \Omega_c h^2, 100\theta_{MC}, \tau, \ln(10^{10} A_s), n_s)$$

$$\Theta_{\text{Planck}} = (y_{\text{cal}}, A_{217}^{\text{CIB}}, \xi^{tSZ-\text{CIB}}, A_{143}^{tSZ}, A_{100}^{PS}, A_{143}^{PS}, A_{143 \times 217}^{PS}, A_{217}^{PS}, \\ A_{100}^{kSZ}, A_{100}^{\text{dust } TT}, A_{143}^{\text{dust } TT}, A_{143 \times 217}^{\text{dust } TT}, A_{217}^{\text{dust } TT}, c_{100}, c_{217})$$

$$\Theta_{\text{extensions}} = (n_{\text{run}}, n_{\text{run,run}}, w, \Sigma m_\nu, m_{\nu, \text{sterile}}^{\text{eff}})$$


- ▶ Parameter estimation:  $L, \pi \rightarrow \mathcal{P}$ : model parameters
- ▶ Model comparison:  $L, \pi \rightarrow Z$ : how good model is

# Sampling

How to describe a high-dimensional posterior

# Sampling


How to describe a high-dimensional posterior



# Sampling

How to describe a high-dimensional posterior


- In high dimensions, posterior  $\mathcal{P}$  occupies a vanishingly small region of the prior  $\pi$ .






# Sampling

How to describe a high-dimensional posterior




- ▶ In high dimensions, posterior  $\mathcal{P}$  occupies a vanishingly small region of the prior  $\pi$ .
- ▶ *Sampling* the posterior is an excellent compression scheme.






# Sampling



# Sampling






# Sampling





# Sampling



# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$

# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{1}{N} \sum_{i=1}^N m_1^{(i)}$$

# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{1}{N} \sum_{i=1}^N m_1^{(i)}$$

- ▶ Mass covariance:

$$\text{Cov}(m_1, m_2) \approx \frac{1}{N} \sum_{i=1}^N (m_1^{(i)} - \bar{m}_1)(m_2^{(i)} - \bar{m}_2)$$

# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{1}{N} \sum_{i=1}^N m_1^{(i)}$$

- ▶ Mass covariance:

$$\text{Cov}(m_1, m_2) \approx \frac{1}{N} \sum_{i=1}^N (m_1^{(i)} - \bar{m}_1)(m_2^{(i)} - \bar{m}_2)$$

- ▶ Marginalised samples: Just ignore the other coordinates.

# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{1}{N} \sum_{i=1}^N m_1^{(i)}$$

- ▶ Mass covariance:

$$\text{Cov}(m_1, m_2) \approx \frac{1}{N} \sum_{i=1}^N (m_1^{(i)} - \bar{m}_1)(m_2^{(i)} - \bar{m}_2)$$

- ▶ Marginalised samples: Just ignore the other coordinates.
- ▶ N.B. Typically have *weighted* samples

# Why do sampling?

Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{1}{N} \sum_{i=1}^N m_1^{(i)}$$

- ▶ Mass covariance:

$$\text{Cov}(m_1, m_2) \approx \frac{1}{N} \sum_{i=1}^N (m_1^{(i)} - \bar{m}_1)(m_2^{(i)} - \bar{m}_2)$$

- ▶ Marginalised samples: Just ignore the other coordinates.
- ▶ N.B. Typically have *weighted* samples

# Why do sampling?

## Marginalisation over the posterior

- ▶ Set of  $N$  samples  $S = \{\Theta^{(i)} : i = 1, \dots, N : \Theta^{(i)} \sim \mathcal{P}\}$
- ▶ Mean mass:

$$\bar{m}_1 \equiv \langle m_1 \rangle_{\mathcal{P}} \approx \frac{\sum_{i=1}^N w^{(i)} m_1^{(i)}}{\sum_{i=1}^N w^{(i)}}$$

- ▶ Mass covariance:

$$\text{Cov}(m_1, m_2) \approx \frac{\sum_{i=1}^N (m_1^{(i)} - \bar{m}_1)(m_2^{(i)} - \bar{m}_2)}{\sum_{i=1}^N w^{(i)}}$$

- ▶ Marginalised samples: Just ignore the other coordinates.
- ▶ N.B. Typically have *weighted* samples



# Parameter estimation

# Parameter estimation

- ▶ The name of the game is therefore drawing samples  $S$  from the posterior  $\mathcal{P}$  with the minimum number of likelihood calls.

# Parameter estimation

- ▶ The name of the game is therefore drawing samples  $S$  from the posterior  $\mathcal{P}$  with the minimum number of likelihood calls.
- ▶ Gridding is doomed to failure in high dimensions.

# Parameter estimation

- ▶ The name of the game is therefore drawing samples  $S$  from the posterior  $\mathcal{P}$  with the minimum number of likelihood calls.
- ▶ Gridding is doomed to failure in high dimensions.
- ▶ Enter Metropolis Hastings.

## Current sampling approaches

# Current sampling approaches

1. Metropolis Hastings.

# Current sampling approaches

1. Metropolis Hastings.
2. Hamiltonian Monte-Carlo (HMC).

# Current sampling approaches

1. Metropolis Hastings.
2. Hamiltonian Monte-Carlo (HMC).
3. Ensemble sampling (e.g. emcee).



# Metropolis Hastings

# Metropolis Hastings

- ▶ Turn the  $N$ -dimensional problem into a one-dimensional one.

# Metropolis Hastings

- ▶ Turn the  $N$ -dimensional problem into a one-dimensional one.
  1. Pick random direction

# Metropolis Hastings

- ▶ Turn the  $N$ -dimensional problem into a one-dimensional one.
  1. Pick random direction
  2. Choose step length


# Metropolis Hastings

- ▶ Turn the  $N$ -dimensional problem into a one-dimensional one.
  1. Pick random direction
  2. Choose step length
  3. If uphill, make step. . .

# Metropolis Hastings

- ▶ Turn the  $N$ -dimensional problem into a one-dimensional one.
  1. Pick random direction
  2. Choose step length
  3. If uphill, make step. . .
  4. . . otherwise sometimes make step.


# Metropolis Hastings



# Metropolis Hastings



# Metropolis Hastings



# Metropolis Hastings

Struggles with...

# Metropolis Hastings

Struggles with...

1. Burn in

# Metropolis Hastings

Struggles with...

1. Burn in
2. Multimodality

# Metropolis Hastings

Struggles with...

1. Burn in
2. Multimodality
3. Correlated Peaks

# Metropolis Hastings

Struggles with...

1. Burn in
2. Multimodality
3. Correlated Peaks
4. Phase transitions

# Hamiltonian Monte-Carlo

# Hamiltonian Monte-Carlo

- ▶ Key idea: Treat  $\log L(\Theta)$  as a potential energy



# Hamiltonian Monte-Carlo

- ▶ Key idea: Treat  $\log L(\Theta)$  as a potential energy
- ▶ Guide walker under “force”:

$$F(\Theta) = \nabla \log L(\Theta)$$

# Hamiltonian Monte-Carlo

- ▶ Key idea: Treat  $\log L(\Theta)$  as a potential energy
- ▶ Guide walker under “force”:

$$F(\Theta) = \nabla \log L(\Theta)$$

- ▶ Walker is naturally “guided” uphill

# Hamiltonian Monte-Carlo

- ▶ Key idea: Treat  $\log L(\Theta)$  as a potential energy
- ▶ Guide walker under “force”:

$$F(\Theta) = \nabla \log L(\Theta)$$

- ▶ Walker is naturally “guided” uphill
- ▶ Conserved quantities mean efficient acceptance ratios.

# Hamiltonian Monte-Carlo

- ▶ Key idea: Treat  $\log L(\Theta)$  as a potential energy
- ▶ Guide walker under “force”:

$$F(\Theta) = \nabla \log L(\Theta)$$

- ▶ Walker is naturally “guided” uphill
- ▶ Conserved quantities mean efficient acceptance ratios.
- ▶ stan is a fully fledged, rapidly developing programming language with HMC as a default sampler.

# Ensemble sampling

# Ensemble sampling

- ▶ Instead of one walker, evolve a set of  $n$  walkers.

# Ensemble sampling

- ▶ Instead of one walker, evolve a set of  $n$  walkers.
- ▶ Can use information present in ensemble to guide proposals.

# Ensemble sampling

- ▶ Instead of one walker, evolve a set of  $n$  walkers.
- ▶ Can use information present in ensemble to guide proposals.
- ▶ emcee: affine invariant proposals.



# Ensemble sampling

- ▶ Instead of one walker, evolve a set of  $n$  walkers.
- ▶ Can use information present in ensemble to guide proposals.
- ▶ emcee: affine invariant proposals.
- ▶ emcee is not the only (or even best) affine invariant approach.

## The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.
  - ▶ Inspired by thermodynamics.
  - ▶ Suffers from similar issues to MCMC.
  - ▶ Unclear how to choose correct annealing schedule

# The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\mathcal{Z} = P(D|M)$$

$$= \langle \mathcal{L} \rangle_{\pi}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.
  - ▶ Inspired by thermodynamics.
  - ▶ Suffers from similar issues to MCMC.
  - ▶ Unclear how to choose correct annealing schedule

## The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

## The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.

# The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.

## The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.
  - ▶ Inspired by thermodynamics.

# The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.
  - ▶ Inspired by thermodynamics.
  - ▶ Suffers from similar issues to MCMC.



## The fundamental issue with all of the above

- ▶ They don't give you evidences!

$$\begin{aligned}\mathcal{Z} &= P(D|M) \\ &= \int P(D|\Theta, M)P(\Theta|M)d\Theta \\ &= \langle \mathcal{L} \rangle_{\pi}\end{aligned}$$

- ▶ MCMC fundamentally explores the posterior, and cannot average over the prior.
- ▶ Simulated annealing gives one possibility for computing evidences.
  - ▶ Inspired by thermodynamics.
  - ▶ Suffers from similar issues to MCMC.
  - ▶ Unclear how to choose correct annealing schedule

What is nested sampling?

# What is nested sampling?

- ▶ Nested sampling is an alternative way of sampling posteriors.

# What is nested sampling?

- ▶ Nested sampling is an alternative way of sampling posteriors.
- ▶ Uses ensemble sampling to compress prior to posterior.

# What is nested sampling?

- ▶ Nested sampling is an alternative way of sampling posteriors.
- ▶ Uses ensemble sampling to compress prior to posterior.
- ▶ In doing so, it circumvents many issues (dimensionality, topology, geometry) that beset standard approaches.

# Nested Sampling

John Skilling's alternative to traditional MCMC!

# Nested Sampling

John Skilling's alternative to traditional MCMC!

New procedure:

# Nested Sampling

John Skilling's alternative to traditional MCMC!

New procedure:

Maintain a set  $S$  of  $n$  samples, which are sequentially updated:



# Nested Sampling

John Skilling's alternative to traditional MCMC!

New procedure:

Maintain a set  $S$  of  $n$  samples, which are sequentially updated:

$S_0$ : Generate  $n$  samples uniformly over the space (from the prior  $\pi$ ).

# Nested Sampling

John Skilling's alternative to traditional MCMC!

New procedure:

Maintain a set  $S$  of  $n$  samples, which are sequentially updated:

- $S_0$ : Generate  $n$  samples uniformly over the space (from the prior  $\pi$ ).
- $S_{n+1}$ : Delete the lowest likelihood sample in  $S_n$ , and replace it with a new uniform sample with higher likelihood

# Nested Sampling

John Skilling's alternative to traditional MCMC!

New procedure:

Maintain a set  $S$  of  $n$  samples, which are sequentially updated:


$S_0$ : Generate  $n$  samples uniformly over the space (from the prior  $\pi$ ).

$S_{n+1}$ : Delete the lowest likelihood sample in  $S_n$ , and replace it with a new uniform sample with higher likelihood

Requires one to be able to uniformly within a region, subject to a *hard likelihood constraint*.


# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid



# Nested Sampling


Graphical aid






# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid



# Nested Sampling


Graphical aid






# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid



# Nested Sampling


Graphical aid






# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid



# Nested Sampling


Graphical aid






# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid




# Nested Sampling

Graphical aid



# Nested Sampling


Graphical aid





# Nested Sampling

Graphical aid



## Nested sampling

## Nested sampling

- ▶ The set of dead points are posterior samples with an appropriate weighting factor

## Nested sampling

- ▶ The set of dead points are posterior samples with an appropriate weighting factor
- ▶ They can also be used to calculate evidences, since it sequentially updates the priors.

## Sampling from a hard likelihood constraint

## Sampling from a hard likelihood constraint

*"It is not the purpose of this introductory paper to develop the technology of navigation within such a volume. We merely note that exploring a hard-edged likelihood-constrained domain should prove to be neither more nor less demanding than exploring a likelihood-weighted space."*

*— John Skilling*

## Sampling from a hard likelihood constraint

*"It is not the purpose of this introductory paper to develop the technology of navigation within such a volume. We merely note that exploring a hard-edged likelihood-constrained domain should prove to be neither more nor less demanding than exploring a likelihood-weighted space."*

*— John Skilling*

- ▶ Most of the work in NS to date has been in attempting to implement a hard-edged sampler in the NS meta-algorithm.

# Sampling within an iso-likelihood contour

Previous attempts



# Sampling within an iso-likelihood contour

Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

# Sampling within an iso-likelihood contour

Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

# Sampling within an iso-likelihood contour

Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

Hamiltonian M.J. Betancourt (2010)

# Sampling within an iso-likelihood contour

## Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- Suffers in high dimensions

Hamiltonian M.J. Betancourt (2010)

Galilean F. Feroz & J. Skilling (2013)

# Sampling within an iso-likelihood contour

## Previous attempts

**Rejection Sampling** MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

**Hamiltonian** M.J. Betancourt (2010)

**Galilean** F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

# Sampling within an iso-likelihood contour

## Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

Hamiltonian M.J. Betancourt (2010)

Galilean F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

Diffusive Nested Sampling B. Brewer et al. (2009,2016).

# Sampling within an iso-likelihood contour

## Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

Hamiltonian M.J. Betancourt (2010)

Galilean F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

Diffusive Nested Sampling B. Brewer et al. (2009,2016).

- ▶ Very promising

# Sampling within an iso-likelihood contour

## Previous attempts

Rejection Sampling MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

Hamiltonian M.J. Betancourt (2010)

Galilean F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

Diffusive Nested Sampling B. Brewer et al. (2009,2016).

- ▶ Very promising
- ▶ Still needs tuning.



# Sampling within an iso-likelihood contour

## Previous attempts

**Rejection Sampling** MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

**Hamiltonian** M.J. Betancourt (2010)

**Galilean** F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

**Diffusive Nested Sampling** B. Brewer et al. (2009,2016).

- ▶ Very promising
- ▶ Still needs tuning.

**Slice Sampling** PolyChord; Handley et al. (2015).

# Sampling within an iso-likelihood contour

## Previous attempts

**Rejection Sampling** MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

**Hamiltonian** M.J. Betancourt (2010)

**Galilean** F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

**Diffusive Nested Sampling** B. Brewer et al. (2009,2016).

- ▶ Very promising
- ▶ Still needs tuning.

**Slice Sampling** PolyChord; Handley et al. (2015).

- ▶ Current “state-of-the-art”.

# Sampling within an iso-likelihood contour

## Previous attempts

**Rejection Sampling** MultiNest; F. Feroz & M. Hobson (2009).

- ▶ Suffers in high dimensions

**Hamiltonian** M.J. Betancourt (2010)

**Galilean** F. Feroz & J. Skilling (2013)

- ▶ Requires gradients and tuning

**Diffusive Nested Sampling** B. Brewer et al. (2009,2016).


- ▶ Very promising
- ▶ Still needs tuning.

**Slice Sampling** PolyChord; Handley et al. (2015).

- ▶ Current “state-of-the-art”.
- ▶ PolyChord 2.0 imminent.


# Object detection

Toy problem




# Object detection

Toy problem




# PolyChord in action

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




# PolyChord in action

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction



# PolyChord in action


Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction






# PolyChord in action

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




# PolyChord in action

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




# 0 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




# 1 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




## 2 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




### 3 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




## 4 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction



## 5 internal knots


Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction






## 6 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




## 7 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




## 8 internal knots

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction




# Bayes Factors

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction



# Marginalised plot

Primordial power spectrum  $\mathcal{P}_{\mathcal{R}}(k)$  reconstruction



# Dark energy equation of state reconstruction

# Dark energy equation of state reconstruction

- ▶ Same thing, but for Dark energy equation of state  $w(z)$  (quintessence).


# Dark energy equation of state reconstruction

- ▶ Same thing, but for Dark energy equation of state  $w(z)$  (quintessence).
- ▶ Data used is Planck 2015, BOSS DR 11, JLA supernovae and BOSS Ly $\alpha$  data




# Flat, variable $w$

Dark energy equation of state reconstruction




# Tilted

Dark energy equation of state reconstruction




# 1 internal node

Dark energy equation of state reconstruction




## 2 internal nodes

Dark energy equation of state reconstruction




# 3 internal nodes

Dark energy equation of state reconstruction




# Marginalised plot - just extension models

Dark energy equation of state reconstruction



# Marginalised plot - including LCDM

Dark energy equation of state reconstruction



# Useful links

My email: [wh260@cam.ac.uk](mailto:wh260@cam.ac.uk)

My room: Room 104, Tuesday-Thursday this week

PolyChord: [ccpforge.cse.rl.ac.uk/gf/project/polychord](http://ccpforge.cse.rl.ac.uk/gf/project/polychord)

MultiNest: [ccpforge.cse.rl.ac.uk/gf/project/multinest](http://ccpforge.cse.rl.ac.uk/gf/project/multinest)

Stan: [mc-stan.org/](http://mc-stan.org/)

emcee: [dan.iel.fm/emcee/current/](http://dan.iel.fm/emcee/current/)