UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

General Certificate of Education O Level

MARK SCHEME for the June 2005 question paper

5070 CHEMISTRY

5070/02 **Paper 2 (Theory 1), maximum mark 75**

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

- CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the June 2005 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

UNIVERSITY of CAMBRIDGE
International Examinations

JUNE 2005

GCE O Level

MARK SCHEME

MAXIMUM MARK: 75

SYLLABUS/COMPONENT: 5070/02

CHEMISTRY
Paper 2 (Theory 1)

UNIVERSITY of CAMBRIDGE
International Examinations

Section A

Maximum 45 marks

A1 four names at **(1)** each: penalise correct formulae once only

(a) nitrogen dioxide

(b) silicon dioxide

(c) aluminium oxide

(d) lead(II) iodide **[Total: 4]**

A2 **(a)** iron has positive ions and delocalised electrons **(1)**
the electrons are free to move **(1)**
moving electrons is an electric current **(1)** **[3]**

(b) high carbon steels are strong or are brittle (allow harder) **(1)**
low carbon steels are soft or are more easily shaped
(allow more malleable) **(1)** **[2]**

(c) (i) conditions are air (oxygen) and water or moist air **(1)**

(ii) magnesium is above iron in the reactivity series
(or is more reactive) **(1)**
hence it corrodes before the iron **(1)** **[3]**

(d) any two from:
coloured compounds/variable oxidation states/can act as
catalysts/valency/form complex ions **[2]**

(e) calculation
for idea of dividing by correct A_r **(1)**
dividing by the smallest **(1)**
for final formula only if first 2 fully correct **(1)**

K	Fe	C	N
0.547/39	0.195/56	0.252/12	0.294/14
0.0140	0.00348	0.0210	0.0210
4	1	6	6

i.e. $K_4FeC_6N_6$ or $K_4Fe(CN)_6$ **[3]**

[Total: 13]

© University of Cambridge International Examinations 2005

A3 **(a)** Group 0 or the noble gas group or Group 8 **[1]**

(b) Any two sensible suggestions at **(1)** each
e.g: Mendeleev's table has:
Groups and periods reversed (only allow once)
no A_r
no atomic numbers
no transition metals
periods 4 and/or 5 and all or a specific group has two elements
group numbers Arabic rather than Roman **[2]**

(c) any two observations at **(1)** each
fizzes/runs on the surface/flame/dissolves/explodes/melts
equation **(1)**
$2\ Rb + 2\ H_2O \rightarrow 2\ RbOH + H_2$ **[3]**

[Total:6]

A4 **(a)** boiling point **[1]**

(b) (i) making chemicals or feedstock or make petrol
not make plastics **(1)**

(ii) for road surfaces **(1)** **[2]**

(c) (i) saturated is single bonds or no double/triple bonds
or maximum number of hydrogen atoms **(1)**
hydrocarbon is carbon and hydrogen only **(1)**

(ii) correct methane structure (all dots = 1) **(2)** **[4]**

(d) any two ideas at **(1)** each:
enables supply to match demand (allow more useful)
make more petrol
make hydrogen
make alkenes e.g. ethene **[2]**

[Total: 9]

A5 **(a) (i)** hydrogen is below sodium in the reactivity series **(1)**

(ii) chloride ions are removed (leaving hydroxide ions) **(1)**
[2]

(b) (i) chlorine bleaches litmus or turns starch/iodide paper blue **(1)**

(ii) hydrogen pops with a burning splint **(1)** **[2]**

(c) chlorine kills bacteria
(not just sterilises the water) **[1]**

(d) burning hydrogen does not produce pollutants or only forms
water or hydrogen is not a finite resource, is renewable **[1]**

© University of Cambridge International Examinations 2005

(e) **(i)** no products or no reaction **(1)**

(ii) sodium chloride and bromine, both needed for **(1)**
(allow NaC*l* and Br_2) **[2]**

[Total: 8]

A6 **(a)** sodium ion shown as 2.8 **(1)**
chloride ion shown as 2.8.8 **(1)**
(charges not needed. Outer shell only = 0) **[2]**

(b) **(i)** strong attraction between oppositely charged ions **(1)**

(ii) higher charges on the ions **(1)**
hence stronger attraction **(1)**
(independent marks) **[3]**

(c) ions cannot move in the solid but can move in the melt **[1]**

[Total: 6]

[Section A: score any 45 from 46]

© University of Cambridge International Examinations 2005

Section B

Answer any three questions

B7 **(a)** ozone is formed by photochemical reactions
(or sparks in air, u.v on O_2) **[1]**

(b) ozone removed by reaction with chlorine (atoms) **(1)**
derived from CFC's **(1)**
ozone loss causes skin cancers or cataracts or crop damage
or skin diseases or eye damage **(1)**
(allow O_3 + CFC for **(1)**) **[3]**

(c) **(i)** bond breaking is endothermic/absorbs energy **(1)**
and bond forming is exothermic/releases energy
more energy released than absorbed (only if first point scored) **(1)**

(ii) as temperature increases molecules move faster
or increased k.e. **(1)**
hence more frequent collisions
or more molecules energy exceeds the activation energy **(1)**

(iii) calculation
48 g ozone releases 143 kJ **(1)**
16 g ozone releases 47.66 kJ or 47.7 kJ **(1)**
(answer alone **(1)**, units needed)
(if 6 x 16 = 96 g ozone used, then **(0)**)
(if 0.33 used, answer = 47.2) **[6]**

[Total: 10]

B8 **(a)** calculation **(2)**
143.5 g AgC*l* contains 108 g Ag
0.287 g AgC*l* contains 0.216 g Ag
(answer alone **(1)**, units needed) **[2]**

(b) oxidation is electron loss or an increase in O.N. **(1)**
copper(I) is oxidised because it loses an electron
or its O.N. increases **(1)**
chlorine is reduced because it gains an electron
or its O.N. decreases **(1)** **[3]**

(c) equation **(1)**
$Ag + CuCl_2 \rightarrow AgCl + CuCl$ **[1]**

(d) **(i)** equation **(1)** state symbols **(1)**
$CuCl_2(aq) + 2\,NaOH(aq) \rightarrow Cu(OH)_2(s) + 2\,NaCl(aq)$,
(or ionic, $Cu^{2+} + 2OH^- \rightarrow Cu(OH)_2$)
(scores **(1)** for states)

(ii) name is copper(II) hydroxide (allow copper hydroxide) **(1)**
colour is blue or blue-green **(1)**
(colour only for correct name) **[4]**

[Total: 10]

© University of Cambridge International Examinations 2005

B9 **(a)** **(i)** the catalyst is iron or Fe_2O_3 **(1)**

(ii) equation
$N_2 + 3H_2 \rightarrow 2NH_3$ **(1)**

(iii) the temperature is 280 °C **(1)**
the pressure is 400 atmos **(1)**

(iv) higher temperature gives faster reaction **(1)**
(higher yield = -1) **[5]**

(b) a catalyst increases reaction rate **(1)**
(not alters the rate)
a lower activation energy **(1)**
hence saves energy **(1)**
(third mark only if E_a given) **[3]**

(c) equation **(1)**
$Ca(OH)_2 + 2\ NH_4NO_3 \rightarrow Ca(NO_3)_2 + 2\ H_2O + 2\ NH_3$
ammonia lost as a gas **(1)**
[2]

[Total: 10]

B10 **(a)** name is butanoic acid (not butenoic) **(1)**

(b) formula is $C_5H_{11}CO_2H$ (not $C_6H_{12}O_2$) **(1)**

(c) structure of ethyl ethanoate **(1)**
allow full structure or condensed version, $CH_3CO_2C_2H_5$

(d) allow any suitable named oxidising reagent **(1)**
e.g. (acidified) potassium dichromate(VI) or air or oxygen
(allow formula) **[(a) to (d) 4]**

(e) equation **(1)**
$Mg + 2\ CH_3CO_2H \rightarrow Mg(CH_3CO_2)_2 + H_2$
calculation **(2)**
50 cm^3 acid is 0.05 mol 0.025 mol Mg needed
24 x 0.025 = 0.60 g
(answer alone **(1)**, unit needed) **[3]**

(f) ethanoic acid is weak and hydrochloric is strong **(1)**
lower $[H^+]$ concentration in ethanoic acid **(1)**
[2]

(g) ionic equation **(1)**
$H^+ + OH^- \rightarrow H_2O$ **[1]**

[Total: 10]

© University of Cambridge International Examinations 2005