www.PapaCambridge.com

O

ADDITIONAL MATHEMATICS **4037/12**

Paper 1 **May/June 2010**

2 hours

Additional Materials: Answer Booklet/Paper
Graph paper (2 sheets)

9495576362

This document consists of **6** printed pages and **2** blank pages.

DCA (KN) 12935/4
© UCLES 2010

UNIVERSITY of CAMBRIDGE
International Examinations

[Turn over

www.PapaCambridge.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$

$$a^2 = b^2 + c^2 - 2bc \cos A.$$

$$\Delta = \frac{1}{2} bc \sin A.$$

www.PapaCambridge.com

1 Find the coordinates of the points of intersection of the curve $y^2 + y = 10x - 8x^2$ and the line $y + 4x + 1 = 0$.


2 The expression $6x^3 + ax^2 - (a + 1)x + b$ has a remainder of 15 when divided by $x + 2$ and a remainder of 24 when divided by $x + 1$. Show that $a = 8$ and find the value of b. [5]

3 Given that $\overrightarrow{OA} = \begin{pmatrix} -17 \\ 25 \end{pmatrix}$ and $\overrightarrow{OB} = \begin{pmatrix} 4 \\ 5 \end{pmatrix}$, find

(i) the unit vector parallel to $\overrightarrow{AB}$, [3]

(ii) the vector $\overrightarrow{OC}$, such that $\overrightarrow{AC} = 3\overrightarrow{AB}$. [2]

4




Variables x and y are such that, when y^2 is plotted against $\sec x$, a straight line graph passing through the points (2.4, 1.6) and (1.3, 3.8) is obtained.

(i) Express y^2 in terms of $\sec x$. [3]

(ii) Hence find the exact value of $\cos x$ when $y = 2$. [2]

www.PapaCambridge.com

5



The diagram shows part of the curve $y = 6 - \frac{3}{x}$ which passes through the point A where $x = 3$. The normal to the curve at the point A meets the x-axis at the point B. Find the coordinates of the point B. [5]

6 **(a)** **(i)** On the same diagram, sketch the curves $y = \cos x$ and $y = 1 + \cos 2x$ for $0 \leqslant x \leqslant 2\pi$. [3]

(ii) Hence state the **number** of solutions of the equation

$$\cos 2x - \cos x + 1 = 0 \text{ where } 0 \leqslant x \leqslant 2\pi.$$ [1]

(b) The function f is given by $f(x) = 5\sin 3x$. Find

(i) the amplitude of f, [1]

(ii) the period of f. [1]

7 The table shows values of the variables p and v which are related by the equation $p = kv^n$, where k and n are constants.

v	10	50	110	230
p	1412	151	53	19

(i) Using graph paper, plot $\lg p$ against $\lg v$ and draw a straight line graph. [3]

Use your graph to estimate

(ii) the value of n, [2]

(iii) the value of p when $v = 170$. [2]

www.PapaCambridge.com


8 Given that $\mathbf{A} = \begin{pmatrix} 4 & 3 \\ 1 & 2 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} -2 & 0 \\ 1 & 4 \end{pmatrix}$, find

(i) $3\mathbf{A} - 2\mathbf{B}$,

(ii) $\mathbf{A}^{-1}$, [2]

(iii) the matrix $\mathbf{X}$ such that $\mathbf{X}\mathbf{B}^{-1} = \mathbf{A}$. [3]

9



The diagram shows a sector OXY of a circle centre O, radius 3 cm and a sector OAB of a circle centre O, radius 8 cm. The point X lies on the line OA and the point Y lies on the line OB. The perimeter of the region $XABYX$ is 15.5 cm. Find

(i) the angle AOB in radians, [3]

(ii) the ratio of the area of the sector OXY to the area of the region $XABYX$ in the form $p : q$, where p and q are integers. [4]

10 A music student needs to select 7 pieces of music from 6 classical pieces and 4 modern pieces. Find the number of different selections that she can make if

(i) there are no restrictions, [1]

(ii) there are to be only 2 modern pieces included, [2]

(iii) there are to be more classical pieces than modern pieces. [4]

www.PapaCambridge.com

11 A particle moves in a straight line such that its displacement, x m, from a fixed point O at time t seconds is given by $x = 12\{\ln(2t + 3)\}$. Find


(i) the value of t when the displacement of the particle from O is 48 m,

(ii) the velocity of the particle when $t = 1$, [3]

(iii) the acceleration of the particle when $t = 1$. [3]

12 Answer only **one** of the following two alternatives.

EITHER



The diagram shows part of a curve for which $\frac{dy}{dx} = 8\cos 2x$. The curve passes through the point $B\left(\frac{\pi}{4}, 7\right)$. The line $y = 5$ meets the curve at the points A and C.

(i) Show that the curve has equation $y = 3 + 4\sin 2x$. [3]

(ii) Find the x-coordinate of the point A and of the point C. [4]

(iii) Find the area of the shaded region. [5]

OR

A curve is such that $\frac{dy}{dx} = 6e^{3x} - 12$. The curve passes through the point (0, 1).

(i) Find the equation of the curve. [4]

(ii) Find the coordinates of the stationary point of the curve. [3]

(iii) Determine the nature of the stationary point. [2]

(iv) Find the coordinates of the point where the tangent to the curve at the point (0, 1) meets the x-axis. [3]

BLANK PAGE

www.PapaCambridge.com

www.PapaCambridge.com

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of