www.PapaCambridge.com

O

ADDITIONAL MATHEMATICS **4037/22**

Paper 2 **May/June 2010**

2 hours

Additional Materials: Answer Booklet/Paper
Graph paper (2 sheets)

9126272629

This document consists of **6** printed pages and **2** blank pages.

DCA (KN) 12939/4
© UCLES 2010

UNIVERSITY of CAMBRIDGE
International Examinations

[Turn over

www.PapaCambridge.com

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}.$$

Binomial Theorem

$$(a + b)^n = a^n + \binom{n}{1} a^{n-1} b + \binom{n}{2} a^{n-2} b^2 + \ldots + \binom{n}{r} a^{n-r} b^r + \ldots + b^n,$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!r!}$.

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1.$$

$$\sec^2 A = 1 + \tan^2 A.$$

$$\operatorname{cosec}^2 A = 1 + \cot^2 A.$$

Formulae for ΔABC

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}.$$


$$a^2 = b^2 + c^2 - 2bc \cos A.$$

$$\Delta = \frac{1}{2} bc \sin A.$$

www.PapaCambridge.com


1 Find $\int\left(2 + 5x - \frac{1}{(x-2)^2}\right)\mathrm{d}x$.

2 **(a)**



Copy the diagram and shade the region which represents the set $A \cup (B \cap C')$. [1]

(b)




Express, in set notation, the set represented by the shaded region. [1]

(c) The universal set $\mathscr{E}$ and the sets P and Q are such that $\mathrm{n}(\mathscr{E}) = 30$, $\mathrm{n}(P) = 18$ and $\mathrm{n}(Q) = 16$. Given that $\mathrm{n}(P \cup Q)' = 2$, find $\mathrm{n}(P \cap Q)$. [2]

3 The volume $V\,\mathrm{cm}^3$ of a spherical ball of radius $r\,\mathrm{cm}$ is given by $V = \frac{4}{3}\pi r^3$. Given that the radius is increasing at a constant rate of $\frac{1}{\pi}\,\mathrm{cm\,s^{-1}}$, find the rate at which the volume is increasing when $V = 288\pi$. [4]

www.PapaCambridge.com

4




The diagram shows a right-angled triangle ABC in which the length of AB is $\frac{16}{\sqrt{2}}$, the length of BC is $7\sqrt{3}$ and angle BCA is θ.

(i) Find $\tan\theta$ in the form $\frac{a\sqrt{6}}{b}$, where a and b are integers. [2]

(ii) Calculate the length of AC, giving your answer in the form $c\sqrt{d}$, where c and d are integers and d is as small as possible. [3]

5 Solve the equation $2x^3 - 3x^2 - 11x + 6 = 0$. [6]

6



The diagram shows part of the line $y = 12 - 2x$. The point $Q(x, y)$ lies on this line and the points P and R lie on the coordinate axes such that $OPQR$ is a rectangle.

(i) Write down an expression, in terms of x, for the area A of the rectangle $OPQR$. [2]

(ii) Given that x can vary, find the value of x for which A has a stationary value. [3]

(iii) Find this stationary value of A and determine its nature. [2]

www.PapaCambridge.com

7 **(i)** Sketch the graph of $y = |3x + 9|$ for $-5 < x < 2$, showing the coordinates of the points where the graph meets the axes.

(ii) On the same diagram, sketch the graph of $y = x + 6$.

(iii) Solve the equation $|3x + 9| = x + 6$. [3]

8 **(a)** **(i)** Write down the first 4 terms, in ascending powers of x, of the expansion of $(1 - 3x)^7$. [3]

(ii) Find the coefficient of x^3 in the expansion of $(5 + 2x)(1 - 3x)^7$. [2]

(b) Find the term which is independent of x in the expansion of $\left(x^2 + \frac{2}{x}\right)^9$. [3]

9 **(i)** Given that $y = \frac{x + 2}{(4x + 12)^{\frac{1}{2}}}$, show that $\frac{dy}{dx} = \frac{k(x + 4)}{(4x + 12)^{\frac{3}{2}}}$, where k is a constant to be found. [5]

(ii) Hence evaluate $\int_1^{13} \frac{x + 4}{(4x + 12)^{\frac{3}{2}}}\,dx$. [3]

10 **(a)** Given that $\log_p X = 6$ and $\log_p Y = 4$, find the value of

(i) $\log_p \left(\frac{X^2}{Y}\right)$, [2]

(ii) $\log_Y X$. [2]

(b) Find the value of 2^z, where $z = 5 + \log_2 3$. [3]

(c) Express $\sqrt{512}$ as a power of 4. [2]

11 **(a)** Solve, for $0 < x < 3$ radians, the equation $4\sin x - 3 = 0$, giving your answers correct to 2 decimal places. [3]

(b) Solve, for $0° < y < 360°$, the equation $4\operatorname{cosec} y = 6\sin y + \cot y$. [6]

www.PapaCambridge.com

12 Answer only **one** of the following two alternatives.

EITHER

It is given that $\mathrm{f}(x) = 4x^2 + kx + k$.

(i) Find the set of values of k for which the equation $\mathrm{f}(x) = 3$ has no real roots. [5]

In the case where $k = 10$,

(ii) express $\mathrm{f}(x)$ in the form $(ax + b)^2 + c$, [3]

(iii) find the least value of $\mathrm{f}(x)$ and the value of x for which this least value occurs. [2]

OR

The functions f, g and h are defined, for $x \in \mathbb{R}$, by

$$\mathrm{f}(x) = x^2 + 1,$$

$$\mathrm{g}(x) = 2x - 5,$$

$$\mathrm{h}(x) = 2^x.$$

(i) Write down the range of f. [1]

(ii) Find the value of gf(3). [2]

(iii) Solve the equation $\mathrm{fg}(x) = \mathrm{g}^{-1}(15)$. [5]

(iv) On the same axes, sketch the graph of $y = \mathrm{h}(x)$ and the graph of the inverse function $y = \mathrm{h}^{-1}(x)$, indicating clearly which graph represents h and which graph represents h^{-1}. [2]

BLANK PAGE

www.PapaCambridge.com

www.PapaCambridge.com

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of