UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

General Certificate of Education O Level

MARK SCHEME for the November 2004 question paper

5070 CHEMISTRY

5070/02 **Paper 2 (Theory 1), maximum mark 75**

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

- CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the November 2004 question papers for most IGCSE and GCE Advanced Level syllabuses.

UNIVERSITY of CAMBRIDGE
International Examinations

NOVEMBER 2004

GCE O Level

MARK SCHEME

MAXIMUM MARK: 75


SYLLABUS/COMPONENT: 5070/02

CHEMISTRY
Paper 2 (Theory 1)

UNIVERSITY of CAMBRIDGE
International Examinations

A1	**(a)**		rises then falls <u>only</u>; **NOT** references to B and C	**1**
	(b)	**(i)**	less than 12/any number <12;	**1**
		(ii)	eutrophication; weed/algae grows more/faster; <u>rotting/decomposition/bacteria</u> uses up oxygen; 'use up oxygen' alone does not score	**any 2**
	(c)		decreases; decreases; increases;	**3**
				7 marks
A2			**a to d accept correct formulae, use list principle**	
	(a)		bromine and methane/(both needed)	**1**
	(b)		lithium	**1**
	(c)		iodine and bromine/Br_2 and I_2 (both needed)	**1**
	(d)		lithium and lead (II) bromide (both needed)	**1**
	(e)		methane has a <u>simple</u> (covalent) structure (not discussion of breaking bonds in methane); silicon dioxide has a <u>giant/lattice/macromolecular</u> (covalent) structure;	**2**
	(f)		electrolysis; of <u>molten</u> lead bromide; **allow:** (metal) displacement; by more reactive metal/named more reactive metal (magnesium, zinc, iron);	**2**
				8 marks

© University of Cambridge International Examinations 2005

A3	**(a)**	**(i)**	(conc) H_2SO_4; **not** dilute H_2SO_4, **accept** phosphoric acid heat/reflux/50 - 150 °C; ignore pressure	**2**
		(ii)	 (**allow** condensed OH as shown)	**1**
		(iii)	$CH_3COOH + C_3H_7OH \rightarrow CH_3COOC_3H_7 + H_2O$; **e.c.f.** from **(ii)** **allow** molecular formulae LHS = 1 RHS = 1	**2**
	(b)	**(i)**	pH meter/universal indicator/electrical conductivity test; shows different pH/orange for carboxylic acid, red for hydrochloric/different colours (if colours stated, must be correct)/electrical conductivity different/electrical conductivity higher in HC*l* **1 mark max** for chemical reactions: add reactive/named solid (as in **(iii)**) and compare rates/test for chloride ion using silver nitrate;	**2**
		(ii)	metal carbonate/metal oxide/named metal carbonate or named oxide (**not** Group I oxide or CaO)/magnesium metal, zinc metal	**1**
			consequential on <u>correct</u> substance- carbonate or metal – see bubbles metal oxide – solid disappears, **accept** dissolves	**1**
				9 marks
A4	**(a)**		blocks oxygen uptake in blood; **not** 'breathing difficulties'	**1**
	(b)	**(i)**	H_2O;	**1**

© University of Cambridge International Examinations 2005

		(ii)	Pd oxidation states (+)2 to 0; C oxidation states (+)2 to (+)4;	**2**
		(iii)	palladium has been reduced and C has been oxidised; palladium ox state has fallen, C has increased/palladium accepted electrons from carbon; e.c.f. from **(ii)**	**2**
	(c)		extraction of iron, zinc, lead or tin/blast furnace	**1**
				7 marks
A5	**(a)**		$Zn + Cu^{2+} \rightarrow Cu + Zn^{2+}$ check equation is <u>correct direction</u> ignore state symbols	**1**
	(b)		arrow in external circuit from zinc to copper (to the left)	**1**
	(c)		zinc iron lead copper Zn and Cu correct = 1 iron lead correct = 1	**2**
	(d)		magnesium/aluminium	**1**
				5 marks
A6	**(a)**	**(i)**	(aqueous) lithium hydroxide/lithium carbonate; **not** lithium oxide evaporation/(allow to) crystallise;	**2**

© University of Cambridge International Examinations 2005

		(ii)	(aqueous) barium chloride/barium nitrate/barium hydroxide; filtration	**2**
		(iii)	copper oxide or copper carbonate; **accept** copper hydroxide <u>partial</u> evaporation (owtte)/leave to crystallise	**2**
	(b)		relative molecular mass $(NH_4)_2SO_4$ = 132; 34 g NH_3 makes 132 g $(NH_4)_2SO_4$ owtte; mass formed = 132/34 x 51 = 198 g usual calculation rules apply.	**3**
				9 marks
				Total Section A = 45

Section B

B7	**(a)**		**Diagram** standard rate curve shape; labels on axes 'volume' against 'time' (owtte); **Explanation** reaction stops when magnesium carbonate <u>used up</u>;	**2** **1**
	(b)		M_r $MgCO_3$ = 84; no mols CO_2 = 10.5/84 (=0.125 mols); volume = 0.125 x 24 = 3 dm^3 usual calculation rules apply	**3**
	(c)	**(i)**	faster; because zinc carbonate is less (thermally) stable than magnesium carbonate ORA ignore references to metal reactivity	**2**

© University of Cambridge International Examinations 2005

		(ii)	less carbon dioxide; because there are <u>fewer moles</u> of zinc carbonate. **Calculation** leading to 2.02 dm^3 = 2 marks <u>different</u> amount of CO_2 because 10.5 g zinc carbonate contains a <u>different</u> number of moles = 1 mark	**2**
				10 marks
B8	**(a)**		**A** diesel oil **B** paraffin **C** naphtha	**1**
	(b)		fractions vaporise/evaporate/boil; <u>condense</u> at <u>different temperatures</u>; lowest boiling points come out at highest point of tower/ temp of tower higher at bottom	**3**
	(c)	**(i)**	correct <u>method</u> Mass of C/mass of compound x 100; correct <u>masses</u> used octane 96/114 x 100 = 84.2 %; <u>both fully correct</u> hexadecane 192/226 x 100 = 85.0 % Guidance: one calculation fully correct scores 2; both calculations fully correct scores 3; allow e.c.f. for minor arithmetical errors.	**3**
		(ii)	$2C_{16}H_{34} + 49O_2 \rightarrow 32CO_2 + 34H_2O$ ignore state symbols	**1**
		(iii)	less oxygen is needed (per molecule) to combust octane ORA/more carbon <u>atoms</u> in hexadecane/more <u>carbon per molecule/</u>higher percentage C by mass; 'more carbon' alone is not enough	**1**
	(d)		hydrogen and ethanol/alcohol ignore solar	**1**
				10 marks

© University of Cambridge International Examinations 2005

B9	**(a)**		reaction is exothermic/gives out heat/gives out energy	**1**
	(b)		$4Fe + 3O_2 \rightarrow 2Fe_2O_3$ (1) for Fe_2O_3; CONSEQUENTIAL (1) for rest of equation correct	**2**
	(c)	**(i)**	rises slowly then faster; melting the scrap is endothermic/uses energy /temperature of molten iron changes less when scrap is melting.	**2**
		(ii)	saving metal ores/saving energy for extraction/saves need to dispose of scrap iron.	**1**
	(d)	**(i) and(ii)**	(mark together) more carbon in high carbon steel; both alloys contain more iron than carbon;	**2**
		(iii)	**Property**: low C steel softer/weaker/more easily shaped/less brittle than high carbon steel; ORA **Structure**: properties change because carbon atoms are smaller than iron atoms (may come from reference to diagram)/metallic bonding is disrupted/lattice is disrupted/alloy structure is less regular/layers need to slip when steel changes shape	**2**
				10 marks
B10	**(a)**		correct set-up showing battery and two electrodes dipping in an electrolyte; nickel at cathode and silver at anode; named electrolyte: silver nitrate.	**3**
	(b)		anode reaction: $Ag(s) \rightarrow Ag^+(aq) + e^-$; cathode reaction: $Ag^+(aq) + e^- \rightarrow Ag(s)$; **incorrect state symbols** on fully correct equations (1) mark **electrodes reversed** with fully correct equations (1) mark	**2**

© University of Cambridge International Examinations 2005

	(c)	**(i)**	**Salt A:** Gp1 sulphate or Gp 1/2 nitrate or zinc sulphate or nitrate or magnesium sulphate/<u>dilute</u> (aqueous) sodium chloride; **Salt B:** Gp 1/2 chloride or zinc chloride;	**2**
		(ii)	oxygen relights glowing spill; hydrogen pops when lit; chlorine bleaches (damp) litmus/indicator paper OR mix with Group I iodide/bromide, solution goes yellow/brown;	**3**
				10 marks
				Total Section B = 30

© University of Cambridge International Examinations 2005