

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1

GENVASC**GP Practice:**

- With consent, please draw 3x EDTA tubes 3ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

02/01/2018 vs.1