

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1

GENVASC

GP Practice:

- With consent, please draw 2 x EDTA tubes 4ml (purple)
- Attach the GENVASC consent form to the bag
- Send to pathology separate to clinical samples using routine pathology transit

15/03/2017 v1