

DCT v3.13: recorded training variants

Variants	BLCA	KIRC	Mean (2)
Exp0: patient NLL only	0.7001 $\pm$ 0.0503	<u>0.8120 $\pm$ 0.0394</u>	0.7560
Exp1: + IPCW ranking	0.7097 $\pm$ 0.0584	0.7935 $\pm$ 0.0343	0.7516
Exp2: + per-slot NLL	0.7189 $\pm$ 0.0537	0.7921 $\pm$ 0.0206	0.7555
Exp3: + diversity, no reconstruction	<u>0.7205 $\pm$ 0.0403</u>	0.8049 $\pm$ 0.0416	<u>0.7627</u>
Exp4: self only (0.025)	0.7101 $\pm$ 0.0434	0.8046 $\pm$ 0.0329	0.7574
Exp5: cross only (0.025)	0.7188 $\pm$ 0.0474	0.8027 $\pm$ 0.0278	0.7607
Full: self + cross (0.05 each)	0.7238 $\pm$ 0.0467	0.8224 $\pm$ 0.0163	0.7731
Direct: cross reconstruction from WSI slots	0.7029 $\pm$ 0.0474	0.8030 $\pm$ 0.0302	0.7530
Independent: product coupling	0.7093 $\pm$ 0.0408	0.8054 $\pm$ 0.0171	0.7573

Five-fold best-validation C-index; UNI2-h; seed 3. Red: first; underline: second among available variants.

Exp4/Exp5 retain weight 0.025; Full uses 0.05 each. These historical recipes are not pure matched-weight removals.

Two-cohort mean is descriptive BLCA/KIRC macro-average, not the ten-cancer Overall. $\pm$: fold sample SD.