

Divide and Conquer

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Divide-and-Conquer: Overview

Divide-and-conquer.

- Break up problem into several parts.
- Solve each part recursively.
- Combine solutions to sub-problems into overall solution.

Most common usage.

- Break up problem of size n into **two** equal parts of size $\frac{1}{2}n$.
- Solve two parts recursively.
- Combine two solutions into overall solution in **linear time**.

Remark.

- Partition the problem into **disjoint** subproblems.
- Each subproblem is the same type as the original problem.
- A simple partition of the problem and combination of solutions to subproblems may not beat brute force.

The phrase is attributed to Julius Caesar, Philip II, king of Macedon (382-336 BC), describing his political policy.

Divide-and-Conquer: Overview

Examples we have seen so far.

- Binary search
- Mergesort
- Quicksort

The running times of Divide-and-conquer algorithms are often characterized by recurrences.

- Binary search: $T(n) = 2T(n/2) + \Theta(1)$
- Mergesort: $T(n) = 2T(n/2) + \Theta(n)$
- Quicksort: not always partition the subproblems evenly!

$$T(n) = 2T(n/2) + \Theta(n),$$

$$T(n) = T(0) + T(n-1) + \Theta(n),$$

$$T(n) = T(n/4) + T(n/2) + \Theta(n)$$

.....

Solving recurrences.

Iteration, master theorem, recursion tree, etc.

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Master method

Goal. Recipe for solving common divide-and-conquer recurrences:

$$T(n) = a T\left(\frac{n}{b}\right) + f(n)$$


where $a \geq 1$, $b > 1$, and f is asymptotically positive.

Terms.

- $a \geq 1$ is the (integer) number of subproblems.
- $b \geq 2$ is the (integer) factor by which the subproblem size decreases.
- $f(n) \geq 0$ is the work to divide and combine subproblems.

Running time by analyzing the recursion tree.

- Dominated by cost at leaves
(for solving the minimum subproblems)
- Evenly distributed throughout the tree
- Dominated by cost at the root
(for dividing the problem and combining the results)



Master method

Master theorem. Suppose that $T(n)$ is a function on the non-negative integers that satisfies the recurrence

$$T(n) = a T\left(\frac{n}{b}\right) + f(n)$$

where $a \geq 1$, $b > 1$, and f is asymptotically positive.

Case 1. If $f(n) = O(n^k)$ for some constant $k < \log b^a$, then $T(n) = \Theta(n^{\log b^a})$.

Case 2. If $f(n) = \Theta(n^k \log^p n)$ for $p \geq 0$, $k = \log b^a$, then $T(n) = \Theta(n^k \log^{p+1} n)$.

Case 3. If $f(n) = \Omega(n^k)$ for some constant $k > \log b^a$, and if $a f(n/b) \leq c f(n)$ for some constant $c < 1$ and all sufficiently large n , then $T(n) = \Theta(f(n))$.

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Counting Inversions


Music site tries to match your song preferences with others.

- You rank n songs.
- Music site consults database to find people with **similar** tastes.

Similarity metric: number of inversions between two rankings.

- My rank: $1, 2, \dots, n$.
- Your rank: a_1, a_2, \dots, a_n .
- Songs i and j **inverted** if $i < j$, but $a_i > a_j$.

Songs					
	A	B	C	D	E
Me	1	2	3	4	5
You	1	3	4	2	5



Inversions

3-2, 4-2

Brute force: check all $\Theta(n^2)$ pairs i and j .

Applications

Applications.

- Voting theory.
- Collaborative filtering.
- Measuring the "sortedness" of an array.
- Sensitivity analysis of Google's ranking function.
- Rank aggregation for meta-searching on the Web.
- Nonparametric statistics (e.g., Kendall's Tau distance).

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- **Divide**: separate list into two pieces.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- Divide: separate list into two pieces.
- **Conquer**: recursively count inversions in each half.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

5 blue-blue inversions

8 green-green inversions

Conquer: $2T(n / 2)$

5-4, 5-2, 4-2, 8-2, 10-2

6-3, 9-3, 9-7, 12-3, 12-7, 12-11, 11-3, 11-7

Counting Inversions: Divide-and-Conquer

Divide-and-conquer.

- Divide: separate list into two pieces.
- Conquer: recursively count inversions in each half.
- **Combine**: count inversions where a_i and a_j are in different halves, and return sum of three quantities.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

Divide: $O(1)$.

1	5	4	8	10	2	6	9	12	11	3	7
---	---	---	---	----	---	---	---	----	----	---	---

5 blue-blue inversions

8 green-green inversions

Conquer: $2T(n / 2)$

9 blue-green inversions

5-3, 4-3, 8-6, 8-3, 8-7, 10-6, 10-9, 10-3, 10-7

Combine: ???

Total = $5 + 8 + 9 = 22$.

Counting Inversions: Combine

Combine: count blue-green inversions

- Assume each half is **sorted**.
- Count inversions where a_i and a_j are in different halves.
- **Merge** two sorted halves into sorted whole.

↖ to maintain sorted invariant

3	7	10	14	18	19
---	---	----	----	----	----

2	11	16	17	23	25
6	3	2	2	0	0

13 blue-green inversions: $6 + 3 + 2 + 2 + 0 + 0$

Count: $O(n)$

2	3	7	10	11	14	16	17	18	19	23	25
---	---	---	----	----	----	----	----	----	----	----	----

Merge: $O(n)$

$$T(n) \leq T(\lfloor n/2 \rfloor) + T(\lceil n/2 \rceil) + O(n) \Rightarrow T(n) = O(n \log n)$$

Counting Inversions: Implementation

Pre-condition. [Merge-and-Count] A and B are sorted.

Post-condition. [Sort-and-Count] L is sorted.

```
Sort-and-Count(L) {  
    if list L has one element  
        return 0 and the list L  
  
    Divide the list into two halves A and B  
    ( $r_A$ , A)  $\leftarrow$  Sort-and-Count(A)  
    ( $r_B$ , B)  $\leftarrow$  Sort-and-Count(B)  
    ( $r$ , L)  $\leftarrow$  Merge-and-Count(A, B)  
  
    return  $r = r_A + r_B + r$  and the sorted list L  
}
```


Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

--	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total:

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

6

2											
---	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total: 6

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



3	7	10	14	18	19
---	---	----	----	----	----



2	11	16	17	23	25
---	----	----	----	----	----

two sorted halves

6

2											
---	--	--	--	--	--	--	--	--	--	--	--

auxiliary array

Total: 6

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.

$i = 6$



two sorted halves

6




auxiliary array

Total: 6

Merge and Count

Merge and count step.


- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6

Merge and Count

Merge and count step.


- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and Count

Merge and count step.


- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: 6 + 3

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2 + 0$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2 + 0$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.




Total: $6 + 3 + 2 + 2 + 0 + 0$

Merge and Count

Merge and count step.

- Given two sorted halves, count number of inversions where a_i and a_j are in different halves.
- Combine two sorted halves into sorted whole.



Total: $6 + 3 + 2 + 2 + 0 + 0 = 13$

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Closest Pair of Points

Closest pair. Given n points in the plane, find a pair with smallest Euclidean distance between them.

Fundamental geometric primitive.

- Graphics, computer vision, geographic information systems, molecular modeling, air traffic control.
- Special case of nearest neighbor, Euclidean MST, Voronoi.

↖ fast closest pair inspired fast algorithms for these problems

Brute force. Check all pairs of points p and q with $\Theta(n^2)$ comparisons.

1-D version. $O(n \log n)$ easy if points are on a line.


Assumption. No two points have same x coordinate.

↑
to make presentation cleaner

Closest Pair of Points

Algorithm.


- **Divide:** draw vertical line L so that roughly $\frac{1}{2}n$ points on each side.



Closest Pair of Points

Algorithm.


- Divide: draw vertical line L so that roughly $\frac{1}{2}n$ points on each side.
- **Conquer**: find closest pair in each side recursively.



Closest Pair of Points


Algorithm.

- Divide: draw vertical line L so that roughly $\frac{1}{2}n$ points on each side.
- Conquer: find closest pair in each side recursively.
- **Combine**: find closest pair with one point in each side. ← seems like $\Theta(n^2)$
- Return best of 3 solutions.



Closest Pair of Points


Find closest pair with one point in each side, assuming that distance $< \delta$.



Closest Pair of Points

Find closest pair with one point in each side, **assuming that distance $< \delta$** .


- Observation: only need to consider points within δ of line L .



Closest Pair of Points

Find closest pair with one point in each side, **assuming that distance $< \delta$** .


- Observation: only need to consider points within δ of line L .
- Sort points in 2δ -strip by their y coordinate.



Closest Pair of Points

Find closest pair with one point in each side, **assuming that distance $< \delta$** .

- Observation: only need to consider points within δ of line L .
- Sort points in 2δ -strip by their y coordinate.
- Only check distances of those within 11 positions in sorted list!



Closest Pair of Points


Def. Let s_i be the point in the 2δ -strip, with the i^{th} smallest y -coordinate.

Claim. If $|i - j| \geq 12$, then the distance between s_i and s_j is at least δ .

Pf.

- No two points lie in same $\frac{1}{2}\delta$ -by- $\frac{1}{2}\delta$ box.
- Two points at least 2 rows apart have distance $\geq 2(\frac{1}{2}\delta)$. ▪

Fact. Still true if we replace 12 with 7.



Closest Pair Algorithm

```
Closest-Pair( $p_1, \dots, p_n$ ) {  
  Compute separation line  $L$  such that half the points  
  are on one side and half on the other side.  $O(n \log n)$   
  
   $\delta_1 = \text{Closest-Pair}(\text{left half})$   
   $\delta_2 = \text{Closest-Pair}(\text{right half})$   $2T(n / 2)$   
   $\delta = \min(\delta_1, \delta_2)$   
  
  Delete all points further than  $\delta$  from separation line  $L$   $O(n)$   
  
  Sort remaining points by  $y$ -coordinate.  $O(n \log n)$   
  
  Scan points in  $y$ -order and compare distance between  
  each point and next 11 neighbors. If any of these  
  distances is less than  $\delta$ , update  $\delta$ .  $O(n)$   
  
  return  $\delta$ .  
}
```

Closest Pair of Points: Analysis

Running time.

$$T(n) \leq 2T(n/2) + O(n \log n) \Rightarrow T(n) = O(n \log^2 n)$$

Q. Can we achieve $O(n \log n)$?

A. Yes. Don't sort points in strip from scratch each time.

- Each recursive returns two lists: all points sorted by y coordinate, and all points sorted by x coordinate.
- Sort by **merging** two pre-sorted lists.

$$T(n) \leq 2T(n/2) + O(n) \Rightarrow T(n) = O(n \log n)$$

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Integer Addition

Addition. Given two n -bit integers a and b , compute $a + b$.

Grade-school. $\Theta(n)$ bit operations.

	1	1	1	1	1	1	0	1	
		1	1	0	1	0	1	0	1
+	0	1	1	1	1	1	1	0	1
<hr/>									
	1	0	1	0	1	0	0	1	0

Remark. Grade-school addition algorithm is optimal.

Integer Multiplication

Multiplication. Given two n -bit integers a and b , compute $a \times b$.

Grade-school. $\Theta(n^2)$ bit operations.

[illegible]

Q. Is grade-school multiplication algorithm optimal?

Divide-and-Conquer Multiplication: Warmup

To multiply two n -bit integers a and b :

- Multiply four $\frac{1}{2}n$ -bit integers, recursively.
- Add and shift to obtain result.

$$\begin{aligned}a &= 2^{n/2} \cdot a_1 + a_0 \\b &= 2^{n/2} \cdot b_1 + b_0 \\ab &= (2^{n/2} \cdot a_1 + a_0)(2^{n/2} \cdot b_1 + b_0) = 2^n \cdot a_1 b_1 + 2^{n/2} \cdot (a_1 b_0 + a_0 b_1) + a_0 b_0\end{aligned}$$

Ex. $a = \underbrace{1000}_{a_1} \underbrace{1101}_{a_0} \quad b = \underbrace{1110}_{b_1} \underbrace{0001}_{b_0}$

$$T(n) = \underbrace{4T(n/2)}_{\text{recursive calls}} + \underbrace{\Theta(n)}_{\text{add, shift}} \Rightarrow T(n) = \Theta(n^2)$$

↖ assumes n is a power of 2

Conjecture. [Kolmogorov 1952] Grade-school algorithm is optimal.

Theorem. [Karatsuba 1960] Conjecture is wrong.

Karatsuba Multiplication

To multiply two n -bit integers a and b :

- Add two $\frac{1}{2}n$ bit integers.
- Multiply **three** $\frac{1}{2}n$ -bit integers, recursively.
- Add, subtract, and shift to obtain result.

$$a = 2^{n/2} \cdot a_1 + a_0$$

$$b = 2^{n/2} \cdot b_1 + b_0$$

$$ab = 2^n \cdot a_1 b_1 + 2^{n/2} \cdot (a_1 b_0 + a_0 b_1) + a_0 b_0$$

$$= 2^n \cdot a_1 b_1 + 2^{n/2} \cdot ((a_1 + a_0)(b_1 + b_0) - a_1 b_1 - a_0 b_0) + a_0 b_0$$

1

2

1

3

3

Karatsuba Multiplication

To multiply two n -bit integers a and b :

- Add two $\frac{1}{2}n$ bit integers.
- Multiply **three** $\frac{1}{2}n$ -bit integers, recursively.
- Add, subtract, and shift to obtain result.

$$\begin{aligned}
 a &= 2^{n/2} \cdot a_1 + a_0 \\
 b &= 2^{n/2} \cdot b_1 + b_0 \\
 ab &= 2^n \cdot a_1 b_1 + 2^{n/2} \cdot (a_1 b_0 + a_0 b_1) + a_0 b_0 \\
 &= 2^n \cdot a_1 b_1 + 2^{n/2} \cdot ((a_1 + a_0)(b_1 + b_0) - a_1 b_1 - a_0 b_0) + a_0 b_0
 \end{aligned}$$

1
2
1
3
3

Theorem. [Karatsuba-Ofman 1962] Can multiply two n -bit integers in $O(n^{1.585})$ bit operations.

$$T(n) \leq \underbrace{3T(n/2)}_{\text{recursive calls}} + \underbrace{\Theta(n)}_{\text{add, subtract, shift}} \Rightarrow T(n) = O(n^{\lg 3}) = O(n^{1.585})$$

Outline

- Overview
- Recap master theorem
- Counting inversions
- Closest pair of points
- Integer multiplication
- Matrix multiplication
- FFT (evaluate a polynomial of $n-1$ -th degree)

Matrix Multiplication

Matrix multiplication. Given two n -by- n matrices A and B , compute $C = AB$.

$$c_{ij} = \sum_{k=1}^n \underbrace{a_{ik} b_{kj}}$$

$$\begin{bmatrix} c_{11} & c_{12} & \cdots & c_{1n} \\ c_{21} & c_{22} & \cdots & c_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ c_{n1} & c_{n2} & \cdots & c_{nn} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix} \times \begin{bmatrix} b_{11} & b_{12} & \cdots & b_{1n} \\ b_{21} & b_{22} & \cdots & b_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ b_{n1} & b_{n2} & \cdots & b_{nn} \end{bmatrix}$$

Grade-school. $\Theta(n^3)$ arithmetic operations.

Q. Is grade-school matrix multiplication algorithm optimal?

Block Matrix Multiplication

$$\begin{bmatrix} 152 & 158 & 164 & 170 \\ 504 & 526 & 548 & 570 \\ 856 & 894 & 932 & 970 \\ 1208 & 1262 & 1316 & 1370 \end{bmatrix} = \begin{bmatrix} 0 & 1 & 2 & 3 \\ 4 & 5 & 6 & 7 \\ 8 & 9 & 10 & 11 \\ 12 & 13 & 14 & 15 \end{bmatrix} \times \begin{bmatrix} 16 & 17 & 18 & 19 \\ 20 & 21 & 22 & 23 \\ 24 & 25 & 26 & 27 \\ 28 & 29 & 30 & 31 \end{bmatrix}$$

Labels with arrows pointing to the blocks:

- C_{11} points to the first two columns of the result matrix.
- A_{11} points to the top-left 2x2 block of matrix A.
- A_{12} points to the top-right 2x2 block of matrix A.
- B_{11} points to the top-left 2x2 block of matrix B.
- B_{21} points to the bottom-left 2x2 block of matrix B.

$$C_{11} = A_{11} \times B_{11} + A_{12} \times B_{21} = \begin{bmatrix} 0 & 1 \\ 4 & 5 \end{bmatrix} \times \begin{bmatrix} 16 & 17 \\ 20 & 21 \end{bmatrix} + \begin{bmatrix} 2 & 3 \\ 6 & 7 \end{bmatrix} \times \begin{bmatrix} 24 & 25 \\ 28 & 29 \end{bmatrix} = \begin{bmatrix} 152 & 158 \\ 504 & 526 \end{bmatrix}$$

Matrix Multiplication: Warmup


To multiply two n -by- n matrices A and B :

- Divide: partition A and B into $\frac{1}{2}n$ -by- $\frac{1}{2}n$ blocks.
- Conquer: multiply 8 pairs of $\frac{1}{2}n$ -by- $\frac{1}{2}n$ matrices, recursively.
- Combine: add appropriate products using 4 matrix additions.

$$\begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$\begin{aligned} C_{11} &= (A_{11} \times B_{11}) + (A_{12} \times B_{21}) \\ C_{12} &= (A_{11} \times B_{12}) + (A_{12} \times B_{22}) \\ C_{21} &= (A_{21} \times B_{11}) + (A_{22} \times B_{21}) \\ C_{22} &= (A_{21} \times B_{12}) + (A_{22} \times B_{22}) \end{aligned}$$

Assume n is a power of 2



$$T(n) = \underbrace{8T(n/2)}_{\text{recursive calls}} + \underbrace{\Theta(n^2)}_{\text{add, form submatrices}} \Rightarrow T(n) = \Theta(n^3)$$


Fast Matrix Multiplication

Key idea. multiply 2-by-2 blocks with only **7 multiplications**. [Strassen 1969]

$$\begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \times \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$\begin{aligned} C_{11} &= P_5 + P_4 - P_2 + P_6 \\ C_{12} &= P_1 + P_2 \\ C_{21} &= P_3 + P_4 \\ C_{22} &= P_5 + P_1 - P_3 - P_7 \end{aligned}$$

$$\begin{aligned} P_1 &= A_{11} \times (B_{12} - B_{22}) \\ P_2 &= (A_{11} + A_{12}) \times B_{22} \\ P_3 &= (A_{21} + A_{22}) \times B_{11} \\ P_4 &= A_{22} \times (B_{21} - B_{11}) \\ P_5 &= (A_{11} + A_{22}) \times (B_{11} + B_{22}) \\ P_6 &= (A_{12} - A_{22}) \times (B_{21} + B_{22}) \\ P_7 &= (A_{11} - A_{21}) \times (B_{11} + B_{12}) \end{aligned}$$



Assume n is a power of 2

$$T(n) = \underbrace{7T(n/2)}_{\text{recursive calls}} + \underbrace{\Theta(n^2)}_{\text{add, subtract}} \Rightarrow T(n) = \Theta(n^{\log_2 7}) = O(n^{2.81})$$

Strassen's algorithm beats the ordinary algorithm on today's machines for $n \geq 32$ or so.

Fast Matrix Multiplication: Theory




FIG. 1. $\omega(t)$ is the best exponent announced by time τ .

Best known. $O(n^{2.376})$ [Coppersmith-Winograd, 1987]

Conjecture. $O(n^{2+\epsilon})$ for any $\epsilon > 0$.

Caveat. Theoretical improvements to Strassen are progressively less practical.

Homework

Exercises 2, 3, 7 in Chapter 5 "Divide and Conquer" by Jon Kleinberg and Eva Tardos. Addison-Wesley, 2005.