

ATLANTIC SMALL-MAMMAL: A DATASET OF COMMUNITIES OF RODENTS AND MARSUPIALS OF THE ATLANTIC FORESTS OF SOUTH AMERICA

Ricardo S. Bovendorp^{1, *}, Nacho Villar¹, Edson F. de Abreu-Junior², Carolina Bello¹, André L. Regolin¹, Alexandre R. Percequillo² and Mauro Galetti^{1, *}

1. Universidade Estadual Paulista (UNESP), Instituto de Biociências, Departamento de Ecologia, CP. 199, 13506-900 Rio Claro, São Paulo, Brazil
2. Departamento de Ciências Biológicas, Escola Superior de Agricultura “Luiz de Queiroz” (ESALQ), Universidade de São Paulo, Piracicaba, São Paulo, 13418-900, Brazil

* Correspondence and requests for materials should be addressed to Ricardo S. Bovendorp or Mauro Galetti (email: bovendorp@rc.unesp.br; mgaletti@rc.unesp.br).

Introduction

Elton’s seminal work on vole, mice and lemmings ([Elton 1924](#), [1942](#)) has inspired many generations of animal ecologists fascinated by the factors regulating small mammal populations and community dynamics. Since Elton’s work, small mammal ecology has been at the epicenter of the advancement of general theories on top-down vs. bottom up control, trophic cascades and population cycles ([Oksanen et al. 1981](#), [Lambin et al. 2002](#), [Korpimäki et al. 2003](#)), coexistence and habitat selection ([Brown and Davidson 1977](#), [Abramsky et al. 1990](#), [Morris 1999](#), [Morris et al. 2000](#)), kin selection and maternal effects ([Chitty 1960](#), [Boonstra and Boag 1987](#), [Lambin and Krebs 1991](#), [1993](#)), the evolution of social and mating systems ([Ostfeld 1990](#), [Sherman and Wolff 2007](#), [Odden et al. 2014](#)), and many others. Ecologists have also used small mammals as model systems for the development of new field and statistical tools ([Otis et al. 1978](#), [Nichols and Pollock 1990](#), [Kendall et al. 1995](#)), but also for applied ecological issues such as the impact of habitat fragmentation and destruction on vertebrate populations and communities ([Andreassen et al. 1998](#), [Pardini et al. 2010](#)) and the ecology of infectious diseases ([Allan et al. 2003](#), [LoGiudice et al. 2003](#)).

One area where small mammal ecology has been less informative is in understanding macroecological patterns of biodiversity, population dynamics and community assembly. For such studies, a large quantity of data from communities across ecosystems is needed so as

to inquire questions about relationships between organisms and their environment at large spatial scales and clarify generalizations in pattern and process ([Brown 1971](#), [May 1988](#), [Purvis and Hector 2000](#), [Karp et al. 2012](#)). Large community databases are also important for biodiversity hotspots regions because of the urgency in conservation actions. However, perhaps because historically temperate regions have been the stronghold of small mammal ecologists, information on the ecology of small mammals from tropical regions has received less attention. For example, Brown's 1995 notorious book on Macroecology ([Brown 1995](#)) begins with an example of predicted range shifts in small mammal communities on mountain ranges, but, like most of the rest of the examples in the book, patterns are almost exclusively derived from small mammal desert communities in North America. Recently some large datasets have become available for mammal and bird traits ([Wilman et al. 2014](#)) and fruit-frugivore interactions ([Bello et al. 2017](#)) on biodiversity hotspots, but no publicly available database exists on non-volant small mammals for such regions.

One of the region with the highest diversity of small mammals is the Atlantic forest of South America ([Galindo-Leal and Câmara 2003](#)). The Atlantic forest hosts 119 species of small mammals represented by 22 species of marsupials (Family Didelphidae) and 97 species of small rodents (families Caviidae, Cricetidae, Ctenomyidae, Echimyidae, Cricetidae [=Muridae] and Sciuridae) ([Costa et al. 2000](#), [Paglia et al. 2012](#)). The Atlantic forest was one of the largest rainforests of South America, originally covering around 150 million hectares along the Brazilian coast; today fragmented patches cover less than 12% of its original area ([Ribeiro et al. 2009](#)). Nonetheless, the remaining forest hosts a larger number of endemic species and is considered one of the five major hotspots in the world for conservation ([Myers et al. 2000](#), [Visconti et al. 2011](#)).

Here we compiled a dataset comprising 136 studies covering 300 locations, that sampled small mammals in the Atlantic forest of Brazil and Paraguay (Figure 1). Ecological inventories of small mammals in the Atlantic forest started in 1945 ([Davis 1945](#)) and have been carried out in all major biogeographical region in the Atlantic forest (Figure 2). The ATLANTIC-SMALL MAMMAL represents the largest dataset of inventories of small mammal communities for the Neotropical region. We present in this dataset the species composition, richness, and relative abundance (captures/trap nights), which are amenable for population and community ecology studies.

METADATA

CLASS I. DATA SET DESCRIPTORS

A. Data set identity:

Title: ATLANTIC-SMALL MAMMAL: a dataset of communities of rodents and marsupials of the Atlantic Forests of South America

B. Data set and metadata identification codes:

Suggested Data Set Identity Codes: ATLANTIC-SM_Capture.csv, ATLANTIC-SM_Reference.csv, ATLANTIC-SM_Study_site.csv

C. Data set description

Principal Investigators:

1. Ricardo S. Bovendorp, Carolina Bello, André L. Regolin, Nacho Villar, and Mauro Galetti,

Universidade Estadual Paulista (UNESP), Instituto de Biociências, Departamento de Ecologia, CP. 199, Rio Claro, São Paulo, 13506-900, Brazil

2. Edson F. de Abreu-Jr. and Alexandre R. Percequillo

Departamento de Ciências Biológicas, Escola Superior de Agricultura “Luiz de Queiroz” (ESALQ), Universidade de São Paulo, Piracicaba, São Paulo, 13418-900, Brazil

Abstract: The contribution of small mammal ecology to the understanding of macroecological patterns of biodiversity, population dynamics and community assembly has been hindered by the absence of large datasets of small mammal communities from tropical regions. Here we compile the largest dataset of inventories of small mammal communities for the Neotropical region. The dataset reviews small mammal communities from the Atlantic forest of South America, one of the regions with the highest diversity of small mammals and a global biodiversity hotspot, though currently covering less than 12% of its original area due to anthropogenic pressures. The dataset comprises 136 references from 300 locations covering

seven vegetation types of tropical and subtropical Atlantic forests of South America, and presents data on species composition, richness, and relative abundance (captures/trap-nights). One paper was published more than 70 years ago but 80% of them were published after 2000. The dataset comprises 53,518 individuals of 124 species of small mammals, including 30 species of marsupials and 94 species of rodents. Species richness averaged 8.2 species (1 - 21) per site. Only two species occurred in more than 50% of the sites (the common opossum, *Didelphis aurita* and black-footed pigmy rice rat *Oligoryzomys nigripes*). Mean species abundance varied 430-fold, from 4.3 to 0.01 individuals/trap-night. The dataset also revealed a hyper-dominance of 22 species that comprised 78.29% of all individuals captured, with only seven species representing 44% of all captures. The information contained on this dataset can be applied in the study of macroecological patterns of biodiversity, communities and populations, but also to evaluate the ecological consequences of fragmentation and defaunation, and predict disease outbreaks, trophic interactions and community dynamics in this biodiversity hotspot.

D. Key words: biodiversity hotspot, tropical forest, biodiversity dataset, rodent, marsupial, small mammals, species richness, communities, hyper-dominance, live traps.

Description

The dataset is restricted to seven vegetation types in the domain of the Atlantic Forest that cover tropical and subtropical forests in Brazil, Paraguay and Argentina ([Ribeiro et al. 2009](#), [Morrone 2014](#)) (Figure 1). Seventy percent of the references are articles in peer reviewed journals and 27% are theses. The duration of the studies varied from one to 75 months. Details of methods can be found in [Bovendorp et al. \(2017\)](#).

We here report the raw data from 300 small mammal communities present in 136 studies and 300 locations. Of these locations, 46% are located in protected areas. Most studies were carried out in Ombrophilous (66%) and Semideciduous (21%) forests, which are the dominant vegetation type of this biome ([Ribeiro et al. 2009](#)). The forest sizes varied from 0.15 to 791,652 ha (average= 11,886 ha, median = 390 ha). Altitude varied from sea level to 2,700 m (Parque Nacional de Caparó, Rio de Janeiro). Rainfall varied from 188 mm/year to 4,200 mm/year.

The dataset included 53,518 marked individuals, representing 124 species of non-volant small mammal classified in two orders: Didephimorphia, with 13 genera and 30 species; and

Rodentia with 37 genera and 94 species. Species richness ranged from one species to 23 species. Only two species occurred in more than 50% of the localities (Figure 2).

Most of species are classified as less concern, but four species are considered endangered (Table 1). We recorded that 24% of the sites have at least one invasive species (*Rattus rattus*, *R. norvegicus* and *Mus musculus*). One invasive species (*Rattus rattus*) comprised 1.4% of all individuals captured.

CLASS II. RESEARCH ORIGIN DESCRIPTORS

A. Overall project description

Identity: A compilation of small mammals captured in the Atlantic Forest of Brazil, Argentina and Paraguay.

Period of Study: Dates of source publications range from 1945–2016.

Objectives: Our objectives for compiling the data were: (1) to summarize information about small mammal inventories in the Atlantic Forest biome, focusing on species richness and abundance; and (2) to identify the gaps on the knowledge of small mammal communities of the Atlantic Forest biome to guide future sampling efforts. Our dataset represents a first attempt to obtain a large-scale inventory of small mammals, with potential applications in macroecological studies ([Galetti et al. 2013](#)), conservation strategies ([Banks-Leite et al. 2014](#)), and population and community ecology research ([Pacheco et al. 2013](#)).

Abstract: Same as above.

Sources of funding: The compilation of this data set was supported by grants and scholarships from Programa BIOTA from Fundação de Amparo à Pesquisa do Estado de São Paulo (São Paulo Research Foundation; FAPESP), numbers 2008/55483-8, 2013/25441-0, 2014/18800-6 and 2014/01986-0 (RBS); 2013/50421-2, 2014/01986-0 and 2014/50434-0 (MG); 2015/11521-7 (NV); 2013/22492-2(CB) and CNPq 153423/2016-1 (ALR). MG and ARP receive research grants and fellowships from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazilian Research Council, CNPq).

B. Specific subproject description

Site description: The Atlantic Forest is an important biodiversity hotspot in South America (Willis et al. 2007). It comprises tropical and subtropical evergreen and semideciduous forests with highly heterogeneous environmental conditions (Morellato and Haddad 2000). It supports up to 8% of the world's total species richness and one of the highest rates of endemism in the world (Myers et al. 2000). The Atlantic Forest supports at least 15,519 plant species (3,343 trees) (BFG 2015), 891 bird species (Moreira-Lima 2014), 543 amphibians (Haddad et al. 2013), 200 reptiles (Bérnils and Costa 2015), 350 fishes (MMA 2010), and 298 mammals (Paglia et al. 2012) including 22 species of marsupials (Paglia et al. 2012) and 105 species of rodents (Patton et al. 2015).

Seventy two percent of the Brazilian population lives in areas of the Atlantic Forest domain (~145 million people) (IBGE 2013). Therefore, many past and present anthropogenic activities such as logging, sugarcane and coffee plantations, agribusiness, industrialization and unplanned urban expansion have contributed to the ecosystem deterioration. By now, the conservation of Atlantic Forest is critical, the natural remnants accounting for only 12% of the original biome and above 80% of these remnants are within < 50 ha fragments (Ribeiro et al. 2009), and 88% of the fragments are defaunated from large mammals (Jorge et al. 2013).

Data compilation: Data was obtained from published literature including 136 papers, theses, scientific conferences, technical reports, and our own unpublished data. We searched for potential studies in the following sources: (i) online academic databases (e.g., ISI Web of Knowledge, Google Scholar, Scielo, Scopus, JStore) (ii) digital libraries of state and federal Brazilian universities, (iii) references cited in literature, and (iv) email contacts with local experts. The searches were performed with the following key words: small mammal(s), survey(s), inventory(ies), and Atlantic forest; searches were conducted in English, Portuguese and Spanish.

Research Methods: We included in this database, studies that reported sampling of small mammal species, trapping methods, trap size, sampling design, strata sampled, effort and abundance. Lack of information were filled with NA. However, some studies that did not provide a substantial detailed information about captures were excluded at early stages of the

database. We also included information from the geographical location (when provided, we tabulated latitude, longitude, locality, municipality, state and country). Small mammals information (occurrence and abundance) were compiled from (Davis 1945, Fonseca and Robinson 1990, Herrmann 1991, Stallings et al. 1991, Bergallo 1994, Paglia et al. 1995, Grelle 1996, Palma 1996, Graipel et al. 1997, Stevens and Husband 1998, D'Andrea et al. 1999, Gentile and Fernandez 1999, Lessa et al. 1999, Moura 1999, Vieira 1999, Abel et al. 2000, Barros-Battesti et al. 2000, Passamani 2000, Passamani et al. 2000, Briani et al. 2001, Silva 2001, Bonvicino et al. 2002, Feliciano et al. 2002, Gheler-Costa et al. 2002, Olifiers 2002, Pires et al. 2002, Rosa 2002, Graipel et al. 2003, Grelle 2003, Marinho 2003, Moura 2003, Vieira and Monteiro Filho 2003, Castro and Fernandez 2004, Nieri-Bastos et al. 2004, Pardini 2004, Paresque et al. 2004, Rocha 2004, Santos et al. 2004, Vieira et al. 2004, Cherem 2005, Gaspar 2005, Horn 2005, Oliveira et al. 2005, Passamani et al. 2005, Pedó 2005, Atique 2006, Carlos 2006, Graipel et al. 2006, Graipel and Filho 2006, Pardini and Umetsu 2006, Puttker et al. 2006, Uchôa 2006, Umetsu et al. 2006, Vera Y Conde and Rocha 2006, Caceres et al. 2007, Caldara-Junior and Leite 2007, D'Andrea et al. 2007, Finokiet 2007, Iob 2007, Langone 2007, Pena 2007, Scheibler and Christoff 2007, Silva et al. 2007, Umetsu and Pardini 2007, Amaral 2008, Bezerra and de Moraes 2008, Bressiani and Graipel 2008, Cademartori et al. 2008, Modesto et al. 2008, Moreira et al. 2008, Pinheiro and Geise 2008, Prevedello et al. 2008, Salvador and Fernandez 2008, Steiner-Souza et al. 2008, Umetsu et al. 2008, Antunes et al. 2009, Asfora and Pontes 2009, Moreira et al. 2009, Naxara et al. 2009, Passamani and Ribeiro 2009, Pereira and Geise 2009, Pessôa et al. 2009, Pinto et al. 2009, Vieira et al. 2009, Abreu et al. 2010, Antunes et al. 2010, Galiano 2010, Lima et al. 2010, Paise 2010, Pedó et al. 2010, Peters et al. 2010, Pinotti 2010, Andreazzi et al. 2011, Martins 2011, Melo et al. 2011, Passamani and Fernandez 2011a, Passamani and Fernandez 2011b, Rocha et al. 2011, Amaral et al. 2012, Cerboncini 2012, Dantas-Torres et al. 2012, Leiner and Silva 2012, Loretto 2012, Martin et al. 2012, Mesquita and Passamani 2012, Rosa 2012, Silveira 2012, Albuquerque et al. 2013, Behs 2013, Ferracioli 2013, Gheler-Costa et al. 2013, Lima et al. 2013, Luza et al. 2013, Machado et al. 2013, Milagres et al. 2013, Pacheco et al. 2013, Pereira et al. 2013, Prevedello 2013, Quintela et al. 2013, Rodarte 2013, Siqueira et al. 2013, Abreu and de Oliveira 2014, Balieiro et al. 2014, Cerboncini et al. 2014, de la Sancha 2014, Grazzini 2014, Maestri and Marinho 2014, Mochi-Jr 2014, Oliveira et al. 2014, Rosalino et al. 2014, Teixeira et al. 2014, Tortato et al. 2014, Balieiro et al. 2015, Regolin et al. 2015, Rocha-Mendes et al. 2015, Bergallo 1991, Regolin and Cherem Unpubl. Data-b, a, Souza and Galetti) and our own unpublished data.

Taxonomic data: We used animal taxonomic information for marsupials according to Voss and Jansa (2009) and for the rodent species Patton et al. (2015). We added one column with the current scientific name based on Patton et al. (2015) where this association was possible (*e.g.*, changes on generic attribution, species group name synonymy, species identified as sp. or spp. were sometimes identifiable by geography). In addition, we recorded the basic publishing information of each study (author, title, year, journal, volume, number of issue, pages, place of publication, DOI to the document and others relevant information).

C. Data Limitations and Potential Enhancements

We recognize that documenting all species of small mammals present in megadiverse ecosystems is a challenging task for several reasons, mostly related to species biology and their sampling. In general, although small bodied animals, species of rodents and marsupials occupy quite different niches and exhibit distinct and diversified life-histories, paralleling other more conspicuous medium and large size mammals. As small size mammals, these species are prone to exhibit seasonal population fluctuations in response to wet and dry seasons: during wet seasons, populations are usually higher, benefiting from the higher productivity of forests. Population sizes of these mammals also respond to more drastic events, either climatic (as El Niño and La Niña) or biological (as the flowering and fructification of bamboo), presenting explosive responses and attaining huge sizes (Jaksic and Lima 2003). Therefore, there are marked population fluctuations, with periods of high abundance and likely density, and periods with small abundances. Ideally then, sampling should be performed over different periods to characterize these fluctuations, but on our dataset sampling is quite heterogeneous, limiting controlled comparisons.

Being diverse on their life-histories and habits, small mammals are challenging to sample and survey (Voss and Emmons 1996, Bovendorp et al. 2017), thus field sampling strongly constrained data assembly. Usual sampling methods with conventional live-traps (such as Sherman and Tomahawk traps) are quite selective, as they employ baits that attract only a subset of the assemblage, usually terrestrial/scansorial omnivorous species. These traps are usually set on the ground level and on the understory, with very few surveys of the canopy (Vieira and Monteiro 2003, Lambert et al. 2006) or aquatic habitats (as river courses, river and lake). Pitfall traps are quite effective and less selective (Hice and Schmidly 2002), but if buckets are small only young individuals will be sampled, and young are more difficult to identify than adults. These methods (conventional and pitfall traps) should be used obligatorily,

to enhance trapping success, not only quantitatively but also qualitatively, along with active search (night census, along with night hunting; see (Voss and Emmons 1996, Voss et al. 2001), an aspect often neglected by mammalogists. Therefore, sampling of small mammals is complex and demands a combination of survey methods, along with consistent sampling effort, that can be obtained only with large number of traps and/or long-term surveys (see for a review on length of survey): a large number of traps set for a long number of days in a standardized design will produce a high sampling effort, that usually produce more complete lists of species, along with information on abundance and density. Most surveys in our database are heterogeneous, with sampling efforts quite unstandardized (e.g. sampling design, number of traps, sampling days and sampling period being highly variable), precluding more comprehensive diagnosis of diversity throughout the Atlantic Forest. In addition, the length of studies is highly variable, and most of them are conducted through one or two years, limiting the ability to obtain reliable population estimates.

Some aspects of the biology of small mammals also make them difficult to sample, as some species are more elusive, exploring their surroundings cautiously and in a small radius, and generally avoiding traps (trap-shy), especially conventional traps. On the other hand some species are quite inquisitive, actively prospecting the surrounding and being positively attracted to the traps; in some cases the reward offered by the bait is so valuable that in capture-recapture studies, previously captured individuals kept returning to the trap. This precludes from other individuals from being caught, biasing the sample.

Another consistent limitation is the uneven sampling throughout the Atlantic Forest, as extensive areas of the biome remain understudied. States of Santa Catarina, Paraná, Bahia, Alagoas, Pernambuco and Paraíba have been subject to very few studies, while Sergipe and Rio Grande do Norte are virtually not studied. The Atlantic Forest of Argentina and Paraguay also present few available studies on small mammal assemblages. One major aspect is the definition of the Atlantic Forest employed by several authors recovered on this database. Several sampling areas are located on transitional landscapes, and the lists of species include species that are typical of other biomes, such as Pampas in southern Brazil, Cerrado in southeastern Brazil, and Caatinga in northeastern Brazil.

Considering that a study was performed in Atlantic Forest, with an adequate sampling, both qualitative and quantitative, at the appropriate time, some limitation also is apparent on species identification, the most important aspect on the diagnosis of an assemblage. Being quite similar externally, small mammals are difficult to identify. The application of a specific name to an individual or a population is a hypothesis, and as such must be testable under the scientific

paradigm: other scientists should be able to examine the same specimens and confirm or refute previous hypothesis. However, many studies did not report the collection of voucher specimens or genetic samples for a posterior species confirmation, employing new conceptual and/or methodological approaches. Another issue is the use of the appropriate nomenclature: scientific names change through time, with names of the species group taxa being replaced by other names or transferred to other genera. These efforts should be acknowledged by specialists working on the ecology of small mammals, to follow the most current taxonomic conventions. Despite these impediments, we recognize the massive effort of biologists who carried out these studies. Collectively, it represents the best available dataset of the communities of small mammals in a tropical forest biome. We expect that, based on this database, the community of scientists will be able to recognize some patterns and be able to determine: i) priority areas to sample small mammals; ii) minimum trapping effort to obtain consistent information on small mammal assemblages; iii) most adequate periods for sampling these species along the latitudinal and longitudinal extension of the biome; iv) lists of voucher specimens and reference material, and the institutions where they are housed; v) a minimal sampling protocol that will standardize the study of small mammals in the Atlantic Forest and other tropical biomes.

CLASS III. DATA SET STATUS AND ACCESSIBILITY

A. Status

Latest update: March 2017

Latest Archive date: March 2017

Metadata status: Last update 30 March 2017, version submitted

Data verification: Data is mostly from published sources. We searched for extreme values, corrected any transcription errors and homogenized the taxonomic information.

B. Accessibility

Contact person: Ricardo S. Bovendorp or Mauro Galetti, Departamento de Ecologia, Universidade Estadual Paulista, Rio Claro, São Paulo, 13506-900, Brazil. E-mail: bovendorp@rc.unesp.br; mgaletti@rc.unesp.br

Copyright restrictions: None.

Proprietary restrictions: Please cite this data paper when the data are used in publications. We also request that researchers and teachers inform us of how they are using the data.

Costs: None.

CLASS IV. DATA STRUCTURAL DESCRIPTORS

A. Data Set File

Identity: ATLANTIC_SM_Study_Site.csv

ATLANTIC_SM_Reference.csv

ATLANTIC_SM_Capture.csv

Size: 2620 records, 496 KB

Format and storage mode: comma-separated values (.csv)

Header information: See column descriptions in section B.

Alphanumeric attributes: Mixed.

Data Anomalies: If no information is available for a given record, this is indicated as 'NA'.

B. Variable information

1) Table 2. Site Information

2) Table 3. Reference Information

3) Table 4. Capture Information

CLASS V. SUPPLEMENTAL DESCRIPTORS

A. Data acquisition

1. **Data request history:** None

2. **Data set updates history:** None

3. **Data entry/verification procedures**

G. History of data set usage

Bovendorp et al. (2017) used the small mammal dataset in order to determine the influence of trap configuration, trap type, and sampling effort on measures of species richness and abundance.

ACKNOWLEDGMENTS

RSB, MG and NV received FAPESP fellowships and grants (2008/55483-8, 2013/25441-0, 2014/18800-6 and 2014/01986-0 (RBS); 2013/50421-2, 2014/01986-0 and 2014/50434-0

(MG); 2015/11521-7 (NV), 2013/22492-2(CB)) and CNPq 153423/2016-1 (ALR). MG and ARP received research grants and fellowships from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (Brazilian Research Council, CNPq). The authors would like to thank Laís Rodrigues and Thaís Labecca for helping in the early phase of this dataset, and M.C. Ribeiro and C. Bello for comments on the manuscript. This work is dedicated to R. Cerqueira for his contribution on the advancement of Neotropical mammals.

TABLES

Table 1. IUCN conservation status of small mammal species reported in the ATLANTIC dataset. CR= Critically, DD= Data Deficient, EN=Endangered, LC= Least Concern, NT= Near Threatened, VU=Vulnerable

IUCN conservation status	Small mammals
Critically endangered (CR)	1
Endangered (EN)	4
Vulnerable (VU)	2
Near Threatened (NT)	3
Least Concern (LC)	79
Data Deficient (DD)	10
Not evaluated (NE)	7

Table 2. Site information. Description of the fields related with the study sites information.

Type of information	Field	Description	Levels	Example
SITE INFORMATION	ID	Field that link the study site to the complete information table	Sm01-Sm305	Sm01
	Reference_number	The reference number which report small mammal communities	1-166	1
	Country	Country of the study		Brazil
	State	State of the study		SP
	Municipality	Municipality of the study		Cananéia
	Study_location	Specific location of the study		Parque Estadual Ilha do Cardoso, Cananéia, SP
	Latitude	Decimal coordinates		-25.06930556
	Longitude	Decimal coordinates		-47.9200356
	Precision	Precision of the given coordinate	Precise Not-Precise	Precise
	Size	Forest size (area in hectares)		15100
	Altitude	Altitude (m) above sea level		60

	Annual_rainfall	Anual precipitation (mm)		500
	Vegetation type	Atlantic forest vegetation type	Ombrophilous Semideciduous Deciduous Araucaria Restinga Stepe	Ombrophilous Forest
	Protect_area	If the study is inside to the protected areas		Yes
	Matrix	Predominant vegetation surrounding the forest patch		Sugarcane plantations

Table 3. Reference information. Description of the fields related with reference information of each study.

Type of information	Field	Description	Levels	Example
REFERENCE INFORMATION	Reference_number	The reference number which report small mammal communities	1-136	10
	Reference	Extended information of the reference		Cáceres, N. C., Ghizoni-Jr, I. R., & Graipel, M. E. (2002). Diet of two marsupials, <i>Lutreolina crassicaudata</i> and <i>Micoureus demerarae</i> , in a coastal Atlantic Forest island of Brazil. <i>Mammalia</i> , 66(3), 331–340.
	Publication_year	Year of publication		2001
	Type of publication	Type of publication	Article Book Thesis Unpublished	Article

Table 4. Capture information. Description of the fields related with the complete capture information.

Type of information	Field	Description	Levels	Example
CAPTURE INFORMATION	ID	Field that links the study site to the complete information table	Sm01-Sm305	sm1
	Reference_number	The reference number which reports small mammal communities	1-136	10
	Month_start	Month when the survey started	January-December	March
	Year_start	Year when the survey started		1999
	Month_finish	Month when the survey finished	January-December	July
	Year_finish	Year when the survey finished		2000
	Total_of_months	Number of months used to sample small mammal		16
	Sampling_habitat	Habitat location that was sampled in the patch or site	Interior Edge Rivers Etc....	Interior
	Sampling_design	Design used to capture small mammal	Aleatory Grid Line transect	Grid
	Effort	Total traps-type per night used to capture small mammal		2692
	Method	Trap type used to capture small mammal	Live traps Pitfall traps Catwalks Manual	Live traps

Trap_size	Size of traps used to capture small mammal	S - small - ($\leq 23 \times 7 \times 8$ cm) M - medium - ($> 23 \times 7 \times 8$ cm and $< 38 \times 10 \times 12$ cm) L - large - ($\geq 38 \times 10 \times 12$ cm) 20, 30, 35, 40, 60, 100L-liters	M
Strata_sampling	Trap height for sampling small mammal	Ground Understory Canopy	Ground
Order	Order taxonomic classification		Didelphi morphia
Genus	Genus taxonomic classification		<i>Metachirus</i>
Species_name_on_paper	Species taxonomic classification reported on paper		<i>Metachirus nudicaudatus</i>
Actual_species_name	Species taxonomic classification reviewed by specialist		<i>Metachirus nudicaudatus</i>
Species_origin	If the species is native or introduced in Brazil	Native Invasive	Native
Individuals_captured	Total of individuals captured for every trap type	Individuals/effort (trap-nights)	1
Abundance_(ind/trap-nights)	Number of individuals captured by trap-night		0,03
Voucher_Specimens	Paper related voucher specimens in collection or museum	Yes No	Yes

Figures:




Fig 1. Distribution of the small mammal assemblage records on the Atlantic Forest biome. The gray colors show the domain of the Atlantic Forest and the green colors show the remaining Atlantic forest patches (Huang et al. 2007, Ribeiro et al. 2009). The dots show the locations of the original studies reporting the small mammal assemblages.




Fig 2. Total number of sites occupied by every small mammal species. Blue bars indicate species from the Order Rodentia and orange bars species from the Order Didephimorphia.




Fig 3. Mean abundance per site (mean number of individuals captured per trap night) for every small mammal species. The narrow bars show standard errors, blue bars indicate species from the Order Rodentia and orange bars species from the Order Marsupialia.




Fig 5. Number of studies conducted by major vegetation type and year in the Atlantic forest. The intensity of the color shade indicates the number of studies. All studies were carried out between 1981 and 2014, with a single exception of the 1942 Davis' study. From the 7 major vegetation types, the Ombrophilous forest is the best studied.

Literature Cited

- Abel, I. S., G. Marzagao, N. H. Yoshinari, and T. T. Schumaker. 2000. Borrelia-like spirochetes recovered from ticks and small mammals collected in the Atlantic Forest Reserve, Cotia county, State of Sao Paulo, Brazil. *Mem Inst Oswaldo Cruz* **95**:621-624.
- Abramsky, Z., M. L. Rosenzweig, B. Pinshow, J. S. Brown, B. Kotler, and W. A. Mitchell. 1990. Habitat selection: an experimental field test with two gerbil species. *Ecology* **71**:2358-2369.
- Abreu, M., A. Wieliczko, A. Mesquita, and E. Vieira. 2010. Consumo de pequenos mamíferos por canídeos simpátricos do sul do Brasil: sobreposição de nichos e seleção de presas. *Neotropical Biology and Conservation* **5**:16-23.
- Abreu, M. S. L., and L. R. de Oliveira. 2014. Patterns of arboreal and terrestrial space use by non-volant small mammals in an Araucaria forest of southern Brazil. *Anais Da Academia Brasileira De Ciencias* **86**:807-819.
- Albuquerque, H. G., P. F. Martins, F. S. Pessôa, T. C. Modesto, J. L. Luz, D. S. L. Raíces, N. C. Ardente, I. C. M. Lessa, N. Attias, T. Jordão-Nogueira, M. C. Enrici, and H. d. G. Bergallo. 2013. Mammals of a forest fragment in Cambuci municipality, State of Rio de Janeiro, Brazil. *Check List* **9**:1505-1509.
- Allan, B. F., F. Keesing, and R. S. Ostfeld. 2003. Effect of forest fragmentation on Lyme disease risk. *Conservation Biology* **17**:267-272.
- Amaral, H. H. D. O. M. 2008. Ecologia de Phthiraptera, Siphonaptera e Acari (Ixodidae) de pequenos roedores e marsupiais do Parque Estadual da Pedra Branca, Rio de Janeiro, Brasil. Universidade Federal Rural do Rio de Janeiro.
- Amaral, H. H. d. O. M., A. B. Silva, V. Gomes, I. P. F. Quinelato, V. F. Morelli-Amara, N. Serra-Freire, R. W. Carvalho, and A. G. Carvalho. 2012. Roedores e marsupiais capturados no Parque Estadual da Pedra Branca, Rio de Janeiro: distribuição e relação com o ambiente. *Uniaubeu* **5**:158-180.
- Andreassen, H. P., K. Hertzberg, and R. A. Ims. 1998. Space-use responses to habitat fragmentation and connectivity in the root vole *Microtus oeconomus*. *Ecology* **79**:1223-1235.
- Andreazzi, C. S. d., V. Rademaker, R. Gentile, H. M. Herrera, A. M. Jansen, and P. S. D'Andrea. 2011. Population ecology of small rodents and marsupials in a semi-deciduous tropical forest of the southeast Pantanal, Brazil. *Zoologia (Curitiba)* **28**:762-770.
- Antunes, P. C., M. A. A. Campos, L. G. R. Oliveira-Santos, and M. E. Graipel. 2009. Population dynamics of *Euryoryzomys russatus* and *Oligoryzomys nigripes* (Rodentia, Cricetidae) in an Atlantic forest area, Santa Catarina Island, southern Brazil. *Biotemas* **22**:143-151.
- Antunes, P. C., M. A. A. Campos, L. G. R. Oliveira-Santos, and M. E. Graipel. 2010. Population dynamics of *Akodon montensis* (Rodentia, Cricetidae) in the Atlantic forest of southern Brazil. *Mammalian Biology* **75**:186-190.
- Asfora, P. H., and A. R. M. Pontes. 2009. The small mammals of the highly impacted North-eastern Atlantic Forest of Brazil, Pernambuco Endemism Center. *Biota Neotrop.* **9**:31-35.
- Atique, M. S. 2006. Estudo de uma comunidade de pequenos mamíferos em uma área de Floresta Atlântica. Universidade Regional de Blumenau.
- Balieiro, P., D. Behs, M. E. Graipel, S. S. Dornelles, L. M. Tiepolo, and M. J. Cremer. 2015. Riqueza de mamíferos não voadores em florestas de restinga do Sul do Brasil. *Mastozoologia Neotropical* **22**:367-373.
- Balieiro, P., M. L. P. Miranda, S. S. Dornelles, A. Soares, I. Hass, and I. J. Sbalqueiro. 2014. Pequenos mamíferos não voadores (Rodentia e Didelphimorphia) do Parque Estadual Acaraí, São Francisco do Sul, Santa Catarina, Brasil. *Boletim da Sociedade Brasileira de Mastozoologia* **71**:37-41.

- Banks-Leite, C., R. Pardini, L. R. Tambosi, W. D. Pearse, A. A. Bueno, R. T. Bruscagin, T. H. Condez, M. Dixo, A. T. Igari, A. C. Martensen, and J. P. Metzger. 2014. Conserving Brazil's Atlantic forests response. *Science* **346**.
- Barros-Battesti, D. M., R. Martins, C. R. Bertim, N. H. Yoshinari, V. L. N. Bonoldi, E. P. Leon, M. Miretzki, and T. T. S. Schumaker. 2000. Land fauna composition of small mammals of a fragment of Atlantic forest in the State of Sao Paulo, Brazil. *Revista Brasileira De Zoologia* **17**:241-249.
- Behs, D. 2013. Influência da heterogeneidade ambiental na composição de comunidades de roedores e marsupiais em áreas de Mata Atlântica, no sul de Santa Catarina. Universidade do Extremo Sul Catarinense, Criciúma.
- Bello, C., M. Galetti, D. Montan, M. A. Pizo, T. C. Mariguela, L. Culot, F. Bufalo, F. Labecca, F. Pedrosa, R. Constantini, C. Emer, W. R. Silva, F. R. da Silva, O. Ovaskainen, and P. Jordano. 2017. ATLANTIC-FRUGIVORY: A plant-frugivore interaction dataset for the Atlantic Forest. *Ecology*.
- Bergallo, H. D. 1994. Ecology of a small mammal community in an Atlantic forest area in southeastern Brazil. *Studies On Neotropical Fauna And Environment* **29**:197-217.
- Bergallo, H. G. 1991. Dinâmica populacional, área de vida, parasitismo e mutualismo de pequenos mamíferos da Estação Ecologica da Juréia, SP. Universidade Estadual de Campinas, Campinas.
- Bérnils, R. S., and H. C. Costa. 2015. Brazilian reptiles. *in* R. S. Bérnils and H. C. Costa, editors. List of species. Sociedade Brasileira de Herpetologia, São Paulo.
- Bezerra, A. C., and D. A. a. de Moraes. 2008. Alimentação de pequenos mamíferos em um fragmento de Mata Atlântica nordestina (Mata do Engenho de Coimbra, Usina Serra Grande, Ibateguara, AL). XVI CONIC - Congresso de iniciação científica da UFPE, Pernambuco.
- BFG, T. B. F. g. 2015. Growing knowledge: an overview of seed plant diversity in Brazil. *Rodriguésia* **66**:1085-1113.
- Bonvicino, C. R., S. M. Lindbergh, and L. S. Maroja. 2002. Small non-flying mammals from conserved and altered areas of Atlantic forest and Cerrado: comments on their potencial use for monitoring environment. *Braz. J. Biol.* **62**:765-774.
- Boonstra, R., and P. T. Boag. 1987. A test of the Chitty hypothesis: inheritance of life-history traits in meadow voles *Microtus pennsylvanicus*. *Evolution* **41**:929-947.
- Bovendorp, R. S., R. McCleery, and M. Galetti. 2017. Optimising sampling methods for small mammal communities in Neotropical rainforests. *Mammal Review* **47**:148-158.
- Bressiani, V. B., and M. E. Graipel. 2008. Comparação de métodos para captura da cuíca-d'água, *Chironectes minimus* (Zimmerman, 1780) (Mammalia, Dipelphidae) no sul do Brasil. *Mastozoologia Neotropical* **15**:33-39.
- Briani, D. C., R. T. Santori, M. V. Vieira, and N. Gobbi. 2001. Mamíferos não-voadores de um fragmento de mata mesófila semidecídua, do interior do Estado de São Paulo, Brasil. *HOLOS Environment* **1**:141-149.
- Brown, J. H. 1971. Mammals on mountaintops: nonequilibrium insular biogeography. *The American Naturalist* **105**:467-574.
- Brown, J. H. 1995. *Macroecology*. University of Chicago Press, Chicago.
- Brown, J. H., and D. W. Davidson. 1977. Competition between seed-eating rodents and ants in desert ecosystems. *Science* **196**:880-882.
- Caceres, N. C., M. R. Bornschein, W. H. Lopes, and A. R. Percequillo. 2007. Mammals of the Bodoquena mountains, southwestern Brazil: an ecological and conservation analysis. *Revista Brasileira De Zoologia* **24**:426-435.
- Cademartori, C. V., M. Saraiva, C. Saraiva, and J. Andrades-Miranda. 2008. Nota sobre a fauna de pequenos roedores em mosaico antropogênico com remanescente florestal do domínio Mata Atlântica, sul do Brasil. *Biodiversidade Pampeana* **6**:34-38.

- Caldara-Junior, V., and Y. L. R. Leite. 2007. Uso de habitats por pequenos mamíferos no Parque Estadual da Fonte Grande, Vitória, Espírito Santo, Brasil. *Bol. Mus. Biol. Mello Leitão* **21**:57-77.
- Carlos, H. S. A. 2006. Uso de corredores florestais e matriz de pasto por pequenos mamíferos em Mata Atlântica do Estado do Rio de Janeiro. Universidade Federal de Minas Gerais.
- Castro, E. B. V., and F. A. S. Fernandez. 2004. Determinants of differential extinction vulnerabilities of small mammals in Atlantic forest fragments in Brazil. *Biological conservation* **119**:73-80.
- Cerboncini, R. A. S. 2012. Respostas de pequenos mamíferos ao efeito de borda da ferrovia Paranaguá-Curitiba no Parque Estadual Pico do Marumbi, Morretes – PR. Universidade Federal do Paraná.
- Cerboncini, R. A. S., M. B. G. Rubio, I. P. Bernardi, T. V. Braga, J. J. Roper, and F. C. Passos. 2014. Small mammal community structure and vertical space use preferences in nonfragmented Atlantic forest. *Mammalia* **0**:1-8.
- Cherem, J. J. 2005. Registros de mamíferos não voadores em estudos de avaliação ambiental no sul do Brasil. *Biotemas* **18**:169-202.
- Chitty, D. 1960. Population processes in the vole and their relevance to general theory. *Canadian Journal of Zoology* **38**:99-113.
- Costa, L. P., Y. L. R. Leite, G. A. B. da Fonseca, and M. T. da Fonseca. 2000. Biogeography of South American forest mammals: endemism and diversity in the Atlantic forest. *Biotropica* **32**:872-881.
- D'Andrea, P. S., R. Gentile, L. S. Maroja, F. A. Fernandes, R. Coura, and R. Cerqueira. 2007. Small mammal populations of an agroecosystem in the Atlantic forest domain, southeastern Brazil. *Braz J Biol* **67**:179-186.
- D'Andrea, R. S., R. Gentile, R. Cerqueira, C. E. V. Grelle, C. Horta, and L. Rey. 1999. Ecology of small mammals in a Brazilian rural area. *Revista Brasileira De Zoologia* **16**:611-620.
- Dantas-Torres, F., F. M. Alessio, D. B. Siqueira, J. F. Mauffrey, M. F. Marvulo, T. F. Martins, J. Moraes-Filho, M. C. Camargo, S. R. D'Auria, M. B. Labruna, and J. C. Silva. 2012. Exposure of small mammals to ticks and rickettsiae in Atlantic forest patches in the metropolitan area of Recife, North-eastern Brazil. *Parasitology* **139**:83-91.
- Davis, D. E. 1945. The annual cycle of plants, mosquitoes, birds, and mammals in two Brazilian forests. *Ecological Society of America* **15**:243-295.
- de la Sancha, N. U. 2014. Patterns of small mammal diversity in fragments of subtropical interior Atlantic forest in eastern Paraguay. *Mammalia* **78**:1-13.
- Elton, C. S. 1924. Periodic fluctuations in the numbers of animals: their causes and effects. *Journal of Experimental Biology* **2**:119-163.
- Elton, C. S. 1942. Voles, mice and lemmings: problems in population dynamics. Clarendon Press, Oxford.
- Feliciano, B. R., F. A. S. Fernandez, D. De Freitas, and M. S. L. Figueiredo. 2002. Population dynamics of small rodents in a grassland between fragments of Atlantic forest in southeastern Brazil. *Mammalian Biology* **67**:304-314.
- Ferracioli, P. 2013. Interação entre uma comunidade de pequenos mamíferos e a espécie *Araucaria angustifolia* em uma floresta ombrófila mista do sul do Brasil. Universidade Federal de Lavras, Lavras, MG.
- Finokiet, M. 2007. Efeito de borda sobre uma assembléia de pequenos mamíferos em um fragmento de floresta estacional decidual de Santa Maria, RS, sul do Brasil. Universidade Federal De Santa Maria.
- Fonseca, G. A. B., and J. G. Robinson. 1990. Forest size and structure: competitive and predatory effects on small mammal communities. *Biological conservation* **53**:265-294.

Galetti, M., R. Guevara, M. C. Côrtes, R. Fadini, S. Von Matter, A. B. Leite, F. Labecca, T. Ribeiro, C. S. Carvalho, R. G. Collevatti, M. M. Pires, P. R. Guimarães, P. H. Brancalion, M. C. Ribeiro, and P. Jordano. 2013. Functional extinction of birds drives rapid evolutionary changes in seed size. *Science* **340**:1086-1090.

Galiano, D. 2010. Dinâmica populacional e efeitos de variáveis ambientais sobre a fauna de pequenos mamíferos em um fragmento de floresta com Araucária no sul do Brasil. Universidade Federal do Rio Grande do Sul, Porto Alegre.

Galindo-Leal, C., and G. I. Câmara. 2003. The Atlantic forest of South America: biodiversity status, threats, and outlook. [State of the Hotspots.]. The Atlantic forest of South America: biodiversity status, threats, and outlook. [State of the Hotspots.]:i-xxi, 1-488.

Gaspar, D. A. 2005. Comunidade de mamíferos não voadores de um fragmento de floresta Atlântica semidecídua do Município de Campinas, SP. Universidade Estadual de Campinas.

Gentile, R., and F. Fernandez. 1999. A field comparison of two capture-mark-recapture estimators of small mammal populations. *Revista Brasileira De Zoologia* **16**:1109 - 1114.

Gheler-Costa, C., G. Sabino-Santos, L. S. Amorim, L. M. Rosalino, L. T. M. Figueiredo, and L. M. Verdade. 2013. The effect of pre-harvest fire on the small mammal assemblage in sugarcane fields. *Agriculture Ecosystems & Environment* **171**:85-89.

Gheler-Costa, C., L. M. Verdade, and A. I. F. de Almeida. 2002. Mamíferos não-voadores do campus Luiz de Queiroz , Universidade de São Paulo, Piracicaba, Brasil. *Zoologia (Curitiba, Impr.)* **19**:203-214.

Graipel, M. c. E., J. J. Cherem, E. L. A. Monteiro-Filho, and L. Glock. 2006. Dinâmica populacional de marsupiais e roedores no Parque Municipal da Lagoa do Peri, Ilha de Santa Catarina, sul do Brasil. *Mastozoología Neotropical* **13**:31-49.

Graipel, M. E., J. J. Cherem, D. A. Machado, P. C. Garcia, M. E. Menezes, and M. Soldateli. 1997. Vertebrados da Ilha de Ratoes Grande, Santa Catarina, Brasil. *Biotemas* **10**:105-122.

Graipel, M. E., J. J. Cherem, P. R. M. Miller, and L. Glock. 2003. Trapping small mammals in the forest understory: a comparison of three methods. *Mammalia* **67**:551-558.

Graipel, M. E., and M. d. S. Filho. 2006. Reprodução e dinâmica populacional de *Didelphis aurita* Wied-Neuwied (Mammalia : Didelphimorphia) em ambiente periurbano na Ilha de Santa Catarina , sul do Brasil. *Biotemas* **19**:65-73.

Grazzini, G. 2014. Identidade e diversidade dos pequenos mamíferos não voadores da Floresta Nacional do Piraí do Sul, Paraná, Brasil. Universidade Federal do Paraná.

Grelle, C. E. V. 1996. Análise tridimensional de uma comunidade de pequenos mamíferos. Universidade Federal de Minas Gerais.

Grelle, C. E. V. 2003. Forest structure and vertical stratification of small mammals in a secondary Atlantic forest, southeastern Brazil. *Studies On Neotropical Fauna And Environment* **38**:81-85.

Haddad, C. F. B., L. F. Toledo, C. P. A. Prado, D. Loebmann, J. L. Gasparini, and I. Sazima. 2013. Guia dos Anfíbios da Mata Atlântica - Diversidade e Biologia. Anolis Books, São Paulo.

Herrmann, G. 1991. Estrutura da comunidade de pequenos mamíferos em áreas secundárias de Mata Atlântica. Universidade Federal de Mina Gerais.

Hice, C. L., and D. J. Schmidly. 2002. The effectiveness of pitfall traps for sampling small mammals in the Amazon Basin. *Mastozoología Neotropical* **9**:85-89.

Horn, G. B. 2005. A assembléia de pequenos mamíferos da floresta paludosa do Faxinal, Torres-RS: sua relação com a borda e o roedor *Akodon montensis* (Rodentia, Muridae) como potencial dispersos de sementes endozoocóricas. Universidade Federal do Rio Grande do Sul.

Huang, C. Q., S. Kim, A. Altstadt, J. R. G. Townshend, P. Davis, K. Song, C. J. Tucker, O. Rodas, A. Yanosky, R. Clay, and J. Musinsky. 2007. Rapid loss of Paraguay's Atlantic forest and the status of protected areas - A Landsat assessment. *Remote Sensing of Environment* **106**:460-466.

- IBGE. 2013. Atlas do censo demográfico 2011. Ministério do Planejamento, Orçamento e Gestão. Fundação Instituto Brasileiro de Geografia e Estatística, Rio de Janeiro.
- Iob, G. 2007. Influência de frutos e sementes na abundância de pequenos mamíferos e a relação com a predação e dispersão de sementes da araucária (*Araucaria angustifolia*). Universidade Federal do Rio Grande do Sul.
- Jaksic, F. M., and M. Lima. 2003. Myths and facts on ratadas: bamboo blooms, rainfall peaks and rodent outbreaks in South America. Pages 237-251 *Austral Ecology*.
- Jorge, M. L. S. P., M. Galetti, M. C. Ribeiro, and K. M. P. M. B. Ferraz. 2013. Mammal defaunation as surrogate of trophic cascades in a biodiversity hotspot. *Biological conservation* **163**:49-57.
- Karp, D. S., A. J. Rominger, J. Zook, J. Ranganathan, P. R. Ehrlich, and G. C. Daily. 2012. Intensive agriculture erodes beta-diversity at large scales. *Ecology Letters* **15**:963-970.
- Kendall, W. L., K. H. Pollock, and C. Brownie. 1995. A likelihood-based approach to capture-recapture estimation of demographic parameters under the robust design. *Biometrics* **51**:293-308.
- Korpimäki, E., T. Klemola, K. Norrdahl, L. Oksanen, T. Oksanen, P. B. Banks, G. O. Batzli, and H. Henttonen. 2003. Vole cycles and predation. *Trends in Ecology & Evolution* **18**:494-495.
- Lambert, T. D., J. R. Malcolm, and B. L. Zimmerman. 2006. Amazonian small mammal abundances in relation to habitat structure and resource abundance. *Journal of Mammalogy* **87**:766-776.
- Lambin, X., and C. J. Krebs. 1991. Can changes in female relatedness influence microtine population dynamics. *Oikos* **61**:126-132.
- Lambin, X., and C. J. Krebs. 1993. Influence of female relatedness on the demography of Townsend's vole populations in spring. *Journal of Animal Ecology* **62**:536-550.
- Lambin, X., C. J. Krebs, R. Moss, and N. G. Yoccoz. 2002. Population cycles: inferences from experimental, modelling, and time series approaches. Pages 155-176 in A. A. Berryman, editor. *Population cycles: the case for trophic interactions*. Oxford University Press, New York.
- Langone, P. Q. 2007. Importância da matriz e das características do hábitat sobre a assembléia de pequenos mamíferos em fragmento de mata de restinga no sul do Brasil. Universidade Federal do Rio Grande Do Sul.
- Leiner, N. O., and W. R. Silva. 2012. Non-volant small mammals at an Atlantic forest area situated nearby a limestone quarry (Limeira quarry), State of São Paulo, Brazil. *Biota Neotrop.* **12**:1-7.
- Lessa, G., P. R. Gonçalves, M. M. M. Júnior, F. M. Costa, R. F. Pereira, and A. P. Paglia. 1999. Caracterização e monitoramento da fauna de pequenos mamíferos terrestres de um fragmento de mata secundária em Viçosa, Minas Gerais. *Cadernos do Departamento de Ciências Biológicas da PUC Minas* **7**:41-49.
- Lima, B. S., F. Dantas-Torres, M. R. de Carvalho, J. F. Marinho-Junior, E. L. de Almeida, M. E. F. Brito, F. Gomes, and S. P. Brandao-Filho. 2013. Small mammals as hosts of *Leishmania* spp. in a highly endemic area for zoonotic leishmaniasis in north-eastern Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **107**:592-597.
- Lima, D. O., B. O. Azambuja, V. L. Camilotti, and N. C. Caceres. 2010. Small mammal community structure and microhabitat use in the austral boundary of the Atlantic forest, Brazil. *Zoologia* **27**:99-105.
- LoGiudice, K., R. S. Ostfeld, K. A. Schmidt, and F. Keesing. 2003. The ecology of infectious disease: Effects of host diversity and community composition on Lyme disease risk. *Proceedings of the National Academy of Sciences* **100**:567-571.

- Loretto, D. 2012. Ecologia de pequenos mamíferos arborícolas: estado do conhecimento, métodos de amostragem e estudo populacional, com ênfase no bioma da Mata Atlântica. Universidade Federal do Rio de Janeiro.
- Luza, A. L., N. Zanella, A. U. Christoff, A. F. Barreto-Lima, and J. V. Grando. 2013. Relação entre fatores exógenos e a abundância de roedores em remanescente de Floresta Ombrófila Mista, Rio Grande do Sul. *Revista Brasileira de Biociências* **11**:263-268.
- Machado, F. S., R. Gregorin, and P. S. B. Mouallem. 2013. Small mammals in high altitude phytophysognomies in southeastern Brazil: are heterogeneous habitats more diverse? *Biodivers Conserv* **22**:1769-1782.
- Maestri, R., and J. R. Marinho. 2014. Singing in the rain. Rainfall and moonlight affect daily activity patterns of rodents in a Neotropical forest. *Acta Theriologica* **59**:427-433.
- Marinho, J. R. 2003. Estudo da comunidade e do fluxo gênico de roedores silvestres em um gradiente altitudinal de Mata Atlântica na área de influência da RST-453/RS-486 - Rota-do-Sol. Universidade Federal do Rio Grande do Sul, Porto Alegre.
- Martin, P. S., C. Gheler-Costa, P. C. Lopes, L. M. Rosalino, and L. M. Verdade. 2012. Terrestrial non-volant small mammals in agro-silvicultural landscapes of southeastern Brazil. *Forest Ecology And Management* **282**:185-195.
- Martins, T. K. 2011. Determinantes ecológicos do risco de extinção: abundância local, amplitude de nicho, capacidade de dispersão e a resposta das espécies de pequenos mamíferos à fragmentação florestal no Planalto Atlântico Paulista. University of São Paulo, São Paulo.
- May, R. M. 1988. How many species are there on earth? *Science* **241**:1441-1449.
- Melo, G. L., J. Sponchiado, A. F. Machado, and N. C. Caceres. 2011. Small-mammal community structure in a South American deciduous Atlantic forest. *Community Ecology* **12**:58-66.
- Mesquita, A. O., and M. Passamani. 2012. Composition and abundance of small mammal communities in forest fragments and vegetation corridors in southern Minas Gerais, Brazil. *Revista De Biologia Tropical* **60**:1335-1343.
- Milagres, B. S., A. F. Padilha, C. E. Montandon, R. N. Freitas, R. Pacheco, D. H. Walker, M. B. Labruna, C. L. Mafra, and M. A. M. Galvao. 2013. Spotted fever group Rickettsia in small rodents from areas of low endemicity for Brazilian spotted fever in the eastern region of Minas Gerais State, Brazil. *American Journal of Tropical Medicine and Hygiene* **88**:937-939.
- MMA, M. d. M. A. d. E. d. S. P. 2010. Inventário da fauna do Município de São Paulo 2010. Diário Oficial da Cidade de São Paulo, São Paulo.
- Mochi-Jr, C. M. 2014. Composição taxonômica e avaliação da diversidade da fauna de pequenos mamíferos não-voadores na formação submontana do Parque Nacional Saint-Hilaire/Lage, Mata Atlântica costeira do Paraná. Universidade Federal do Paraná.
- Modesto, T. C., F. S. Pessoa, M. C. Enrici, N. Attias, T. Jordão-Nogueira, L. d. M. Costa, H. G. Albuquerque, and H. d. G. Bergallo. 2008. Mamíferos do Parque Estadual do Desengano, Rio de Janeiro, Brasil. *Biota Neotrop.* **8**:153-159.
- Moreira, J. C., E. G. Manduca, P. R. Gonçalves, M. M. d. Moraes Júnior, R. F. Pereira, G. Lessa, and J. A. Dergam. 2009. Small mammals from Serra do Brigadeiro State Park, Minas Gerais, southeastern Brazil: species composition and elevational distribution. *Arquivos do Museu Nacional* **67**:103-118.
- Moreira, J. C., E. G. Manduca, P. R. Goncalves, R. Stumpp, C. G. C. Pinto, and G. Lessa. 2008. Mammals, Volta Grande environmental unity, triângulo mineiro, States of Minas Gerais and São Paulo, southeastern Brazil Check List **4**:349-357.
- Moreira-Lima, L. 2014. Aves da Mata Atlântica: riqueza, composição, status, endemismos e conservação. Universidade de São Paulo, São Paulo.
- Morellato, L. P. C., and C. F. B. Haddad. 2000. Introduction: the Brazilian Atlantic forest. *Biotropica* **32**:786-792.

- Morris, D. W. 1999. A haunting legacy from isoclines: mammal coexistence and the ghost of competition. *Journal of Mammalogy* **80**:375-384.
- Morris, D. W., B. J. Fox, J. Luo, and V. Monamy. 2000. Habitat-dependent competition and the coexistence of Australian heathland rodents. *Oikos* **91**:294-306.
- Morrone, J. J. 2014. Biogeographical regionalisation of the Neotropical region. *Zootaxa* **3782**:1-110.
- Moura, R. T. 1999. Análise comparativa da estrutura de comunidades de pequenos mamíferos em remanescente de Mata Atlântica e plantio de cacau em sistema de Cabruca no sul da Bahia. Universidade de Minas Gerais.
- Moura, R. T. 2003. Distribuição e ocorrência de mamíferos na Mata Atlântica do sul da Bahia. Page 22 in P. I. Prado, E. C. Landau, R. T. Moura, L. P. S. Pinto, G. A. B. Fonseca, and K. N. Alger, editors. Corredor de Biodiversidade da Mata Atlântica do Sul da Bahia. programas.inema.ba.gov.br, Ilhéus, IESB / CI / CABS / UFMG / UNICAMP.
- Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. da Fonseca, and J. Kent. 2000. Biodiversity hotspots for conservation priorities. *Nature* **403**:853-858.
- Naxara, L., B. T. Pinotti, and R. Pardini. 2009. Seasonal microhabitat selection by terrestrial rodents in an old-growth Atlantic forest. *Journal of Mammalogy* **90**:404-415.
- Nichols, J. D., and K. H. Pollock. 1990. Estimation of recruitment from immigration versus *in situ* reproduction using Pollock's robust design. *Ecology* **71**:21-26.
- Nieri-Bastos, F. A., D. M. Barros-Battesti, P. M. Linardi, M. Amaku, A. Marcili, A. E. Favorito, and R. Pinto-Da-Rocha. 2004. Ectoparasites of wild rodents from Parque Estadual da Cantareira (Pedra Grande Nuclei), São Paulo, Brazil. *Rev. Bras. Parasitol. Vet* **13**:29-35.
- Odden, M., R. A. Ims, O. G. Stoen, J. E. Swenson, and H. P. Andreassen. 2014. Bears are simply voles writ large: social structure determines the mechanisms of intrinsic population regulation in mammals. *Oecologia* **175**:1-10.
- Oksanen, L., S. D. Fretwell, J. Arruda, and P. Niemela. 1981. Exploitation ecosystems in gradients of primary productivity. *American Naturalist* **118**:240-261.
- Olifiers, N. 2002. Fragmentação, habitat e as comunidades de pequenos mamíferos da Bacia do Rio Macacu, RJ. Universidade Federal de Minas Gerais, Belo Horizonte.
- Oliveira, F. F., S. F. Ferrari, and S. D. B. Silva. 2005. Mamíferos não-voadores. Page 131 in C. M. d. Carvalho and J. C. Vilar, editors. Parque Nacional Serra de Itabaiana Levantamento da Biota
- Oliveira, R. C., R. Gentile, A. Guterres, J. Fernandes, B. R. Teixeira, V. Vaz, F. P. Valdez, L. H. B. Vicente, S. F. da Costa-Neto, C. Bonvicino, P. S. DAndrea, and E. R. S. Lemos. 2014. Ecological study of hantavirus infection in wild rodents in an endemic area in Brazil. *Acta Tropica* **131**:1-10.
- Ostfeld, R. S. 1990. The ecology of territoriality in small mammals. *Trends in Ecology & Evolution* **5**:411-415.
- Otis, D. L., K. P. Burnham, G. C. White, and D. R. Anderson. 1978. Statistical inference from capture data on closed animal populations. *Wildlife Monographs*:7-135.
- Pacheco, M., M. Kajin, R. Gentile, P. L. Zangrandi, M. V. Vieira, and R. Cerqueira. 2013. A comparison of abundance estimators for small mammal populations. *Zoologia (Curitiba, Impresso)* **30**:182-190.
- Paglia, A. P., G. A. B. da Fonseca, A. B. Rylands, G. Herrmann, L. M. S. Aguiar, A. G. Chiarello, Y. L. R. Leite, L. P. Costa, S. Siciliano, M. C. M. Kierulff, S. L. Mendes, V. C. Tavares, R. A. Mittermeier, and J. Patton. 2012. Lista anotada dos mamíferos do Brasil. *Occasional Papers in Conservation Biology* **6**:1-76.
- Paglia, A. P., P. De Marco Júnior, F. M. Costa, R. F. Pereira, and G. Lessa. 1995. Heterogeneidade estrutural e diversidade de pequenos mamíferos em um fragmento de mata secundária de Minas Gerais, Brasil. *Revista Brasileira De Zoologia* **12**:67-79.

- Paise, G. 2010. Efeitos da fragmentação de habitat sobre a comunidade de pequenos mamíferos de Mata Atlântica no Estado de São Paulo. Universidade Estadual de Campinas.
- Palma, A. R. T. 1996. Separação de nichos entre pequenos mamíferos de Mata Atlântica. Universidade Estadual de Campinas.
- Pardini, R. 2004. Effects of forest fragmentation on small mammals in an Atlantic forest landscape. *Biodiversity and Conservation* **13**:2567-2586.
- Pardini, R., A. d. A. Bueno, T. A. Gardner, P. I. c. Prado, and J. P. Metzger. 2010. Beyond the fragmentation threshold hypothesis: regime shifts in biodiversity across fragmented landscapes. *PLoS ONE* **5**:e13666.
- Pardini, R., and F. Umetsu. 2006. Pequenos mamíferos não-voadores da Reserva Florestal do Morro Grande: distribuição das espécies e da diversidade em uma área de Mata Atlântica. *Biota Neotropica* **6**:1-22.
- Paresque, R., W. P. d. Souza, S. L. Mendes, and V. Fagundes. 2004. Composição cariotípica da fauna de roedores e marsupiais de duas áreas de Mata Atlântica do Estado do Espírito Santo, Brasil. *Boletim do Museu de Biologia Mello Leitão* **17**:5-33.
- Passamani, M. 2000. Análise da comunidade de marsupiais em Mata Atlantica de Santa Teresa, Espírito Santo. *Bol. Mus. Biol. Mello Leitão* **11**:215-228.
- Passamani, M., J. Dalmaschio, and S. A. Lopes. 2005. Mamíferos não-voadores em áreas com predomínio de Mata Atlântica da Samarco Mineração S.A., município de Anchieta, Espírito Santo. *Biotemas* **18**:135-149.
- Passamani, M., and F. A. S. Fernandez. 2011a. Abundance and richness of small mammals in fragmented Atlantic forest of southeastern Brazil. *Studies On Neotropical Fauna And Environment* **45**:553-565.
- Passamani, M., and F. A. S. Fernandez. 2011b. Movements of small mammals among Atlantic forest fragments in Espírito Santo, southeastern Brazil. *Mammalia* **75**:83-86.
- Passamani, M., S. r. L. Mendes, and A. G. Chiarello. 2000. Non-volant mammals of the Estação Biológica de Santa Lúcia and adjacent areas of Santa Teresa, Espírito Santo, Brazil. *Bol. Mus. Biol. Mello Leitão* **11**:201-214.
- Passamani, M., and D. Ribeiro. 2009. Small mammals in a fragment and adjacent matrix in southeastern Brazil. *Braz. J. Biol.* **69**:305-309.
- Patton, J. L., U. F. J. Pardiñas, and G. D'Elia. 2015. *Mammals of South America. Volume 2: Rodents*, University of Chicago Press.
- Pedó, E. 2005. Assembléia de pequenos mamíferos não-voadores em áreas de ecótono campo-floresta com Araucária na região dos Campos de Cima da Serra, Rio Grande do Sul. Universidade Federal do Rio Grande do Sul.
- Pedó, E., T. R. O. d. Freitas, and S. M. Hartz. 2010. The influence of fire and livestock grazing on the assemblage of non-flying small mammals in grassland-Araucaria forest ecotones, southern Brazil. *Zoologia (Curitiba, Impr.)* **27**:533-540.
- Pena, D. C. H. 2007. Epidemiologia das riquetsioses em área de foco silencioso para febre maculosa brasileira, município de Santa Cruz do Escalvado, Minas Gerais. Universidade Federal de Ouro Preto, Ouro Preto.
- Pereira, L. G., and L. Geise. 2009. Non-flying mammals of Chapada Diamantina (Bahia, Brazil). *Biota Neotrop.* **9**:185-196.
- Pereira, S. N., D. Dias, I. P. Lima, A. C. I. S. Maas, M. A. Martins, D. P. Bolzan, D. b. Souza França, M. B. Oliveira, A. L. Peracchi, and M. F. t. S. Ferreira. 2013. Mamíferos de um fragmento florestal em Volta Redonda, Estado do Rio de Janeiro. *Biosci. J.* **29**:1017-1027.
- Pessôa, F. S., T. C. Modesto, H. G. Albuquerque, N. Attias, and H. G. Bergallo. 2009. Non-volant mammals, Reserva Particular do Patrimônio Natural (RPPN) Rio das Pedras, municipality of Mangaratiba, State of Rio de Janeiro, Brazil. *Check List* **5**:577-586.

- Peters, F. B., P. R. d. O. Roth, L. F. Machado, E. D. L. Coelho, D. M. H. Jung, and A. U. Christoff. 2010. Assembléia de mamíferos dos agroecossistemas constituintes da Bacia Hidrográfica do rRo da Várzea, Rio Grande do Sul. *Biotemas*:91-107.
- Pinheiro, P. S., and L. Geise. 2008. Non-volant mammals of Picinguaba, Ubatuba, State of São Paulo, southeastern Brazil. *Bol. Mus. Biol. Mello Leitão* **23**:51-59.
- Pinotti, B. T. 2010. Pequenos mamíferos terrestres e a regeneração da Mata Atlântica: influência da estrutura do hábitat e da disponibilidade de alimento na recuperação da fauna. Universidade de São Paulo.
- Pinto, I. d. S., A. C. C. Loss, A. Falqueto, and Y. L. R. Leite. 2009. Pequenos mamíferos não voadores em fragmentos de Mata Atlântica e áreas agrícolas em Viana, Espírito Santo, Brasil. *Biota Neotrop.* **9**:355-360.
- Pires, A. S., P. K. Lira, F. A. S. Fernandez, G. M. Schittini, and L. C. Oliveira. 2002. Frequency of movements of small mammals among Atlantic coastal forest fragments in Brazil. *Biological conservation* **108**:229-237.
- Prevedello, J. A. 2013. A importância dos recursos alimentares para populações e comunidades animais: uma meta-análise global e um estudo experimental na Mata Atlântica. Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ.
- Prevedello, J. A., P. Ferreira, B. S. Papi, D. Loretto, and M. V. Vieira. 2008. Uso do espaço vertical por pequenos mamíferos no Parque Nacional Serra dos Órgãos, RJ: um estudo de 10 anos utilizando três métodos de amostragem. *Espaço & Geografia* **11**:35-58.
- Purvis, A., and A. Hector. 2000. Getting the measure of biodiversity. *Nature* **405**:212-219.
- Puttker, T., Y. Meyer-Lucht, and S. Sommer. 2006. Movement distances of five rodent and two marsupial species in forest fragments of the coastal Atlantic rainforest, Brazil. *Ecotropica* **12**:131-139.
- Quintela, F. M., B. I. Goncalves, G. E. Trindade, M. B. d. Santos, A. M. Tozetti, F. M. Quintela, B. I. Goncalves, G. E. Trindade, M. B. d. Santos, and A. M. Tozetti. 2013. Pequenos mamíferos não voadores (Didelphimorphia, Rodentia) em campos litorâneos do extremo sul do Brasil. *Biota Neotropica* **13**:284-289.
- Regolin, A. L., and J. J. Cherem. Unpubl. Data-a. Reference 144.
- Regolin, A. L., and J. J. Cherem. Unpubl. Data-b. Reference 145.
- Regolin, A. L., N. Furnari, F. de Castro Jacinavicius, P. M. Linardi, and C. J. de Carvalho-Pinto. 2015. Ectoparasites of the critically endangered insular cavy, *Cavia intermedia* (Rodentia: Caviidae), southern Brazil. *International Journal for Parasitology: Parasites and Wildlife* **4**:37-42.
- Ribeiro, M. C., J. P. Metzger, A. C. Martensen, F. J. Ponzoni, and M. M. Hirota. 2009. The Brazilian Atlantic forest: how much is left, and how is the remaining forest distributed? Implications for conservation. *Biological conservation* **142**:1141-1153.
- Rocha, F. S. 2004. A conservação de pequenos mamíferos no Pontal do Paranapanema (SP): diagnóstico de padrões de comunidades e sugestões de manejo. Universidade de São Paulo, São Paulo
- Rocha, M. F., M. Passamani, and J. Louzada. 2011. A small mammal community in a forest fragment, vegetation corridor and coffee matrix system in the Brazilian Atlantic forest. *PLoS ONE* **6**:e23312.
- Rocha-Mendes, F., C. L. Neves, R. A. Nobre, R. M. Marques, and M. Galetti. 2015. Mamíferos não voadores do Núcleo Santa Virgínia, Parque Estadual da Serra do Mar, São Paulo, Brasil. *Biota neotrop.*(Online, Ed. port.) **15**.
- Rodarte, R. R. D. P. 2013. Ecologia trófica de pequenos mamíferos não voadores em uma área contínua de Mata Atlântica. Universidade Estadual Paulista “Júlio de Mesquita Filho” Instituto de Biociências – Rio Claro.

- Rosa, A. O. 2002. Comparação da diversidade de mamíferos não-voadores em áreas de floresta de restinga e áreas reflorestadas com *Pinus elliottii* no Sul do Brasil. Universidade do Vale do Rio Dos Sinos– UNISINOS.
- Rosa, C. A. 2012. Efeito de borda de rodovias em pequenos mamíferos de fragmentos florestais tropicais. Universidade Federal de Lavras.
- Rosalino, L. M., P. S. Martin, C. Gheler-Costa, P. C. Lopes, and L. M. Verdade. 2014. Neotropical small mammals' diversity in the early cycle of commercial *Eucalyptus* plantations. *Agroforest Syst* **88**:427-436.
- Salvador, C. H., and F. A. S. Fernandez. 2008. Population dynamics and conservation status of the insular cavy *Cavia intermedia* (Rodentia: Caviidae). *Journal of Mammalogy* **89**:721-729.
- Santos, A. u. B., S. Lóss, and Y. L. Leite. 2004. Padrões de uso de estratos da floresta por pequenos mamíferos no Parque Estadual da Fonte Grande, Vitória, Espírito Santo. *Natureza on line* **2**:27-33.
- Scheibler, D. R., and A. U. Christoff. 2007. Habitat associations of small mammals in southern Brazil and use of regurgitated pellets of birds of prey for inventorying a local fauna. *Brazilian Journal of Biology* **67**:619-625.
- Sherman, P. W., and J. O. Wolff. 2007. Rodent societies: an ecological & evolutionary perspective. University of Chicago Press, Chicago.
- Silva, C. R. 2001. Riqueza e diversidade de mamíferos não-voadores em um mosaico formado por plantios de *Eucalyptus saligna* e remanescentes de floresta Atlântica no município de Pilar do Sul, SP. Escola Superior de Agricultura Luiz de Queiroz, Universidade de São Paulo,.
- Silva, M. A., A. Ronconi, N. Cordeiro, D. E. Bossi, H. G. Bergallo, M. C. Costa, J. C. Balieiro, and F. L. Varzim. 2007. Blood parasites, total plasma protein and packed cell volume of small wild mammals trapped in three mountain ranges of the Atlantic forest in southeastern Brazil. *Braz J Biol* **67**:531-535.
- Silveira, F. 2012. Estrutura populacional de pequenos mamíferos na Reserva do Cachoeira, APA de Guaraqueçaba, Paraná.
- Siqueira, D. B., F. M. Aléssio, J. F. Mauffrey, M. F. V. Marvulo, V. O. Ribeiro, R. L. Oliveira, H. F. J. Pena, S. M. Gennari, R. A. Mota, M. A. G. Faustino, L. C. Alves, J. P. Dubey, and J. C. R. SILVA. 2013. Seroprevalence of *Toxoplasma gondii* in wild marsupials and rodents from the Atlantic forest of Pernambuco State, northeastern region, Brazil. *Journal Of Parasitology* **99**:1140-1143.
- Souza, R., and M. Galetti. Unpubl. Data. Reference 01
- Stallings, J. R., G. A. B. Fonseca, L. P. S. Pinto, L. Aguiar, and E. L. Sábato. 1991. Mamíferos do Parque Florestal Estadual do Rio Doce, Minas Gerais, Brasil. *Revista Brasileira De Zoologia* **7**:663-677.
- Steiner-Souza, F., P. Cordeiro-Estrela, A. R. Percequillo, A. F. Testoni, and S. L. Althoff. 2008. New records of *Rhagomys rufescens* (Rodentia: Sigmodontinae) in the Atlantic forest of Brazil. *Zootaxa* **1824**:28-34.
- Stevens, S. M., and T. P. Husband. 1998. The influence of edge on small mammals: evidence from Brazilian Atlantic forest fragments. *Biological conservation* **85**:1-8.
- Teixeira, B. R., N. Loureiro, L. Strecht, R. Gentile, R. C. Oliveira, A. Guterres, J. Fernandes, L. H. Mattos, S. M. Raboni, G. Rubio, C. R. Bonvicino, C. N. dos Santos, E. R. Lemos, and P. S. D'Andrea. 2014. Population ecology of hantavirus rodent hosts in southern Brazil. *American Journal of Tropical Medicine and Hygiene* **91**:249-257.
- Tortato, F. R., A. F. Testoni, and S. L. Althoff. 2014. Mastofauna terrestre da Reserva Biológica Estadual do Sassafrás, Doutor Pedrinho, Santa Catarina, Sul do Brasil. *Biotemas* **27**:123-129.

- Uchôa, T. 2006. Comunidades dos pequenos mamíferos em dois estágios sucessionais de floresta Atlântica e suas implicações à ecologia e conservação. Universidade Federal do Paraná.
- Umetsu, F., J. P. Metzger, and R. Pardini. 2008. Importance of estimating matrix quality for modeling species distribution in complex tropical landscapes: a test with Atlantic forest small mammals. *Ecography* **31**:359-370.
- Umetsu, F., L. Naxara, and R. Pardini. 2006. Evaluating the efficiency of pitfall traps for sampling small mammals in the neotropics. *Journal of Mammalogy* **87**:757-765.
- Umetsu, F., and R. Pardini. 2007. Small mammals in a mosaic of forest remnants and anthropogenic habitats-evaluating matrix quality in an Atlantic forest landscape. *Landscape Ecology* **22**:517-530.
- Vera Y Conde, C. F., and C. F. D. Rocha. 2006. Habitat disturbance and small mammal richness and diversity in an Atlantic rainforest area in southeastern Brazil. *Brazilian Journal of Biology* **66**:983-990.
- Vieira, E. M. 1999. Estudo comparativo de comunidades de pequenos mamíferos em duas áreas de Mata Atlântica situadas a diferentes altitudes no sudoeste do Brasil. Universidade Estadual de Campinas.
- Vieira, E. M., and E. L. A. Monteiro. 2003. Vertical stratification of small mammals in the Atlantic rain forest of south-eastern Brazil. *Journal Of Tropical Ecology* **19**:501-507.
- Vieira, E. M., and E. L. A. Monteiro Filho. 2003. Vertical stratification of small mammals in the Atlantic rain forest of south-eastern Brazil. *Journal Of Tropical Ecology* **19**:501-507.
- Vieira, M. V., C. E. Grelle, and R. Gentile. 2004. Differential trappability of small mammals in three habitats of southeastern Brazil. *Braz J Biol* **64**:895-900.
- Vieira, M. V., N. Olifiers, A. C. Delciellos, V. Z. Antunes, L. R. Bernardo, C. E. V. Grelle, and R. Cerqueira. 2009. Land use vs. fragment size and isolation as determinants of small mammal composition and richness in Atlantic forest remnants. *Biological conservation* **142**:1191-1200.
- Visconti, P., R. L. Pressey, D. Giorgini, L. Maiorano, M. Bakkenes, L. Boitani, R. Alkemade, A. Falcucci, F. Chiozza, and C. Rondinini. 2011. Future hotspots of terrestrial mammal loss. *Philosophical Transactions Of The Royal Society Of London Series B-Biological Sciences* **366**:2693-2702.
- Voss, R. S., and L. Emmmons. 1996. Mammalian diversity in neotropical lowland rainforests: a preliminary assessment. *Bulletin of the American Museum of Natural History, New York*.
- Voss, R. S., and S. A. Jansa. 2009. Phylogenetic relationships and classification of didelphid marsupials, an extant radiation of New World metatherian mammals. *Bulletin Of The American Museum Of Natural History* **322**:1-177.
- Voss, R. S., D. P. Lunde, and N. B. Simmons. 2001. The mammals of Paracou, French Guiana: a neotropical lowland rainforest fauna - part 2. Nonvolant species. *Bulletin Of The American Museum Of Natural History*:3-236.
- Willis, K. J., M. B. Araujo, K. D. Bennett, B. Figueroa-Rangel, C. A. Froyd, and N. Myers. 2007. How can a knowledge of the past help to conserve the future? Biodiversity conservation and the relevance of long-term ecological studies. *Philosophical Transactions of the Royal Society B-Biological Sciences* **362**:175-186.
- Wilman, H., J. Belmaker, J. Simpson, C. de la Rosa, M. M. Rivadeneira, and W. Jetz. 2014. EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology* **95**:2027-2027.