


People in Ecosystems/Watershed Integration (PEWI)
Advanced Level Index Notes



PEWI team

August 8, 2017

Summary

This is the document of index. This document includes all advanced tab description.

Acknowledgements

Thanks to so many people.

Contents

Summary	i
Acknowledgements	ii
1 Land Use	1
1.1 Conventional Corn	1
1.2 Conventional Soy	1
1.3 Conservation Corn	1
1.4 Conservation Soy	2
1.5 Alfalfa	2
1.6 Mixed Fruit and Vegetables	2
1.7 Grass Hay	3
1.8 Switchgrass	3
1.9 Permanent Pasture	3
1.10 Rotational Grazing	4
1.11 Wetland	4
1.12 Prairie	4
1.13 Conventional Forest	5
1.14 Conservation Forest	5
1.15 Short-Rotation Woody Bioenergy	5
2 Physical Features	6
2.1 Topographic Relief	6
2.2 Flood Frequency	6
2.3 Strategic Wetlands	6
2.4 Subwatershed Boundaries	7
2.5 Drainage Class	7
2.6 Soil Class	7
2.6.1 Hydrologic Group	8
2.7 Soil Texture	8
2.8 Corn Suitability Rating	9
3 Precipitation	10
4 Management Practices	11
4.1 Crop Rotation	11
5 Modules	12
5.1 Habitat	12
5.1.1 Biodiversity	12
5.1.2 Game Wildlife	12
5.2 Soil Quality	13
5.2.1 Carbon Sequestration	14
5.2.2 Erosion Control	15
5.3 Water Quality	15

5.3.1	Nitrate Retention	16
5.3.2	Phosphorus Retention	16
5.3.3	Sediment Control	17
5.4	Yield	17
5.4.1	Corn Yield	18
5.4.2	Soybean Yield	18
5.4.3	Alfalfa Hay Yield	19
5.4.4	Grass Hay Yield	20
5.4.5	Cattle Yield	20
5.4.6	Mixed Fruit and Vegetable Yield	21
5.4.7	Wood Yield	22
5.4.8	Switch Grass Yield	22
5.4.9	Woody Biomass	22
6	Functionality	23
6.1	Navigating the PEWI Workspace	23
6.1.1	Tool Panel	23
6.1.2	Results Panel	23
6.1.3	Comment Panel	24
6.1.4	Index Information Box	24
6.1.5	View Perspective	24
6.1.6	Save Watershed	25
6.2	Main Menu	25
6.3	Guided Play	25
6.4	Multiplayer Mode	25
6.5	Pause Menu	25
6.6	Hover Information	26
6.7	Hotkeys	26
6.8	Hotkey Customization	26
6.9	Randomization	27
6.10	Undo Function	27
6.11	Print Function	27
6.12	Record	27
6.13	About	27

1 Land Use

Upland grid cells, those not containing part of the river, are 10 acres (4 hectares) in area. Grid cells along the river decrease in area to account for the percent of the grid cell underwater. Land use for a particular land cover is calculated by the sum of the area of each grid cell using that land cover. To find percent land use, this sum is divided by the total area of the watershed and multiplied by 100.

1.1 Conventional Corn

Corn performs best on gentle terrain with high drainage and low flooding probability. The conventional corn land cover in PEWI assumes conventional tillage and no use of cover crops, terraces, contours, grassed waterways, or stream buffers.

Strategic placement can maximize your per-unit yield: Muscatine 119, Tama 120B, Nicollet 55, and Clarion 138B soils are highly productive for annual crops in general, while Buckney 1636 and Gara-Armstrong 993E2 soils are unproductive.

With a conventional corn-dominated land cover, yield, erosion control, and nitrate, phosphorus, and sediment retention vary significantly under different precipitation scenarios.

To improve ecosystem scores in extreme precipitation conditions, try switching from conventional to conservation crops.

Conventional crops also do not provide points in the wildlife habitat or carbon sequestration categories. To improve habitat scores, try adding some perennial or native habitat. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.2 Conventional Soy

Soybeans perform best on gentle terrain with high drainage and low flooding probability. The conventional soybean land cover in PEWI assumes conventional tillage and no use of cover crops, terraces, contours, grassed waterways, or stream buffers. Strategic placement can maximize your per-unit yield: Muscatine 119, Tama 120B, Nicollet 55, and Clarion 138B soils are highly productive for annual crops in general, while Buckney 1636 and Gara-Armstrong 993E2 soils are unproductive.

With a conventional soy-dominated land cover, yield, erosion control, and nitrate, phosphorus, and sediment retention vary significantly under different precipitation scenarios.

To improve ecosystem scores in extreme precipitation conditions, try switching from conventional to conservation crops.

Conventional crops also do not provide points in the wildlife habitat or carbon sequestration categories. To improve habitat scores, try adding some perennial or native habitat. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.3 Conservation Corn

Corn performs best on gentle terrain with high drainage and low flooding probability.

Strategic placement can maximize your per-unit yield: Muscatine 119, Tama 120B, Nicollet 55, and Clarion 138B soils are highly productive for annual crops in general, while Buckney 1636 and Gara-Armstrong 993E2 soils are unproductive.

Conservation corn base yield is equivalent to that of conventional corn.

One hundred percent conservation corn cover gives sediment retention scores above 90, and phosphorus retention and erosion control scores above 85 in all precipitation scenarios. Nitrate retention varies substantially under different precipitation scenarios. To improve nitrate retention scores in extreme precipitation conditions, try switching to perennial land uses.

Conservation crops provide minimal carbon sequestration and moderate habitat points. To improve carbon sequestration scores, try adding some woody land covers or wetlands. To improve habitat scores, try adding some native vegetation.

1.4 Conservation Soy

Soybeans perform best on gentle terrain with high drainage and low flooding probability.

Strategic placement can maximize your per-unit yield: Muscatine 119, Tama 120B, Nicollet 55, and Clarion 138B soils are highly productive for annual crops in general, while Buckney 1636 and Gara-Armstrong 993E2 soils are unproductive.

Conservation soybean base yield is equivalent to that of conventional soybean.

One hundred percent conservation soybean cover gives sediment retention scores above 90, and phosphorus retention and erosion control scores above 80 in all precipitation scenarios. Nitrate retention varies substantially under different precipitation scenarios. To improve nitrate retention scores in extreme precipitation conditions, try switching to perennial crops.

Conservation crops provide minimal points in carbon sequestration, and moderate habitat points. To improve carbon sequestration scores, try adding some woody land covers or wetlands. To improve habitat scores, try adding some native vegetation.

1.5 Alfalfa

Alfalfa performs best on soils with excessive to moderate drainage.

Strategic placement can maximize your per-unit yield: Tama 120B, Muscatine 119, Tama 120C2, Nicollet 55, Nodaway 220, and Clarion 138B soils are highly productive for alfalfa and grass hay, while Buckney 1636, Okoboji 90, and Gara-Armstrong 993E2 soils are approximately half as productive.

Yield varies only slightly in extreme precipitation scenarios.

One hundred percent alfalfa land cover gives the maximum score in nitrate retention, erosion control and sediment retention scores above 90, and phosphorus retention above 80 in all precipitation scenarios. Phosphorus contribution is decreased for areas with an elevation grade of less than 9

Alfalfa does not provide points in the habitat categories and provides only minimal carbon sequestration benefit. To improve habitat scores, try switching to grass hay or adding some native vegetation or stream buffers. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.6 Mixed Fruit and Vegetables

Mixed fruit and vegetables perform best on fine sandy loam, silt loam, or loam.

Strategic placement can maximize your per-unit yield: Buckney 1636, Downs 162D2, Nodaway 220, Ackmore-Colo 5B, Muscatine 119, and Gara-Armstrong 993E2 soils are highly productive for mixed fruit and vegetables, while all other soils are less than half as productive.

With a mixed fruit and vegetable land cover, erosion control, and nitrate, phosphorus, and sediment retention vary significantly under different precipitation scenarios. Yield is decreased in wet years. To improve these scores in extreme precipitation conditions, try switching to perennial or conservation crops.

As a collection of row crops, the mixed fruit and vegetable land cover attains nitrate retention scores similar to those of conventional and conservation corn and soybean. Similarly, it contributes little to carbon sequestration and therefore provides no points in that category. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

As a high-input, high diversity land cover, mixed fruit and vegetables provide moderate points in biodiversity and high points in game wildlife.

1.7 Grass Hay

Grass hay performs best on soils with excessive to moderate drainage.

Strategic placement can maximize your per-unit yield: Tama 120B, Muscatine 119, Tama 120C2, Nicollet 55, Nodaway 220, and Clarion 138B soils are highly productive for alfalfa and grass hay, while Buckney 1636, Okoboji 90, and Gara-Armstrong 993E2 soils are approximately half as productive.

Yield varies only slightly in extreme precipitation scenarios.

One hundred percent grass hay land cover gives the maximum score in nitrate retention, while erosion control, and sediment and phosphorus retention score above 95 in all precipitation scenarios.

As a stream buffer and low input land use, grass hay provides moderate points in the habitat categories. To improve habitat scores, try adding some native vegetation. Grass hay provides minimal carbon sequestration benefit. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.8 Switchgrass

Switchgrass is highly adaptable to soil type. Muscatine 119, Tama 120B, Nicollet 55, Nodaway 220, Clarion 138B, Canisteo 507, and Coland 135 soils are all highly productive for switchgrass. Switchgrass is also productive on soil types that have minimal yield in corn or soy land covers, including Buckney 1636, Okoboji 90, Downs 162D2, and Gara-Armstrong 993E2 soils.

Yield varies only slightly in extreme precipitation scenarios.

One hundred percent grass hay land cover gives the maximum score in sediment retention, erosion control, and nitrate retention, with phosphorus retention scores above 95 in all precipitation scenarios.

As a stream buffer, grassland and low input land use, switchgrass provides points in the habitat categories. To improve habitat scores, try adding some native vegetation including wetland and conservation forest. Switchgrass provides moderate carbon sequestration benefit. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.9 Permanent Pasture

Permanent pasture performs best on soils with excessive to moderate drainage.

Strategic placement can maximize your per-unit yield: Tama 120B, Muscatine 119, Tama 120C2, Nicollet 55, Nodaway 220, and Clarion 138B soils are highly productive for alfalfa and grass hay,

while Buckney 1636, Okoboji 90, and Gara-Armstrong 993E2 soils are approximately half as productive.

Yield varies only slightly in extreme precipitation scenarios, but is much higher for rotational grazing land cover than for permanent pasture land cover. To improve yield, try switching from permanent pasture to rotational grazing.

One hundred percent permanent pasture cover gives the maximum score in nitrate retention, with erosion control scores and sediment retention scores above 90, and phosphorus retention scores above 80 in all precipitation scenarios.

Permanent pasture does not provide points in the habitat categories. To improve habitat scores, try switching to the rotational grazing land cover. Permanent pasture provides minimal carbon sequestration benefit. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.10 Rotational Grazing

Rotational grazing performs best on soils with excessive to moderate drainage.

Strategic placement can maximize your per-unit yield: Tama 120B, Muscatine 119, Tama 120C2, Nicollet 55, Nodaway 220, and Clarion 138B soils are highly productive for rotational grazing and permanent pasture, while Buckney 1636, Okoboji 90, and Gara-Armstrong 993E2 soils are approximately half as productive.

Yield varies only slightly in extreme precipitation scenarios, but is much higher for rotational grazing land cover than for permanent pasture land cover.

One hundred percent rotational grazing cover gives the maximum score in nitrate retention, with erosion control scores above 95, sediment retention scores above 90, and phosphorus retention scores above 85 in all precipitation scenarios.

As a stream buffer, grassland, and low input, high diversity land use, rotational grazing provides the highest level of points in the game wildlife category and relatively high points in the biodiversity category. To improve habitat scores, try diversifying the watershed cover by adding some native vegetation. Rotational grazing provides minimal carbon sequestration benefit. To improve carbon sequestration scores, try adding some woody land covers or wetlands.

1.11 Wetland

Wetlands have a high rate of carbon sequestration in PEWI that applies to all land in the wetland cover type. The biodiversity benefits and nitrate reduction associated with wetland use are optimized by using Strategic Wetlands, located in flat, poorly drained areas of the watershed. Strategic Wetlands may be prairie potholes, or low spots in the left lobe of the watershed, or shallow areas along the river. As a native, high diversity, low input land cover, it provides the highest level of habitat benefits. Game Wildlife benefits from wetlands are derived from percent area of land coverage, as well as using wetland or another stream buffer in appropriate cells.

1.12 Prairie

One hundred percent prairie land cover gives the maximum score in sediment retention, erosion control, phosphorus retention, and nitrate retention.

Prairie has a moderate rate of carbon sequestration in PEWI that applies to all land in the prairie cover type. To improve carbon sequestration scores, try incorporating woody land covers or wetlands.

As a native, high diversity, low input land cover, it provides the highest level of habitat benefits. Game Wildlife benefits from prairie are derived from percent area of land coverage, as well as using prairie or another stream buffer in appropriate cells. Use a diverse mix of prairie and other native vegetation to maximize habitat scores.

1.13 Conventional Forest

Forest land covers perform best on well drained, moderately deep, or moderately wet soils, and are least productive in wet, sandy, saline, or sodic soils.

Strategic placement can maximize your per-unit yield: Gara-Armstrong 993E2 and Downs 162D2 soils are highly productive for forest land covers but perform poorly for annual crops. The conventional forest land cover maximizes wood yield.

As a perennial crop, the conventional forest land cover attains maximum nitrate retention scores. It achieves erosion control, sediment retention, and phosphorus retention scores above 97

Conventional forest land cover provides high points in the carbon sequestration category. To improve carbon sequestration further, include some short-rotation woody land cover.

1.14 Conservation Forest

Forest land covers perform best on well drained, moderately deep, or moderately wet soils, and are least productive in wet, sandy, saline, or sodic soils.

Strategic placement can maximize your per-unit yield: Gara-Armstrong 993E2 and Downs 162D2 soils are highly productive for forest land covers but perform poorly for annual crops. The conventional forest land cover maximizes wood yield.

As a perennial crop, the conservation forest land cover attains maximum nitrate scores. It achieves erosion, sediment, and phosphorus control scores above 97% in all precipitation scenarios.

Conservation forest land cover provides high points in the carbon sequestration category. To improve carbon sequestration further, include some short-rotation woody land cover.

As a native, high diversity, low input land cover, the conservation forest land use provides the highest level of habitat benefits. To maximize habitat scores, try diversifying the watershed by including some other native vegetation covers.

1.15 Short-Rotation Woody Bioenergy

In PEWI, short rotation woody crops perform equally well in all soil, topography, and precipitation scenarios. One hundred percent short-rotation land cover gives the maximum score in carbon sequestration, water and soil quality scores of over 95, and moderate habitat scores.

2 Physical Features

In the Physical Features tab, you'll find information on topography, soil properties, subwatershed boundaries, and strategic wetland areas. These properties can help you strategize the placement of land uses. Physical features influence the Ecosystem Services gained from each land cover choice, from soil and water quality improvement to yield.

2.1 Topographic Relief

Topography influences water runoff patterns, which in turn influences gross erosion, sediment retention, and phosphorus retention scores.

Slope length steepness is also contingent on topography and can range from a factor of .05 to 2.20. See the Erosion Control tab for information about Slope length steepness factor and its role in erosion calculations. Topography also affects the support practice factor in erosion calculations for conservation corn and soy, since terracing and contouring are assumed for elevation grades above 2%. [\[1\]](#)

The runoff factor for phosphorus incorporates Runoff Curve Numbers, which in turn depend on topographic relief. For more information on these calculations, please see the Sediment and Phosphorus tabs.

2.2 Flood Frequency

This feature shows the frequency of flooding for each PEWI grid cell, as found in the Iowa Soil Properties and Interpretations Database (ISPAID) Soil Survey.[\[2\]](#) Flooding categories are defined below, according to the ISPAID manual. In PEWI, flooding frequency is incorporated into yield calculations: yield is determined as per ISPAID calculations as a function of CSR2 results. The CSR2 formula is $CSR2 = S-M-W-F-D \pm EJ$, where F is the field condition including ponding and flooding properties, among other factors.[\[3\]](#) See the Yield tab for more information.

- NONE = 0 = Flooding is not probable.
- RARE = 10 = Flooding is unlikely but possible under unusual weather conditions.
- OCCAS = 20 = Flooding occurs on an average of 50 times or less in 100 years.
- FREQ = 40 = Flooding occurs on an average of more than 50 times in 100 years.
- PONDED = 50 = Standing water on soils in closed depressions. Unless the soils are artificially drained, the water can be removed only by percolation or evapotranspiration. (Ponded is for short duration unless otherwise specified).

2.3 Strategic Wetlands

There are 20 strategic wetlands in PEWI. These include “prairie potholes” in the lower lobe of the watershed that are prone to flooding, but have poor drainage, and shallow areas along the river or a tributary that would regularly have high moisture content in the soil. Utilizing strategic wetlands by planting Wetland in these areas leads to a significant improvement in ecological services.

Wetlands including prairie pothole wetlands and stream buffer wetlands provide native habitat for amphibians, invertebrates, and birds including ring-necked pheasants and mallards,[\[4\]](#) increasing

biodiversity and game wildlife services. It also curbs nitrate pollution; for subwatersheds with at least one strategic wetland, nitrate loss is cut by 52%.^[5]

2.4 Subwatershed Boundaries

This feature depicts the boundaries of each subwatershed, which are important in water runoff patterns influencing nitrate loss. Subwatershed boundaries were realistically calculated based on topographic modeling for the watershed.

2.5 Drainage Class

This feature shows the relative drainage of each pixel in PEWI. Drainage classes are defined below, according to the Iowa Soil Properties and Interpretations Database (ISPAID) manual.^[6] In PEWI, drainage class influences phosphorus, and ISPAID yield values incorporate drainage class information; for more information see the Yield tab.^[7] The subsurface drainage component (SDC) of the phosphorus delivery calculation is a function of flow factor (FF), which is dependent on drainage class. Additionally, the runoff component of the phosphorus calculation incorporates runoff curve numbers based on hydrologic group. See the Phosphorus Tab and Hydrologic Group Tab for more information.

- E = 10 = Excessive
- E-SE = 15 = Excessive-Somewhat excessive
- SE = 20 = Somewhat excessive
- SE-W = 25 = Somewhat excessive-Well
- W = 30 = Well
- W-MW = 35 = Well-Moderately well
- MW = 40 = Moderately well
- MW-SP = 45 = Moderately well-Somewhat poor
- SP = 50 = Somewhat poor
- SP-P = 55 = Somewhat poor-Poor
- P = 60 = Poor
- P-VP = 65 = Poor-Very poor
- VP = 70 = Very poor

2.6 Soil Class

There are thirteen soil classes in PEWI. Each soil class has its own set of properties, including percent slope, flood frequency, drainage class, hydrologic group, soil texture, Corn Suitability Rating, and taxonomic classification. The class of a soil can tell you about its physical and chemical composition, which influences its behavior and production capacity in various situations.

2.6.1 Hydrologic Group

The following chart includes each hydrologic group as defined by the ISPAID Manual.

- Group A = Soils having a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravely sands. These soils have a high rate of water transmission.
- Group B = Soils having a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.
- Group C = Soils having a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.
- Group D = Soils having a very slow infiltration rate (high runoff potential) when thoroughly wet. These consist chiefly of clays that have a high shrink-swell potential, soils with a permanent high water table, soils that have a clay pan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

2.7 Soil Texture

Texture refers to the composition of the soil, including organic matter content and particle size distribution. Abbreviations for texture types used in the ISPAID are below.

- C = Clay
- CL = Clay loam
- CN-SIL = Channery silt loam
- CO-SICL = Cobbly silty clay loam
- CS = Coarse sand
- CSL = Coarse sandy loam
- FS = Fine sand
- FSL = Fine sandy loam
- GR-SL = Gravelly sandy loam
- L = Loam
- LS = Loamy sand
- LFS = Loamy fine sand
- MK = Muck

- MK-SICL = Mucky silty clay loam
- MK-SIL = Mucky silt loam
- S = Sand
- SIC = Silty clay
- SICL = Silty clay loam
- SIL = Silt loam
- SL = Sandy loam
- SP = Sapric
- VFSL = Very fine sandy loam

A table of soil texture by soil type is below.

County	ISPAID Soil Type	Texture
Boone County	Clarion 138B	L
	Buckney 1636	FSL
	Canisteo 507	SICL
	Coland 135	CL
	Nicollet 55	L
	Okoboji 90	MK-SIL
Jasper County	Downs 162D2	SIL
	Gara-Armstrong 993E2	L
	Ackmore-Colo 5B	SIL
	Tama 120C2	SICL
	Tama 120B	SICL
	Muscatine 119	SICL
	Nodaway 220	SIL

2.8 Corn Suitability Rating

The CSR2 calculates the corn suitability value as: $CSR2 = S-M-W-F-D \pm EJ$

- Where S is the taxonomic subgroup class of the soil
- M is the family particle size class (soil particle composition)
- W is a function of the soil's water holding capacity
- F is the field condition factor, including slope, erosion class, flooding, and topsoil thickness
- D incorporates soil depth and "T factor," or soil loss tolerance
- EJ is an expert judgment factor accounting for series abnormalities such as textural extremes or high bulk density.[\[8\]](#)

3 Precipitation

Precipitation is based on historical annual precipitation data from Iowa to simulate climate variability. Scenarios are broken into three categories.

- The Dry category includes the 62.4 cm/yr (24.58 in/yr) scenario, with a probability of 5%, and the 71.6 cm/yr (28.18 in/yr) scenario, with a probability of 15%.
- The Normal category includes the 77.2 cm/yr (30.39 in/yr) scenario, with a probability of 15%, the 81.7 cm/yr (32.16 in/yr) scenario, with a probability of 15%, and the 87.2 cm/yr (34.34 in/yr) scenario, with a probability of 15%.
- The Wet category includes the 92.6 cm/yr (36.47 in/yr) scenario, with a probability of 15%, and the 114.6 cm/yr (45.1 in/yr) scenario, with a probability of 5%.

In PEWI, the level of precipitation influences water quality and soil quality metrics including nitrate and phosphorus runoff, gross erosion, and sediment transport. This is because improved water flow carries a greater quantity of soil and nutrients downstream. Extremes in precipitation also decrease yield for annual crops, mixed fruits and vegetables, alfalfa, grass hay, switchgrass, permanent pasture, and rotational grazing.^[9] Calculations for yield for each land use can be found in the corresponding yield tab.

4 Management Practices

Here you'll find information on management practices used in PEWI's calculations.

4.1 Crop Rotation

A complete chart for Cover Management Factor is found below:

Cover Man- age- ment Factor	C_{ij}	Conventional Corn preceding annual row crop: 0.085, <i>Conservation Corn</i> 0.116, <i>Conservation Soybean</i> 0.150, <i>Conventional Corn</i> 0.200, <i>Conventional Soybean</i> 0.200, <i>Mixed Fruit and Vegetables</i>	0.085 - 0.200
		Conservation Corn preceding annual row crop: 0.020, <i>Conservation Corn</i> 0.031, <i>Conservation Soybean</i> 0.085, <i>Conventional Corn</i> 0.116, <i>Conventional Soybean</i> 0.116, <i>Mixed Fruit and Vegetables</i>	0.020-0.116
		Conventional Soybean/Mixed Fruit and Vegetables preceding annual row crop: 0.156, <i>Conservation Corn</i> 0.178, <i>Conservation Soybean</i> 0.260, <i>Conventional Corn</i> 0.300, <i>Conventional Soybean</i> 0.300, <i>Mixed Fruit and Vegetables</i>	0.156-0.300
		Conservation Soybean preceding annual row crop: 0.052, <i>Conservation Corn</i> 0.055, <i>Conservation Soybean</i> 0.156, <i>Conventional Corn</i> 0.178, <i>Conventional Soybean</i> 0.178, <i>Mixed Fruit and Vegetables</i>	0.052-0.178

5 Modules

The scientific modules in PEWI display ecosystem service scores, the benefits that the watershed provides to people. PEWI tracks ecosystem services in four categories:

- Habitat
- Soil Quality
- Water Quality
- Yield

5.1 Habitat

This category includes two Habitat metrics. Providing habitat to a multitude of species is important for water quality, nutrient recycling, pollination, and biocontrol, including suppression of pests that decrease crop production. A highly diverse ecosystem is also more highly adaptable to regional and global stressors and extreme disturbances, such as fires or flooding.

5.1.1 Biodiversity

Biodiversity is expressed on a 0-10 scale and calculated from a composite score of three nested categories and two location-specific categories.

- 1.5 points maximum awarded for percentage of watershed area in low-input and/or high diversity land uses, conservation annual crops, or permanent pasture.
- 1.5 points maximum awarded for percentage of watershed area in high-diversity land uses.
- 4 points maximum awarded for percentage of watershed area in native land uses.
- 1.5 points maximum awarded for wetland coverage and strategic use.
- 1.5 points maximum awarded for using a stream buffer with an appropriate land use type (conservation crop, forest, grass hay, switchgrass, mixed fruit and vegetable, prairie, rotational grazing, short-rotation woody bioenergy, or wetland). Biodiversity is maximized when each of these criteria is fulfilled and minimized in conventional annual crop scenarios.

5.1.2 Game Wildlife

Game wildlife is expressed on a 0-10 scale and calculated from six categories.

- 1.5 points maximum awarded for percentage of watershed area in low-input and/or high diversity land uses, conservation annual crops, or permanent pasture.
- 4 points maximum awarded for percentage of watershed area in high diversity land uses.
- 1.0 points maximum awarded for at least 5
- 1.0 points maximum awarded for at least 5

- 1.0 points maximum awarded for at least 5
- 1.5 points maximum awarded for percentage of stream buffers with appropriate land use types (conservation corn, conservation soybean, forest, grass hay, switchgrass, mixed fruit and vegetable, prairie, rotational grazing, short-rotation woody bioenergy, or wetland).

5.2 Soil Quality

Soil quality is important for agricultural production and ecosystem health. PEWI monitors two metrics of soil quality: carbon sequestration and erosion prevention. Soils have an immense carbon-holding capacity and, through the actions of plants and microbes, can reduce air pollutant levels. Protecting soils can also minimize erosion, which can diminish crop productivity and deteriorate aquatic stream habitat.[\[10\]](#)

5.2.1 Carbon Sequestration

The annual carbon sequestration of the watershed is calculated as the sum, as land-use type i goes from 1 to n , of the product of land-use type area and the associated rate of sequestration, based on Fissore et al. and Al-Kaisi, Yin, and Licht.^[11] A complete chart for Cover Management Factor is found below:

Source	Annual Row Crop Types	Measured Unit	PEWI Land-use Types	Values (Mg ha ⁻¹ yr ⁻¹)
Fissore et al. (2010)	Forests	Total biomass and soils	Conservation Forest, Conventional Forest	3.67
	Incorporation of cover crops	Soils	Conservation Corn, Conservation Soybean	0.40
	Perennial Grassland	Soils	Prairie	1.07
	Pasture or hay land	Soils	Grass hay, Permanent pasture, Rotational grazing	0.29
	Prairie Potholes	Soils	Wetland	3.05
	Short-Rotation Woody Crops	Total Biomass and Soils	Short-Rotation	4.69
Al-Kaisi et al. (2005)	Switchgrass	SOC, 0-15 cm soil depth	Herbaceous Perennial Bioenergy	1.20
	Corn-Soy Alfalfa Rotation	SOC, 0-15 cm soil depth	Alfalfa	0.50

5.2.2 Erosion Control

Gross erosion is divided into two subcomponents: calculations for ephemeral gully erosion and calculations based on the Revised Universal Soil Loss Equation (RUSLE) for rill and interrill erosion.[12] Ephemeral gully erosion is modeled through a modification to USDA NRCS estimates of 6.7 Mg/ha for Iowa, increased by 50% to 10.1 Mg/ha for conventional annual row-crops and decreased by 50% to 3.4 Mg/ha for conservation annual row-crop practices. This is directionally consistent with current understanding of the relationship between conservation practices and ephemeral gully erosion.[13]

The RUSLE rill and interrill calculation is expressed as $A = R * K * LS * C * P$, where

- A is the estimated net soil loss,
- R is the erosivity factor,
- K is the soil erodibility factor,
- L is the slope length factor,
- S is the slope steepness factor,
- C is the cover-management factor, and
- P is the supporting practices factor (ars.gov).

The R-factor estimates the influence of climate on erosion. In PEWI, the R-factor varies according to changes in precipitation.[14] The K-factor, representing the tendency of soil to become displaced through erosion, is dependent on soil type and is provided by ISPAID.[15] The relationship between L and S is estimated by an NRCS graph depicting slope length as a function of slope grade.[16] C-factor values are based on estimates for the Squaw Creek Watershed in central Iowa, which depends on current and prior land use (see Crop Rotation tab for full chart). [17] P-factor is the product of a contour subfactor and a terrace subfactor, where contouring and terracing is assumed for conservation corn and soybean on slopes above 2%. The P-factor is estimated according to the USLE Erosion Prediction guidelines, using median row grades, low ridge or oriented roughness height, 10 year EI (storm intensity) equal to 80, and maximum recommended terrace interval spacing.[18] The gross erosion in PEWI ranges from $1.36 * 10^2$ Mg/year to $1.1 * 10^5$ Mg/year.

5.3 Water Quality

Water quality is a salient issue for Corn Belt states and other agricultural regions. PEWI tracks three metrics of water quality: nitrate, sediment, and phosphorus. Nitrate-N concentrations in drinking water exceeding the Environmental Protection Agency’s National Regulation limit of 10 mg/L can lead to negative health effects, including blue baby syndrome.[19] Downstream, nitrate-N loading also creates hypoxic (low-oxygen) zones in gulf regions, leading to loss of marine life.[20] Sediment delivery impacts water quality both locally and downstream.[21] Phosphorus loss impacts local producers who incur expenses of phosphorus fertilizer application, regional aquatic ecosystems and human economy through freshwater eutrophication, and distant scales through gulf hypoxia.[22]

5.3.1 Nitrate Retention

In PEWI, Nitrate Retention is the indexed inverse of annual mean nitrate-N concentration in surface water at the watershed outlet. The nitrate-N contribution for the watershed is derived as follows:

$$\sum_{n=1}^n \left[\max \left\{ 100 * PW_i \sum_{j=1}^{m_i} R_{ij} C_{ij}, 2 \right\} * \frac{A(i.)}{A(..)} \right]$$

where i represents the watershed, j represents the pixel, P is a precipitation multiplier, W_i is a strategic wetland multiplier. The multipliers can take the following values:

- P = 0.86 in dry years; 1.69 in normal years following dry years; 2.11 in wet years after dry years; 1.00 in all other years. [23]
- $W_i = 0.48$ for subwatersheds with at least one strategic wetland; 1.00 otherwise. [24]
- $R_{ij} = 0.14 * A_{ij} / A_i$ for conservation corn, conservation soy, conventional corn, conventional soy, and mixed fruit and vegetables, and 0.00 otherwise, where A_{ij} is the area of the grid cell and A_i is the area of the subwatershed (Schilling Libra, 2000). [25]
- $C_{ij} = 0.69 * A_{ij} / A_i$ for conservation corn and conservation soy in the Des Moines Lobe, $0.62 * A_{ij} / A_i$ for conservation corn and conservation soy in the Southern Iowa Drift Plain, and $1.00 * A_{ij} / A_i$ otherwise. [26]

5.3.2 Phosphorus Retention

Phosphorus loading for each grid cell is found by $EC_{ij} + RC_{ij} + SDC_{ij}$

where EC_{ij} represents erosion component, RC_{ij} is a runoff component, and SDC_{ij} is a subsurface drainage component. The erosion component, which measures delivery with sediment, is broken down as:

$$\frac{B_{ij} * E_{ij} * SDR_{ij} * EF_{ij} * STPE_{ij}}{2,000}$$

where B_{ij} is buffer factor, E_{ij} is gross erosion, SDR_{ij} is sediment delivery ratio, EF_{ij} is enrichment factor [27], and $STPE_{ij}$ is soil test phosphorus erosion factor. Information for gross erosion can be found under the erosion control tab. Information for buffer factor and SDR can be found under the sediment control tab. The enrichment factor accounts for the impact of management techniques on soil phosphorus levels. [28] The STP Erosion factor estimates the phosphorus release over time from STP in eroded sediment [29] and is found by: $0.7 * (500 + STP_{ij}) * \frac{2,000}{1,000,000}$

where STP_{ij} is averaged over the major landform region area (MLRA) at 30 ppm for the Des Moines Lobe and 27 ppm for the Southern Iowa Drift Plain. [30]

The runoff component, measuring delivery with runoff, is broken down as:

$$RF_{ij} * PF * (STPR_{ij} + PA_{ij})$$

where RF_{ij} is the runoff factor, PF is the precipitation factor, $STPR_{ij}$ is the soil test phosphorus runoff factor, and PA_{ij} is the phosphorus application factor. RF_{ij} is a function of Runoff Curve Number (RCN) representing the fraction of precipitation that runs off the field. [31] PF is the annual precipitation in million Mg/ha. $STPR_{ij}$ is found by:

$$0.05 + 0.005 * STP_{ij}$$

according to Technical Note 25.[32] The subsurface drainage component, measuring delivery with subsurface drainage, is broken down as FFij*PF*STPDij where FFij is the flow factor and STPDij is the soil test phosphorus drainage factor. Flow factor is either 0 or 0.1 according to P-Index Technical Note 25.[33] STPDij is 0.2 for STP values over 100 and 0.1 for all other values.[34] STP values in PEWI are always under 100.

5.3.3 Sediment Control

Calculations were based on modified P-index guidelines.[35] PEWI's sediment delivery is determined as a product of three factors: gross erosion (Eij), sediment delivery ratio (SDRij), and buffer factor (Bij).

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} (B_{ij} * E_{ij} * SDR_{ij})$$

The buffer factor is equal to 0.5 for land uses considered vegetative buffers: conservation forest, conventional forest, grass hay, herbaceous perennial bioenergy, prairie, short-rotation woody bioenergy, and wetland. Buffer factor is 1.0 for all other land uses.[36]

Information for Gross Erosion can be found in the Erosion Control tab.

The Sediment Delivery Ratio is calculated as a function of drainage area. SDR for cells in the Des Moines Lobe is 0.044 and for cells in the Southern Iowa Drift Plain is 0.278.[37]

5.4 Yield

The General equation is:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} [YieldType] * YP_{ij}$$

where Aij is the area of the grid cell, YBij[Yield Type] is the yield base rate for the type of product, and YPij is the yield precipitation factor. Yield precipitation factors for each precipitation level are shown below:

Landuse Types	Precipitation Factors						
	24.58	24.58 in	30.39 in	32.16 in	34.34 in	36.47 in	45.1 in
Alfalfa	0.75	0.90	1.00	1.00	1.00	0.90	0.75
Grass Hay	0.75	0.90	1.00	1.00	1.00	0.90	0.75
Herbaceous Perennial	0.75	0.90	1.00	1.00	1.00	0.90	0.75
Permanent Pasture	0.75	0.90	1.00	1.00	1.00	0.90	0.75
Rotational Grazing	0.75	0.90	1.00	1.00	1.00	0.90	0.75
Conservation Corn	0.95	1.00	1.00	1.00	1.00	1.00	0.95
Conservation Soybean	0.95	1.00	1.00	1.00	1.00	1.00	0.95
Conventional Corn	0.95	1.00	1.00	1.00	1.00	1.00	0.95
Conventional Soybean	0.95	1.00	1.00	1.00	1.00	1.00	0.95
Mixed Fruit and Vegetables	1.00	1.00	1.00	1.00	1.00	0.90	0.75
All Others	1.00	1.00	1.00	1.00	1.00	1.00	1.00

5.4.1 Corn Yield

The yield equation for corn is given by

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} \left[Corn \right] * YP_{ij}$$

Corn production in PEWI is maximized in a normal precipitation scenario, with all grid cells in either conventional or conservation corn land use types. Yield on a per-unit basis is maximized in Muscatine 119 soil, followed closely by Tama 120B, Nicollet 55, and Clarion 138B. Per-unit yield is minimized in Buckney 1636 soils and Gara-Armstrong 993E2 soils, and is decreased in dry and wet precipitation years. A complete yield base rate chart by soil type can be found below:

County	ISPAID Soil Type	Yield Base Rate Mg/ha-1year
Boone County	Clarion 138B	14.0
	Buckney 1636	0
	Canisteo 507	13.4
	Coland 135	13.2
	Nicollet 55	14.3
	Okoboji 90	11.2
Jasper County	Downs 162D2	12.9
	Gara-Armstrong 993E2	0
	Ackmore-Colo 5B	12.6
	Tama 120C2	13.9
	Tama 120B	14.7
	Muscatine 119	15.1
	Nodaway 220	13.1

5.4.2 Soybean Yield

The yield equation for soybeans is given by:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} \left[Soybean \right] * YP_{ij}$$

Soybean production in PEWI is maximized in a normal precipitation scenario, with all grid cells in either conventional or conservation soybean land use types. Yield on a per-unit basis is maximized in Muscatine 119 soil, followed closely by Tama 120B, Nicollet 55, and Clarion 138B. Per-unit yield is minimized in Buckney 1636 soils and Gara-Armstrong 993E2 soils, and is decreased in dry and wet precipitation years. A complete yield base rate chart by soil type can be found below:

County	ISPAID Soil Type	Yield Base Rate Mg/ha-1year
Boone County	Clarion 138B	4.37
	Buckney 1636	0
	Canisteo 507	4.17
	Coland 135	4.10
	Nicollet 55	4.44
	Okoboji 90	3.49
Jasper County	Downs 162D2	4.03
	Gara-Armstrong 993E2	0
	Ackmore-Colo 5B	3.90
	Tama 120C2	4.30
	Tama 120B	4.57
	Muscatine 119	4.70
	Nodaway 220	4.10

5.4.3 Alfalfa Hay Yield

The yield equation for alfalfa is given by:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * Y B_{ij} \left[\text{Alfalfa} \right] * Y P_{ij}$$

Alfalfa production in PEWI is maximized in a normal precipitation scenario, with all grid cells in alfalfa hay land use type. Yield on a per-unit basis is maximized in Tama 120B, followed closely by Muscatine 119, Tama 120C2, and Nicollet 55. Per-unit yield is minimized in Buckney 1636 soils, Gara-Armstrong 993E2 soils, and Okoboji 90 soils, and it is decreased in extreme precipitation years of 62.4 cm/yr (24.58 in/yr) and 114.6 cm/year (45.1 in/yr). A complete yield base rate chart by soil type can be found below:

County	ISPAID Soil Type	Yield Base Rate Mg/ha-1year
Boone County	Clarion 138B	14.1
	Buckney 1636	8.07
	Canisteo 507	9.64
	Coland 135	9.42
	Nicollet 55	14.4
	Okoboji 90	8.07
Jasper County	Downs 162D2	12.6
	Gara-Armstrong 993E2	8.07
	Ackmore-Colo 5B	9.19
	Tama 120C2	14.6
	Tama 120B	15.5
	Muscatine 119	15.0
	Nodaway 220	14.1

5.4.4 Grass Hay Yield

The yield equation for grass hay is given by:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} \left[GrassHay \right] * YP_{ij}$$

Grass Hay production in PEWI is maximized in a normal precipitation scenario, with all grid cells in grass hay land use type. Yield on a per-unit basis is maximized in Tama 120B, followed closely by Muscatine 119, Tama 120C2, and Nicollet 55. Per-unit yield is minimized in Buckney 1636 soils, Gara-Armstrong 993E2 soils, and Okoboji 90 soils, and it is decreased in extreme precipitation years of 62.4 cm/yr (24.58 in/yr) and 114.6 cm/year (45.1 in/yr). A complete yield base rate chart by soil type can be found below:

County	ISPAID Soil Type	Yield Base Rate Mg/ha-1year
Boone County	Clarion 138B	14.1
	Buckney 1636	8.07
	Canisteo 507	9.64
	Coland 135	9.42
	Nicollet 55	14.4
	Okoboji 90	8.07
Jasper County	Downs 162D2	12.6
	Gara-Armstrong 993E2	8.07
	Ackmore-Colo 5B	9.19
	Tama 120C2	14.6
	Tama 120B	15.5
	Muscatine 119	15.0
	Nodaway 220	14.1

5.4.5 Cattle Yield

The yield equation for head of cattle is given by:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} \left[Cattle \right] * YP_{ij}$$

where $YB_{ij} \left[Cattle \right] = YB_{ij} \left[Alfalfa \right] * \frac{SU}{DI * GS}$

- SU is the seasonal utilization rate, which is equal to .35 in permanent pasture and .55 in rotational grazing
- DI is average daily intake, estimated at 3% of live bodyweight of .544 Mg, equal to .0163 Mg/day/animal
- GS is the grazing season length, April 15 to November 1, which is 200 days

Cattle production in PEWI is maximized in any of the middle five precipitation scenarios, with all grid cells in rotational grazing land use type. Yield on a per-unit basis is maximized

in Tama 120B, followed closely by Muscatine 119, Tama 120C2, and Nicollet 55. Per-unit yield is minimized in Buckney 1636 soils, Gara-Armstrong 993E2 soils, and Okoboji 90 soils, and it is low in extreme precipitation years of 62.4 cm/year or 114.6 cm/year. A complete yield base rate chart by soil type can be found below:

County	ISPAID Soil Type	Yield Base Rate Mg/ha-1year
Boone County	Clarion 138B	14.1
	Buckney 1636	8.07
	Canisteo 507	9.64
	Coland 135	9.42
	Nicollet 55	14.4
	Okoboji 90	8.07
Jasper County	Downs 162D2	12.6
	Gara-Armstrong 993E2	8.07
	Ackmore-Colo 5B	9.19
	Tama 120C2	14.6
	Tama 120B	15.5
	Muscatine 119	15.0
	Nodaway 220	14.1

5.4.6 Mixed Fruit and Vegetable Yield

The yield equation for mixed fruit and vegetables is:

$$\sum_{i=1}^n \sum_{j=1}^{m_i} A_{ij} * YB_{ij} \left[Cattle \right] * YP_{ij}$$

where $YB_{ij} \left[FruitVeg \right] = 0.25 * STM * (GRY + GBY + SQY + STY)$ and ranges from 6.58 to 16.5 Mg/ha/yr

- GRY = Grape Yield = 17.6 Mg/ha/yr
- GBY = Green Bean Yield = 6.73 Mg/ha/yr
- SQY = Squash Yield = 33.6 Mg/ha/yr
- STY = Strawberry Yield = 7.85 Mg/ha/yr

Soil Texture Multiplier	Fine Sandy Loam (FSL)	1.00
	Silt loam (SIL)	0.90
	Loam (L)	0.85
	Clay loam (CL), Mucky silt loam (MK-SIL), Silty clay loam (SICL)	0.40

A chart for Soil Texture Multiplier (STM) for each soil type can be found in the Texture tab. Green bean, squash, and strawberry yield values are from Iowa State University Extension, with adjustments for unfavorable soil texture.[?] Due to lack of published data for Iowa grapes, regionally

similar data was used from New York State indicating 17.6 Mg/ha.[?] Mixed fruit and vegetable production is maximized in a dry or normal precipitation year. Yield on a per-unit basis is maximized in fine sandy loam (Buckney 1636 soils), followed by silt loam (Downs 162D2, Ackmore-Colo 5B, and Nodoway 220) soils. It is minimized in clay loam (Coland 135), mucky silt loam (Okoboji 90), and silty clay loam (Tama 120C2, Tama 120B, and Muscatine 119).

5.4.7 Wood Yield

Wood production is based on estimates generated using the 2007 Iowa Woodland Suitability Composite.[17] Yield estimates assume appropriate selection of tree species for a given location. A 30% reduction in wood production estimates for conservation forest is applied to account for differences in management.

5.4.8 Switch Grass Yield

Switchgrass biomass yields are based on a modification to Iowa Soil Properties and Interpretations Database (ISPAID) corn suitability ratings.[18]

5.4.9 Woody Biomass

Woody biomass yield estimates are based on a 10-year aspen biomass of 224 Mg/ha.[19] Annual production assumes temporally spaced plantings such that a constant 22.4 Mg/ha is harvested each year.

6 Functionality

This section describes how to navigate the PEWI application, including play mode options and watershed editing tools.

6.1 Navigating the PEWI Workspace

Opening Sandbox or playing a PEWI Level brings you to the PEWI Workspace. In the workspace, there are many tools that you can use to edit the watershed and monitor ecosystem services.

6.1.1 Tool Panel

The tool panel has seven features to use in designing your watershed. The Land Use tab includes 15 land cover options to use in your watershed, ranging from annual and perennial crops to pasture, forests, and native land cover.

The Precipitation tab shows the precipitation for each year. PEWI simulates real life by choosing randomly from seven precipitation levels. To override the presets, select a precipitation level from the drop-down menu.

The Years tab allows you to add another year of simulation to your watershed. Some of the impacts of land use choices vary with weather or previous practices. You can explore these effects by playing across multiple years.

The Indicator Tab includes overlay maps for Nitrate, Erosion, and Phosphorus risk. You can edit the land use while in overlay mode to target a specific risk area.

The Physical Features tab includes five watershed behavior overlay maps displaying flood frequency, strategic wetland areas, subwatershed boundaries, drainage class, and soil class. You can edit the land use while in overlay mode to target a specific region.

The Yield Tab allows you to strategically place landuse types on the map, by providing information for each yield type and their respective yield base rates based off soil type. To learn more about this feature, check out the Yield Tab video.

The buttons located above the Toolbar are for placing landuse types on the map. The leftmost button is the individual brush, allowing you to paint one landuse type at a time. The middle button allows you to paint multiple tiles at once by dragging and clicking the corners. The arrow button is the Undo button, allowing you to undo previous changes. Any amount of previous changes can be undone. You can also undo changes by pressing the "U" hotkey on the keyboard.

6.1.2 Results Panel

The results page provides data visualization and numerical results for your watershed design.

Land Use Graph The first circle graph gives the percent of the watershed you've devoted to each of the land uses. Scroll over the graph or the legend on the right to view the quantitative data for that land use. Use the up and down arrows to switch between List view and Category view.

List view displays each land use individually. Category view groups land uses into seven categories: annual grain, annual legume, mixed fruit and vegetables, pasture, non-pasture perennial herbs, perennial legume, and perennial wetland. Use the arrows to switch between years.

Ecosystem Services Graph The ecosystem services radar plot shows the zero-to-100 score of ecosystem services pertaining to water quality, soil health, and habitat. A watershed design with high scores in many ecosystem services will produce a radar plot shape with a large area, while a watershed design with low ecosystem service scores will produce a smaller shape.

Precipitation Graph The precipitation graph shows the amount of precipitation for each year of the simulation.

Data Sheet Click on the abacus icon at the top of the Results page to access the data tab with two numerical tables. Both tables provide three metrics: the zero-to-100 scale used in the Scores tab, English units, and metric units. The first table gives the percent area of each land use by year, and the second gives ecosystem service scores by year. At the bottom of the page are precipitation data and strategic wetland use for each year.

6.1.3 Comment Panel

Click the comment bubble to show or hide the comment panel. In Level mode, the comment panel will show instructions for completing the next objective in the level.

6.1.4 Index Information Box

The Index Information Box is located at the bottom right portion of the screen. Each tab is given particular information about their intended usage and each individual feature for each tab is given some information in the Index Box and where to learn more, by directing the User to its respective Index entry. Every tab, from the Landuse Tab to the Yield Tab has information. Check them out!

6.1.5 View Perspective

Press or hold D to rotate the watershed clockwise. Press or hold A to rotate the watershed counterclockwise. Press or hold W to pivot your viewpoint toward the horizon (flat) Press or hold S to pivot your viewpoint toward the perpendicular Right click to rotate the canvas with your mouse.

Use the up and down arrow keys to zoom in and out. Use the left and right arrow keys to pan left and right.

Press T to toggle 3-D topographical view.

Press E to reset the camera position to default.

6.1.6 Save Watershed

Click the download icon in the upper right-hand corner to save your watershed to your computer. Upload to continue with your design.

6.2 Main Menu

Select Sandbox to explore the watershed on your own. Select Play to play an interactive PEWI level. Select Utilities to design a level, set up Multiplayer Mode, or aggregate Multiplayer Maps. For more information, see the Levels tab and Multiplayer Mode tab.

6.3 Guided Play

PEWI comes with predesigned, thematic levels, ranging from introductory to complex. The N-Factor (Nitrogen) set focuses on the interaction between nitrate levels, physical features, land uses, and other ecosystem service scores. The P-Factor (Phosphorus) focuses on improving phosphorus, sediment, and erosion levels. The B-Factor (Biodiversity) focuses on habitat variety and suitability.

To create your own level, choose Design a Level from the Utilities menu. You can write a customized lesson with your own objectives and board presets.

6.4 Multiplayer Mode

PEWI offers a Multiplayer Mode. Under Utilities, click on the "Set-Up Multiplayer" button. In Multiplayer Mode, up to 6 Players can be added. Click on the Player numbers to draw out boundaries for each Player on the map with the brush tools.

You can also merge and combine Player boundaries and add and remove Players. Once finished, press "V" to download a Multi-Map file. Return to sandbox and click on the down arrow on the upper right part of the screen. Click on Upload and select a Player from your map files. In this case, Player 1 will be given only their boundaries drawn out for them to paint landuse types on. Scores in the Results are adjusted for these sections.

You can continue to adjust any Player maps as seen fit. Once finished, hit download to update the maps. To see all of the maps put together at once, go to "Utilities" and "Aggregate Multi-Maps". Add the files that have been changed and download the aggregate maps. Go back to "Sandbox" and upload the combined map file.

6.5 Pause Menu

The Pause Menu can be accessed by pressing the "Escape Key" Clicking on the Index will take you to the various Index Entries, where you can learn about different features of PEWI with videos The "Instructor Options" feature can also be accessed where customizations can be made and are further explored in the "Instructor Options" video The final part of the Pause Menu is the "Main Menu" feature, where you can go back to PEWI's main menu.

6.6 Hover Information

Specific information can be shown when hovered over tiles on the map. Each map when hovered over will display the tile's coordinates, the Land Cover that is currently being placed on the map, Precipitation level, and Soil Type. On some maps, there are some additional bits of information when hovered over. For the Nitrate, Erosion, and Phosphorus Maps, raw scores will show up that correspond to their map keys. Strategic Wetlands and Subwatershed Boundaries have their sections appear, while Flood Frequency, Drainage and Soil Class have their map key information appear. Yield maps all have their raw scores appear as well. Hover information displayed can be customized in the Instructor Options page and can be toggled on and off accordingly.

6.7 Hotkeys

- Shift + Click Fill the watershed with the selected land type
- E Reset camera position
- Escape Access Pause Menu
- R Randomize all land use types
- T Toggle topographical view
- U Undo previous land change
- O Toggle previous overlay
- B Toggle recording feature
- V Create multiplayer maps in multiplayer utility
- D Rotate clockwise
- A Rotate counterclockwise
- W Pivot toward the horizon (flat)
- S Pivot toward the perpendicular
- Q Toggle Flying Mode
- Use the up and down arrow keys to zoom in and out
- Use the left and right arrow keys to pan left and right
- Right Click Rotate watershed with your mouse

6.8 Hotkey Customization

By pressing the Esc Key and going to Instructor Options, a list of Hotkeys is available with their default settings. However, the hotkeys can be customized and adjusted as much as possible, with the option of having two hotkeys per feature. You can also reset the hotkeys to bring them back to their default settings.

6.9 Randomization

Press "R" to randomize the landuse tiles on the watershed. The watershed can be randomized an infinite number of times. Press the Esc key to go to Instructor Options, where you can turn off landuse types. By turning off Conventional Corn, the landuse type will no longer appear when randomized. This applies for any other landuse types that are turned off. Landuse types can be turned back on and put back into randomization through the Instructor Options as well.

6.10 Undo Function

In PEWI, there is an undo function that allows you to undo changes to the map, such as painting tiles or randomizing the map. Once you want to make changes to your previous edits to the map, click on the undo button, located next to the paint brush buttons above the tool box. You can also undo your changes by pressing the "Undo Hotkey" which is "U" on the keyboard. You can undo up to your oldest change to the map. The undo hotkey also extends to painting multiple tiles at once.

6.11 Print Function

You can print your maps, results, and much more with Pewi. Activate the Print function by pressing "P" on the keyboard and select what you want to be printed out on a PDF. There are plenty of categories to choose from to print, including: Landuse Type Maps for set years, results, levels maps, and much more. Buttons such as "Select All" and "Reset" can help you choose. Once finished, you can click "Preview PDF" to see what will be on your PDF and then "Download" when ready. You can then print out your PDF once you save it. When you're finished, click "Exit".

6.12 Record

By pressing B, you can record your actions with PEWI. The record feature will keep track of all actions. When you're done recording, press B again to save what was recorded. If you want to see your recording, go back to the main menu and click on "Utilities", then "Run User Simulation" where you can upload your recording. Once clicking "Click here to begin Simulation", you can view your recording. A bar at the bottom left of the screen will show the recording's progress and it's possible to fast forward and rewind. Once finished, you can replay or return to the Main Menu.

6.13 About

People in Ecosystems/Watershed Integration (PEWI) is a simple web-based educational game designed to provide a scientific platform for teaching, discussing, and evaluating the tradeoffs associated with agricultural land use and management. Players iteratively manipulate land use annually for three years in a virtual 6000 acre watershed to meet a variety of goals. The tool computes a variety of results, including agricultural yields, soil erosion, stream pollution, and wildlife habitat.

PEWI v3 is developed and maintained to be compatible with Google Chrome (versions 49+), Internet Explorer (versions 11+), Mozilla Firefox (versions 47+), and Safari (versions 9.1+).

PEWI is a product of Iowa State University Department of Natural Resource Ecology and Management. The product license can be found [here](#). Members of the PEWI development team were Lisa Schulte Moore, Assata Caldwell, Carrie Chennault, Justin Choe, Ryan Frahm, Noah Hagen, Jake Hill, Charles Labuzzetta, Laura Roy, Nancy Shryock, John Tyndall, Robert Valek, and John VanDyk.

PEWI v3 was funded through grants from the Iowa State University Department of Agronomy, McKnight Foundation, and USDA Forest Service.

More information on PEWI, including information on tool development, learning exercises, additional acknowledgements, and how you can contribute can be found at the project website: www.nrem.iastate.edu/pewi. Questions can be sent to pewi@iastate.edu.

References

- [1] K. G. Renard et al., Predicting Soil Erosion by Water: A Guide to Conservation Planning with RUSLE (Washington, D.C: U.S. Department of Agriculture Agriculture Research Service, 1997); USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “Section I FOTG: USLE Erosion Prediction” (Des Moines, Iowa, September 2002), http://kaleita.public.iastate.edu/Iowa_USLE_Guide.pdf.
- [2] Iowa State University, “Iowa Soil Properties and Interpretations Database,” 2010, <http://www.extension.iastate.edu/soils/ispaid>.
- [3] C. L. Burras et al., “Corn Suitability Rating 2 (CSR2) Equation and Component Values” (Iowa State University, March 8, 2015), <http://www.extension.iastate.edu/soils/sites/www.extension.iastate.edu/files/soils/ISU%20CSR2%20formula%20with%20values%20V1.1%2009Mar15.pdf>.
- [4] Carrie Chennault, “People in Ecosystems / Watershed Integration: Visualizing Ecosystem Services Tradeoffs in Agricultural Landscapes” (Master’s thesis, Iowa State University, 2014), 26, 30.
- [5] Chennault, “People in Ecosystems / Watershed Integration,” p. 85 (Table 2-8).
- [6] G. A. Miller et al., “ISPAID 7.3 Manual” (Iowa State University, Iowa Agriculture and Home Economics Experiment Station and University Extension, 2010), www.extension.iastate.edu/Documents/soils/ISPAID_73man.doc
- [7] Miller et al., “ISPAID 7.3 Manual.”
- [8] C.L. Burras et al., “Corn Suitability Rating 2 (CSR2) Equation and Component Values” (Iowa State University, March 8, 2015), <http://www.extension.iastate.edu/soils/sites/www.extension.iastate.edu/files/soils/ISU%20CSR2%20formula%20with%20values%20V1.1%2009Mar15.pdf>.
- [9] G. A. Miller et al., “ISPAID 7.3 Manual” (Iowa State University, Iowa Agriculture and Home Economics Experiment Station and University Extension, 2010), www.extension.iastate.edu/Documents/soils/ISPAID_73man.doc; Emily Heaton and Matt Liebman

- (Iowa State University, personal communication, 2014); Emily Heaton (Iowa State University, personal communication, 2014).
- [10] J. Lyons and C.C. Courtney, “A Review of Fisheries Habitat Improvement Projects in Warmwater Streams with Recommendations for Wisconsin USA,” Wisconsin Department of Natural Resources Technical Bulletin, no. 169 (1990): 1–34; E.G. Smith et al., “Soil Quality Attribute Time Paths: Optimal Levels and Values,” *Journal of Agricultural and Resource Economics* 25, no. 1 (2000): 307–24.
 - [11] Cinzia Fissore et al., “Limited Potential for Terrestrial Carbon Sequestration to Offset Fossil-Fuel Emissions in the Upper Midwestern US,” *Frontiers in Ecology and the Environment* 8, no. 8 (October 2010): 409–13, doi:10.1890/090059; Mahdi M. Al-Kaisi, Xinhua Yin, and Mark A. Licht, “Soil Carbon and Nitrogen Changes as Influenced by Tillage and Cropping Systems in Some Iowa Soils,” *Agriculture, Ecosystems Environment* 105, no. 4 (March 5, 2005): 635–47, doi:10.1016/j.agee.2004.08.002.
 - [12] Carrie Chennault, “People in Ecosystems / Watershed Integration: Visualizing Ecosystem Services Tradeoffs in Agricultural Landscapes” (Master’s thesis, Iowa State University, 2014), 44.
 - [13] Ibid.
 - [14] Kenneth G. Renard and Jeremy R. Freimund, “Using Monthly Precipitation Data to Estimate the R-Factor in the Revised USLE,” *Journal of Hydrology* 157, no. 1 (May 1, 1994): 287–306, doi:10.1016/0022-1694(94)90110-4.
 - [15] G. A. Miller et al., “ISPAID 7.3 Manual” (Iowa State University, Iowa Agriculture and Home Economics Experiment Station and University Extension, 2010), www.extension.iastate.edu/Documents/soils/ISPAID_73man.doc.
 - [16] USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “Iowa Technical Note No. 29: Choosing the Planning Area of a Field By ‘dominant Critical Area’.” (Des Moines, Iowa, 2008), <http://efotg.sc.egov.usda.gov/references/public/IA/AgronomyTECHnote29Nov08.pdf>.
 - [17] A.A. Wendt, “Watershed Planning in Central Iowa: An Integrated Assessment of the Squaw Creek Watershed for Prioritization of Conservation Practice Establishment” (Master’s thesis, 2007), ProQuest Dissertations and Theses Full Text database (UMI No. 1446134).
 - [18] Chennault, “People in Ecosystems / Watershed Integration,” 44-46.
 - [19] National Primary Drinking Water Regulations, 40 C.F.R. § 141 (2010).
 - [20] US EPA [United States Environmental Protection Agency], “Hypoxia in the Northern Gulf of Mexico: An Update by the EPA Science Advisory Board.” (Washington, D.C, December 2007), http://water.epa.gov/type/watersheds/named/msbasin/upload/2008_1_31_msbasin_sab_report_2007.pdf.
 - [21] J. Lyons and C.C. Courtney, “A Review of Fisheries Habitat Improvement Projects in Warmwater Streams with Recommendations for Wisconsin USA,” Wisconsin Department of Natural Resources Technical Bulletin, no. 169 (1990): 1–34; E.G. Smith et al., “Soil Quality Attribute Time Paths: Optimal Levels and Values,” *Journal of Agricultural and Resource Economics* 25, no. 1 (2000): 307–24; K.B. Vache, J.M. Eilers, and M.V. Santelmann, “Water Quality Modeling of Alternative Agricultural Scenarios in the US Corn

- Belt,” *Journal of the American Water Resources Association* 38, no. 3 (2002): 773–87, doi:10.1111/j.1752-1688.2002.tb00996.x.; Iowa, “Iowa Nutrient Reduction Strategy. A Science and Technology-Based Framework to Assess and Reduce Nutrients to Iowa Waters and the Gulf of Mexico” (Iowa Department of Agriculture and Land Stewardship, Iowa Department of Natural Resources, and Iowa State University College of Agriculture and Life Sciences, 2013); R.B. Alexander et al., “Differences in Phosphorus and Nitrogen Delivery to the Gulf of Mexico from the Mississippi River Basin,” *Environmental Science Technology* 42, no. 3 (2008): 822–30, doi: 10.1021/es0716103.
- [22] Alexander et al., “Phosphorus and Nitrogen Delivery to the Gulf of Mexico”; Iowa, “Iowa Nutrient Reduction Strategy”; US EPA, “Hypoxia in the Northern Gulf of Mexico”; Linda M. Jacobson, Mark B. David, and Laurie E. Drinkwater, “A Spatial Analysis of Phosphorus in the Mississippi River Basin,” *Journal of Environment Quality* 40, no. 3 (2011): 931, doi:10.2134/jeq2010.0386.
- [23] G.W. Randall and D.J. Mulla, “Nitrate Nitrogen in Surface Waters as Influenced by Climatic Conditions and Agricultural Practices,” *Journal of Environmental Quality* 30, no. 2 (2001): 337–44.
- [24] Thomas Isenhardt (Iowa State University, personal communication, 2013).
- [25] K.E. Schilling and R.D. Libra, “The Relationship of Nitrate Concentrations in Streams to Row Crop Land Use in Iowa. *Journal of Environmental Quality*,” *Journal of Environmental Quality* 29, no. 6 (2000): 1846–51.
- [26] Iowa, “Iowa Nutrient Reduction Strategy. A Science and Technology-Based Framework to Assess and Reduce Nutrients to Iowa Waters and the Gulf of Mexico” (: Iowa Department of Agriculture and Land Stewardship, Iowa Department of Natural Resources, and Iowa State University College of Agriculture and Life Sciences, 2013).
- [27] USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “Iowa Technical Note No. 25: Iowa Phosphorus Index” (Des Moines, Iowa, 2004a), http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_007643.pdf.
- [28] USDA NRCS, “Technical Note 25”; Carrie Chennault, “People in Ecosystems / Watershed Integration: Visualizing Ecosystem Services Tradeoffs in Agricultural Landscapes,” Table 2-12 (Master’s thesis, Iowa State University, 2014).
- [29] USDA NRCS, “Technical Note 25.”
- [30] Iowa, “Iowa Nutrient Reduction Strategy. A Science and Technology-Based Framework to Assess and Reduce Nutrients to Iowa Waters and the Gulf of Mexico” (: Iowa Department of Agriculture and Land Stewardship, Iowa Department of Natural Resources, and Iowa State University College of Agriculture and Life Sciences, 2013).
- [31] A.P. Mallarino et al., “Background and Basic Concepts of the Iowa Phosphorus Index: A Support Document to the USDA-NRCS Field Office Technical Note 25.” (Des Moines, Iowa, 2005), Iowa State University, USDA Natural Resource Conservation Service, and USDA Agricultural Research Service National Soil Tilth Laboratory, Retrieved from <ftp://ftp-fc.sc.egov.usda.gov/IA/technical/BackgroundBasics.pdf>; USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “National Engineering Handbook. Part 630, Hydrology.” (Washington, D.C, 2004b), <ftp://ftp.wcc.nrcs.usda.gov/wntsc/HH/NEHhydrology/ch9.pdf>.

- [32] USDA NRCS, “Technical Note 25.”
- [33] Ibid.
- [34] Ibid.
- [35] 1. USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “Iowa Technical Note No. 25: Iowa Phosphorus Index” (Des Moines, Iowa, 2004a), http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_007643.pdf; A.P. Mallarino et al., “Background and Basic Concepts of the Iowa Phosphorus Index: A Support Document to the USDA-NRCS Field Office Technical Note 25.” (Des Moines, Iowa, 2005), Iowa State University, USDA Natural Resource Conservation Service, and USDA Agricultural Research Service National Soil Tilth Laboratory, Retrieved from <ftp://ftp-fc.sc.egov.usda.gov/IA/technical/BackgroundBasics.pdf>.
- [36] USDA NRCS, “Technical Note 25.”
- [37] USDA NRCS [United States Department of Agriculture Natural Resources Conservation Service], “Erosion and Sediment Delivery” (Des Moines, Iowa, 1998), http://efotg.sc.egov.usda.gov/references/public/IA/Erosion_and_sediment_delivery.pdf.
- [38] H. G. Taber, “Commercial Vegetable Production – A Small Farm Opportunity?” (Iowa State University Extension and Outreach, 2009), <http://ia.foodmarketmaker.com/uploads/0a88216bd7a96e10152b7c7a6891a667.pdf>.
- [39] R.M. Post and J.A. Robinson, “The SARE - Cornell Organic Grape Project” (The Third N.J. Shaulis Organic Grape and Wine Production Symposium, New York, 1995); K. Delate and H. Friedrich, “Organic Apple and Grape Performance in the Midwestern US,” *Sustainability of Horticultural Systems in the 21st Century* 638 (2004): 309–20.