

Dark Matter Benchmark Models for Early LHC Run-2 Searches: Report of the ATLAS/CMS Dark Matter Forum

Daniel Abercrombie^a, Nural Akchurin^b, Ece Akilli^c, Juan Alcaraz Maestre^d,
Brandon Allen^a, Barbara Alvarez Gonzalez^e, Jeremy Andrea^f, Alexandre Arbey^g,
Georges Azuelos^h, Patrizia Azziⁱ, Mihailo Backović^j, Yang Bai^k, Swagato Banerjee^k,
James Beacham^l, Alexander Belyaev^m, Antonio Boveia (Forum organizer and
editor)^e, Amelia Jean Brennanⁿ, Oliver Buchmueller^o, Matthew R. Buckley^p,
Giorgio Busoni^q, Michael Buttignol^r, Giacomo Cacciapaglia^s, Regina Caputo^t,
Linda Carpenter^u, Nuno Filipe Castro^v, Guillelmo Gomez Ceballos^a, Yangyang
Cheng^w, John Paul Chou^x, Arely Cortes Gonzalez^y, Chris Cowden^z, Francesco
D'Eramo^{aa}, Annapaola De Cosa^{ab}, Michele De Gruttola^e, Albert De Roeck^e, Andrea
De Simone^{ac}, Aldo Deandrea^{ad}, Zeynep Demiragli^a, Anthony DiFranzo^{ae}, Caterina
Doglioni (Forum organizer and editor)^{af}, Tristan du Pree^e, Robin Erbacher^{ag},
Johannes Erdmann^{ah}, Cora Fischer^{ai}, Henning Flaecher^{aj}, Patrick J. Fox^{ak},
Benjamin Fuks^{al}, Marie-Helene Genest^{am}, Bhawna Gomber^k, Andreas Goudelis^{an},
Johanna Gramling^c, John Gunion^{ao}, Kristian Hahn^{ap}, Ulrich Haisch¹, Roni
Harnik^{aq}, Philip C. Harris^e, Kerstin Hoepfner^{ar}, Siew Yan Hoh^{as}, Dylan George
Hsu^a, Shih-Chieh Hsu^{at}, Yutaro Iiyama^a, Valerio Ippolito^{au}, Thomas Jacques^{av},
Xiangyang Ju^k, Felix Kahlhoefer^{aw}, Alexis Kalogeropoulos^{ax}, Laser Seymour
Kaplan^k, Lashkar Kashi^k, Valentin V. Khoze^{ay}, Raman Khurana^{az}, Khristian
Kotov^{ba}, Dmytro Kovalskyi^a, Suchita Kulkarni (editor)^{bb}, Shuichi Kunori^{bc}, Viktor
Kutzner^{bd}, Hyun Min Lee^{be}, Sung-Won Lee^{bf}, Seng Pei Liew^{bg}, Tongyan Lin^{bh},
Steven Lowette (Forum organizer and editor)^{bi}, Romain Madar^{bj}, Sarah Malik
(Forum organizer and editor)^{bk}, Fabio Maltoni^{bl}, Mario Martinez Perez^{bm}, Olivier
Mattelaer^{bn}, Kentarou Mawatari^{bo}, Christopher McCabe^{bp}, Théo Megy^{bq}, Enrico
Morgante^{br}, Stephen Mrenna (Forum organizer and editor)^{bs}, Chang-Seong
Moon^{bt}, Siddharth M. Narayanan^a, Andy Nelson^{bu}, Sérgio F. Novaes^{bt}, Klaas Ole
Padeken^{bv}, Priscilla Pani (editor)^{bw}, Michele Papucci^{bx}, Manfred Paulini^{by},
Christoph Paus^a, Jacopo Pazzini^{bz}, Björn Penning^{ca}, Michael E. Peskin^{cb}, Deborah
Pinna^{ab}, Massimiliano Procura^{cc}, Shamona F. Qazi^{cd}, Davide Racco^{ce}, Emanuele
Re¹, Antonio Riotto^{cf}, Thomas G. Rizzo^{cg}, Rainer Roehrig^{ch}, David Salek^{ci}, Arturo
Sanchez Pineda^{cj}, Subir Sarkar^{ck}, Alexander Schmidt^{cl}, Steven Randolph
Schramm^c, William Shepherd^{cm}, Gurpreet Singh^{cn}, Livia Soffi^{co}, Norraphat
Srimanobhas^{cp}, Kevin Sung^{cq}, Tim M. P. Tait^{cr}, Timothee Theveneaux-Pelzer^{cs},
Marc Thomas^{ct}, Mia Tosi^{cu}, Daniele Trocino^{cv}, Sonaina Undleeb^{cw}, Alessandro
Vichi^{cx}, Fuquan Wang^k, Lian-Tao Wang^{cy}, Ren-Jie Wang^{cz}, Nikola Whallon^{da},
Steven Worm^{db}, Mengqing Wu^{dc}, Sau Lan Wu^k, Hongtao Yang^k, Yong Yang^{ab},
Shin-Shan Yu^{dd}, Bryan Zaldivar^{de}, Marco Zanetti^{bz}, Zhiqing Zhang^{df}, Alberto
Zucchetta^{bz}

^aMIT, USA

^bTexas Tech University, USA

^cUniversité de Genève, DPNC, Switzerland

^dCentro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Spain

^eCERN, Switzerland

- ^fInstitut Pluridisciplinaire Hubert Curien/Département Recherches Subatomiques, Université de Strasbourg/CNRS-IN2P3, France
- ^gUniversité de Lyon and Centre de Recherche Astrophysique de Lyon, CNRS and Ecole Normale Supérieure de Lyon, France and CERN Theory Division, Switzerland
- ^hUniversity of Montreal and TRIUMF, Canada
- ⁱINFN Padova, Italy
- ^jCentre for Cosmology, Particle Physics and Phenomenology (CP3), Université catholique de Louvain, Belgium
- ^kUniversity of Wisconsin-Madison, USA
- ^lOhio State University, USA
- ^mRutherford Appleton Laboratory and University of Southampton, United Kingdom
- ⁿThe University of Melbourne, Australia
- ^oImperial College London, United Kingdom
- ^pDepartment of Physics and Astronomy, Rutgers University, USA
- ^qSISSA and INFN, Sezione di Trieste, Italy
- ^rInstitut Pluridisciplinaire Hubert Curien/Département Recherches Subatomiques, Université de Strasbourg/CNRS-IN2P3, France
- ^sUniversité de Lyon and Université Lyon 1, CNRS/IN2P3, UMR5822, IPNL, France
- ^tSanta Cruz Institute for Particle Physics, Department of Physics and Department of Astronomy and Astrophysics, University of California at Santa Cruz, USA
- ^uOhio State University, USA
- ^vLIP-Minho, Braga, and Departamento de Física e Astronomia, Faculdade de Ciências da Universidade do Porto, Portugal
- ^wUniversity of Chicago, USA
- ^xRutgers University, USA
- ^yIFAE Barcelona, Spain
- ^zTexas Tech University, USA
- ^{aa}University of California and LBNL, Berkeley, USA
- ^{ab}University of Zurich, Switzerland
- ^{ac}SISSA and INFN, Sezione di Trieste, Italy
- ^{ad}Université de Lyon and Université Lyon 1, CNRS/IN2P3, UMR5822, IPNL, France
- ^{ae}Department of Physics and Astronomy, University of California, Irvine and Theoretical Physics Department, Fermilab, USA
- ^{af}Lund University, Sweden
- ^{ag}University of California, Davis, USA
- ^{ah}Institut für Experimentelle Physik IV, Technische Universität Dortmund, Germany
- ^{ai}IFAE Barcelona, Spain
- ^{aj}H.H. Wills Physics Laboratory, University of Bristol, United Kingdom
- ^{ak}Fermilab, USA
- ^{al}Institut Pluridisciplinaire Hubert Curien/Département Recherches Subatomiques, Université de Strasbourg/CNRS-IN2P3, France
- ^{am}LPSC, Université Grenoble-Alpes, CNRS/IN2P3, France
- ^{an}Institut für Hochenergiephysik, Österreichische Akademie der Wissenschaften, Austria
- ^{ao}University of California, Davis, USA
- ^{ap}Northwestern University, USA
- ^{aq}Theoretical Physics Department, Fermilab, USA
- ^{ar}RWTH Aachen University, III. Physikalisches Institut A, Germany
- ^{as}National Centre for Particle Physics, Universiti Malaya, Malaysia
- ^{at}Physics, University of Washington, Seattle, USA
- ^{au}Laboratory for Particle Physics and Cosmology, Harvard University, USA
- ^{av}Department of Theoretical Physics, University of Geneva, Switzerland
- ^{aw}DESY, Germany
- ^{ax}Deutsches Elektronen-Synchrotron (DESY), Germany
- ^{ay}Institute of Particle Physics Phenomenology, Durham University, United Kingdom
- ^{az}National Central University, Taiwan
- ^{ba}The Ohio State University, USA
- ^{bb}Institut für Hochenergiephysik, Österreichische Akademie der Wissenschaften, Austria

^{bc}Texas Tech University, USA
^{bd}RWTH Aachen University, III. Physikalisches Institut A, Germany
^{be}Department of Physics, Chung-Ang University, Korea
^{bf}Texas Tech University, USA
^{bg}Department of Physics, University of Tokyo, Japan
^{bh}Kavli Institute for Cosmological Physics, University of Chicago, USA
^{bi}Vrije Universiteit Brussel - IIHE, Belgium
^{bj}Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France
^{bk}Imperial College London, United Kingdom
^{bl}Centre for Cosmology, Particle Physics and Phenomenology (CP3), Université catholique de Louvain, Belgium
^{bm}IFAE Barcelona, Spain
^{bn}IPPP Durham, United Kingdom
^{bo}Theoretische Natuurkunde and IIHE/ELEM, Vrije Universiteit Brussel, and International Solvay Institutes, Belgium
^{bp}GRAPPA, University of Amsterdam, Netherlands
^{bq}Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France
^{br}Department of Theoretical Physics, University of Geneva, Switzerland
^{bs}FNAL, USA
^{bt}Universidade Estadual Paulista, Brazil
^{bu}University of California, Irvine, USA
^{bv}RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany
^{bw}Stockholm University, Sweden
^{bx}Theoretical Physics Group, Lawrence Berkeley National Laboratory, and Berkeley Center for Theoretical Physics, University of California, Berkeley, USA
^{by}Carnegie Mellon University, USA
^{bz}Università di Padova, Italy
^{ca}Imperial College London, United Kingdom
^{cb}SLAC, Stanford University, USA
^{cc}Universität Wien, Austria
^{cd}National Centre for Physics, Quaid-i-Azam University, Pakistan
^{ce}Department of Theoretical Physics, University of Geneva, Switzerland
^{cf}Department of Theoretical Physics, University of Geneva, Switzerland
^{cg}SLAC, USA
^{ch}Max-Planck-Institut fuer Physik, Germany
^{ci}Nikhef and GRAPPA, Netherlands
^{cj}INFN Sezione di Napoli, and Dipartimento di Fisica, Università di Napoli, Italy
^{ck}Rudolf Peierls Centre for Theoretical Physics, University of Oxford, United Kingdom, and Niels Bohr Institute, Copenhagen, Denmark
^{cl}University of Hamburg, Germany
^{cm}University of California Santa Cruz Department of Physics and Santa Cruz Institute for Particle Physics, USA, and Niels Bohr International Academy, University of Copenhagen, Denmark
^{cn}Chulalongkorn University, Thailand
^{co}Cornell University, USA
^{cp}Chulalongkorn University, Faculty of Science, Department of Physics, Thailand
^{cq}Northwestern University, USA
^{cr}Department of Physics and Astronomy, University of California, Irvine, USA
^{cs}Laboratoire de Physique Corpusculaire, Clermont Université and Université Blaise Pascal and CNRS/IN2P3, Clermont-Ferrand, France
^{ct}Southampton University, United Kingdom
^{cu}University of Padova and INFN, Italy
^{cv}Northeastern University, Boston, USA
^{cw}Texas Tech University, USA
^{cx}Theory division, CERN, Switzerland
^{cy}Enrico Fermi Institute and Department of Physics and Kavli Institute for Cosmological Physics, University of Chicago, USA
^{cz}Department of Physics, Northeastern University, USA

^{da}Physics, University of Washington, Seattle, USA
^{db}Particle Physics Department, Rutherford Appleton Laboratory, United Kingdom
^{dc}Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS/IN2P3, France
^{dd}National Central University, Taiwan
^{de}Université Libre de Bruxelles, Belgium
^{df}Laboratoire de l'Accélérateur Linéaire, Univ. Paris-Sud 11 et IN2P3/CNRS, France

Abstract

This document is the final report of the ATLAS-CMS Dark Matter Forum, a forum organized by the ATLAS and CMS collaborations with the participation of experts on theories of Dark Matter, to select a minimal basis set of dark matter simplified models that should support the design of the early LHC Run-2 searches. A prioritized, compact set of benchmark models is proposed, accompanied by studies of the parameter space of these models and a repository of generator implementations. This report also addresses how to apply the Effective Field Theory formalism for collider searches and present the results of such interpretations.

Keywords: Dark Matter, Simplified Models, EFT, LHC

1. Introduction

Dark matter (DM) ¹ has not yet been observed in particle physics experiments, and there is not yet any evidence for non-gravitational interactions between DM and the Standard Model (SM) particles. If such interactions exist, particles of DM
5 could be produced at the LHC. Since DM particles themselves do not produce signals in the LHC detectors, one way to observe them is when they are produced in association with a visible SM particle $X(=g, q, \gamma, Z, W, \text{ or } h)$. Such reactions, which are observed at colliders as particles or jets recoiling against an invisible state, are

¹Many theories of physics beyond the Standard Model predict the existence of stable, neutral, weakly-interacting and massive particles that are putative DM candidates. In the following, we refer to such matter as DM, even though the observation of such matter at a collider could only establish that it is neutral, weakly-interactive, massive and stable on the distance-scales of tens of meters.

called “mono-X” or $\cancel{E}_T + X$ reactions (see e.g Refs. [1, 2, 3, 4, 5]), where \cancel{E}_T is the
10 missing transverse momentum observable in the detector.

Early Tevatron and LHC Run-1 searches for $\cancel{E}_T + X$ signatures at CDF [6], ATLAS [7, 8, 9, 10, 11, 12, 13, 14] and CMS [15, 16, 17, 18, 19, 20, 21], employed a basis of contact interaction operators in effective field theories (EFTs) [22, 23] to calculate the possible signals. These EFTs assume that production of DM takes place
15 through a contact interaction involving a quark-antiquark pair, or two gluons, and two DM particles. In this case, the missing energy distribution of the signal is determined by the nature and the mass of the DM particles and the Lorentz structure of the interaction. Only the overall production rate is a free parameter to be constrained or measured. Provided that the contact interaction approximation holds,
20 these EFTs provide a straightforward way to compare the results from different collider searches with non-collider searches for DM.

The EFT describes the case when the mediator of the interaction between SM and DM particles are very heavy; if this is not the case, models that explicitly include these mediators are needed [22, 24, 5, 25, 26, 27, 24, 28]. Some “simplified
25 models” [29, 30, 31] of DM production were constructed, including particles and interactions beyond the SM. These models can be used consistently at LHC energies, and provide an extension to the EFT approach. Many proposals for such models have emerged (see, for example Refs. [32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48]). At the LHC, the kinematics of mono-X reactions occurring
30 via a TeV-scale mediator can differ substantially from the prediction of the contact interaction. The mediator may also produce qualitatively different signals, such as decays back into the SM particles. Thus, appropriate simplified models are an important component of the design, optimization, and interpretation of DM searches at ATLAS and CMS. This has already been recognized in the CDF, ATLAS and CMS
35 searches quoted above, where both EFT and selected simplified model results are presented.

1.1. The ATLAS/CMS Dark Matter Forum

To understand what signal models should be considered for the upcoming LHC Run-2, groups of experimenters from both ATLAS and CMS collaborations have
40 held separate meetings with small groups of theorists, and discussed further at the DM@LHC workshop [40, 39, 49]. These discussions identified overlapping sets of simplified models as possible benchmarks for early LHC Run-2 searches. Following the DM@LHC workshop, ATLAS and CMS organized a forum, called the *ATLAS-CMS Dark Matter Forum*, to form a consensus on the use of these simplified models
45 and EFTs for early Run-2 searches with the participation of experts on theories of DM. This is the final report of the ATLAS-CMS Dark Matter Forum.

One of the guiding principles of this report is to channel the efforts of the ATLAS and CMS collaborations towards a minimal basis of dark matter models that should influence the design of the early Run-2 searches. At the same time, a thorough
50 survey of realistic collider signals of DM is a crucial input to the overall design of the search program.

The goal of this report is such a survey, though confined within some broad assumptions and focused on benchmarks for kinematically-distinct signals which are most urgently needed. As far as time and resources have allowed, the assumptions
55 have been carefully motivated by theoretical consensus and comparisons of simulations. But, to achieve such a consensus in only a few months before the start of Run-2, it was important to restrict the scope and timescale to the following:

1. The forum should propose a prioritized, compact set of benchmark simplified models that should be agreed upon by both collaborations for Run-2 searches.
60 The values for the scan on the parameters of the models for which experimental results are provided should be specified, to facilitate theory reinterpretation beyond the necessary model-independent limits that should be provided by all LHC DM searches.
2. The forum should recommend the use of the state of the art calculations for these benchmark models. Such a recommendation will aid the standardiza-
65 tion the event generator implementation of the simplified models and the

harmonization of other common technical details as far as practical for early Run-2 LHC analyses. It would be desirable to have a common choice of leading order (LO) and next-to-leading order (NLO) matrix elements corresponding to the state of the art calculations, parton shower (PS) matching and merging, factorization and renormalization scales for each of the simplified models. This will also lead to a common set of theory uncertainties, which will facilitate the comparison of results between the two collaborations.

3. The forum should discuss how to apply the EFT formalism and present the results of EFT interpretations.
4. The forum should prepare a report summarizing these items, suitable both as a reference for the internal ATLAS and CMS audiences and as an explanation of early Run-2 LHC benchmark models for theory and non-collider readers. This report represents the views of its endorsers, as participants of the forum.

1.2. Grounding Assumptions

We assume that interactions exist between the SM hadrons and the particles that constitute cosmological DM. If this is not the case, then proton collisions will not directly produce DM particles, and DM will not scatter off nuclei in direct detection experiments.

The DM itself is assumed to be a single particle, a Dirac fermion WIMP, stable on collider timescales and non-interacting with the detector. The former assumption is reductionistic. The rich particle content of the SM is circumstantial evidence that the DM sector, which constitutes five times as much of the mass of the universe, may be more complex than a single particle or a single interaction. But, as was often the case in the discoveries of the SM, here only one mediator and one search channel might play a dominant role in the opening stages of an LHC discovery. The latter assumption focuses our work on early LHC searches, where small kinematic differences between models will not matter in a discovery scenario, and with the imminent re-start of the LHC our report relies heavily on a large body of existing theoretical work which assumed Dirac fermionic DM.

Different spins of DM particles will typically give similar results. Exceptions exist: For example, the choice of Majorana fermions forbids some processes that are allowed for Dirac fermions [22]. Aside from these, adjusting the choice of Dirac or Majorana fermions or scalars will produce only minor changes in the kinematic distributions of the visible particle and is expected to have little effect on cut-and-count² analysis. Thus the choice of Dirac fermion DM should be sufficient as benchmarks for the upcoming Run-2 searches.

One advantage of collider experiments lies in their ability to study and possibly characterize the mediator. A discovery of an anomalous \cancel{E}_T signature at the LHC would not uniquely imply discovery of dark matter, while at the same time e.g. discovery of an anomalous and annually-modulated signal in a direct-detection experiment would leave unanswered many questions about the nature of the interaction that could be resolved by the simultaneous discovery of a new mediator particle. Collider, direct, and indirect detection searches provide complementary ways to approach this problem [50], and it is in this spirit that much of our focus is on the mediator.

We systematically explore the basic possibilities for mediators of various possible spins and couplings. All models considered are assumed to produce a signature with pairs of DM particles. Though more varied and interesting possibilities are added to the literature almost daily, these basic building blocks account for much of the physics studied at hadron colliders in the past three decades.

We also assume that Minimal Flavor Violation (MFV) [51, 52, 53, 54] applies to the models included in this report. This means that the flavor structure of the couplings between DM and ordinary particles follows the same structure as the SM. This choice is simple, since no additional theory of flavor is required, beyond what is already present in the SM, and it provides a mechanism to ensure that the models do not violate flavor constraints. As a consequence, spin-0 resonances must

²Cut-and-count refers to an analysis that applies a certain event selection and checks the inclusive number of events which pass. This is to be contrasted with a shape analysis, which compares the distribution of events.

have couplings to fermions proportional to the SM Higgs couplings. Flavor-safe models can still be constructed beyond the MFV assumption, for example [55],
125 and deserve further study. For a discussion of MFV in the context of the simplified models included in this report, see Ref. [49].

In the parameter scan for the models considered in this report, we make the assumption of a minimal decay width for the particles mediating the interaction between SM and DM. This means that only decays strictly necessary for the self-
130 consistency of the model (e.g. to DM and to quarks) are accounted for in the definition of the mediator width. We forbid any further decays to other invisible particles of the Dark Sector that may increase the width or produce striking, visible signatures. Studies within this report show that, for cut-and-count analyses, the kinematic distributions of many models, and therefore the sensitivity of these
135 searches, do not depend significantly on the mediator width, as long as the width remains smaller than the mass of the particle and that narrow mediators are sufficiently light.

The particle content of the models chosen as benchmarks is limited to one single kind of DM whose self-interactions are not relevant for LHC phenomenology, and
140 to one type of SM/DM interaction at a time. These assumptions only add a limited number of new particles and new interactions to the SM. These simplified models, independently explored by different experimental analyses, can be used as starting points to build more complete theories. Even though this factorized picture does not always lead to full theories and leaves out details that are necessary for the
145 self-consistency of single models (e.g. the mass generation for mediator particles), it is a starting point to prepare a set of distinct but complementary collider searches for DM, as it leads to benchmarks that are easily comparable across channels.

1.3. Choices of benchmarks considered in this report and parameter scans

Contact interaction operators have been outlined as basis set of theoretical build-
150 ing blocks representing possible types of interactions between SM and DM particles in [23]. The approach followed by LHC searches (see e.g. Refs. [15, 7] for recent jet+ \cancel{E}_T Run-1 searches with the 8 TeV dataset) so far has been to simulate only a pri-

oritized set of the possible operators with distinct kinematics for the interpretation
 of the constraints obtained, and provide results that may be reinterpreted in terms
 155 of the other operators. This report intends to follow this strategy, firstly focusing
 on simplified models that allow the exploration of scenarios where the mediating
 scale is not as large. In the limit of large mediator mass, the simplified models
 map onto the EFT operators. Secondly, this report considers specific EFT bench-
 marks whenever neither a simplified model completion nor other simplified models
 160 yielding similar kinematic distributions are available and implemented in one of the
 event generators used by both collaborations. This is the case for dimension-5 or
 dimension-7 operators with direct DM-electroweak boson couplings³. Considering
 these models as separate experimental benchmarks will allow to target new signal
 regions and help validate the contact interaction limit of new simplified models de-
 165 veloped to complete these specific operators. Results from these EFT benchmarks
 should include the condition that the momentum transfer does not probe the scale
 of the interaction; whenever there is no model that allows a direct mapping between
 these two quantities, various options should be tested to ensure a given fraction of
 events within the range of applicability of the EFT approach. Experimental searches
 170 should in any case deliver results that are independent from the specific benchmark
 tested, such as fiducial cross-sections that are excluded in a given signal region.

When choosing the points to be scanned in the parameter space of the mod-
 els, this report does not quantitatively consider constraints that are external to the
 MET+X analyses. This is the case also for results from LHC experiments searching
 175 for mediator decays. The main reason for not doing so in this report is the diffi-
 culty of incorporating these constraints in a rigorous quantitative way within the
 timescale of the Forum. However, even if the parameter scans and the searches
 are not optimized with those constraints in mind, we intend to make all informa-
 tion available to the community to exploit the unique sensitivity of colliders to all

³An example of a dimension-5 operator for scalar DM is described in Appendix A. Dimension-7 oper-
 ators of DM coupling to gauge bosons exist in the literature, but they require a larger particle spectrum
 with respect to the models studied in this report.

180 possible DM signatures.

1.4. Structure of this report and dissemination of results

The report provides a brief theoretical summary of the models considered, starting from the set of simplified models and contact interactions put forward in previous discussions and in the literature cited above. Its main body documents the studies done within this Forum to identify a kinematically distinct set of model parameters to be simulated and used as benchmarks for early Run-2 searches. The implementation of these studies according to the state of the art calculations is detailed, including instructions on how to estimate theoretical uncertainties in the generators used for these studies. The presentation of results for EFT benchmarks is also covered.

Chapter 2 of this report is dedicated to simplified models with radiation of a hard object either from the initial state or from the mediator. These models produce primarily monojet signatures, but should be considered for all $\cancel{E}_T + X$ searches. Chapter 3 contains studies on the benchmark models for final states specifically containing an electroweak boson ($W/Z/\gamma/H$). In this case, both simplified models leading to mono-boson signatures and contact interaction operators are considered. Details of the state of the art calculations and on the implementation of the simplified models in Monte Carlo generators are provided in Chapter 4. Chapter 5 is devoted to the treatment of the presentation of results for the benchmark models from contact interaction operators. Chapter 6 prescribes how to estimate theoretical uncertainties on the simulation of these models. Chapter ?? concludes the report.

Further models that could be studied beyond early searches and their implementation are described in Appendix A. For these models, either the implementation could not be fully developed by the time of this report, or some of the grounding assumptions were not fully met. Some of these models have been used in previous ATLAS and CMS analyses and discussed thoroughly within the Forum. They are therefore worth considering for further studies and for Run-2 searches, since they lead to unique $\cancel{E}_T + X$ signatures that are not shared by any other of the models included in this report. Appendix B contains the necessary elements that should be

210 included in the results of experimental searches to allow for further reinterpretation.

It is crucial for the success of the work of this Forum that these studies can be employed as cross-check and reference to the theoretical and experimental community interested in early Run-2 searches. For this reason, model files, parameter cards,
215 and cross-sections for the models considered in these studies are publicly available. The SVN repository of the Forum [56] contains the models and parameter files necessary to reproduce the studies within this report. Details and cross-sections for these models, as a function of their parameters, will be published on HEPData [57].

2. Simplified models for all $\cancel{E}_T + X$ analyses

220 In this Chapter we review models that yield $X + \cancel{E}_T$ signatures, where X is a QCD parton or γ, W, Z or h .

The primary simplified models for Dirac fermion DM studied and recommended by this Forum for early LHC Run-2 searches are detailed in this Chapter, comprising spin-0 and spin-1 mediators. Section 2.1 covers the s -channel exchange of a vector
225 mediator⁴, while we consider both s -channel and t -channel exchange for scalar mediators in Section 2.2 and 2.3 respectively. Spin-2 mediators are briefly mentioned in Section 2.4. While these models are general and cover a broad set of signatures, the discussion and studies are focused on the monojet final state. Details on final states with electroweak (EW) boson radiation and with heavy flavor quarks from
230 diagrams arising within these models are also discussed in this Chapter.

A summary of the state of the art calculations and implementations for these models is provided in Table 6.1. Section 4 details the implementation of these models that have been used for the studies in this Chapter and that will be employed for the simulation of early Run-2 benchmark models for LHC DM searches.

⁴Colored vector mediators can be exchanged in the t -channel, but there are no examples in literature so far.

235 *2.1. Vector and axial vector mediator, s-channel exchange*

A simple extension of the SM is an additional $U(1)$ gauge symmetry, where a DM candidate particle has charges only under this new group. Assuming that some SM particles are also charged under this group, a new gauge boson can mediate interactions between the SM and DM.

240 We consider the case of a DM particle χ of mass m_χ that is a Dirac fermion and where the production proceeds via the exchange of a spin-1 mediator of mass M_{med} in the s-channel, illustrated in Fig. 1.




Figure 1: Representative Feynman diagram showing the pair production of DM particles in association with a parton from the initial state via a vector or axial-vector mediator. The cross section and kinematics depend upon the mediator and DM masses, and the mediator couplings to DM and quarks respectively: $(M_{\text{med}}, m_\chi, g_\chi, g_q)$.

We consider two models with vector and axial-vector couplings between the spin-1 mediator Z' and SM and DM fields, with the corresponding interaction La-

245 grangians:

$$\mathcal{L}_{\text{vector}} = g_q \sum_{q=u,d,s,c,b,t} Z'_\mu \bar{q} \gamma^\mu q + g_\chi Z'_\mu \bar{\chi} \gamma^\mu \chi \quad (1)$$

$$\mathcal{L}_{\text{axial-vector}} = g_q \sum_{q=u,d,s,c,b,t} Z'_\mu \bar{q} \gamma^\mu \gamma^5 q + g_\chi Z'_\mu \bar{\chi} \gamma^\mu \gamma^5 \chi. \quad (2)$$

The coupling g_q is assumed to be universal to all quarks. It is also possible to consider other models in which mixed vector and axial-vector couplings are considered,

for instance the couplings to the quarks are axial-vector whereas those to DM are vector.

250 As mentioned in the Introduction, when no additional visible or invisible decays contribute to the width of the mediator, the minimal width is fixed by the choices of couplings g_q and g_χ . The effect of larger widths is discussed in Section 2.5.2. For the vector and axial-vector models, the minimal width is:

$$\Gamma_{\min}^V = \frac{g_\chi^2 M_{\text{med}}}{12\pi} \left(1 + \frac{2m_\chi^2}{M_{\text{med}}^2} \right) \beta_{DM} \theta(M_{\text{med}} - 2m_\chi) \quad (3)$$

$$+ \sum_q \frac{3g_q^2 M_{\text{med}}}{12\pi} \left(1 + \frac{2m_q^2}{M_{\text{med}}^2} \right) \beta_q \theta(M_{\text{med}} - 2m_q),$$

$$\Gamma_{\min}^A = \frac{g_\chi^2 M_{\text{med}}}{12\pi} \beta_{DM}^3 \theta(M_{\text{med}} - 2m_\chi) \quad (4)$$

$$+ \sum_q \frac{3g_q^2 M_{\text{med}}}{12\pi} \beta_q^3 \theta(M_{\text{med}} - 2m_q).$$

$\theta(x)$ denotes the Heaviside step function, and $\beta_f = \sqrt{1 - \frac{4m_f^2}{M_{\text{med}}^2}}$ is the velocity of
 255 the fermion f with mass m_f in the mediator rest frame. Note the color factor 3 in the quark terms. Figure 2 shows the minimal width as a function of mediator mass for both vector and axial-vector mediators assuming the coupling choice $g_q = g_\chi = 1$. With this choice of the couplings, the dominant contribution to the minimal width comes from the quarks, due to the combined quark number and color factor
 260 enhancement. We specifically assume that the vector mediator does not couple to leptons. If such a coupling were present, it would have a minor effect in increasing the mediator width, but it would also bring in constraints from measurements of the Drell-Yan process that would unnecessarily restrict the model space.

Therefore, the minimal set of parameters under consideration for these two
 265 models is

$$\{ g_q, g_\chi, m_\chi, M_{\text{med}} \} \quad (5)$$

together with the spin structure of their couplings.




Figure 2: Minimal width as a function of mediator mass for vector and axial-vector mediator assuming couplings of 1. The total width is shown as solid lines for DM masses of 10 GeV, 30 GeV, 100 GeV and 300 GeV in black, red, brown and green, respectively. The individual contributions from DM are indicated by dotted lines with the same colors. The contribution from all quarks but top is shown as magenta dotted line and the contribution from top quarks only is illustrated by the dotted blue line. The dotted black line shows the extreme case $\Gamma_{\text{min}} = M_{\text{med}}$.

A thorough discussion of these models and their parameters can also be found in [58].

These simplified models are known and available in event generators at NLO +
 270 PS accuracy, as detailed in Section 4.1.1. Results in this Section have been obtained using the model implementation within the POWHEG generator (v3359) [59], interfaced to PYTHIA 8 [60] for the parton shower.

In addition, for the vector models considered, initial and final state radiation of a Z' can occur which can appear as a narrow jet if it decays hadronically and
 275 may not be distinguishable from a QCD jet, thus accounting for some fraction of the monojet signal. The ISR and FSR of Z' becomes more important at large values of the couplings [61].

2.1.1. Parameter scan

In order to determine an optimal choice of the parameter grid for the simulation of early Run-2 benchmark models, dependencies of the kinematic quantities
 280 and cross sections on the model parameters have been studied. Only points that are kinematically distinct will be fully simulated, while instructions on how to rescale the results according to models with different cross sections are presented in Section 2.5. The following paragraphs list the main observations from the scans over
 285 the parameters that support the final proposal for the benchmark signal grid.

Scan over the couplings. To study the dependence of kinematic distributions on the coupling strength, samples were generated where a pair of $m_\chi = 10$ GeV DM particles is produced on-shell from the mediator of $M_{\text{med}} = 1$ TeV. Figure 3 compares the shapes of the \cancel{E}_T distribution for the different choices of the coupling strength.
 290 This is a generator-level prediction with no kinematic selections or detector simulation. Coupling values in the scan range 0.1–1.45, fixing $g_q = g_\chi$, correspond to a rough estimate of the lower sensitivity of mono-jet analyses and a maximum coupling value such that $\Gamma_{\text{min}} < M_{\text{med}}$. We observe that the shapes of the \cancel{E}_T or jet p_T distributions do not depend on the couplings (and consequently the width) in
 295 the ranges considered. A large width of the mediator implies a broad integral over




Figure 3: Scan over couplings. The \cancel{E}_T distribution is compared for the vector mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively. All figures in this Section have been obtained using the model implementation within the POWHEG generator (v3359) [59], interfaced to PYTHIA 8 [60] for the parton shower.

the contributing parton distributions, which might not be well approximated by the midpoint of this integral. This study shows that the effect, in the p_T distribution of the observed gluon, is not important.

Based on similar findings for different choices of M_{med} and m_χ , we conclude that the shapes of kinematic distributions are not altered by coupling variations, neither for the on-shell mediator case where $M_{\text{med}} > 2m_\chi$, nor for the off-shell case where $M_{\text{med}} < 2m_\chi$. Only the production cross sections change. Differences in kinematic distributions are expected only close to the transition region between on-shell and off-shell mediators.

Special care needs to be taken when coupling strengths are combined with extremely heavy mediators. Figure 4 suggests a change in the shape of the \cancel{E}_T distribution for a $M_{\text{med}} = 5$ TeV mediator once $\Gamma_{\text{min}}/M_{\text{med}}$ is of the order of a percent or lower.

Such heavy mediators, although inaccessible with early LHC data, are interesting since they provide a good approximation for benchmark EFT models. The observed difference among the simplified models in the plot arises from the fact






Figure 4: Comparison of the E_T^{miss} distributions from the D5 EFT sample and the vector models with 5 TeV heavy mediator of various widths. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $E_T > 300$ GeV and $E_T > 500$ GeV cut, respectively.

that the region of low invariant masses of the DM pair, $m_{\tilde{\chi}\chi}$, is suppressed due to narrow Breit-Wigner peak that only probes a narrow window of parton distribution functions. For wider mediators, the low mass region is significantly enhanced by
315 parton distribution functions at low Bjorken x , as illustrated in Fig. 5(a). This explains why the sample with the narrowest mediator in Fig. 4 is heavily suppressed in terms of production cross section and also gives different E_T shape. Furthermore, Fig. 4 compares the vector model with 5 TeV mediator to the D5 EFT sample and reveals that the simplified models with larger mediator widths (e.g. for couplings
320 of 1 where $\Gamma_{\text{min}}/M_{\text{med}} \sim 0.5$) are the ones resembling the kinematics of contact interactions. This reflects the fact that in an EFT there is no enhancement due to on-shell mediators, leading to a closer resemblance to an off-shell regime where no peak in the $m_{\tilde{\chi}\chi}$ distribution is present. In case of narrow width mediators, e.g. $\Gamma_{\text{min}}/M_{\text{med}} \sim 0.05$, even larger mediator masses need to be chosen in order to sig-
325 nificantly suppress the peak in the $m_{\tilde{\chi}\chi}$ distribution and reproduce the kinematic shapes of an EFT model. Figure 5(b) verifies that the choice of 10 TeV mediator mass is sufficient to achieve that.

Since kinematic distributions are robust to changes in the specific values of cou-



(a)



(b)

Figure 5: Invariant mass of the DM pair in the vector mediator samples with $m_{\chi} = 10$ GeV, $M_{\text{med}} = 5$ TeV and different coupling strengths (a). A similar comparison is shown for the samples with different mediator masses considering $\Gamma_{\text{min}}/M_{\text{med}} = 0.05$ and 0.1 (b). An EFT sample is also displayed in the latter case. The distributions are normalised to unit area.

pling ⁵, the choice of $g_q = 0.25$ and $g_\chi = 1$ is reasonable to reduce the parameter
 330 space to be scanned. There are no complications associated with small couplings,
 but, also, the early part of Run 2 will not be sensitive to them. The range of cou-
 plings we recommend to generate limit the calculated width of the mediator to be
 near or below M_{med} .

For direct mediator searches, such as $q\bar{q} \rightarrow Z' \rightarrow q\bar{q}$, different couplings ($g_q \neq$
 335 g_χ) might also be considered. A scan in g_χ vs g_q can then be performed for a fixed
 mediator mass. Such searches may restrict g_q to a greater degree than g_χ .

Scan over m_χ . For a fixed mediator mass M_{med} and couplings, the DM mass falls
 into three regimes:

On-shell: When $M_{\text{med}} \gg 2m_\chi$, most mediators are on-shell. The hardness of the ISR is
 340 set by M_{med} , and the kinematic distributions do not strongly depend on m_χ .
 This is illustrated in Fig. 6 for an example of $M_{\text{med}} = 1$ TeV, $10 \text{ GeV} < m_\chi <$
 300 GeV . The cross section decreases as the m_χ approaches $M_{\text{med}}/2$. A coarse
 binning along m_χ is sufficient.

Threshold: When $M_{\text{med}} \approx 2m_\chi$, the production is resonantly enhanced, and both the cross
 345 section and kinematic distributions change more rapidly as a function of the
 two masses, and finer binning is needed in order to capture the changes.

Off-shell: When $M_{\text{med}} \ll 2m_\chi$, the DM pair is produced by an off-shell mediator. The
 mediator propagator gives an explicit suppression of $(M_{\text{med}}/Q)^2$ that sup-
 presses hard ISR. The $m_\chi = 1$ TeV case, shown in Fig. 6, and Figure 7 demon-
 350 strates that the \cancel{E}_T spectrum hardens with increasing m_χ , accompanied by the
 gradual decrease of the cross section. Due to the significant cross section sup-
 pression, it is not necessary to fully populate the parameter space. Imminent
 LHC searches are not expected to be sensitive to these signals.

⁵This applies as long as heavy narrow mediators are generated without any truncation of low-mass
 tails at the generator-level.




Figure 6: Scan over DM mass. The \cancel{E}_T distribution is compared for the vector mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.




Figure 7: Scan over DM mass. The \cancel{E}_T distribution is compared for the vector mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.




Figure 8: Scan over mediator mass. The \cancel{E}_T distribution is compared for the vector mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.

Scan over the mediator mass. Changing the mediator mass for fixed DM mass and couplings leads to significant differences in cross section and shapes of the kinematic variables for the on-shell regime, as shown in Fig. 8. As expected, higher mediator masses lead to harder \cancel{E}_T spectra. On the other hand, the \cancel{E}_T shapes are similar for off-shell mediators. This is illustrated in Fig. 9. Therefore, a coarse binning in M_{med} is sufficient in the off-shell regime.

Spin structure of the couplings. This section compares the kinematic properties of vector, axial-vector and mixed vector/axial-vector models. The samples with pure vector and pure axial-vector couplings are compared for $M_{\text{med}} = 100$ GeV and different DM masses in Fig. 10. No differences in the shape of the \cancel{E}_T distributions are observed between the samples with coincident masses. In the case of the on-shell mediators, where $2m_\chi \ll M_{\text{med}}$, the cross sections of the pure vector and pure axial-vector models are similar. With increasing DM mass towards the $2m_\chi = M_{\text{med}}$ transition and further into the off-shell regime, the relative difference between the cross sections of the two samples is increasing, with the vector ones having larger cross sections.

Figure 11 shows the samples generated with pure and mixed couplings for $m_\chi =$




Figure 9: Scan over mediator mass. The \cancel{E}_T distribution is compared for the vector mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.




Figure 10: Comparison of the pure vector and pure axial-vector couplings. The \cancel{E}_T distribution is shown for the samples generated with $M_{\text{med}} = 100$ GeV and different DM masses. Ratios of the normalized distributions are shown for between the samples with coincident masses. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.

100 GeV and $M_{\text{med}} = 1$ TeV, i.e. where the mediator is on-shell. The mediator width between the pure vector and pure axial-vector couplings differ only by 2% in this case, and $< 10\%$ agreement between the cross sections is found. The mediator widths for the samples with the same type coupling to quarks agree at better than
 375 1% since the width is dominated by the quark contribution, as expected from Eq. 3. No significant differences between the samples with same type DM coupling are seen, given the statistical precision of the generated samples. This is expected since the mediator is on-shell, and the details of the invisible decay are unimportant in cut-and-count searches.

380 For the off-shell case, shown in Fig. 12 for $m_\chi = 100$ GeV and $M_{\text{med}} = 100$ GeV, there is approximately a factor 2 difference between the cross-sections of the samples with pure couplings is observed. As in the previous case, the samples with the same type coupling to DM are similar both in terms of cross sections and \cancel{E}_T shape. Since the contribution to the mediator width from DM is closed in this case, only
 385 the quark couplings define the width. Only couplings to light quarks are opened in the case of $M_{\text{med}} = 100$ GeV for which the differences between the partial widths of vector and axial-vector couplings are marginal. This explains the similar minimal widths for all four samples stated in Fig. 12.

In general, the coupling to quarks is not expected to play an important role in
 390 the kinematics as it is only needed to produce the mediator which is confirmed by the observations above. Based on this argument and on the observations above, we recommend to consider only the models with pure vector couplings or pure axial-vector couplings for simulation.

Proposed parameter grid. The final step in proposing a parameter grid is to evalu-
 395 ate the sensitivity of Run-2 LHC data with respect to rate and/or kinematics. The parameter scan focuses on two important regions, the light mediator region and the heavy mediator limit to reproduce the EFT limit, and takes into account the projected sensitivities for the mono-jet analysis.

Considering simplified models also allows to discuss constraints from different
 400 search channels. In the case of the s -channel exchange, the results from the mono-




Figure 11: Comparison of the pure vector, V-V, and pure axial-vector, A-A, couplings with mixed couplings, A-V and V-A where the first (second) letter indicates the SM (dark sector) vertex. The \cancel{E}_T distribution is shown for the samples generated with $m_\chi = 100$ GeV and $M_{\text{med}} = 1$ TeV. Ratios of the normalized distributions are shown for A-V over V-V and for V-A over A-A. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.




Figure 12: Comparison of the pure vector, V-V, and pure axial-vector, A-A, couplings with mixed couplings, A-V and V-A where the first (second) letter indicates the SM (dark sector) vertex. The \cancel{E}_T distribution is shown for the samples generated with $m_\chi = 100$ GeV and $M_{\text{med}} = 100$ GeV. Ratios of the normalized distributions are shown for A-V over V-V and for V-A over A-A. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively. The suppression by β^3 for $m_\chi \sim M_{\text{med}}$ can be seen for the curves representing axial DM coupling.

jet final states, where the mediator decays to a DM pair, one can also take into account dijet constraints on the processes where the mediator decays back to the SM particles. The importance of the dijet results depend on the magnitude of the coupling g_q . We recommend to keep the two channels rather independent by choosing $g_q = 0.25$ and $g_\chi = 1$, based on the findings given in Ref. [62]. Furthermore, it is also important to mention this choice leads to $\Gamma_{\min}/M_{\text{med}} \lesssim 0.06$. Note that the usual choice of $g_q = g_\chi = 1$ used in literature leads to $\Gamma_{\min}/M_{\text{med}} \sim 0.5$, questioning the applicability of the narrow width approximation.

The expected upper limit at 95% confidence level on the product of cross section, acceptance and efficiency, $\sigma \times A \times \epsilon$, in the final Run-1 ATLAS mono-jet analysis [7] is 51 fb and 7.2 fb for $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV, respectively. Projected sensitivities for a 14 TeV mono-jet analysis are available from ATLAS [14]. These ATLAS studies estimate a factor of two increase in sensitivity with the 2015 data. The generator level cross section times efficiency times acceptance at $\cancel{E}_T > 500$ GeV for the model with couplings $g_q = 0.25$ and $g_\chi = 1$, a light DM particle of $m_\chi = 10$ GeV and a $M_{\text{med}} = 1$ TeV vector mediator is at the order of 100 fb, i.e. the early Run-2 mono-jet analysis is going to be sensitive to heavier mediators than this. The value of $\sigma \times \epsilon \times A$ at $\cancel{E}_T > 500$ GeV for a 5 TeV vector mediator is at the order of 0.1 fb, therefore this model lies beyond the reach of the LHC in the early Run-2. However, models with high enough mediators are still useful to reproduce the EFT result.

Following these arguments, M_{med} grid points are chosen, roughly equidistant in a logarithmic scale: 10 GeV, 20 GeV, 50 GeV, 100 GeV, 200 GeV, 300 GeV, 500 GeV, 1000 GeV and 2000 GeV. In the threshold regime $M_{\text{med}} = 2m_\chi$, the m_χ grid points are taken at approximately $M_{\text{med}}/2$, namely: 10 GeV, 50 GeV, 150 GeV, 500 GeV and 1000 GeV. Points on the on-shell diagonal are always chosen to be 5 GeV away from the threshold, to avoid numerical instabilities in the event generation. The detailed studies of the impact of the parameter changes on the cross section and kinematic distributions presented earlier in this section support removing some of the grid points and relying on interpolation. The optimized grids proposed for the vector and axial-vector mediators are given in Table. 1. One point at very

high mediator mass (10 TeV) is added for each of the DM masses scanned, to aid the reinterpretation of results in terms of contact interaction operators (EFTs), as discussed in Section 5.2.

m_χ/GeV	$M_{\text{med}}/\text{GeV}$									
1	10	20	50	100	200	300	500	1000	2000	10000
10	10	15	50	100						10000
50	10		50	95	200	300				10000
150	10				200	295	500	1000		10000
500	10						500	995	2000	10000
1000	10							1000	1995	10000

Table 1: Simplified model benchmarks for s -channel simplified models (spin-1 mediators decaying to Dirac DM fermions in the V and A case, taking the minimum width for $g_q = 0.25$ and $g_\chi = 1$)

435 Tables 2 and 3 give the $\Gamma_{\text{min}}/M_{\text{med}}$ ratio for the parameter grid proposed for vector and axial-vector s -channel models, respectively. The numbers range from ~ 0.02 in the off-shell regime at $2m_\chi > M_{\text{med}}$ to ~ 0.06 in the on-shell regime for heavy mediators where all coupling channels contribute.

2.1.2. Additional considerations for $V+\cancel{E}_T$ signatures

440 All models detailed in this Section are applicable to signatures where a photon, a W boson, a Z boson or a Higgs boson is radiated from the initial state partons instead of a gluon. The experimental signature is identified as $V+\cancel{E}_T$ and it has been sought by ATLAS and CMS in Refs. [16, 8, 17, 9, 11, 10]. This signature is also produced by the models described in Section 3.

445 Monojet searches are generally more sensitive with respect to final states including EW bosons, due to the much larger rates of signal events featuring quark or gluon radiation with respect to radiation of bosons [63], in combination with the low branching ratios if leptons from boson decays are required in the final state. The rates for the Higgs boson radiation is too low for these models to be considered
450 a viable benchmark [48]. However, the presence of photons, leptons from W and Z

m_χ / GeV	$M_{\text{med}} / \text{GeV}$									
	10	20	50	100	200	300	500	1000	2000	10000
1	0.049	0.051	0.051	0.051	0.051	0.051	0.056	0.056	0.056	0.056
10	0.022	0.024	0.054	0.052						0.056
50	0.022		0.025	0.025	0.055	0.053				0.056
150	0.022				0.025	0.025	0.061	0.058		0.056
500	0.022						0.029	0.030	0.060	0.057
1000	0.022							0.030	0.030	0.057

Table 2: Minimal width of the vector mediator exchanged in s -channel divided by its mass, assuming $g_q = 0.25$ and $g_\chi = 1$. The numbers tabulated under $2m_\chi = M_{\text{med}}$ correspond to the width calculated for $M_{\text{med}} - 5 \text{ GeV}$.

m_χ / GeV	$M_{\text{med}} / \text{GeV}$									
	10	20	50	100	200	300	500	1000	2000	10000
1	0.045	0.049	0.051	0.051	0.051	0.051	0.053	0.055	0.056	0.056
10	0.020	0.022	0.047	0.050						0.056
50	0.020		0.025	0.025	0.045	0.048				0.056
150	0.020				0.025	0.025	0.044	0.053		0.056
500	0.020						0.027	0.029	0.050	0.056
1000	0.020							0.029	0.030	0.055

Table 3: Minimal width of the axial-vector mediator exchanged in s -channel divided by its mass, assuming $g_q = 0.25$ and $g_\chi = 1$. The numbers tabulated under $2m_\chi = M_{\text{med}}$ correspond to the width calculated for $M_{\text{med}} - 5 \text{ GeV}$.

decays, and W or Z bosons decaying hadronically allow backgrounds to be rejected more effectively, making $Z/\gamma/W+\cancel{E}_T$ searches still worth comparing with searches in the jet+ \cancel{E}_T final state (see e.g. Ref. [64]).

In the case of a spin-1 mediator, an example Feynman diagram for these processes can be constructed by taking Fig. 1 and replacing the gluon with γ, W or Z .

When the initial state radiation is a W boson, Run-1 searches have considered three benchmark cases, varying the relative coupling of the W to u and d quarks. The simplified model with a vector mediator exchanged in the s-channel includes only the simplest of these cases, in which the W coupling to u and d quarks is identical, as required naively by SU(2) gauge invariance. With some more complex model building, other cases are possible. The case in which the u and d couplings have opposite sign is particularly interesting, since this enhances the $W + \cancel{E}_T$ signal over the jet+ \cancel{E}_T signal [65, 44, 66]. An example of a model of this type is discussed in Appendix A.4.

Simulations for the models in this Section have been done at the LO+PS level using MADGRAPH5_AMC@NLO 2.2.3 interfaced to PYTHIA 8, and therefore no special runtime configuration is needed for pythia 8. Even though merging samples with different parton multiplicities is possible, this has not been deemed necessary as the visible signal comes from the production of a heavy SM boson whose transverse momentum distribution is sufficiently well described at LO+PS level.

In these $V+\cancel{E}_T$ models, as in the case of the jet+ \cancel{E}_T models, p_T of the boson or the \cancel{E}_T does not depend strongly on the width of the mediator. An example of the particle-level analysis acceptance using the generator-level cuts from Ref. [8] for the photon+ \cancel{E}_T analysis, but raising the photon p_T cut to 150 GeV, is shown in Figure 4, comparing a width that is set to $\Gamma = M_{med}/3$ to the minimal width (the ratio between the two widths ranges from 1.05 to 1.5 with increasing mediator masses).

Examples of relevant kinematic distributions for selected benchmark points are shown in Fig. 13.

Acceptance ratio: $\Gamma = \Gamma_{\min}$ vs $\Gamma = M_{\text{med}}/3$				
	m_χ/GeV			
$M_{\text{med}}/\text{GeV}$	10	50	200	400
50	0.96	0.99		0.95
100	0.97			
300	1.00	1.02		
600			0.96	
1000	1.01	1.02	1.03	
3000	1.02	1.03		1.01


Table 4: Analysis acceptance ratios for the photon+ \cancel{E}_T analysis when varying the mediator width, in the case of a vector mediator exchanged in the s -channel. The figures shown in this Section have been obtained using a LO UFO model in MADGRAPH5_AMC@NLO 2.2.3 interfaced to PYTHIA 8 for the parton shower.

2.2. Scalar and pseudoscalar mediator, s -channel exchange


In this section, we consider a parallel situation to the vector and axial-vector mediators in the previous sections: a real scalar or a pseudoscalar where the associated scalar is decoupled at higher energies⁶. This section is largely based on
485 Refs. [42, 41, 43] which contain a thorough discussion of these models.

Assuming MFV, spin-0 resonances behave in a similar fashion as the SM Higgs boson. If the mediators are pure singlets of the SM, their interactions with quarks are not $SU(2)_L$ invariant. To restore this invariance, one could include the mixing of such mediators with the Higgs sector. This leads to extra interactions and a more
490 complex phenomenology with respect to what considered in this Section (for a more complete discussion, see Refs. [42, 43]). In the interest of simplicity, we do not study models including those interactions in this report as early Run-2 benchmark models, but we give an example of a model of this kind in Appendix A.6.

⁶This assumption does not hold in a UV-complete model where the two components of the complex scalar mediator would be approximately degenerate. The complex scalar case could be studied separately in the case of heavy flavor final states given the sufficiently different kinematics.



(a) Leading photon transverse momentum distribution for the photon+ E_T final state, for different mediator mass choices, for $m_\chi = 10$ GeV. (b) Leading photon transverse momentum distribution for the photon+ E_T final state, for different DM mass choices, with $M_{\text{med}} = 1$ TeV.



(c) Missing transverse momentum distribution for the leptonic $Z+E_T$ final state, for different mediator mass choices, for $m_\chi = 15$ GeV. (d) Missing transverse momentum distribution for the hadronic $W+E_T$ final state.

Figure 13: Kinematic distributions relevant for searches with W, Z and photons in the final state, for the simplified model with a vector mediator exchanged in the s -channel.

Relative to the vector and axial-vector models discussed above, the scalar models are distinguished by the special consequences of the MFV assumption: the very narrow width of the mediator and its extreme sensitivity to which decays are kinematically available, and the loop-induced coupling to gluons. The interaction Lagrangians are




Figure 14: One-loop diagrams of processes exchanging a scalar (S) or pseudoscalar (P) mediator, leading to a mono-jet signature.

$$\mathcal{L}_\phi = g_\chi \phi \bar{\chi} \chi + \frac{\phi}{\sqrt{2}} \sum_i (g_u y_i^u \bar{u}_i u_i + g_d y_i^d \bar{d}_i d_i + g_\ell y_i^\ell \bar{\ell}_i \ell_i), \quad (6)$$

$$\mathcal{L}_a = i g_\chi a \bar{\chi} \gamma_5 \chi + \frac{ia}{\sqrt{2}} \sum_i (g_u y_i^u \bar{u}_i \gamma_5 u_i + g_d y_i^d \bar{d}_i \gamma_5 d_i + g_\ell y_i^\ell \bar{\ell}_i \gamma_5 \ell_i) \quad (7)$$

where ϕ and a are respectively the scalar and pseudoscalar mediators, and the
500 Yukawa couplings y_i^f are normalized to the Higgs vev as $y_i^f = \sqrt{2} m_i^f / v$.

The couplings to fermions are proportional to the SM Higgs couplings, yet one is still allowed to adjust an overall strength of the coupling to charged leptons and the relative couplings of u - and d -type quarks. As in the preceding sections, for the sake of simplicity and straightforward comparison, we reduce the couplings to the
505 SM fermions to a single universal parameter $g_q \equiv g_u = g_d = g_\ell$. Unlike the vector and axial-vector models, the scalar mediators are allowed to couple to leptons.⁷

The relative discovery and exclusion power of each search can be compared in this framework. However, we again emphasize the importance of searching the full

⁷This contribution plays no role for most of the parameter space considered. The choice to allow lepton couplings follows Refs. [42, 41].

set of allowed channels in case violations of these simplifying assumptions lead to
 510 significant modifications of the decay rates that unexpectedly favor different chan-
 nels than the mix obtained under our assumptions. The coupling g_χ parametrizes
 the entire dependence on the structure between the mediator and the dark sector.

Given these simplifications, the minimal set of parameters under consideration
 is

$$\{m_\chi, m_{\phi/a} = M_{\text{med}}, g_\chi, g_q\}. \quad (8)$$

515 Fig. 14 shows the one-loop diagrams producing a jet+X signature. The full calcu-
 lation of the top loop is available at LO for DM pair production in association with
 one parton.

The minimal mediator width (neglecting the small contributions from quarks
 other than top in the loop) is given by

$$\begin{aligned} \Gamma_{\phi,a} = \sum_f N_c \frac{y_f^2 g_q^2 m_{\phi,a}}{16\pi} \left(1 - \frac{4m_f^2}{m_{\phi,a}^2}\right)^{x/2} &+ \frac{g_\chi^2 m_{\phi,a}}{8\pi} \left(1 - \frac{4m_\chi^2}{m_{\phi,a}^2}\right)^{x/2} \\ &+ \frac{\alpha_s^2 g_q^2 m_{\phi,a}^3}{32\pi^3 v^2} \left|f_{\phi,a}\left(\frac{4m_t^2}{m_{\phi,a}^2}\right)\right|^2 \end{aligned} \quad (9)$$

where $x = 3$ for scalars and $x = 1$ for pseudoscalars. The loop integrals, with f as
 complex functions, are

$$f_\phi(\tau) = \tau \left[1 + (1 - \tau) \arctan^2 \left(\frac{1}{\sqrt{\tau - 1}} \right) \right], \quad (10)$$

$$f_a(\tau) = \tau \arctan^2 \left(\frac{1}{\sqrt{\tau - 1}} \right) \quad (11)$$

520 where $\tau = 4m_t^2/m_{\phi,a}^2$.

The minimal widths for scalar and pseudo-scalar mediators with $g_q = g_\chi = 1$
 are shown in Fig. 20, illustrating the effect of choosing the SM Higgs-like Yukawa
 couplings for the SM fermions. For the mediator mass above twice the top quark
 mass m_t , the minimal width receives the dominant contribution from the top quark.
 525 For lighter mediator masses, DM dominates as the couplings to lighter quarks are
 Yukawa suppressed.

As shown in the diagram of Fig. 14, the lowest order process of these models al-
 ready involves a one-loop amplitude in QCD, and only LO predictions are currently

available. The generator used for the studies for the jet+ \cancel{E}_T signature is POWHEG
530 [59, 43, 67, 68, 69], with PYTHIA 8 [60] for the parton shower; within this imple-
mentation, the scalar and pseudoscalar mediator benchmark models are known at
LO+PS accuracy.

2.2.1. Parameter scan

Similarly as in the case of the vector and axial-vector couplings of spin-1 medi-
535 ators, scans in the parameter space are performed also for the scalar and pseudo-
scalar couplings of the spin-0 mediators in order to decide on the optimized param-
eter grid for the presentation of Run-2 results. Figures 15- 19 show the scans over
the couplings, DM mass and mediator mass and the same conclusions apply as in
Section 2.1.

A scan over the mediator mass is shown in Fig. 19 where $M_{\text{med}} = 300$ GeV and
540 500 GeV are chosen to be below and above $2m_t$. The off-shell case is assumed by
taking an extreme limit ($m_\chi = 1$ TeV) in order to study solely the effects of the
couplings to quarks. No differences in the kinematic distributions are observed and
also the cross sections remain similar in this case. No significant changes appear
545 for mediator masses around the $2m_t$ threshold.

It can be seen in Fig. 21 that the kinematics for the scalar and pseudoscalar
models coincides when considering the diagrams in Fig. 14. For this reason, we
recommend to fully simulate only one of the two models. No preference is given
between the two models as they have the same kinematics, although it is worth
550 noting that the pseudo-scalar model has been used for a DM interpretation of the
DAMA signal and of the galactic center excess [70]. Like in the case of the vector
and axial-vector models described in Section 2.1.1, the differences between the
cross sections for the scalar and pseudo-scalar samples with the same m_χ and M_{med}
are increasing with the DM mass for fixed mediator mass, with the pseudo-scalar
555 model yielding larger cross sections. There is an increasing difference between the
minimal widths close to the $2m_\chi = M_{\text{med}}$ threshold.

Proposed parameter grid. The optimized parameter grid in the $M_{\text{med}}-m_\chi$ plane for
scalar and pseudo-scalar mediators is motivated by similar arguments as in the pre-




Figure 15: Scan over couplings. The \cancel{E}_T distribution is compared for the scalar mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300 \text{ GeV}$ and $\cancel{E}_T > 500 \text{ GeV}$ cut, respectively. Studies in all figures for the jet+ \cancel{E}_T signature is POWHEG, with PYTHIA 8 for the parton shower;




Figure 16: Scan over DM mass. The \cancel{E}_T distribution is compared for the scalar mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300 \text{ GeV}$ and $\cancel{E}_T > 500 \text{ GeV}$ cut, respectively.




Figure 17: Scan over DM mass. The \cancel{E}_T distribution is compared for the scalar mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.




Figure 18: Scan over mediator mass. The \cancel{E}_T distribution is compared for the scalar mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.

vious section. Therefore, a similar pattern is followed here, with the exception of
 560 taking $g_q = g_\chi = 1$. The choice of $g_q = 0.25$ for the vector and axial-vector models
 is motivated by suppressing constraints from di-jets, which is not a concern in the
 scalar and pseudo-scalar mediator case. Here a di-jet signal emerges only at the 2-
 loop level through diagrams where the mediator is produced via gluon-gluon fusion
 and decays back into two gluons through a top loop. The strong loop suppression
 565 renders such signals unobservable at the LHC. Further constraints on the scalar and
 pseudo-scalar mediators may emerge from searches in $t\bar{t}$ final states. Studies of the
 electroweak effects to $t\bar{t}$ production suggest that one can only expect percent level
 contributions for $g_q \sim O(1)$ [71]. Therefore, keeping $g_q = g_\chi = 1$ is a reasonable
 choice in the case of the scalar and pseudo-scalar mediators. Contrary to the vector
 570 and axial-vector models, note that couplings of 1 lead to $\Gamma_{\min}/M_{\text{med}} \lesssim 0.1$, ensuring
 the narrow width approximation is applicable. Furthermore, the sensitivity to the
 highest mediator masses has to be re-evaluated. The generator level cross section
 times the acceptance at $\cancel{E}_T > 500$ GeV for the model with couplings $g_q = g_\chi = 1$,
 light DM of $m_\chi = 10$ GeV and a $M_{\text{med}} = 500$ GeV scalar mediator is at the order of
 575 10 fb, i.e. just at the edge of the early Run-2 sensitivity. Increasing the mediator
 mass to 1 TeV pushes the product $\sigma \times A$ down to approximately 0.1 fb, below the
 LHC sensitivity. Therefore, we choose to remove the 2 TeV mediator mass from
 the grid and present the final grid with 33 mass points only, as shown in Tab. 5.
 One point at very high mediator mass (10 TeV) is added for each of the DM masses
 580 scanned, to aid the reinterpretation of results in terms of contact interaction oper-
 ators (EFTs).

For the parameter grid for scalar and pseudo-scalar mediator s -channel exchange,
 the $\Gamma_{\min}/M_{\text{med}}$ ratio is given in Tables 6 and 7, respectively. In the on-shell regime,
 the ratio is between 0.04 and 0.1. Very narrow resonances with $\Gamma_{\min}/M_{\text{med}} < 0.001$
 585 correspond to the mass points where the mediator is off-shell. Note that the loop-
 induced contribution from gluons is ignored in the width calculation.

m_χ (GeV)	M_{med} (GeV)									
1	10	20	50	100	200	300	500	1000	10000	
10	10	15	50	100						10000
50	10		50	95	200	300				10000
150	10				200	295	500	1000		10000
500	10						500	995		10000
1000	10							1000		10000

Table 5: Simplified model benchmarks for s -channel simplified models (spin-0 mediators decaying to Dirac DM fermions in the scalar and pseudoscalar case, taking the minimum width for $g_q = 1$ and $g_\chi = 1$)

2.2.2. Additional considerations for $V + \cancel{E}_T$ signatures

The discussion of parameters for the model with a color-singlet, spin-0 mediator parallels that in Section 2.

590 Even though the sensitivity of mono-boson searches to this model is low and it may not be in reach of early LHC searches, this model can be generated for W, Z and photon searches in order to reproduce the kinematics of contact interaction operators that are further described in Section 3.2.1, to aid later reinterpretation.

Other models of DM that couple dominantly to electroweak gauge bosons through
595 either pseudo-scalar or vector mediators can be found in Ref. [72].

2.2.3. Additional considerations for $t\bar{t}$ and $b\bar{b} + \cancel{E}_T$ signatures

With the MFV assumption, the top and bottom quark can play an important role in the phenomenology. The scalar and pseudoscalar mediator models predict not only the monojet process described in Section 2.2, but also production of Dark
600 Matter in association with top (or bottom) pairs, as illustrated in Fig. 22. Dedicated searches including jets from heavy flavor quarks in the final state can be designed for this signature. Another class of simplified models, which includes a Dark Matter interpretation among many others, and yields a single top quark in the final state, is detailed in Appendix A.1.




Figure 19: Scan over mediator mass. The \cancel{E}_T distribution is compared for the scalar mediator models using the parameters as indicated. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $\cancel{E}_T > 300$ GeV and $\cancel{E}_T > 500$ GeV cut, respectively.

m_χ / GeV	$M_{\text{med}} / \text{GeV}$								
	10	20	50	100	200	300	500	1000	10000
1	0.040	0.040	0.040	0.040	0.040	0.040	0.062	0.089	0.099
10	<0.001	<0.001	0.040	0.040					0.099
50	<0.001		<0.001	<0.001	0.040	0.040			0.099
150	<0.001				<0.001	<0.001	0.062	0.089	0.099
500	<0.001						0.022	0.049	0.099
1000	<0.001							0.049	0.099

Table 6: Minimal width of the scalar mediator exchanged in s -channel divided by its mass, assuming $g_q = g_\chi = 1$. The loop-induced gluon contribution is ignored. The numbers tabulated under $2m_\chi = M_{\text{med}}$ correspond to the width calculated for $M_{\text{med}} = 5$ GeV.




Figure 20: Minimal width as a function of mediator mass for scalar and pseudo-scalar mediator assuming couplings of 1. The total width is shown as solid lines for DM masses of $m_\chi = 10$ GeV, 30 GeV, 100 GeV and 300 GeV in black, red, brown and green, respectively. The individual contributions from DM are indicated by dotted lines with the same colors. The contribution from all quarks but top is shown as magenta dotted line and the contribution from top quarks only is illustrated by the dotted blue line. The dotted beige line shows the contribution from the coupling to gluons. The dotted black line shows the extreme case $\Gamma_{min} = M_{med}$.




Figure 21: Comparison of the E_T^{miss} distributions for the scalar and pseudoscalar models for different $M_{\text{med}} = 300$ GeV and different DM masses. Ratios of the normalized distributions with respect to the first one are shown. A_{300} and A_{500} in the table denote the acceptance of the $E_T > 300$ GeV and $E_T > 500$ GeV cut, respectively.

m_χ / GeV	$M_{\text{med}} / \text{GeV}$								
	10	20	50	100	200	300	500	1000	10000
1	0.040	0.040	0.040	0.040	0.040	0.040	0.083	0.095	0.099
10	<0.001	<0.001	0.040	0.040					0.099
50	<0.001		<0.001	<0.001	0.040	0.040			0.099
150	<0.001				<0.001	<0.001	0.083	0.095	0.099
500	<0.001						0.043	0.056	0.099
1000	<0.001							0.056	0.099

Table 7: Minimal width of the pseudo-scalar mediator exchanged in s -channel divided by its mass, assuming $g_q = g_\chi = 1$. The loop-induced gluon contribution is ignored. The numbers tabulated under $2m_\chi = M_{\text{med}}$ correspond to the width calculated for $M_{\text{med}} - 5$ GeV.




Figure 22: Representative Feynman diagram showing the pair production of Dark Matter particles in association with $t\bar{t}$ (or $b\bar{b}$).

605 In addition to the $t\bar{t}$ +DM models illustrated in Fig. 22, some theoretically motivated scenario (e.g. for high $\tan\beta$ in 2HDM in the pMSSM) privilege the coupling of spin-0 mediators to down generation quarks. This assumption motivates the study of final states involving b -quarks as a complementary search to the $t\bar{t}$ +DM models, to directly probe the b -quark coupling. An example of such a model can be found in
 610 Ref. [42] and can be obtained by replacing top quarks with b quarks in Fig. 22. Note that, because of the kinematics features of b quark production relative to heavy t quark production, a $b\bar{b}$ +DM final state may only yield one experimentally visible b quark, leading to a mono- b signature in a model that conserves b flavor.

Dedicated implementations of these models for the work of this Forum are avail-
 615 able at LO+PS accuracy, even though the state of the art is set to improve on a timescale beyond that for early Run-2 DM searches as detailed in Section 4.1.5. The studies in this Section have been produced using a leading order UFO model within MADGRAPH5_AMC@NLO 2.2.2 [73, 74, 75] using PYTHIA 8 for the parton shower.

620 *Parameter scan.* The parameter scan for the dedicated $t\bar{t}+\cancel{E}_T$ searches has been studied in detail to target the production mechanism of DM associated with heavy flavor quarks, and shares many details of the scan for the scalar model with a gluon radiation. The benchmark points scanning the model parameters have been se-

lected to ensure that the kinematic features of the parameter space are sufficiently
 625 represented. Detailed studies were performed to identify points in the m_χ , $m_{\phi,a}$,
 g_χ , g_q (and $\Gamma_{\phi,a}$) parameter space that differ significantly from each other in terms
 of expected detector acceptance. Because missing transverse momentum is the key
 observable for searches, the mediator p_T spectra is taken to represent the main kine-
 matics of a model. Another consideration in determining the set of benchmarks is
 630 to focus on the parameter space where we expect the searches to be sensitive during
 the 2015 LHC run. Based on a projected integrated luminosity of 30 fb^{-1} expected
 for 2015, we disregard model points with a cross section times branching ratio
 smaller than 0.1 fb , corresponding to a minimum of one expected event assuming
 a 0.1% efficiency times acceptance.

635 The kinematics is most dependent on the masses m_χ and $m_{\phi,a}$. Figure 23 and 24
 show typical dependencies for scalar and pseudoscalar couplings respectively. Typi-
 cally, the mediator p_T spectrum broadens with larger $m_{\phi,a}$. The kinematics are also
 different between on-shell ($M_{\text{med}} > 2m_\chi$) and off-shell ($M_{\text{med}} < 2m_\chi$) mediators
 as discussed in Section 2.2. Furthermore, the kinematic differences in the \cancel{E}_T spec-
 640 trum between scalar and pseudoscalar are larger for light mediator masses with
 respect to heavier mediators. It is therefore important to choose benchmark points
 covering on-shell and off-shell mediators with sufficient granularity, including the
 transition region between on-shell and off-shell mediators.

Typically only weak dependencies on couplings are observed (see Fig 25) where
 645 the variation with width of the integral over parton distributions is unimportant. As
 shown in Section 2.1.1, for couplings $\sim O(1)$ the width is large enough that the p_T
 of the mediator is determined mainly by the PDF.

At large mediator masses ($\sim 1.5 \text{ TeV}$) or very small couplings ($\sim 10^{-2}$), width
 effects are significant, but these regimes have production cross sections that are too
 650 small to be relevant for 30 fb^{-1} and are not studied here. However, with the full
 Run 2 dataset, such models may be within reach.

Another case where the width can impact the kinematics is when $m_{\phi,a}$ is slightly
 larger than $2m_\chi$. Here, the width determines the relative contribution between on-
 shell and off-shell mediators. An example is given in Fig. 26. As the minimal width




Figure 23: Example of the dependence of the kinematics on the scalar mediator mass in the $t\bar{t} + \cancel{E}_T$ signature. The Dark Matter mass is fixed to be $m_\chi = 1\text{GeV}$.

choice pursued in this document is the most conservative one, this effect can be neglected in order to reduce the number of benchmark points to be generated.

The points for the parameter scan chosen for this model are listed in Table 5, chosen to be harmonized with those for other analyses employing the same scalar model as benchmark. Based on the sensitivity considerations above, DM masses are only simulated up to 500 GeV (but the 5 TeV mediator point is retained) leading to a total of 24 benchmark points. However for these searches we recommend to generate and simulate scalar and pseudoscalar models separately, as the kinematics differs due to the different coupling of the mediator to the final state top quarks in the two cases, as shown in Figs. 23 and 24.

Similar studies were performed in the $b\bar{b}$ case. It was found that they show the same weak dependence of the kinematics of the event on the mediator width. The same benchmark parameters of the $t\bar{t}$ case could then be chosen.




Figure 24: Example of the dependence of the kinematics on the pseudoscalar mediator mass in the $t\bar{t} + \cancel{E}_T$. The Dark Matter mass is fixed to be $m_\chi = 1\text{ GeV}$. All figures concerning the $t\bar{t} + \cancel{E}_T$ signature have been produced using a leading order model within MADGRAPH5_AMC@NLO 2.2.2, using PYTHIA 8 for the parton shower.

2.3. Colored scalar mediator, t -channel exchange

The preceding sections address models with a Dirac fermion coupled to the SM through exchange of a neutral spin-0 or spin-1 particle in an s -channel process. A t -channel process may couple the SM and DM directly, leading to a different phenomenology. For completeness, we examine a model where χ is a Standard Model (SM) singlet, a Dirac fermion; the mediating particle, labeled ϕ , is a charged scalar color triplet and the SM particle is a quark. Such models have been studied in Refs. [38, 76, 36, 34, 77, 46]. However, these models have not been studied as extensively as others in this Forum.

Following the example of Ref. [76], the interaction Lagrangian is written as

$$\mathcal{L}_{\text{int}} = g \sum_{i=1,2} (\phi_{(i),L} \bar{Q}_{(i),L} + \phi_{(i),u,R} \bar{u}_{(i),R} + \phi_{(i),d,R} \bar{d}_{(i),R}) \chi \quad (12)$$




Figure 25: Study of the dependence of kinematics on the width of a scalar mediator $t\bar{t} + \cancel{E}_T$. The width is increased up to four times the minimal width for each mediator and Dark Matter mass combination.

where $Q_{(i),L}$, $u_{(i),R}$ and $d_{(i),R}$ are the SM quarks of the i -th generation and $\phi_{(i),L}$, $\phi_{(i),u,R}$ and $\phi_{(i),d,R}$ are the corresponding mediators, which (unlike the s -channel
680 mediators) must be heavier than χ . These mediators have SM gauge representations under $(SU(3), SU(2))_Y$ of $(3, 2)_{-1/6}$, $(3, 1)_{2/3}$ and $(3, 1)_{-1/3}$ respectively. Variations of the model previously studied in the literature include coupling to the left-handed quarks only [77, 78], to the $\phi_{(i),u,R}$ [34] or $\phi_{(i),d,R}$ [76, 39], or some combination [36, 38].

685 The minimal width of each mediator is expressed, using the example of decay to an up quark, as

$$\Gamma(\phi_{(i)} \rightarrow \bar{u}_{(i)}\chi) = \frac{g_{(i)}^2}{16\pi M_{\phi_{(i)}}^3} (M_{\phi_{(i)}}^2 - m_{u_{(i)}}^2 - m_\chi^2) \times \sqrt{(M_{\phi_{(i)}}^2 - (m_{u_{(i)}} + m_\chi)^2)(M_{\phi_{(i)}}^2 - (m_{u_{(i)}} - m_\chi)^2)}, \quad (13)$$




Figure 26: Dependence of the kinematics on the width of a scalar mediator $t\bar{t} + \cancel{E}_T$. The width is increased up to the mediator mass. Choices of mediator and Dark Matter masses such that $m_{\phi,a}$ is slightly larger than $2m_\chi$ is the only case that shows a sizeable variation of the kinematics as a function of the width.

which reduces to

$$\frac{g_{(i)}^2 M_{\phi_{(i)}}}{16\pi} \left(1 - \frac{m_\chi^2}{M_{\phi_{(i)}}^2} \right)^2 \quad (14)$$

in the limit $M_{\phi_{(i)}}, m_\chi \gg m_{u_{(i)}}$.

The generation index i for $\phi_{(i)}$ is linked to the incoming fermion(s), and it runs
 690 on all three quark generations due to the MFV assumption. Ref. [76] considers
 two extreme cases for this model in terms of cross-sections: the case in which all
 mediator flavors are present, leading to the maximal cross-section, and the case in
 which only right-handed down-type mediators are present. Neither of the models
 in this reference include couplings to the third quark generation, leading to a viola-
 695 tion of the MFV assumption. In the case of purely down-type right-handed squarks
 this is still safe from flavor constraints. Furthermore, reintroducing the third gen-

eration squarks would lead to models that produce qualitatively similar signals in the mono-jet and SUSY squark searches, the main difference being the production cross-section. At the same time the presence of third generation squarks will lead to further constraints from other searches such as those for mono-bjets, for stops and for sbottoms, as discussed in Sec. 2.3.2. The studies in this Section are performed using a model with a mediator coupling to all three generation, following Ref. [46]. Further differences between the two models (hypercharge, chirality) only lead to a change in the cross-section. The LO UFO model is interfaced to MADGRAPH5_AMC@NLO v2.2.3, but it was not possible to go beyond parton-level studies and interface those models to a parton shower in time for the conclusion of this Forum. The state of the art for calculating these models is LO+PS, and the implementation of multi-parton merging has been studied in detail [79, 80, 81, 76], and further studies should be undertaken prior to generating signal samples for early Run-2 LHC searches.

The leading-order processes involved in \cancel{E}_T -jet production are shown in Fig. 27. This model can also give a signal in the \cancel{E}_T + di-jet channel when, for example, the χ is exchanged in the t -channel and the resulting ϕ pair each decay to a jet + χ . Fig. 28 shows the leading order diagrams. Except for the gg induced process, di-jet production through the third-generation mediator $\phi_{(3),u}$ is not possible, and production through $\phi_{(3),d}$ is suppressed. However, if the coupling g includes a Yukawa coupling proportional to the quark mass, and g is sufficiently large, LHC searches will still be sensitive to this model, as explained in Section 2.3.2.

The diagram involving the t -channel exchange of χ is strongly dependent upon the Dirac fermion assumption. For a Majorana fermion, $q\bar{q}$, $\bar{q}q$, and qq production would be possible with the latter having a pronounced enhancement at the LHC.

This model is similar to the simplified model considered in SUSY searches, implemented as the MSSM with only light squarks and a neutralino, except for two distinct points: the χ is a Dirac fermion and the coupling g is not limited to be weak scale ($g \ll 1$). In the MSSM, most of these processes are sub-dominant, even if resonantly enhanced, because the production is proportional to weak couplings. In the more general theories considered here, g is free to take on large values of order

1 or more, and thus diagrams neglected in MSSM simulation can occur at a much higher rate here. While constraints from SUSY jets+ \cancel{E}_T analyses on MSSM models
730 can be recast to apply to the specific model in this report, DM searches should also directly test their sensitivity to the MSSM benchmark models.

The state of the art calculation for these models is LO and they can be interfaced with a parton shower program. The studies in this Section use a LO model implementation within MADGRAPH5_AMC@NLO v2.2.3, but no parton shower could be
735 employed in the time-frame of the conclusions of this Forum. Further implementation details can be found in Section 4.1.3.

2.3.1. Parameter scan

As for the s -channel models, we adopt the simplifying assumption that the mediator masses and couplings are equal for each flavor and handedness. The free
740 parameters are then

$$\{m_\chi, M_\phi, g\}. \quad (15)$$

Ref. [76] studies the parameter space and obtains bounds on this model from LHC Run-1 mono-jet and dijets+ \cancel{E}_T data. The Forum did not exhaustively compare the kinematic distributions of the t -channel models as done in the s -channel case. In particular, the absence of a parton shower simulation can affect some of the conclusions on the points and sensitivity chosen. While this means the conclusions on
745 the parameter scan below should be taken with more caution, the model is plausible and distinctive, and it should be included in the design of early Run-2 LHC searches.

As in the s -channel models, scans should be performed over m_χ and M_ϕ . The
750 viable ranges of both parameters nearly coincide with the scan proposed for the s -channel. For the early Run-2 searches, we recommend to generate and fully simulate a subset of the s -channel mono-jet grid that accounts for the on-shell and off-shell regions. In contrast to the s -channel case, the bounds one obtains from \cancel{E}_T+X searches depend strongly on the width of the mediator, as is visible in Figs. 5 and
755 6 of Ref. [76] and in Fig. 29 (a), except in the heavy mediator limit ($M_\phi \approx 2$ TeV).




Figure 27: Leading order mono-jet t -channel processes, adapted from [76].





Figure 28: Leading order two-jet t -channel processes, adapted from [76].

This figure has been obtained applying a simplified analysis selection (cuts on the leading jet $p_T > 150$ GeV and $\eta < 2.8$, $\cancel{E}_T > 150$ GeV) using MadAnalysis [82, 83]. Figure 29 (b) also shows that, if the DM mass is low and the mediator is produced on-shell and its width is narrow, the cross-section is dominated by $qg \rightarrow q\chi\chi$ diagram. The mediator energy is then split evenly between the light DM particles and the quark, leading to a broad enhancement at $M_{\text{med}}/2$.


Points with distinct kinematic distributions for a preliminary scan in $\{m_\chi, M_\phi, g\}$ are selected taking into account the expected sensitivity of Run-2 searches, and requiring at least 100 events to pass the kinematic cuts outlined for Fig. 29 in 25 fb⁻¹ of collected data, and respect $\Gamma/M_{\text{med}} < 1$. They are outlined in Table 8. The conclusions in this table may change when a parton shower is employed together with multiparton matching.

m_χ / GeV	$M_{\text{med}} / \text{GeV}$	couplings
1	10 50 100 300	0.1, 1, 3, 7
1	500 1000	0.25, 1, 3, 7
1	2000	1, 3, 7
50	55	0.1, 1, 3, 4π
50	200 300	0.1, 1, 3, 7
500	550	1, 3
500	1000	0.25, 1, 3
500	2000	3
1000	1100	3, 4π
1000	2000	3

Table 8: Simplified model benchmark points for t -channel simplified model (spin-0 mediators coupling to Dirac DM fermions, taking the minimum width.)



(a) \cancel{E}_T distribution for a 200 GeV t -channel mediator, when varying the couplings.



(b) Leading jet p_T distribution for a 2 TeV t -channel mediator with small ($g=0.5$) to large ($g=7$) couplings with a DM mass of 1 GeV

Figure 29: [

14pt]Kinematic distributions normalized to unit area from the t -channel model from Ref. [46], using MadAnalysis [84, 82] and simplified analysis cuts on the leading jet $p_T > 150$ GeV and $\eta < 2.8$, $\cancel{E}_T > 150$ GeV. For these models, a LO UFO model is interfaced to MADGRAPH5_AMC@NLO v2.2.3, and studies are at parton-level only.

2.3.2. Additional considerations for $V + \cancel{E}_T$ signatures

The models and parameters with emission of an EW boson generally follow those in Section 2.3. even though different diagrams are involved. A representative Feynman diagram can be constructed by replacing a final-state gluon in Fig. 27 with a γ, W, Z boson, but radiation of electroweak bosons directly from the mediator also leads to a mono-boson signature.

The models considered in Section 2.3 present a relevant difference concerning final states with an electroweak boson. In the model in [46], both right- and left-handed mediators can radiate a Z boson, while only the left-handed mediator in [46] allows for W and Z radiation.

The studies in this Section use the LO+PS UFO model from [46] in MADGRAPH5_AMC@NLO v2.2.3, using PYTHIA 8 for the parton shower. Figure 30 shows the \cancel{E}_T distribution for the hadronic $Z + \cancel{E}_T$ final state, with varying DM and mediator mass, before any selection. The acceptance for a series of basic analysis selections ($\cancel{E}_T > 350$ GeV, leading jet $p_T > 40$ GeV, minimum azimuthal angle between jet and $\cancel{E}_T > 0.4$) applied at the generator level is shown in Figure 31.

The discussion of the parameter scan for the t -channel model in the case of signatures including EW bosons parallels that of the monojet case for mediator and DM masses, but no kinematic dependence on the width is observed, so a coupling scan is not needed.

Additional considerations for signatures with b -quarks + \cancel{E}_T . Models of bottom-flavored Dark Matter that are closely related to the t -channel mediated model from this Section have been proposed in Refs. [85, 86]. We describe the b -FDM model of Ref. [86], created to explain the Galactic Center (GC) gamma-ray excess observed in data collected by the Fermi-LAT collaboration [87, 88]. This model favors couplings to third-generation quarks via Yukawa couplings, therefore respecting the MFV assumption.

The model contains a Dirac fermion transforming as a flavor triplet, exclusively coupling to right-handed down-type quarks. The third component of the triplet χ_b comprises the cosmological DM. Within the MFV framework, the other fermions in




Figure 30: Missing transverse momentum distribution for the hadronic $Z+\cancel{E}_T$ final state, for the simplified model with a colored scalar mediator exchanged in the t -channel.

the flavor triplet can be made sufficiently heavy and weakly-coupled that they can be neglected in the analysis. A flavor singlet, color triplet scalar field Φ mediates the interactions between the DM and the Standard Model quarks. The model is similar to the MSSM with a light bottom squark and neutralino, and is thus a flavor-specific example of a t -channel model. Similar top-flavored models can exist, as e.g. in Refs. [89, 90]. In the case where the top coupling is the main DM coupling, the signal is very similar to a signal from a stop quark, since unlike the other t -channel cases there is no top in the initial state parton distribution functions (PDFs). This is the reason why it wasn't considered as an additional model. More recent literature shows that other flavor states could also contribute to LHC signals, as shown in Ref. [91], but such models will have to be investigated on a longer timescale with respect to that of this Forum.

The Lagrangian considered is given by

$$-\mathcal{L} \supset g \Phi^* \bar{\chi}_b b_R + \text{h.c.} \quad (16)$$




Figure 31: Acceptance for the hadronic $Z+\cancel{E}_T$ final state, for the simplified model with a colored scalar mediator exchanged in the t -channel.

810 This model is known at LO+PS accuracy, and the studies in this Section use a LO model implementation within MADGRAPH5_AMC@NLO v2.2.3 interfaced to PYTHIA 8 for the parton shower. Further implementation details can be found in Section 4.1.5.

Parameter scan. In this model, the interference of diagrams with QCD production 815 of the mediator (which scale as g_s^2) with diagrams that are proportional to the coupling g in the $b+\cancel{E}_T$ and $b\bar{b}+\cancel{E}_T$ final states. In the case of large couplings, this is not conducive to a simple scaling behavior that would allow us to reduce the number of points to be simulated. This can be seen in Fig. 33.

A full study of the parameter scan for this model was not available for this report; 820 thus for early Run-2 searches we recommend scanning a range of possible widths as discussed in a more limited way than for the t -channel mono-jet, spanning from the minimal width to a value approaching the particle limit, e.g. $g = 0.5, 1, 2, 3$. A coupling benchmark such as $g = 1$ should be considered for each mass point

since this would be a distinctive feature of this benchmark from SUSY models with
825 sbottom squarks (see Section 2.3 for further discussion).

A scan of Dark Matter and mediator masses should be done in the on-shell region
 $M_\Phi > m_\chi + m_b$, since the cross-sections in the off-shell region are too small to be
probed with early LHC data, spanning from 10 to 500 GeV in m_χ and from 10 to
1300 GeV in M_Φ . Examples of the kinematic distributions produced by this model
830 are shown in Fig. 32⁸.




Figure 32: E_T^{miss} (left) and jet multiplicity (right) for various DM and mediator masses and couplings normalized to the relic density observed in the early universe. Studies in this section use a LO UFO model implementation within MADGRAPH5_AMC@NLO v2.2.3 interfaced to PYTHIA 8 for the parton shower.

2.4. Spin-2 mediator

In models with extra dimensions, the Kaluza-Klein excitations of the graviton could also serve as a mediator between the SM and dark sector physics. This kind of model was not studied in the forum and is not included in the recommendations,
835 but models such as Ref. [92, 93] may warrant further study on a longer timescale.

2.5. Presentation of results for reinterpretation of s -channel mediator models

The aim of the parameter grid optimization done for the s -channel models in the previous sections is to reduce the parameter space that must be simulated. We then

⁸Following the grounding assumptions in this report, the normalization to the relic density is considered only in these example plots rather than as a necessary ingredient for the parameter scan of this model.




Figure 33: E_T (left) and jet multiplicity (right) for $m_\chi = 35$ GeV and $M_\phi = 500$ GeV for varying couplings of $g = 1, 2$

need a procedure for populating the full parameter space by using the simulated
840 grid points. We recommend doing this as follows:

- When the dependences on parameters are known, the cross sections and efficiencies at general points can be calculated from the grid data.
- In other cases, this information can be obtained by interpolation between the grid points. We have chosen the grid points so that the dependence is
845 sufficiently smooth that this will be possible.

The results of the scan over the couplings presented in the previous sections indicate that there are no changes in kinematic distributions for different choices of the coupling strengths. This means that the acceptance remains the same in the whole g_q - g_χ plane and it is sufficient to perform the detector simulation only for
850 one single choice of g_q, g_χ . The resulting truth-level selection acceptance and the detector reconstruction efficiency can then be applied to all remaining grid points in the g_q - g_χ plane where only the generator-level cross section needs to be known. This significantly reduces the computing time as the detector response is by far the most CPU-intensive part of the Monte Carlo sample production. However, the
855 number of generated samples can be reduced even further if a parameterization of the cross section dependence from one grid point to another exists. In this section, we describe the details of a cross section scaling procedure that can be used to reinterpret results for a fixed coupling for s -channel mediator models. The studies

in this section employ the POWHEG [43] generator.

860 The propagator for the s -channel exchange is written in a Breit-Wigner form as

$$\frac{1}{q^2 - M_{\text{med}}^2 + iM_{\text{med}}\Gamma},$$
 where q is the momentum transfer calculated from the two partons entering the hard process after the initial state radiation, which is equivalent to the momentum of the DM pair⁹. The size of the momentum transfer with respect to the mediator mass allows us to identify three cases:

- 865
- off-shell mediator, when $q^2 \gg M_{\text{med}}^2$ leading to suppressed cross sections,
 - on-shell mediator, when $q^2 \sim M_{\text{med}}^2$ leading to enhanced cross sections,
 - effective field theory (EFT) limit when $q^2 \ll M_{\text{med}}^2$.

In the case of the off-shell mediator and the EFT limit, the first and second term in the propagator dominate, respectively, which reduces the dependence on the
 870 mediator width. Therefore, in these cases one can approximate the cross section as

$$\sigma \propto g_q^2 g_\chi^2. \quad (17)$$

The on-shell regime is the most interesting one as it gives the best chances for a discovery at the LHC given the cross section enhancement. The propagator term with the width cannot be neglected in this case and, in the narrow width approximation which requires $\Gamma \ll M_{\text{med}}$ (this is not necessarily the case in the benchmarks
 875 considered in the scans), one can integrate

$$\int \frac{ds}{(s - M_{\text{med}}^2)^2 + M_{\text{med}}^2 \Gamma^2} = \frac{\pi}{M_{\text{med}} \Gamma} \quad (18)$$

which further implies the cross section scaling

$$\sigma \propto \frac{g_q^2 g_\chi^2}{\Gamma}. \quad (19)$$

The narrow width approximation is important here as it ensures an integration over parton distribution functions (PDFs) can be neglected. In other words, it is assumed

⁹Using a running width and replacing the denominator of the propagator with $q^2 - M_{\text{med}}^2 + iQ^2 \frac{\Gamma}{M_{\text{med}}}$ should be considered in the case of wide mediators [94].

the integrand in Eq. 18 is non-zero only for a small region of s , such that the PDFs
 880 can be taken to be constant in this range. By simplifying the dependence of the
 minimal width on the couplings as $\Gamma \sim g_q^2 + g_\chi^2$, one can approximate this scaling
 rule in the extreme cases as follows

$$\sigma \propto \frac{g_q^2 g_\chi^2}{g_q^2 + g_\chi^2} \xrightarrow{g_q \ll g_\chi} g_q^2 \quad (20)$$

$$\sigma \propto \frac{g_q^2 g_\chi^2}{g_q^2 + g_\chi^2} \xrightarrow{g_q \gg g_\chi} g_\chi^2. \quad (21)$$

However, it is important to keep in mind that this formula omits color and multi-
 plicity factors as well as possible Yukawa suppression, and there is no simple scaling
 885 rule for how the cross section changes with the DM mass and the mediator mass, or
 for mediators with a large width, because PDFs matter in such cases as well. There-
 fore, the scaling procedure outlined above is expected to work only for fixed masses
 and fixed mediator width, assuming the narrow width approximation applies.

Figure 34 shows the minimal width over the mediator mass in the g_q - g_χ plane
 890 for vector and scalar mediators for $M_{\text{med}} = 100$ GeV and 1000 GeV, taking $m_\chi =$
 10 GeV. The individual colors indicate the lines of constant width, along which the
 cross section scaling may work for narrow mediators. The limiting case $\Gamma_{\text{min}} = M_{\text{med}}$
 defines the upper values of the couplings below which the narrow width approxima-
 tion can be considered and provides more stringent constraint than the perturbative
 895 limit $g_q = g_\chi = 4\pi$. For vector and axial-vector mediators, the minimal width is
 predominantly defined by g_q due to the number of quark flavors and the color fac-
 tor. On the contrary, both the SM and DM partial width have comparable contribu-
 tions in case of scalar and pseudo-scalar mediators if the top quark channel is open
 ($M_{\text{med}} > 2m_t$). However, mostly g_χ defines the minimal width for $M_{\text{med}} < 2m_t$ due
 900 to the Yukawa-suppressed light quark couplings.

The performance of the cross section scaling is demonstrated in Fig. 35 where
 two mass points $M_{\text{med}} = 100$ GeV and 1 TeV with $m_\chi = 10$ GeV are chosen and
 rescaled from the starting point $g_q = g_\chi = 1$ according to Eq. 19 to populate the
 whole g_q - g_χ plane. This means the width is not kept constant in this test and this is




Figure 34: Minimal width over the mediator mass for vector (top) and scalar (bottom) mediators as a function of the individual couplings g_q and g_{χ} , assuming $M_{\text{med}} = 100$ GeV (left) and $M_{\text{med}} = 1$ TeV (right). $m_{\chi} = 10$ GeV is considered in all cases. Only the cases with $\Gamma_{\text{min}} < M_{\text{med}}$ are shown.

905 done in purpose in order to point out deviations from the scaling when the width is
 altered. For each mass point, the rescaled cross section is compared to the generator
 cross section and the ratio of the two is plotted. For the given choice of the mass
 points, the scaling seems to work approximately within the precision of $\sim 20\%$ in
 the region where $\Gamma_{\min} < M_{\text{med}}$. Constant colors indicate the lines along which the
 910 cross section scaling works precisely and there is a remarkable resemblance of the
 patterns shown in the plots of the mediator width. To prove the scaling along the
 lines of constant width works, one such line is chosen in Fig. 36 for a scalar mediator,
 defined by $M_{\text{med}} = 300 \text{ GeV}$, $m_\chi = 100 \text{ GeV}$, $g_q = g_\chi = 1$, and the rescaled and
 generated cross sections are found to agree within 3%.

915 2.5.1. Proposed parameter grid for cross-section scaling

We propose to deliver collider results in the g_q - g_χ plane using the following
 prescription, to ease reinterpretation through cross-section scaling:

- Since the shapes of kinematic quantities do not change for different couplings,
 use the acceptance and efficiency for the available $m_\chi = 50 \text{ GeV}$, $M_{\text{med}} =$
 920 300 GeV grid point from the M_{med} - m_χ plane for the scalar and pseudo-scalar
 mediator. In case of the vector and axial-vector mediator, use the grid point
 $m_\chi = 150 \text{ GeV}$, $M_{\text{med}} = 1 \text{ TeV}$.
- Generate additional samples in order to get generator cross sections only. For
 scalar and pseudo-scalar mediator, choose $m_\chi = 50 \text{ GeV}$, $M_{\text{med}} = 300 \text{ GeV}$
 925 with the following values for $g_q = g_\chi$: 0.1, 1, 2, 3. For vector and axial vector
 mediator, choose $m_\chi = 150 \text{ GeV}$, $M_{\text{med}} = 1 \text{ TeV}$ with the following values for
 $g_q = g_\chi$: 0.1, 0.25, 0.5, 0.75, 1, 1.25, 1.5. The upper values are defined by
 the minimal width reaching the mediator mass.
- Rescale the generator cross sections for on-shell resonance production along
 the lines of constant width in order to populate the whole g_q - g_χ plane in the
 region $\Gamma_{\min} < M_{\text{med}}$. The scaling follows from Eq. 19 which for the constant




Figure 35: Ratio of the rescaled and generated cross sections in the g_q - g_χ plane. The point at $g_q = g_\chi = 1$, taken as a reference for the rescaling, is denoted by a star symbol. Scalar model with $M_{\text{med}} = 100$ GeV (left) and 1 TeV (right) is plotted for $m_\chi = 10$ GeV. The limiting case $\Gamma_{\text{min}} = M_{\text{med}}$ is indicated by a black line and no results are shown beyond.




Figure 36: Scaling along the lines of constant width. The line of constant width for $M_{\text{med}} = 300$ GeV and $m_\chi = 100$ GeV, intercepting $g_q = g_\chi = 4$ is shown on left. The generated and rescaled cross sections are compared in the middle, the corresponding ratio is shown on right.

width implies:

$$\sigma' = \sigma \times \frac{g_q'^2 g_\chi'^2}{g_q^2 g_\chi^2}. \quad (22)$$

2.5.2. Rescaling to different mediator width

930 In general it is also important to consider a larger mediator width than Γ_{\min} in order to accommodate additional interactions of the mediator with the visible and hidden sector particles [42, 41]. If the narrow width approximation applies, the cross section scaling method described above can be used to reinterpret the results presented for the minimal width, since multiplying the width by factor n is
935 equivalent to changing the coupling strength by factor \sqrt{n} , i.e.

$$\sigma(g_q, g_\chi, n\Gamma_{\min}(g_q, g_\chi)) \propto \frac{g_q^2 g_\chi^2}{\Gamma_{\min}(\sqrt{n}g_q, \sqrt{n}g_\chi)}. \quad (23)$$

The cross section for the sample with couplings g_q and g_χ and modified mediator width $\Gamma = n\Gamma_{\min}$ can therefore be rescaled from a sample generated with the minimal width corresponding to the couplings scaled by \sqrt{n} as described in the following formula.

$$\sigma(g_q, g_\chi, n\Gamma_{\min}(g_q, g_\chi)) = \frac{1}{n^2} \sigma(\sqrt{n}g_q, \sqrt{n}g_\chi, \Gamma_{\min}(\sqrt{n}g_q, \sqrt{n}g_\chi)) \quad (24)$$

940 The advantage of doing this is in the fact that no event selection and detector response needs to be simulated since the changes in couplings do not have an effect on the shapes of kinematic distributions.

It should be noted again that this procedure is only useful when the narrow width approximation applies. Care must be taken to ensure that is the case. For
945 example, in the vector and axial-vector cases, one quickly breaks this approximation even for small n .

Additional considerations for $t\bar{t}$ and $b\bar{b} + \cancel{E}_T$ signatures. The cross-section scaling considerations shown in Sec. 2.5 still apply for the reactions in the scalar and pseudoscalar models with explicit b and t quarks. Here we detail the specific studies
950 done for the $t\bar{t}$ model.

Given that the kinematics are similar for all couplings $g \simeq 1$, we recommend to generate only samples with $g_\chi = g_q = 1$. It follows from this that these benchmark points should be a good approximation for non-unity couplings and for $g_\chi \neq g_q$, provided that the sample is rescaled to the appropriate cross section times branching
955 ratio.

While the simple scaling function

$$\sigma' \times BR' = [\sigma \times BR] \times \left(\frac{g'_q}{g_q} \right)^2 \times \left(\frac{g'_\chi}{g_\chi} \right)^2 \times \frac{\Gamma}{\Gamma'} \quad (25)$$

is sufficient for a limited range of coupling values (see Fig. 37 for example), this scaling is only approximate (up to 20%) and relies on the narrow width approximation, ignoring PDFs effects.

3. Specific models for signatures with EW bosons

960 $\hat{\mathcal{A}}_{\tilde{a}}$

In this Section, we consider specific models with a photon, a W boson, a Z boson or a Higgs boson in the final state ($V + \cancel{E}_T$ signature), accompanied by DM particles that either couple directly to the boson or are mediated by a new particle. The common feature of those models is that they provide different kinematic distributions
965 with respect to the models described in Section 2.

The models considered in this Section can be divided into two categories:

V-specific simplified models These models postulate direct couplings of new mediators to bosons, e.g. they couple the Higgs boson to a new vector or to a new scalar [48, 96].

970 **Models involving a SM singlet operator including a boson pair that couples to DM through a contact interaction**

Shown on the right-hand side of Figure 38, these models allow for a contact interaction vertex that directly couples the boson to DM [97, 45, 98, 96]. The models of this type included in this report are devoted to simplified models since UV completions for most of these operators proceed through loops and
975 are not available to date. These models provide a benchmark to motivate




Figure 37: An example comparing a simple cross section scaling versus the computation from the MADGRAPH5_AMC@NLO generator, for a scalar $t\bar{t} + \cancel{E}_T$ model with $m_\phi = 400$ GeV, $m_\chi = 1$ GeV and all couplings set to unity. In this example, the scaling relationship holds for Γ_ϕ/m_ϕ below 0.2, beyond which finite width effects become important and the simple scaling breaks down.

signal regions that are unique to searches with EW final states and would otherwise not be studied. However, we recommend to use these models as placeholders and emphasize model-independent results especially in signal regions tailored to these models. Wherever results are interpreted in terms of these operators, a truncation procedure to ensure the validity of the EFT should be employed, as detailed in Section 5.

The following Sections describe the models within these categories, the chosen parameters for each of the benchmark models and the studies towards the choices of the parameters to be scanned.




Figure 38: Sketch of benchmark models including a contact interaction for V+MET searches, adapted from [95].

3.1. Specific simplified models including EW bosons, tailored to Higgs+MET searches

Three benchmark simplified models [48, 96] are recommended for Higgs+MET searches:

- A model where a vector mediator (Z'_B) is exchanged in the s-channel, radiates a Higgs boson, and decays into two DM particles (Fig. 39 (a)). As in Section 2.1, we conservatively omit couplings of the Z'_B to leptons.
- A model where a scalar mediator S is emitted from the Higgs boson and decays to a pair of DM particles (Fig. 40).
- A model where a vector Z' is produced resonantly and decays into a Higgs boson plus an intermediate heavy pseudoscalar particle A^0 , in turn decaying into two DM particles (Fig. 39 (b)).

These models are kinematically distinct from one another, as shown in the comparison of the \cancel{E}_T spectra in Fig. 41 for high and low masses of the pseudoscalar mediator. Figure 41 (a) shows the \cancel{E}_T distribution for models with high mediator masses ($m_S = 1$ TeV, $m_{Z'} = 1$ TeV, $m_{A^0} = 1$ TeV) and DM mass of either 50 (Z'_B and A^0 models) or 65 GeV (scalar mediator model). Figure 41 (b) shows the \cancel{E}_T distribution for models with low pseudoscalar mediator masses ($m_{Z'_B} = 100$ GeV, $m_{Z'} = 1$ TeV, $m_{A^0} = 100$ GeV) and DM mass of 1 TeV for all models.




Figure 39: Examples of Feynman diagrams leading to Higgs+ \cancel{E}_T events: (a) a model with a vector mediator (Z') coupling with DM and with the Higgs boson h , and (b) a 2HDM model with a new invisibly decaying pseudoscalar A^0 from the decay of an on-shell resonance Z' giving rise to a Higgs+ \cancel{E}_T signature






Figure 40: Examples of Feynman diagrams leading to Higgs+ \cancel{E}_T events for a model with a scalar mediator (S) coupling with DM and with the Higgs boson h .



(a) High mediator mass



(b) Low mediator mass

Figure 41: Comparison of the missing transverse momentum distributions at generator level in different simplified models leading to a $Higgs + \cancel{E}_T$ signature. The model parameter settings are detailed in the text. The figures in this Section have been obtained using LO UFO models within MADGRAPH5_AMC@NLO v2.2.2.3, interfaced to PYTHIA 8 for the parton shower.

Predictions for this class of models have been so far considered at LO+ Parton Shower (PS), even though they could be extended to NLO+PS in the near future. The studies in this Section have been performed using a model within MAD-
 1005 GRAPH5_AMC@NLO v5.1.5.12, interfaced to PYTHIA 6 for the parton shower. The implementation details for these models are discussed in Section 4.2.1.

3.1.1. \cancel{E}_T +Higgs from a baryonic Z'

The model shown in Fig. 39 (a) postulates a new gauge boson Z' corresponding
 1010 to a new $U(1)_B$ baryon number symmetry. The stable baryonic states included in this model are the DM candidate particles. The mass of the Z' boson is acquired through a baryonic Higgs h_B , which mixes with the SM Higgs boson.

The interactions between the Z' , the quarks and the DM are described by the following Lagrangian:

$$\mathcal{L} = g_q \bar{q} \gamma^\mu q Z'_\mu + g_\chi \bar{\chi} \gamma^\mu \chi Z'_\mu. \quad (26)$$

The quark couplings g_q are fixed to be equal to one third of the gauge coupling
 1015 g_B , while the DM coupling to the Z' are proportional to the baryon number and to the gauge coupling ($g_\chi = B g_B$). No leptonic couplings of the Z' are allowed, thus evading dilepton constraints. After incorporating the mixing of the baryonic and SM Higgs bosons, this model is described by the following Lagrangian term at
 1020 energies below $m_{Z'}$ ¹⁰:

$$\mathcal{L}_{\text{eff}} = -\frac{g_q g_\chi}{m_{Z'}^2} \bar{q} \gamma^\mu q \bar{\chi} \gamma_\mu \chi \left(1 + \frac{g_{hZ'/Z'}}{m_{Z'}^2} h \right), \quad (27)$$

The first term of this equation is the standard DMV model in the large $M_{Z'}$ limit. This term can lead to a monojet signature, which can be also used to constrain this model. The second term describes the interaction between the Z' and the SM Higgs boson, via the coupling $g_{hZ'/Z'} = \frac{m_{Z'} 2 \sin \theta}{v_B}$, where $\sin \theta$ is the mixing angle between

¹⁰The operator in Eqn. 27 is an effective one, to highlight the two main terms. The full dimension-4 simplified model is used in the model for event generation.

1025 the SM Higgs and the baryonic Higgs h_B , and v_B is the Baryonic Higgs vacuum expectation value.

In its most general form, this model can contribute to mono-Z signals due to the Z' mixing with the Z or photon. Note that EWSB and $U(1)_B$ breaking do not lead to this mixing at tree-level. Instead, kinetic mixing occurs between the $U(1)_Y$ and $U(1)_B$ gauge bosons due to the gauge invariant term $F_Y^{\mu\nu} F_{B\mu\nu}$. This mixing is a free parameter which we assume to be small in order to focus on the mono-Higgs signature. Mixing may also occur due to radiative corrections, however we choose to ignore this here as it is model dependent.

1035 The predictions of the model depend upon the two additional parameters beyond an s-channel simplified model, namely the mixing angle between baryonic Higgs h_B and the SM-like Higgs boson $\sin\theta$ and the coupling of the mediator to SM-like Higgs boson, $g_{hZ'Z'}$. Thus, a full model is specified by:

$$\{M_{\text{med}}, m_\chi, g_\chi, g_q, \sin\theta, g_{hZ'Z'}\}. \quad (28)$$

Parameter scan. The width of the Z' mediator is calculated using all possible decays to SM particles (quarks) and to pairs of DM particles if kinematically allowed as in the DMV model.

1040 The dependence of the missing transverse momentum (\cancel{E}_T) on the model parameters is studied by varying the parameters one at a time. The variation of parameters other than M_{med} and m_χ does not result in significant variations of the \cancel{E}_T spectrum, as shown in Figures 42. Figure 43 shows that for an on-shell mediator, varying m_χ with the other parameters fixed does not affect the \cancel{E}_T distribution, while the distribution broadens significantly in the case of an off-shell mediator. For this reason, the same grid in M_{med}, m_χ as for the vector mediator of the jet+ \cancel{E}_T search (Table 1) is chosen as a starting point. The coupling $g_{hZ'Z'}$, along with g_q and g_χ , are subject to perturbativity bounds:

$$g_q, g_\chi < 4\pi$$

1050 ,

and

$$g_{hZ'Z'} < \sqrt{4\pi} m_{Z'} \sin \theta$$

The value $g_{hZ'Z'}/m_{Z'} = 1$ is chosen as a benchmark value for the generation of Monte Carlo samples since it maximizes the cross section (as shown in the following paragraph) without violating the bounds. The mediator-DM coupling g_χ is fixed to 1, and the mediator-quark g_q coupling is fixed to 1/3. The kinematic distributions do not change as a function of these parameters, so results for other values of $g_{hZ'Z'}/m_{Z'}$, g_χ and g_q can be obtained through rescaling by the appropriate cross sections.

Figs 44 and 45 show the kinematic distributions for the two leading jets in the $H \rightarrow \bar{b}b$ decay channel, for two values of the mediator mass and varying the DM mass.

Experimentalists should perform further studies, beyond those studies performed for the forum, to estimate the reach of the analysis with respect to all points in the grid and therefore decide on a smaller set of grid points to be generated.

3.1.2. $\cancel{E}_T + \text{Higgs}$ from a scalar mediator

A real scalar singlet S coupling to DM can be introduced as a portal between SM and the dark sector through the Higgs field. The most general scalar potential is detailed in Ref. [99], including terms that break \mathbb{Z}_2 . The \mathbb{Z}_2 symmetry, which causes the new scalar to also be a DM candidate, is not covered in this report, but follows Ref. [48] introducing an additional coupling to DM that breaks \mathbb{Z}_2 and leads to a new invisible decay of S . For this reason, no symmetry is broken and no new interactions arise, so there is no dependence on the vacuum expectation value of S : a shift in the field leads to a redefinition of the model couplings. The new scalar S mixes with the SM Higgs boson, and couples to DM through a Yukawa term y_χ . The relevant terms in the scalar potential are:




Figure 42: Missing transverse momentum distributions at generator level in the vector mediator scenario for different values of: the mediator-DM coupling g_χ (left), and the coupling between the mediator and the SM-like Higgs boson, scaled by the mediator mass, $g_{hZ'Z'}/m_{Z'}$ (right).







Figure 43: Missing transverse momentum distributions at generator level in the vector mediator scenario: for different values of the DM mass m_χ and a mediator mass of $M_{\text{med}} = 100 \text{ GeV}$ (left) and $M_{\text{med}} = 1 \text{ TeV}$ (right).



(a) Leading b -jet transverse momentum




(b) Leading b -jet pseudorapidity




(c) Angular distance between the two leading b -jets


Figure 44: Comparison of the kinematic distributions for the two leading b -jets (from the Higgs decay) in the vector Z' simplified model, when fixing the Z' mass to 100 GeV and varying the DM mass.



(a) Leading b -jet transverse momentum



(b) Leading b -jet pseudorapidity



(c) Angular separation of the two leading b -jets

Figure 45: Comparison of the kinematic distributions for the two leading jets from the Higgs decay in the vector Z' simplified model, when fixing the Z' mass to 1000 GeV and varying the DM mass.

$$\begin{aligned}
V &\supset a|H|^2 S + b|H|^2 S^2 + \lambda_h |H|^4 \\
&\longrightarrow \frac{1}{2}a(h+v)^2 S + \frac{1}{2}b(h+v)^2 S^2 + \frac{\lambda_h}{4}(h+v)^4,
\end{aligned} \tag{29}$$

where a, b are new physics couplings and λ_h is the Higgs quartic coupling.

The additional Lagrangian terms for this model are:

$$\mathcal{L} \supset -y_\chi \bar{\chi} \chi (\cos \theta S - \sin \theta h) - \frac{m_q}{v} \bar{q} q (\cos \theta h + \sin \theta S) \tag{30}$$

where θ is the mixing angle between the Higgs boson and the new scalar.

1080 Mono-Higgs signals in this second model arise through processes shown in Fig. 40 (a,b), or through the radiation of a Higgs boson from the t quark in the production loop, in Fig. 40 (c). The first two processes depend on the $h^2 S$ and $h S^2$ cubic terms in Eq. (29). At leading order in $\sin \theta$, these terms are:

$$V_{\text{cubic}} \approx \frac{\sin \theta}{v} (2m_h^2 + m_S^2) h^2 S + b v h S^2 + \dots, \tag{31}$$

1085 with a and λ_h expressed in terms of $\sin \theta$ and m_h^2 , respectively. At leading order of $\sin \theta$, the $h^2 S$ term is fixed once the mass eigenvalues m_h, m_S and mixing angle are specified. The $h S^2$ term is not fixed and remains a free parameter of the model, depending on the new physics coupling b .

This model also has mono-X signatures through h/S mixing. This model is related to the scalar model discussed in Sec. 2.2 in the case of $m_S \gg m_h$ or $m_h \gg m_S$ 1090 and M_{med} equal to the lighter of the two masses, albeit with different mono-Higgs signatures due to the $h S^2$ vertex.

Parameter scan. The model is described by five parameters:

1. the Yukawa coupling of heavy scalar to dark matter, g_χ (also referred to as y_χ)
- 1095 2. the mixing angle between heavy scalar and SM-like Higgs boson, $\sin \theta$;
3. the new physics coupling, b ;
4. mass of heavy scalar, m_S , also termed M_{med} ;

5. mass of DM. m_χ ;

The mixing angle is constrained from current Higgs data to satisfy $\cos \theta = 1$ within 10% and therefore $\sin \theta \lesssim 0.4$. This provides a starting point for the parameter scan in this model: we recommend to set $\sin \theta = 0.3$.

Figure 47 shows that the kinematics does not depend on the value of this angle, and results for different values can be obtained via rescaling the results for this mixing angle according to the relevant cross-section. It can also be observed from Figures 48 and 46 that the kinematics of this model follows that of the equivalent jet+ \cancel{E}_T model: only small changes are observed in the on-shell region, while the relevant distributions diverge when the mediator is off-shell. For this reason, the same grid in M_{med}, m_χ as for the scalar mediator of the jet+ \cancel{E}_T search (Table 5) is chosen as a starting point. The Yukawa coupling to DM y_{DM} is set to 1, the new physics coupling between scalar and SM Higgs $b = 3$. Results for other values can be obtained via a rescaling of the results for these parameters.

Figs. 49 and 50 show the kinematic distributions for the two leading jets in the $H \rightarrow \bar{b}b$ decay channel, for two values of the mediator mass and varying the DM mass.

3.1.3. Higgs+ \cancel{E}_T signal from 2HDM model with a Z' and a new pseudoscalar

In this simplified model [96], a new Z' resonance decays to a Higgs boson h plus a heavy pseudoscalar state A^0 in the 2HDM framework, which in turn decays to a DM pair. This model is represented in the diagram in Fig. 39 (b).

The motivation for coupling the DM to the pseudoscalar is that DM coupling to a Higgs or Z' boson is generically constrained by other signal channels and direct detection. A reason to consider this model is that it has different kinematics due to the on-shell Z' production, where for heavy Z' masses the \cancel{E}_T and p_T spectra are much harder. This model can satisfy electroweak precision tests and constraints from dijet resonance searches, and still give a potentially observable Higgs+ \cancel{E}_T signal.

The model comprises of two doublets, where Φ_u couples to up-type quarks and




Figure 46: Missing transverse momentum distributions at generator level in the scalar mediator scenario, for different values of: the new physics coupling g_b (left), and the mediator-DM coupling g_χ (right).




Figure 47: Missing transverse momentum distributions at generator level in the scalar mediator scenario: for different values of the mixing angle θ .







Figure 48: Missing transverse momentum distributions at generator level in the scalar mediator scenario: for different values of the DM mass m_χ and a mediator mass of $M_{\text{med}} = 100 \text{ GeV}$ (left) and $M_{\text{med}} = 1 \text{ TeV}$ (right).



(a) Leading b -jet transverse momentum




(b) Leading b -jet pseudorapidity




(c) Angular distance between the two leading b -jets


Figure 49: Comparison of the kinematic distributions for the two leading jets from the Higgs decay in the scalar simplified model, when fixing the new scalar mass to 100 GeV and varying the DM mass.



(a) Leading b -jet transverse momentum



(b) Leading b -jet pseudorapidity



(c) Angular distance between the two leading b -jets

Figure 50: Comparison of the kinematic distributions for the two leading jets from the Higgs decay in the scalar simplified model, when fixing the new scalar mass to 1000 GeV and varying the DM mass.

Φ_d couples to down-type quarks and leptons:

$$-\mathcal{L} \supset y_u Q \tilde{\Phi}_u \bar{u} + y_d Q \Phi_d \bar{d} + y_e L \Phi_d \bar{e} + \text{h.c.} \quad (32)$$

After electroweak symmetry breaking, the Higgs doublets attain vacuum expectation values v_u and v_d , and in unitary gauge the doublets are parametrized as

$$\begin{aligned} \Phi_d &= \frac{1}{\sqrt{2}} \begin{pmatrix} -\sin \beta H^+ \\ v_d - \sin \alpha h + \cos \alpha H - i \sin \beta A^0 \end{pmatrix}, \\ \Phi_u &= \frac{1}{\sqrt{2}} \begin{pmatrix} \cos \beta H^+ \\ v_u + \cos \alpha h + \sin \alpha H + i \cos \beta A^0 \end{pmatrix} \end{aligned} \quad (33)$$

where h, H are neutral CP-even scalars, H^\pm is a charged scalar, and A^0 is a neutral CP-odd scalar. In this framework, $\tan \beta \equiv v_u/v_d$, and α is the mixing angle that diagonalizes the $h-H$ mass squared matrix. This model also contains an additional scalar singlet ϕ that leads to spontaneous symmetry breaking. We take $\alpha = \beta - \pi/2$, in the alignment limit where h has SM-like couplings to fermions and gauge bosons as per Ref. [100], and $\tan \beta \geq 0.3$ as implied from the perturbativity of the top Yukawa coupling. The Higgs vacuum expectation values lead to $Z-Z'$ mass mixing, with a small mixing parameter given by

$$\begin{aligned} \epsilon &\equiv \frac{1}{M_{Z'}^2 - M_Z^2} \frac{g g_z}{2 \cos \theta_w} (z_d v_d^2 + z_u v_u^2) \\ &= \frac{(M_Z^0)^2}{M_{Z'}^2 - M_Z^2} \frac{2 g_z \cos \theta_w}{g} z_u \sin^2 \beta, \end{aligned} \quad (34)$$

where z_i are the Z' charges of the two Higgs doublets, and g and g_z related to the mass-squared values in absence of mixing $(M_Z^0)^2 = g^2(v_d^2 + v_u^2)/(4 \cos^2 \theta_w)$ and $(M_{Z'}^0)^2 = g_z^2(z_d^2 v_d^2 + z_u^2 v_u^2 + z_\phi^2 v_\phi^2)$.

The production cross section for this model scales as $(g_z)^2$, as the decay width for this process to leading order in ϵ (Eq. 34) is

$$\Gamma_{Z' \rightarrow h A^0} = (g_z \sin \beta \cos \beta)^2 \frac{|p|}{24\pi} \frac{|p|^2}{M_{Z'}^2}. \quad (35)$$

where the center of mass momentum for the decay products is

$$|p| = \frac{1}{2M_{Z'}} \sqrt{(M_{Z'}^2 - (m_h + m_{A^0})^2)(M_{Z'}^2 - (m_h - m_{A^0})^2)}. \quad (36)$$

The Z' can also decay to Zh , leading to the same signature if the Z decays invisibly.

The partial width for this decay is:

$$\Gamma_{Z' \rightarrow hZ} = (g_z \sin \beta^2)^2 \frac{|p|}{24\pi} \left(\frac{|p|^2}{M_{Z'}^2} + 3 \frac{M_Z^2}{M_{Z'}^2} \right). \quad (37)$$

We recommend to generate these two decays separately and combine them at a later stage.


Parameter scan. The model is described by five parameters:

- 1135 • the pseudoscalar mass M_{A^0} ,
- the DM mass m_χ ,
- the Z' mass, $M_{Z'}$,
- $\tan \beta (\equiv v_u/v_d)$,
- the Z' coupling strength g_z .


1140 To study the signal production and kinematic dependencies on these parameters, we produced signal samples varying each of the five parameters through MADGRAPH5_AMC@NLO for the matrix element, PYTHIA 8 for the parton shower, and DELPHES[101] for a parameterized detector-level simulation.

1145 As seen in Fig. 51, variations of $\tan \beta$ does not lead to any kinematic difference and the production cross section simply scales as a function of $\tan \beta$. Hence we recommend to fix $\tan \beta$ to unity in the signal generation.

Similarly, variations of g_z do not lead to any kinematic changes. The value of g_z for a given $M_{Z'}$ and $\tan \beta$ can be set according to the maximum value allowed by electroweak global fits and dijet constraints, as described in [96]. Since this
1150 parameter does not influence the kinematics, we leave it up to individual analyses to decide whether they generate benchmark points only according to these external constraints.




(a) \cancel{E}_T distribution




(b) $\Delta\phi$ distance between the two b -jets

Figure 51: Kinematic distributions of the signal process varying $\tan\beta$, in the case of a Higgs boson decaying into two b quarks, after parameterized detector simulation: no kinematic dependence is observed on the mixing angle.

Since the DM pair are produced as a result of the decay of A^0 , there are minimal kinematic changes when varying m_χ as long as $m_\chi < M_{A^0}/2$ so that A^0 production is on-shell, as shown in Fig. 52 and 53 (before detector simulation).



(a) \cancel{E}_T distribution



(b) $\Delta\phi$ distance between the two b -jets

Figure 52: Kinematic distributions of the signal process varying m_χ : minimal kinematic dependency on m_χ as expected when A^0 is produced on-shell. Plots shown for $M_{Z'} = 1000$ GeV, $M_{A^0} = 300$ GeV.

We recommend to produce signal events for a fixed $g_z = 0.8$, $\tan\beta = 1$ and $m_\chi = 100$ GeV. For these values, we scan the 2-D parameter space of $M_{Z'}, M_{A^0}$ with $M_{Z'} = 600, 800, 1000, 1200, 1400$ GeV, and $M_{A^0} = 300, 400, 500, 600, 700, 800$ GeV with $M_{A^0} < M_{Z'} - m_h$, for a total of 24 points. The choice of scan is justified by the sensitivity study in [96]: the expected LHC sensitivity for Run-2 is up to $M_{Z'} \sim 1.5$ TeV. For the parameter scan, the DM mass is fixed to 100 GeV.




Figure 53: Missing transverse momentum distributions at generator level in the $Z' + 2\text{HDM}$ scenario for different values of the DM mass m_χ , with $m_{Z'} = 1 \text{ TeV}$ and $m_{A^0} = 300 \text{ GeV}$ (left) and $m_{A^0} = 1 \text{ TeV}$ (right).

For two $M_{Z'}$, M_{A^0} value sets, we vary the DM mass to obtain sample cross section for rescaling results. All LO cross sections for the various parameter scan points are reported on HEPData [57]. The parameter scan excludes the off-shell region,
1165 as the cross-sections are suppressed and the LHC would not have any sensitivity to these benchmark points in early data.

The kinematic distributions with varying $M_{Z'}$ for fixed M_{A^0} are shown in Fig. 54, while the dependency on M_{A^0} is shown in Fig. 55.

This model also allows for an additional source of Higgs plus \cancel{E}_T signal with a
1170 similar kinematics (Fig. 56, shown with detector simulation samples) to the signal process from the decay of $Z' \rightarrow hZ$, where the Z decays invisibly. The partial decay width for the Z' is:

$$\Gamma_{Z' \rightarrow hZ} = (g_z \cos \alpha \sin \beta)^2 \frac{|p|}{24\pi} \left(\frac{|p|^2}{M_{Z'}^2} + 3 \frac{M_Z^2}{M_{Z'}^2} \right). \quad (38)$$


The values for the Z' masses scanned for those samples should follow those of the previous samples, namely values of $M_{Z'} = 600, 800, 1000, 1200, 1400$ GeV. This
1175 signal process has no M_A dependence.

3.2. EFT models with direct DM-boson couplings


The EFT operators considered in this section do not have an implementation of a simplified model completion for Dirac fermion Dark Matter available to date. They provide kinematic distributions that are unique to mono-boson signatures, and that
1180 in most cases are not reproduced by an equivalent simplified model.¹¹

A complete list of effective operators with direct DM/boson couplings for Dirac DM, up to dimension 7, can be found in [97, 45, 98]. Higher dimensional operators, up to dimension 8, leading to Higgs+ \cancel{E}_T signatures, are mentioned in [45, 96]. The first part of this Section outlines the main characteristics for a limited number
1185 of these models that could be considered in early Run-2 searches. However, the EFT approximation made for these operators can be problematic, see Ref. [96] for


¹¹Wherever this is the case, for practical reasons one can only generation a simplified model result in the limiting EFT case, as the results can be rescaled and reinterpreted.



(a) \cancel{E}_T distribution




(b) Leading b -jet p_T distribution




(c) $\Delta\phi$ distance between the two b -jets


Figure 54: Kinematic distributions of the signal process varying $M_{Z'}$, for $m_\chi = 100$ GeV, $M_{A^0} = 300$ GeV.



(a) \cancel{E}_T distribution




(b) Leading b -jet p_T distribution




(c) $\Delta\phi$ distance between the two b -jets


Figure 55: Kinematic distributions of the signal process varying M_{A^0} , for $m_\chi = 100$ GeV, $M_{Z'} = 1000$ GeV.



(a) \cancel{E}_T distribution



(b) Leading b -jet p_T distribution



(c) $\Delta\phi$ distance between the two b -jets

Figure 56: Kinematic distributions of $Z' \rightarrow A^0 h$ exclusive production, $Z' \rightarrow Z h$ exclusive production and Z' inclusive production for $M_{Z'} = 1000$ GeV and $M_{A^0} = 300$ GeV

discussion. For this reason, model-independent results as in Appendix B should be privileged over considering these operators as realistic benchmarks.

However, the Forum discussion highlighted that the EFT approach allows more
1190 model-independence when reinterpreting results, and that it is worth still considering interpretation of the results available in terms of these operators. Furthermore, once simplified models are available for those operators, EFT results can be used as a limiting case for consistency checks. We devote the end of this Section to a discussion on the presentation of results from this model, including an assessment
1195 of their reliability using a conservative procedure that is only dependent on EFT parameters.

The studies in this Section have been performed using a UFO model within MADGRAPH5_AMC@NLO v2.2.3, interfaced to PYTHIA 8 for the parton shower. The implementation of these models is discussed further in Section 4.2.2.

1200 3.2.1. Dimension 5 operators

The lowest dimension benchmark operators we consider are effective dimension 5, such as the one depicted in Figure 57.

Following the notation of [45], models from this category have a Lagrangian that, after electroweak symmetry breaking, includes terms such as:

$$\frac{m_W^2}{\Lambda_5^3} \bar{\chi} \chi W^{+\mu} W_\mu^- + \frac{m_Z^2}{2\Lambda_5^3} \bar{\chi} \chi Z^\mu Z_\mu, \quad (39)$$

1205 where m_Z and m_W are the masses of the Z and W boson, W^μ and Z^μ are the fields of the gauge bosons, χ denotes the Dark Matter fields and Λ_5 is the effective field theory scale. Note that these operators are of true dimension 7, but reduce to effective dimension 5 once the Higgs vacuum expectation values, contained in the W and Z mass terms, are inserted. As such, one expects that these operators would
1210 naturally arise in UV complete models where Dark Matter interacts via a Higgs portal where heavy mediators couple to the Higgs or other fields in an extended Higgs sector. In such models the full theory may be expected to contain additional operators with Higgs-Dark Matter couplings [102]. The above operator also induces




Figure 57: Diagram for EFT operators giving rise to a Higgs+ \cancel{E}_T signature.

signatures with \cancel{E}_T in conjunction with Z and W bosons at tree level, as shown in
 1215 Fig. 38, while at loop level it induces couplings to photon pairs and $Z\gamma$ through W
 loops. In these models, a clear relation exists between final states with photons,
 EW bosons and Higgs boson.

As shown in Fig. 58, the kinematics of this model can be approximated by
 that of a simplified model including a high-mass scalar mediator exchanged in the
 1220 s-channel described in Section 2.2.2. For this reason, the list of benchmark models
 with direct boson-DM couplings for photon, Z and W only includes dimension 7
 operators: if the scalar model with initial state radiation of an EW boson is already
 generated, then its results can be rescaled.

The Higgs+ \cancel{E}_T analysis, however, will not consider the scalar simplified model
 1225 as benchmark, due to the very low sensitivity in early LHC analyses, and will instead
 use this dimension 5 operator.

Parameter scan. The two parameters of this model are the scale of new physics λ
 and the DM particle mass. SM-DM coupling and new physics scale are related by




Figure 58: Comparison of the missing transverse momentum for the simplified model where a scalar mediator is exchanged in the s-channel and the model including a dimension-5 scalar contact operator, in the leptonic $Z+\cancel{E}_T$ final state. All figures in this Section have been performed using a UFO model within MADGRAPH5_AMC@NLO v2.2.3, interfaced to PYTHIA 8 for the parton shower.

$$g_\chi = (246 \text{ GeV})/\lambda.$$

1230 The initial value of the new physics scale λ chosen for the sample generation is 3 TeV. This is a convention and does not affect the signal kinematics: the cross-section of the samples can be rescaled when deriving the constraints on this scale. However, more care should be given when rescaling Higgs+ \cancel{E}_T operators of higher dimensions, as different diagrams have a different λ dependence.

The DM mass values for the benchmark points to be simulated are chosen to span a sufficient range leading to different kinematics, that is within the LHC sensitivity for early searches and that is consistent across the various signatures and EFT operators. We therefore start the mass scan at $m_\chi = 1 \text{ GeV}$, where collider experiments are complementary to direct and indirect detection and choose the last point corresponding to a DM mass of 1.3 TeV. We recommend a scan in the following mass points:

$$m_\chi = 1, 10, 50, 100, 200, 400, 800, 1300 \text{ GeV}.$$

1235 A set of kinematic distributions from the Higgs+ \cancel{E}_T signature where the Higgs decays into two b -quarks is shown in Fig. 59, for points similar to those of the grid scan proposed.

3.2.2. Dimension 7 operators

The dimension-7 benchmark models contain the $SU(2)_L \times U(1)_Y$ gauge-invariant couplings between DM fields and the kinetic terms of the EW bosons. The CP-conserving scalar couplings of this type can be written as


$$\frac{c_1}{\Lambda_S^3} \bar{\chi} \chi B_{\mu\nu} B^{\mu\nu} + \frac{c_2}{\Lambda_S^3} \bar{\chi} \chi W_{\mu\nu}^i W^{i,\mu\nu}. \quad (40)$$

Here $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$ and $W_{\mu\nu}^i = \partial_\mu W_\nu^i - \partial_\nu W_\mu^i + g_2 \epsilon^{ijk} W_\mu^j W_\nu^k$ are the $U(1)_Y$ and $SU(2)_L$ field strength tensor, respectively, and g_2 denotes the weak coupling constant. In the case of the pseudoscalar couplings, one has instead


$$\frac{c_1}{\Lambda_P^3} \bar{\chi} \gamma_5 \chi B_{\mu\nu} \tilde{B}^{\mu\nu} + \frac{c_2}{\Lambda_P^3} \bar{\chi} \gamma_5 \chi W_{\mu\nu}^i \tilde{W}^{i,\mu\nu}, \quad (41)$$

where $\tilde{B}_{\mu\nu} = 1/2 \epsilon_{\mu\nu\lambda\rho} B^{\lambda\rho}$ and $\tilde{W}_{\mu\nu}^i = 1/2 \epsilon_{\mu\nu\lambda\rho} W^{i,\lambda\rho}$ are the dual field strength


1240 tensors. In addition to the CP-conserving interactions (40) and (41), there are also



(a) Leading b -jet transverse momentum



(b) Leading b -jet pseudorapidity



(c) Angular distance between the two leading b -jets

Figure 59: Comparison of the kinematic distributions for the two leading b -jets (from the Higgs decay) in the model with direct interactions between the Higgs boson and the DM particle, when varying the DM mass.

four CP-violating couplings that are obtained from the above operators by the replacement $\tilde{\chi}\chi \leftrightarrow \tilde{\chi}\gamma_5\chi$.

The effective interactions introduced in (40) and (41) appear in models of Rayleigh DM [103]. Ultraviolet completions where the operators are generated through loops of states charged under $U(1)_Y$ and/or $SU(2)_L$ have been proposed in [104] and their LHC signatures have been studied in [105]. If these new charged particles are light, the high- p_T gauge bosons that participate in the \cancel{E}_T processes considered here are able to resolve the substructure of the loops. This generically suppresses the cross sections compared to the EFT predictions [106], and thus will weaken the bounds on the interaction strengths of DM and the EW gauge bosons to some extent. Furthermore, the light charged mediators may be produced on-shell in pp collisions, rendering direct LHC searches potentially more restrictive than \cancel{E}_T searches. Making the above statements precise would require further studies beyond the timescale of this forum.

Since for $\Lambda_S = \Lambda_P$ the effective interactions (40) and (41) predict essentially the same value of the mono-photon, mono- Z and mono- W cross section [45, 98], we consider below only the former couplings. We emphasize however that measurements of the jet-jet azimuthal angle difference in $\cancel{E}_T + 2j$ events may be used to disentangle whether DM couples more strongly to the combination $B_{\mu\nu}B^{\mu\nu}(W_{\mu\nu}^iW^{i,\mu\nu})$ or the product $B_{\mu\nu}\tilde{B}^{\mu\nu}(W_{\mu\nu}^i\tilde{W}^{i,\mu\nu})$ of field strength tensors [97, 98].

After EW symmetry breaking the interactions (40) induce direct couplings between pairs of DM particles and gauge bosons. The corresponding Feynman rule reads:

$$\frac{4i}{\Lambda_S^3} g_{V_1V_2} (p_1^{\mu_2} p_2^{\mu_1} - g^{\mu_1\mu_2} p_1 \cdot p_2), \quad (42)$$

where p_i (μ_i) denotes the momentum (Lorentz index) of the vector field V_i and for simplicity the spinors associated with the DM fields have been dropped. The

couplings $g_{V_i V_j}$ take the form:

$$\begin{aligned}
g_{\gamma\gamma} &= c_w^2 c_1 + s_w^2 c_2, \\
g_{\gamma Z} &= -s_w c_w (c_1 - c_2), \\
g_{ZZ} &= s_w^2 c_1 + c_w^2 c_2, \\
g_{WW} &= c_2,
\end{aligned} \tag{43}$$

with s_w (c_w) the sine (cosine) of the weak mixing angle. Note that our coefficients c_1 and c_2 are identical to the coefficients C_B and C_W used in [98], while they are related via $k_1 = c_w^2 c_1$ and $k_2 = s_w^2 c_2$ to the coefficients k_1 and k_2 introduced in [45].

The coefficients c_1 and c_2 appearing in (43) determine the relative importance of each of the \cancel{E}_T channels and their correlations. For example, one observes that:

- Only c_2 enters the coupling between DM and W bosons, meaning that only models with $c_2 \neq 0$ predict a mono- W signal;
- If $c_1 = c_2$ the mono-photon (mono- Z) signal does not receive contributions from diagrams involving Z (photon) exchange;
- Since numerically $c_w^2/s_w^2 \simeq 3.3$ the mono-photon channel is particularly sensitive to c_1 .

Parameter scan. As stated above and shown in Ref. [95], the kinematic distributions for dimension-7 scalar and pseudoscalar operators only shows small differences. This has been verified from a generator-level study: the signal acceptance after a simplified analysis selection ($\cancel{E}_T > 350$ GeV, leading jet $p_T > 40$ GeV, minimum azimuthal difference between either of the two jets and the \cancel{E}_T direction > 0.4) is roughly 70% for both models, independent from the coefficients c_1 and c_2 . We therefore only suggest to generate one of the two models.

The differences in kinematics for the various signatures are negligible when changing the coefficients c_1 and c_2 , since these coefficient factorize in the matrix

element. Only the case $c_1 = c_2 = 1$ is generated as benchmark; other cases are left for reinterpretation as they will only need a rescaling of the cross-sections.




Figure 60: E_T distribution for the dimension-7 model with a hadronically decaying Z in the final state, for the scalar and pseudoscalar operators representing direct interactions between DM and bosons. The values of the coefficients in the legend are multiplied by 100.

1290 3.2.3. Higher dimensional operators


Many higher dimensional operators can induce signals of photons or $W/Z/H$ bosons in the final state. A complete list can be found in Refs. [48, 96, 47] and references therein.

1295 Although with lower priority with respect to the operators above, a representative dimension-8 operators can be chosen as benchmark, with the form:


$$\frac{1}{\Lambda^4} \bar{\chi} \gamma^\mu \chi B_{\mu\nu} H^\dagger D^\nu H$$

In this case, the new physics scale is Λ is related to the coupling of the DM as $y_\chi = \frac{1}{\Lambda^4}$. An advantage of this operator is that it includes all signatures with EW

1300 bosons, allowing to assess the relative sensitivity of the various channels with the same model. The kinematics for this operator is different with respect to other operators, leading to a harder \cancel{E}_T spectrum, as illustrated by comparing the leading b -jet distribution for the dimension 5 operator to the dimension 8 operator.



(a) Dimension 5 operator



(b) Dimension 7 operator

Figure 61: Comparison of the transverse momentum for the leading b -jet from the Higgs decay for a dimension 5 and dimension 7 operator with direct boson-DM couplings.

3.2.4. Validity of EW contact operators and possible completions

It is important to remember that the operators described in this section may
1305 present problems in terms of the validity of the contact interaction approach for the
energy scales reached at the LHC.

As outlined in [96], designing very high \cancel{E}_T search signal regions that are ex-
clusively motivated by the hard \cancel{E}_T spectra of the dimension 7 and 8 operators will
mean that the momentum transfer in the selected events is larger. This in turn
1310 means that processes at that energy scale (mediators, particles exchanged in loops)
are accessible, and a simple contact interaction will not be able to correctly describe
the kinematics of these signals.

Contact interaction operators like the ones in this section remain useful tools for
comparison of the sensitivity of different search channels, and for reinterpretation of
1315 other models under the correct assumptions. To date, while UV-complete models are
known, their phenomenology has not been studied in full detail as their completion
involves loops.¹²

However, this may be the focus of future theoretical exploration, as discussed
in Ref. [98]. An example of a complete model for scalar DM corresponding to the
1320 dimension-5 operator is provided in the Appendix A. Providing results for the pure
EFT limit of these models will prove useful to cross-check the implementation of
future.

Given these considerations, we recommend to present results for these models
as follows:

- 1325 • Deliver fiducial limits on the cross section of any new physics events, without
any model assumption, according to the guidelines in Appendix B.

¹² An example case for the need of loop completions is a simplified model with an additional scalar
exchanged at tree level. The scalar couples to WW and ZZ in a gauge-invariant way. Integrating out
the mediator does not lead to the Lorentz structure of a dimension-7 operator, so it is not possible to
generate dimension-7 operators that satisfy gauge and Lorentz invariance at the same time. A model
with a spin-1 mediator cannot be considered as an candidate for completion either, since dimension-7
operators only have scalar or pseudoscalar couplings.

- Assess the percentage of events that pass a condition of validity for the EFT approximation that does not depend on a specific completion, and present results removing of the invalid events using the procedure in Section 5 alongside the raw EFT results.

4. Implementation of Models

4.1. Implementation of s -channel and t -channel models for $\cancel{E}_T + X$ analyses

In the studies to date, a number of different Monte Carlo tools have been used to simulate DM signals. In this Chapter, we make recommendations on the accuracy at which simulations should be performed for different final states. We also provide explicit examples of codes and implementations (including specific settings) that have been used to obtain the results in this report. We stress that these recommendations are based on the current status of publicly available codes and users should always check whether new results at a better accuracy have appeared in the meantime. We recommend to update the corresponding analyses directly using the new releases and/or codes respectively, and in case this would not be possible, to at least take into account the new information in the analysis (e.g., via a MC comparison with the latest predictions, or by effectively using global/local K -factors). For all models included in this report, PYTHIA 8 has been used to provide the parton shower simulation. Nevertheless, we note that showering matrix element events with Herwig [107, 108, 109, 110] should be considered as an equally valid alternative.

4.1.1. Implementation of s -channel models for mono-jet signature

These models include those discussed in Secs. 2.1 and 2.2. In monojet analyses, i.e. when final states are selected with a few jets and \cancel{E}_T , observables and in particular the \cancel{E}_T spectrum depend upon the accuracy of the simulation of QCD radiation. For the vector and axial vector models, the current state of the art is NLO+PS. It is particularly simple to obtain simulations for these processes at NLO+PS and even for merged samples at NLO accuracy, starting from SM implementations. We

1355 therefore recommend simulations to be performed at NLO+PS, and in case multi-jet
observables are employed, by merging samples with different multiplicities. Results
at such accuracy can be obtained either in dedicated implementations, such as that
of POWHEG [59], or via general purpose NLO tools like MADGRAPH5_AMC@NLO
employing available UFO models at NLO. A testing version of the full set of these
1360 UFO models has been made available only in June 2015 [111]. For this reason, it
was not used as part of the studies of this Forum on initial Run-2 benchmark mod-
els. Nevertheless, we encourage further study of these UFO models by the ATLAS
and CMS collaborations.

A study using POWHEG [59, 112] has shown that the NLO corrections result
1365 in a substantial reduction in the dependence on the choice of the renormalization
and factorization scales and hence a reduced theoretical uncertainty on the signal
prediction. For the central choice of renormalization and factorization scales, the
NLO corrections also provide a minor enhancement in the cross section due to the
jet veto that has been so far employed in Run-1 analyses.

1370 For the scalar and pseudoscalar models, the lowest order process already in-
volves a one-loop amplitude in QCD. Because of the complexity of performing NLO
calculations for this class of processes and in particular the absence of general meth-
ods for computing two-loop virtual contributions, only LO predictions are currently
available. These can be interfaced to shower programs exactly as usual tree-level
1375 Born computations, i.e. by considering one parton multiplicity at a time or by merg-
ing different parton multiplicities via CKKW or MLM schemes to generate inclu-
sive samples with jet rates at LO accuracy. For spin-0 mediators in the mono-jet
final state, the top-quark loop is the most important consideration. The matrix
element implementation with exact top-loop dependence of the s -channel spin-0
1380 mediated DM production is available in MCFM [112, 41]¹³ at fixed order and in
POWHEG [43] and MADGRAPH5_AMC@NLO [111] for event generation at LO+PS
level. The POWHEG and MCFM implementations include the finite top quark mass
dependence for DM pair production and one extra parton at LO. The same processes

¹³Only the scalar mediator is available in the public release.

are available in MADGRAPH5_AMC@NLO v2.3 and could be made available in the
1385 future in codes like SHERPA+OPENLOOPS/GOSAM, including up to two extra partons
in the final state. Samples can be merged employing CKKW, K_T -MLM procedures.

Most of the results that have been presented in this document for these processes
have been obtained with POWHEG interfaced to PYTHIA 8, matching the state of
the art calculation as of Spring 2015. For future reference, we document the spe-
1390 cific settings needed to run the POWHEG generation for the dark matter models
so they can serve as nominal benchmarks for the early Run-2 ATLAS and CMS DM
analyses. POWHEG parameter cards for all models can be found on the Forum SVN
repository [113, 114, 115, 116].

POWHEG *configuration for s-channel DM models*. The latest POWHEG release is
1395 available for download using the instructions at <http://powhegbox.mib.infn.it/>.
The Forum recommends using at least version 3059.

- POWHEG can generate either unweighted (uniformly-weighted) or weighted events. The relevant keywords in the input card are `bornsuppfact` and `bornktmin`.

1400 1. unweighted events:

`bornsuppfact`: negative or absent

`bornktmin` PT

This runs the program in the most straightforward way, but it is likely
not the more convenient choice, as will be explained below. POWHEG
1405 will generate unweighted events using a sharp lower cut (with value
PT) on the leading-jet p_T . Since this is a generation cut, the user must
check that the choice of `bornktmin` does not change the cross section
for signal events passing analysis selections. It is a good practice to
use 10-20% smaller E_T value in the input card than the final analysis
1410 selection, and check that the final result is independent, by exploring an
even smaller value of `bornktmin`. The drawback of using this mode is
that it is difficult to populate well, and in a single run, both the low- p_T
region as well as the high- p_T tail.

2. weighted events:

1415

`bornsuppfact` PTS

`bornktmin` PT

POWHEG will now produce weighted events, thereby allowing to generate a single sample that provides sufficient statistics in all signal regions.

1420

Events are still generated with a sharp lower cut set by `bornktmin`, but the `bornsuppfact` parameter is used to set the event suppression factor according to

$$F(k_T) = \frac{k_T^2}{k_T^2 + \text{bornsuppfact}^2} . \quad (44)$$

In this way, the events at, for instance, low \cancel{E}_T , are suppressed but receive higher weight, which ensures at the same time higher statistics at high \cancel{E}_T . We recommend to set `bornsuppfact` to 1000.

1425

The `bornktmin` parameter can be used in conjunction with `bornsuppfact` to suppress the low \cancel{E}_T region even further. It is recommended to set `bornktmin` to one-half the value of the lowest \cancel{E}_T selection. For instance, for the event selection used in the CMS/ATLAS monojet analyses, assuming the lowest \cancel{E}_T region being defined above 300 GeV, the proposed value for `bornktmin` is 150. However, this parameter should be set keeping in mind the event selection of all the analyses that will use these signal samples, and hence a threshold lower than 150 may be required.

1430

- The POWHEG monojet implementations can generate events using two expressions for the mediator propagators. The default setup (i.e if the keyword `runningwidth` is absent, commented out or set to 0) is such that a normal Breit-Wigner function is used for the propagator: in this case, the expression

$$Q^2 - M^2 + i M \Gamma$$

is used for the propagator's denominator, where Q is the virtuality of the mediator, and M and Γ are its mass and width, respectively. This is the more

straightforward, simple and transparent option, and it was used for the Forum studies. It should be the method of choice, unless one approaches regions of parameter space where Γ/M starts to approach order 1 values. In those cases, a more accurate modelling (or at least a check of the validity of the fixed width approach) can be achieved by using a running width: by setting the `runningwidth` token to 1, POWHEG uses as the denominator of the mediator propagator the expression

$$Q^2 - M^2 + iQ^2 \frac{\Gamma}{M},$$

which is known to give a more realistic description. See Ref. [94] for a discussion.

1435

- Set the parameters defining the bounds on the invariant mass of the dark matter pair, `mass_low` and `mass_high`, to -1. In this way, POWHEG will assign values internally.

1440

- The minimal values for `ncall1`, `itmx1`, `ncall2`, `itmx2` are 250000, 5, 1000000, 5 for the vector model, respectively.

1445

- The minimal values for `ncall1`, `itmx1`, `ncall2`, `itmx2` are 100000, 5, 100000, 5 for the scalar top-loop model, respectively.
- When NLO corrections are included (as for instance in the vector model), negative-weighted events could occur and should be kept in the event sample, hence `withnegweights` should be set to 1. If needed, their fraction can be decreased by setting `foldsci` and `foldy` to larger values (2 for instance). `foldphi` can be kept to 1.

1450

- One should use the automatic calculation of systematic uncertainties associated with the choice of hard scale and PDFs as described in Section 6.
- `idDM` is the integer that identifies the DM particle in the Monte Carlo event record. This should be chosen so that other tools can process the POWHEG output properly.

POWHEG in itself is not an event generator and must be interfaced with a tool that provides parton showering, hadronization, *etc.* For some time, a PYTHIA 8 [117] interface has existed for POWHEG. The PYTHIA 8 runtime configuration is the following:

```
POWHEG:veto = 1
POWHEG:pTdef = 1
POWHEG:emitted = 0
1455 POWHEG:pTemt = 0
POWHEG:pThard = 0
POWHEG:vetoCount = 100
SpaceShower:pTmaxMatch = 2
TimeShower:pTmaxMatch = 2
```

1465 As always, it is recommended to use the latest PYTHIA 8 release, available at <http://home.thep.lu.se/~torbjorn/Pythia.html>. At the time of this report, the latest version is 8.209.

4.1.2. Merging samples with different parton multiplicities

For the models discussed in the previous section, it is important to calculate the hard process as accurately as possible in QCD. For many other signal models, the \cancel{E}_T signature depends more upon the production and decay of the mediator. In some cases, observables built in terms of the jets present in the final state are considered, relying on the assumption that inclusive samples accurate in higher jet multiplicities are available. In these cases, one can employ LO+PS simulations where different parton multiplicities are merged and then matched to parton shower, using schemes such as CKKW or MLM merging.

Here, we consider the example of an EFT model produced in association with up to 2 additional QCD partons. A Monte Carlo sample based on this method could be used in alternative to a NLO+PS sample for describing shapes and jet distributions (but not for the overall normalisation which would still be at LO). The methodology described here could also be used for the t -channel model discussed in Sec. 2.3.

For the calculation of tree-level merged samples for DM signals, tools that can read UFO files and implement multi-parton merging should be employed, such that MADGRAPH5_AMC@NLO (+PYTHIA 8 or HERWIG++) and SHERPA [118]. In this report we have mostly employed MADGRAPH5_AMC@NLO. MADGRAPH5_AMC@NLO provides a flexible and easy-to-use framework for implementing new models via the FEYNRULES package. MADGRAPH5_AMC@NLO can perform both LO and NLO calculations in QCD, matched/merged to parton showers [81]. For NLO ones, dedicated UFO model implementations at NLO should be used. Several UFO models at NLO are publicly available that although not developed specifically for DM, are suitable to make mode independent simulations at NLO accuracy, including multi-parton merging via the FxFx technique [119]. A dedicated DM UFO implementation has been developed and it has been released as a testing version [111].

Merging events generated via matrix elements with different number of partons in the final state can be achieved by a judicious procedure that avoids double counting of the partons from matrix elements and parton showering. Several merging techniques are available. Based on some comparative studies [120], there is some advantage to using the CKKW-L merging scheme [121] implemented in PYTHIA 8. Alternatively, one can use the k_T -MLM scheme also available in PYTHIA 8.

Generation of the LHE file. The example presented here is a D5 EFT model, and includes tree-level diagrams with $\chi \bar{\chi} + 0, 1, 2$ partons. We stress that MADGRAPH5_AMC@NLO, like POWHEG, is not in itself an event generator, but must be interfaced with an event generator through an LHE file. The production of the LHE file proceeds through setting the process parameters and the run parameters.

The process parameters are:

```
import model MODELNAME
generate p p > chi chi~ [QCD] @0
add process p p > chi chi~ j [QCD] @1
add process p p > chi chi~ j j [QCD] @2
```

The runtime parameters are more numerous, and define the collider properties,

PDF sets, etc. The specific parameters needed for matching are, for the example of CKKW-L matching:

```

    ickkw = 0
    ktdurham = matching scale
1515 dparameter = 0.4
    dokt = T
    ptj=20
    drjj=0
    mmjj=0
1520 ptj1min=0

```

For different kinds of matching, a different choice of ickkw and related parameters would be made.

Implementation of the CKKW-L merging. To illustrate the settings related to merging different multiplicities, the EFT D5 samples were generated with MADGRAPH5_AMC@NLO
1525 version 2.2.2 and showered in PYTHIA 8.201, using the Madgraph parameters in the previous section (Sec. 4.1.2).

The PYTHIA 8 parameters for the CKKW-L k_T -merging scheme are:

```

Merging:ktType          = 1
Merging:TMS              = matching scale
1530 1000022:all = chi chi~ 2 0 0 30.0 0.0 0.0 0.0 0.0
    1000022:isVisible = false
Merging:doKTMerging      = on
Merging:Process           = pp>{chi,1000022}{chi~, -1000022}
Merging:nJetMax           = 2

```

1535 The matching scales should be the same for the generation and parton showering. In the model implementation, the particle data group ID 1000022 is used for weakly interacting dark matter candidates. Since this is a Majorana particle by default (with no corresponding anti-particle), and the model produces a DM Dirac fermion, the particle properties are changed accordingly. Also, the DM mass is set

1540 to 30 GeV. The `Merging:Process` command specifies the lowest parton emission process generated in `MADGRAPH5_AMC@NLO` and `Merging:nJetMax = 2` gives the maximum number of additional parton emissions with respect to the lowest parton emission process.

In general, it is desired to take the hard parton emissions from the matrix element generation in `MADGRAPH5_AMC@NLO` and allow `PYTHIA 8` to take care of soft emissions only. The transition between these two regimes is defined by the matching scale and its optimal value can be determined by studying the cross-section as a function of the number of jets (differential jet rates). The differential rates $\frac{dN_{i \rightarrow j}}{d \log_{10}(k_{\text{cut}})}$ give the number of events which pass from i jets to j jets as the k_T value increases
 1545 beyond k_{cut} . An optimal matching scale should lead to smooth differential jet rates.

Two examples of differential jet rates, using matching scale 30 GeV and 80 GeV, from the EFT D5 sample generated as described in the previous section are given in Fig. 62 and 63, respectively. Although a kink is visible around the matching scale value in both cases, the 80 GeV scale leads to smoother distributions. In order to
 1550 find the optimal matching scale, additional samples with matching scale 50, 70, and 90 GeV are generated as well and a detailed comparison of the differential jet rates close to the transition region is shown in Fig. 64. The largest differences among the samples are visible for the $1 \rightarrow 2$ jets transition where the 30 GeV and 50 GeV scale lead to a drop of the rates around the matching scale values. On the contrary,
 1555 there is a hint of an increased rate around the matching scale value in the sample generated with the 90 GeV scale. Therefore, we recommend to use 80 GeV as the baseline matching scale.

The prescription for the event generation given in Section 4.1.2 starts with the emission of 0 partons and ends with maxim 2 partons in addition. Producing the
 1560 samples separately allows to investigate the relative composition of the individual samples in various parts of the phase space. Figure 65 shows the \cancel{E}_T distribution of the EFT D5 sample with the matching scale at 80 GeV. The plot reveals that the 0-parton sample gives the dominant contribution in the region below the matching scale value that rapidly decreases at higher \cancel{E}_T . Assuming the lowest analysis \cancel{E}_T
 1570 cut in early Run-2 mono-jet analyses at 300 GeV, the generation of the 0-parton




Figure 62: Distributions of differential jet rates $\frac{dN_{i \rightarrow j}}{d \log_{10}(k_{\text{cut}})}$ for EFT D5 sample with CKKW-L matching scale at 30 GeV. The 0-, 1- and 2-parton emission samples are generated separately and indicated in the plots as Prod 1, Prod 2 and Prod 3, respectively. A vertical line is drawn at the matching scale.

emission sample can be safely omitted as it only gives $< 1\%$ contribution at $\cancel{E}_T > 300$ GeV. For the 1- and 2-parton emission samples, one can use a generator cut on the leading parton p_T , ptj1min , in order to avoid generating low \cancel{E}_T events that are irrelevant for the analysis.

1575 In order to describe the signal kinematics correctly and save time during MC production, the parton emissions will only be generated up to a certain multiplicity. The higher multiplicity samples usually have small enough cross sections and the corresponding parts of the phase space can be sufficiently approximated by parton showering in PYTHIA 8. A dedicated study comparing samples generated with up to 1-, 2-, or 3-parton multiplicities was performed, using again the settings for the
 1580 CKKW-L k_T -merging with the 80 GeV matching scale and the `Merging:nJetMax` parameter adjusted accordingly. Figure 66 shows the \cancel{E}_T distribution of the samples




Figure 63: Distributions of differential jet rates $\frac{dN_{i \rightarrow j}}{d \log_{10}(k_{\text{cut}})}$ for EFT D5 sample with CKKW-L matching scale at 80 GeV. The 0-, 1- and 2-parton emission samples are generated separately and indicated in the plots as Prod 1, Prod 2 and Prod 3, respectively. A vertical line is drawn at the matching scale.

at $\cancel{E}_T > 250$ GeV.

With an event selection requiring \cancel{E}_T and the leading jet p_T being larger than
1585 250 GeV, the sample generated with up to 1 parton has 10.3% larger yield compared to the sample with up to 3 partons, while the yield of the sample with up to 2 partons is only 2.3% larger. If an additional cut is applied allowing for up to 3 jets with $p_T > 30$ GeV, the agreement improves to 3.2% larger for up to 1 parton and 0.7% larger for up to 2 partons, compared with up to 3 partons.

1590 A similar comparison is shown in Fig. 67 for the jet multiplicity in the events with the leading jet $p_T > 250$ GeV, where an agreement at the level of $\sim 3\%$ between the samples with up to 2 and 3 parton emissions is observed for number of jets up to 7. This justifies it is sufficient to produce samples with up to 2 parton emissions only at the generator level and ignore generating higher parton emissions.

1595 *4.1.3. Implementation of t -channel models for the $jet+\cancel{E}_T$ final state*

The simulations for t -channel models are available via LO UFO implementations, where events are generated at LO+PS accuracy. The UFO file and parameter cards for the t -channel models with couplings to light quarks only [76] can be found on the Forum SVN repository [122]. The model files from Ref. [46] can also
1600 be found on the repository [123]. The latter is the implementation that has been used for the studies in this report: in the monojet case there are only cross section differences between this model and the model in [122].

Multi-parton simulation and merging are necessary and require particular care for this model: this has not been a topic of detailed studies within the Forum, and
1605 we suggest to follow the procedure outlined in Ref. [76].

4.1.4. Implementation of s -channel and t -channel models with EW bosons in the final state

Currently, simulations for most of these models are available via LO UFO implementations, allowing event generation at the LO+PS accuracy. We note, however,
1610 that inclusion of NLO corrections would be possible. In MADGRAPH5_AMC@NLO, for example, this amounts to simply upgrading the currently employed UFO models to NLO, where the calculations exist for this class of processes. However, this was not available within the timescale of the Forum towards simulation of early Run-2 benchmarks. As a consequence, in this work we have used LO UFO implementa-
1615 tions within MADGRAPH5_AMC@NLO 2.2.3 interfaced to PYTHIA 8 for the parton shower. The corresponding parameter cards used for the Run-2 benchmark models can be found on the Forum SVN repository [124]. This is the implementation that will be used for early Run-2 LHC Dark Matter searches.

None of these models requires merging samples with different parton multiplic-
1620 ities since the visible signal comes from the production of a heavy SM boson whose transverse momentum distribution is sufficiently well described at LO+PS level. As a result, no special runtime configuration is needed for PYTHIA 8.

4.1.5. Implementation of s -channel and t -channel models with heavy flavor quark signatures

1625 Dedicated implementations for DM signals in this final state are available at LO+PS accuracy. However, the state of the art of the simulations for $t\bar{t}$ and $b\bar{b}$ with a generic scalar and vector mediator is NLO+PS accuracy. For example, simulations for $t\bar{t}$ + scalar can be obtained via POWHEG and SHERPA starting from the SM implementations. In MADGRAPH5_AMC@NLO, all final relevant final states, spin-0
1630 (scalar and pseudo scalar) and spin-1, (vector and axial) are available at NLO+PS via the dedicated NLO UFO for DM has been released in June 2015 [111]).

Within the scope of the Forum, simulations for the $t\bar{t}$ and $b\bar{b}$ signatures of the scalar mediator model have been generated starting from a leading order UFO with MADGRAPH5_AMC@NLO 2.2.2, using PYTHIA 8 for the parton shower. The UFO file
1635 and parameter cards that will be used as benchmarks for early Run-2 searches in these final states can be found on the Forum SVN repository [125]. Multi-parton merging has been used for the $b\bar{b}$ case but it has not been studied in detail within this Forum. The b-flavored DM model of Section 2.3.2 is simulated at LO+PS using MADGRAPH5_AMC@NLO v2.2.3 and PYTHIA 8 for the parton shower. The corre-
1640 sponding UFO and parameter files can be found on the Forum SVN repository [126].

Quark flavor scheme and masses. In the case of $b\bar{b}$ final state an additional care should be taken when choosing the flavor scheme generation and whether quarks should be treated as massive or massless.

The production of DM+ $b\bar{b}$, dark matter in association with b jets via a decay of
1645 a (pseudo) scalar boson, is dominated in simplified mediator models by the gluon-gluon initiated production, similar to the production of $Z+b\bar{b}$ at the LHC. The $Z+b\bar{b}$ process has been studied in detail in the $Z(\text{ll})+b$ -jets final state, which can be used to validate both the modelling of DM+bb and, its main background, $Z(\nu\nu)+b\bar{b}$. In this context, the p_T of the Z boson is related to the observed MET, whereas the b -jet
1650 kinematics determines the ratio of mono- b /di- b signatures in the detector.

For basic kinematic criteria applied to $Z+b\bar{b}$ production, this process leads in $\sim 90\%$ of the events to a signature with only 1 b -jet in the acceptance ('Z+1b-

jet production') and only in $\sim 10\%$ of the events to a signature with 2 b -jets in the detector ('Z+2 b -jets production'). The production cross section of the Z+ $b\bar{b}$ process
1655 can be calculated in the 'five-flavor scheme', where b quarks are assumed massless, and the 'four-flavor scheme', where massive b quarks are used [127, 128, 129]. Data slightly favour the cross-section predictions in the five-flavor scheme [130] for the 1 b -jet signature. In this document we have preferred the 5-flavor scheme due to its simplicity and cross section agreement within the SM. The PDF used to
1660 calculate these cross section is NNPDF3.0 (lhaid 263000).

On the other hand, both data [130, 131, 132] and theoretical studies [133, 134] suggest that the best modelling of an inclusive Z+ $b\bar{b}$ sample especially for what concerns b -quark observables, is achieved at NLO+PS using a 4-flavor scheme and a massive treatment of the b -quarks. In Figure 68 we show that, at LO, as expected,
1665 no appreciable difference is visible in the kinematics between either flavor scheme used for DM+ $b\bar{b}$. In our generation we have used NNPDF3.0 set (lhaid 263400).

4.2. Implementation of specific models for $V + \cancel{E}_T$ analyses

4.2.1. Model implementation for mono-Higgs models

Currently, simulations for most of these models are available via LO UFO imple-
1670 mentations, allowing event generation at the LO+PS accuracy. We note, however, that the inclusion of NLO corrections would be possible but not available in time for the conclusion of these studies. In MADGRAPH5_AMC@NLO, for example, this amounts to simply upgrading the currently employed UFO models to NLO. Simulation of loop-induced associated production of DM and Higgs is also possible with
1675 the exact top-quark mass dependence. In MADGRAPH5_AMC@NLO, for example, this can be obtained from the NLO UFO SM and 2HDM implementations.

In this work all three Higgs+ \cancel{E}_T models have been generated at leading order with MADGRAPH5_AMC@NLO 2.2.2, using PYTHIA 8 for the parton shower. No merging procedure has been employed. The LO UFO implementations of the scalar
1680 and vector models that will be used as early Run-2 benchmarks can be found on the Forum SVN repository [135], while the 2HDM model can be found at this link [136].

As a final technical remark, we suggest always to let the shower program handle

the h decay (and therefore to generate a stable h at the matrix element level). In so doing a much faster generation is achieved and the h branching ratios are more accurately accounted for by the shower program.

MADGRAPH5_AMC@NLO *details for scalar mediator Higgs+MET model*. The case of the associated production of a Higgs and scalar mediator via a top-quark loop can be either considered exactly or via an effective Lagrangian where the top-quark is integrated out. While this latter model has been shown not to be reliable [106, 137, 138], for simplicity we have chosen to perform the study in this tree-level effective formulation. A full study of the process including finite top-quark mass and parton shower effects is possible yet left for future work.

MADGRAPH5_AMC@NLO *details for 2HDM Higgs+MET model*. While a 2HDM UFO implementation at NLO accuracy to be used with MADGRAPH5_AMC@NLO has been made available at the end of the work of the Forum [111], in this work we have only considered LO simulations.

The two couplings that can be changed in the implemented model follow the nomenclature below:

- $Tb - \tan \beta$
- $g_Z - g_z$, gauge coupling of Z' to quarks

The other couplings are not changed, including g_x (the $A\tilde{\chi}\chi$ coupling) which has little impact on the signal. $\sin \alpha$ is fixed internally such that $\cos(\beta - \alpha) = 0$. The width of the Z' and A can be computed automatically within MADGRAPH5_AMC@NLO. The couplings here don't affect the signal kinematics, so they can be fixed to default values and then the signal rates can be scaled appropriately.

The nomenclature for the masses in the implemented model is:

- $M_{Z'}$ - PDG ID 32 - Z'
- M_A - PDG ID 28 - A
- M_X - PDG ID 1000022 - dark matter particle

1710 The other masses are unchanged and do not affect the result. Both $Z' \rightarrow hZ(\bar{\nu}\nu)$ and $Z' \rightarrow hA(\tilde{\chi}\chi)$ contribute to the final state, scaling different with model parameters. We recommend to generate them separately, and then add the two signal processes together weighted by cross sections.

4.2.2. Implementation of EFT models for EW boson signatures

1715 The state of the art for these models is LO+PS. NLO+PS can be achieved as well, but the corresponding implementation is not yet available. In our simulations we have implemented the models in the corresponding UFO files and generated events at LO via MADGRAPH5_AMC@NLO 2.2.2, using PYTHIA 8 for the parton shower. UFO files and parameter cards that will be used as early Run-2 benchmarks can
1720 be found on the Forum SVN repository: [135] for operators with Higgs+MET final states and [139] for $W/Z/\gamma$ final states. These models do not require merging.

5. Presentation of EFT results

Most of this report has focused on simplified models of dark matter. In this Chapter, we wish to emphasize the applicability of Effective Field Theories (EFTs)
1725 in the interpretation of DM searches at the LHC. Given our current lack of knowledge about the nature of a DM particle and its interactions, it appears mandatory to provide the necessary information for a model independent interpretation of the collider bounds. This approach should be complemented with an interpretation within a choice of simplified models. We note that, even though EFT benchmarks
1730 are only valid in given conditions, the results provided by the current list of simplified models cannot always characterize the breadth of SM-DM interactions. In at least one case, composite WIMPs [140, 141, 142], the contact interaction framework is the correct one to constrain new confinement scales.

Ideally, experimental constraints should be shown as bounds of allowed signal events in the kinematic regions considered for the search, as detailed in Appendix B. A problematic situation is the attempt to derive a limit on nucleon-dark matter

scattering cross sections from EFT results based on collider data ¹⁴. Experiments that directly probe the nucleon-dark matter scattering cross section are testing the regime of small momentum transfers, where the EFT approximation typically holds. Collider experiments, though, are sensitive to large momentum transfers: We first illustrate the complications that can arise with EFTs at colliders by considering an effective interaction

$$\mathcal{L}_{\text{int}} = \frac{(\bar{q}\gamma_\mu q)(\bar{\chi}\gamma^\mu \chi)}{M_*^2} = (\bar{q}\gamma_\mu q)(\bar{\chi}\gamma^\mu \chi)\frac{g}{\Lambda^2}$$

that couples quarks and DM χ fields.¹⁵ The strength of this interaction is parametrized by $\frac{1}{M_*^2} = \frac{g}{\Lambda^2}$. A monojet signature can be generated from this operator by applying perturbation theory in the QCD coupling. An experimental search will place a limit on M_* . For a fixed M_* , a small value of g will correspond to a small value of Λ . The EFT approximation breaks down if $Q > \Lambda$, where Q is a typical hard scale of the process. The limit on small g can only be reliable if the kinematic region $Q > \Lambda$ is removed from the event generation. However, if a fraction of events is removed from the prediction, the corresponding value of g must increase to match the experimental limit on M_* . On the other hand, if, for the same value of M_* , a large Λ is assumed so that the full set of events fulfill the EFT validity condition, a larger value of g is required. For large enough g , computations based on perturbation theory become unreliable.

In the first part of this Chapter, we summarize two methods that have been advocated to truncate events that do not fulfill the condition necessary for the use of an EFT. These methods are described in detail in Refs. [28, 143, 78, 7, 144, 96]. We then propose a recommendation for the presentation of EFT results for early Run-2 LHC searches.

¹⁴Comparisons between constraints from different experiments meant to highlight their complementarity should be expressed as a function of the model parameters rather than on derived observables; however this is a point that should be developed further after the conclusion of the work of this Forum.

¹⁵The *exact* operator chosen is not important: as detailed in the following, statements concerning the applicability of an EFT can also be made without a specific relation to simplified models.

5.1. Procedures for the truncation of EFT benchmark models

5.1.1. EFT truncation using the momentum transfer and information on UV completion

In the approach described in Ref. [143], the EFT prediction is modified to incorporate the effect of a propagator for a relatively light mediator. For a tree-level interaction between DM and the SM via some mediator with mass M_{med} , the EFT approximation corresponds to expanding the propagator for the mediator in powers of $Q_{\text{tr}}^2/M_{\text{med}}^2$, truncating at lowest order, and combining the remaining parameters into a single parameter M_* (connected to the scale of the interaction Λ in the literature). For an example scenario with a Z' -type mediator (leading to some combination of operators D5 to D8 in the notation of [23] for the EFT limit), this corresponds to setting

$$\frac{g_\chi g_q}{Q_{\text{tr}}^2 - M_{\text{med}}^2} = -\frac{g_\chi g_q}{M_{\text{med}}^2} \left(1 + \frac{Q_{\text{tr}}^2}{M_{\text{med}}^2} + \mathcal{O}\left(\frac{Q_{\text{tr}}^4}{M_{\text{med}}^4}\right) \right) \simeq -\frac{1}{M_*^2}, \quad (45)$$

where Q_{tr} is the momentum carried by the mediator, and g_χ, g_q are the DM-mediator and quark-mediator couplings respectively.¹⁶ A minimal condition that must be satisfied for this approximation to be valid is that $Q_{\text{tr}}^2 < M_{\text{med}}^2 = g_\chi g_q M_*^2$. This requirement avoids the regions: $Q_{\text{tr}}^2 \sim M_{\text{med}}^2$, in which case the EFT misses a resonant enhancement, and it is conservative to ignore this enhancement; and $Q_{\text{tr}}^2 \gg M_{\text{med}}^2$, in which case the signal cross section should fall according to a power of Q_{tr}^{-1} instead of M_{med}^{-1} . The latter is the problematic kinematic region.

The condition $Q_{\text{tr}}^2 < M_{\text{med}}^2 = g_\chi g_q M_*^2$ was applied to restrict the kinematics of the signal and remove events for which the high-mediator-mass approximation made in the EFT would not be reliable. This leads to a smaller effective cross-section, after imposing the event selection of the analysis. This truncated signal was then used to derive a new, more conservative limit on M_* as a function of $(m_\chi, g_\chi g_q)$.

For the example D5-like operator, where the cross section σ scales as M_*^{-4} , there is a simple rule for converting a rescaled cross section into a rescaled con-

¹⁶Here, we ignore potential complications from the mediator width when the couplings are large.

straint on M_* if the original limit is based on a simple cut-and-count procedure. Defining $\sigma_{\text{EFT}}^{\text{cut}}$ as the cross section truncated such that all events pass the condition $\sqrt{g_\chi g_q} M_*^{\text{rescaled}} > Q_{\text{tr}}$, we have

$$M_*^{\text{rescaled}} = \left(\frac{\sigma_{\text{EFT}}}{\sigma_{\text{EFT}}^{\text{cut}}(M_*^{\text{rescaled}})} \right)^{1/4} M_*^{\text{original}}, \quad (46)$$

which can be solved for M_*^{rescaled} via either iteration or a scan. Similar relations exist for a given UV completion of each operator.

This procedure has been proposed in Ref. [143] and its application to ATLAS
 1770 results can be found in Ref. [7] for a range of operators. We reiterate: knowledge of the UV completion for a given EFT operator was necessary for this procedure; this introduces a model-dependence that was not present in the non-truncated EFT results.

Currently, simplified models (including the full effect of the mediator propaga-
 1775 tor) are available for comparison with the data, and since knowledge of the simplified models is needed for the truncation procedure, there is no reason to apply this prescription. Instead, the simplified model limit for large M_* can be presented for interpretation in terms of EFT operators.

5.1.2. EFT truncation using the center of mass energy

The procedure presented in the previous section was predicated on some knowledge of the simplified model. This led to the identification of the mass of the DM pair as the relevant kinematic quantity to use in a truncation procedure. In general, if no assumption is made about the underlying dynamics, it is more conservative to place a limit on the total center of mass energy E_{cm} of the DM production process. Furthermore, the direct connection between the mass scale of the EFT validity, M_{cut} , and the mass scale that normalizes the EFT operator, M_* , is unknown. For such cases, Refs.[144, 96] proposed a procedure to extract model independent and consistent bounds within the EFT that can be applied to any effective Lagrangian describing the interactions between the DM and the SM. This procedure provides conservative limits that can be directly reinterpreted in any completion of the EFT.

The condition ensuring that the EFT approximation is appropriate is:

$$E_{\text{cm}} < M_{\text{cut}}. \quad (47)$$

1780 The relationship between M_{cut} and M_* can be parameterized by an *effective coupling strength* g_* , such that $M_{\text{cut}} = g_* M_*$. A scan over values of g_* provides an indication of the sensitivity of the prediction to the truncation procedure. In the Z' -type model considered above, g_* is equal to $\sqrt{g_\chi g_q}$. The resulting plots are shown in [144] for a particular effective operator.

1785 The advantage of this procedure is that the obtained bounds can be directly and easily recast in any completion of the EFT, by computing the parameters M_* , M_{cut} in the full model as functions of the parameters of the complete theory. On the other hand, the resulting limits will be weaker than those obtained using Q_{tr} and a specific UV completion.

1790 5.1.3. Truncation at the generator level

The conditions on the momentum transfer can also be applied directly at the generator level, by discarding events that are invalid and calculating the limits from this truncated shape. This provides the necessary rescaling of the cross section while keeping the information on the change in the kinematic distributions due to
1795 the removal of the invalid events. This procedure is more general with respect to rescaling the limit in the two sections above, and it should be followed if a search is not simply a counting experiment and exploits the shapes of kinematic distributions.

5.1.4. Sample results of EFT truncation procedures

An example of the application of the two procedures to the limit on M_* from
1800 Ref. [14] as a function of the product of the couplings is shown in Figure 69. Only the region between the dashed and the solid line is excluded. It can be seen that the procedure from [144] outlined in Section 5.1.2, shown in blue, is more conservative than the procedure from Refs. [143, 7], described in Section 5.1.1.

5.1.5. Comments on unitarity considerations

A further consideration applicable to EFT operators at hadron colliders is the potential violation of unitarity. An analysis of the operator $\frac{\bar{q}\gamma_\mu q \tilde{\chi}\gamma^\mu \chi}{M_*^2}$ provides the limit:

$$M_* > \beta(s) \sqrt{s} \sqrt{\frac{\sqrt{3}}{4\pi}}, \quad (48)$$

1805 where \sqrt{s} is (maximally) the collider energy and $\beta(s)$ is the DM velocity [24]. Constraints for other operators have also been derived [145]. This constraint on M_* still is open to interpretation, since the relation to M_{cut} is not resolved, except for a specific simplified model. Derived limits on M_* should be compared to this unitarity bound to check for consistency.

1810 5.2. Recommendation for presentation of EFT results

In this report, we make two recommendations for the presentation of collider results in terms of Effective Field Theories for the upcoming Run-2 searches. A full discussion of the presentation of collider results in relation to other experiments is left to work beyond this Forum, where ATLAS, CMS, the theory community and the
1815 Direct and Indirect Detection communities are to be involved.

We divide the EFT operators in two categories: those that can be mapped to one or more UV-complete simplified models, such as those commonly used in LHC searches so far and detailed in [23], and those for which no UV completion is available to LHC experiments, such as those outlined in Section 3.2.

1820 5.2.1. EFT benchmarks with corresponding simplified models

If a simplified model can be mapped to a given EFT, then the model's high-mediator-mass limit will converge to the EFT. A study of 14 TeV benchmarks for narrow resonances with $g_q = 0.25$ and $g_\chi = 1$ (see Section 2.1.1) shows that a mediator with a mass of at least 10 TeV fully reproduces the kinematics of a contact
1825 interaction and has no remaining dependence on the presence of a resonance. A comparison of the main kinematic variables for the s -channel vector mediator model

with a width of $0.1 M_{\text{med}}$ is shown in Fig. 70.¹⁷

As already observed in Section 2.1.1, varying the DM mass changes the kinematics, both in the simplified model and in the EFT case. This can be seen in Fig. 71.

1830 Based on these studies, the Forum recommends experimental collaborations to add one grid scan point at very high mediator mass (10 TeV) to the scan, for each of the DM masses for the s -channel simplified models described in Section 2. This will allow to reproduce the results of an equivalent contact interaction as a simple extension of the existing parameter scan.

1835 It should be checked that the high-mass mediator case for the simplified model is correctly implemented

5.2.2. EFT benchmarks with no corresponding simplified models

Whenever a UV completion is not available, an EFT still captures a range of possible theories beyond the simplified models that we already consider. However, in
1840 the case of the dimension-7 operators detailed in Section 3.2 we can only roughly control how well the EFT approximation holds, as described in Section 3.2.4. Despite the fact that a propagator was introduced to motivate the truncation procedure for s -channel models, the prescription from Sec. 5.2.1 depends upon the simplified model to derive the energy scaling that is used for the comparison with the momen-
1845 tum transfer. The simple fact remains that the effective coupling of the operator – g/Λ^n – should not allow momentum flow $Q > \Lambda$ or $g > 4\pi$. Given our ignorance of the actual kinematics, the truncation procedure recommended for this purpose is the one described in Section 5.1.2, as it is independent from any UV completion details.

1850 Because there is no UV completion, the parameter M_{cut} can be treated more freely than an explicit function of g and Λ . It makes sense to choose M_{cut} such that we identify the transition region where the EFT stops being a good description of UV complete theories. This can be done using the ratio R , which is defined as the

¹⁷The use of a fixed width rather than the minimal width is exclusive of these plots.

fraction of events for which $\hat{s} > M_{\text{cut}}^2$. For large values of M_{cut} , no events are thrown
 1855 away in the truncation procedure, and $R = 1$. As M_{cut} becomes smaller, eventually
 all events are thrown away in the truncation procedure, i.e. $R = 0$, and the EFT
 gives no exclusion limits for the chosen acceptance.

We propose a rough scan over M_{cut} , such that we find the values of M_{cut} for
 which R ranges from 0.1 to 1. The analysis can then perform a scan over several
 1860 values of M_{cut} , and show the truncated limit for each one of them.

6. Evaluation of signal theoretical uncertainties

A comprehensive and careful assessment of signal theoretical uncertainties plays
 in general a more important role for the background estimations (especially when
 their evaluation is non-entirely data-driven) than it does for signal simulations.
 1865 Nevertheless, also for signal samples theoretical uncertainties are relevant, and may
 become even dominant in certain regions of phase space.

The uncertainties on the factorization and renormalization scales are assessed
 by the experimental collaborations by varying the original scales of the process by
 factors of 0.5 and 2. The evaluation of the uncertainty on the choice of PDF follows
 1870 the PDF4LHC recommendation [146] of considering the envelope of different PDF
 error sets, in order to account for the uncertainty on the various PDFs as well as the
 uncertainty on the choice of the central value PDF. The Forum has not discussed the
 uncertainties related to the merging of different samples, nor the uncertainty due
 to the choice of the modeling of the parton shower. This Chapter provides technical
 1875 details on how scale and PDF uncertainties can be assessed for events generated
 with POWHEG and MADGRAPH5_AMC@NLO.

6.1. POWHEG

When using POWHEG [69, 67, 68], it is possible to study scale and PDF errors
 for the dark matter signals. A fast reweighting machinery is available in POWHEG-
 1880 BOX that allows one to add, after each event, new weights according to different
 scale or PDF choices, without the need to regenerate all the events from scratch.

To enable this possibility, the variable `storeinfo_rwgt` should be set to 1 in the POWHEG input file when the events are generated for the first time¹⁸. After each event, a line starting with

```
1885      #rwgt
```

is appended, containing the necessary information to generate extra weights. In order to obtain new weights, corresponding to different PDFs or scale choice, after an event file has been generated, a line

```
      compute_rwgt 1
```

1890 should be added in the input file along with the desired change in parameters. For instance, `renscfact` and `facscfact` allow one to study scale variations on the renormalization and factorization scales around a central value. By running the program again, a new event file will be generated, named `<OriginalName>-rwgt.lhe`, with one more line at the end of each event of the form

```
1895      #new weight,renfact,facfact,pdf1,pdf2
```

followed by five numbers and a character string. The first of these numbers is the weight of that event with the new parameters chosen. By running in sequence the program in the reweighting mode, several weights can be added on the same file. Two remarks are in order.

- 1900 • The file with new weights is always named
 `<OriginalName>-rwgt.lhe`
 hence care has to be taken to save it as
 `<OriginalName>.lhe`
 before each iteration of the reweighting procedure.
- 1905 • Due to the complexity of the environment where the program is likely to be run, it is strongly suggested as a self-consistency check that the first reweighting is done keeping the initial parameters. If the new weights are not exactly

¹⁸Notice that even if the variable is not present, by default it is set to 1.

the same as the original ones, then some inconsistency must have happened,
or some file was probably corrupted.

1910 It is possible to also have weights written in the version 3 Les Houches format. To
do so, in the original run, at least the token
lhrwgt_id 'ID'
must be present. The reweighting procedure is the same as described above, but
now each new run can be tagged by using a different value for the lhrwgt_id
1915 keyword. After each event, the following lines will appear:

```
<rwgt>
<wgt id='ID'>
<wgt id='ID1'>
</rwgt>
```

1920 A more detailed explanation of what went into the computation of every single
weight can be included in the <header> section of the event file by adding/changing
the line

```
lhrwgt_descr 'some info'
```

in the input card, before each “reweighting” run is performed. Other useful key-
1925 words to group together different weights are lhrwgt_group_name and lhrwgt_group_combine.

More detailed information can be obtained by inspecting the document in /Docs/V2-paper.pdf
under the common POWHEG-BOX-V2 directory.

6.2. The SysCalc package in MADGRAPH5_AMC@NLO

SysCalc is a post-processing package for parton-level events as obtained from
1930 leading-order calculations in MADGRAPH5_AMC@NLO. It can associate to each event
a series of weights corresponding to the evaluation of a certain class of theoretical
uncertainties. The event files in input and output are compliant with the Les
Houches v3 format. For NLO calculations, PDF and scale uncertainties are instead
evaluated automatically by setting corresponding instructions in the run_card.dat
1935 and no post-processing is needed (or possible).

The requirements of the package as inputs are :

- A systematics file (which can be generated by MadGraph 5 v. 1.6.0 or later) [73, 147].
- The Pythia-PGS package (v. 2.2.0 or later) [148]. This is needed only in the case of matching scales variations.
- The availability of LHAPDF5 [149].
- A configuration file (i.e. a text file) specifying the parameters to be varied.

SYS CALC supports all leading order computations generated in MADGRAPH5_AMC@NLO including fixed-order computation and matched-merged computation performed in the MLM scheme [150]. MADGRAPH5_AMC@NLO stores additional information inside the event in order to have access to all the information required to compute the convolution of the PDFs with the matrix element for the various supported systematics.

An example configuration file is as follows:

```

1950
    # Central scale factors
scalefact:
0.5 1 2
    # Scale correlation
1955 # Special value -1: all combination (N**2)
    # Special value -2: only correlated variation
    # Otherwise list of index N*fac_index + ren_index
    # index starts at 0
scalecorrelation:
1960 -1
    #  $\alpha_s$  emission scale factors
alpsfact:
0.5 1 2
    # matching scales
1965 matchscale:

```

```

30 60 120

# PDF sets and number of members (optional)

PDF:

CT10.LHgrid 53

1970 MSTW2008nlo68cl.LHgrid

```

Without matching/merging, SYSCALC is able to compute the variation of renormalisation and factorisation scale (parameter `scalefact`) and the change of PDFs. The variation of the scales can be done in a correlated and/or uncorrelated way, basically following the value of the `scalecorrelation` parameter which can take the following values:

- -1 : to account for all N^2 combinations.
- -2 : to account only for the correlated variations.
- A set of positive values corresponding to the following entries (assuming 0.5, 1, 2 for the `scalefact` entry):

```

1980
0:  $\mu_F = \mu_F^{\text{orig}}/2, \mu_R = \mu_R^{\text{orig}}/2$ 
1:  $\mu_F = \mu_F^{\text{orig}}/2, \mu_R = \mu_R^{\text{orig}}$ 
2:  $\mu_F = \mu_F^{\text{orig}}/2, \mu_R = \mu_R^{\text{orig}} * 2$ 
3:  $\mu_F = \mu_F^{\text{orig}}, \mu_R = \mu_R^{\text{orig}}/2$ 
1985 4:  $\mu_F = \mu_F^{\text{orig}}, \mu_R = \mu_R^{\text{orig}}$ 
5:  $\mu_F = \mu_F^{\text{orig}}, \mu_R = \mu_R^{\text{orig}} * 2$ 
6:  $\mu_F = \mu_F^{\text{orig}} * 2, \mu_R = \mu_R^{\text{orig}}/2$ 
7:  $\mu_F = \mu_F^{\text{orig}} * 2, \mu_R = \mu_R^{\text{orig}}$ 
8:  $\mu_F = \mu_F^{\text{orig}} * 2, \mu_R = \mu_R^{\text{orig}} * 2$ 

```

Without correlation, the weight associated to the renormalisation scale is the following:

$$\mathcal{W}_{\text{new}}^{\mu_R} = \frac{\alpha_S^N(\Delta * \mu_R)}{\alpha_S^N(\mu_R)} * \mathcal{W}_{\text{orig}}, \quad (49)$$

where Δ is the scale variation considered, $\mathcal{W}_{\text{orig}}$ and \mathcal{W}_{new} are respectively the original/new weights associated to the event. N is the power in the strong coupling for

the associated event (interference is not taken account on an event by event basis).

The weight associated to the scaling of the factorisation scale is:

$$\mathcal{W}_{\text{new}}^{\mu_F} = \frac{f_{1,\text{orig}}(x_1, \Delta * \mu_F) * f_{2,\text{orig}}(x_2, \Delta * \mu_F)}{f_{1,\text{orig}}(x_1, \mu_F) * f_{2,\text{orig}}(x_2, \mu_F)} * \mathcal{W}_{\text{orig}}, \quad (50)$$

1990 where $f_{i,\text{orig}}$ are the probabilities from the original PDF set associated to the incoming partons, which hold a proton momentum fraction x_1 and x_2 for the first and second beam respectively.

The variations for the PDF are given by the corresponding weights associated to the new PDF sets:

$$\mathcal{W}_{\text{new}}^{\text{PDF}} = \frac{f_{1,\text{new}}(x_1, \mu_F) * f_{2,\text{new}}(x_2, \mu_F)}{f_{1,\text{orig}}(x_1, \mu_F) * f_{2,\text{orig}}(x_2, \mu_F)} * \mathcal{W}_{\text{orig}}, \quad (51)$$

where $f_{i,\text{new}}$ is the new PDF probability associated to parton i .

In presence of matching, MADGRAPH5_AMC@NLO associates one history of radiation (initial and/or final state radiation) obtained by a k_T clustering algorithm, and calculates α_s at each vertex of the history to a scale given by the aforementioned clustering algorithm. Furthermore, MADGRAPH5_AMC@NLO reweights the PDF in a fashion similar to what a parton shower would do. SYSCALC can perform the associated re-weighting (parameter `alpsfact`) by dividing and multiplying by the associated factor.

For each step in the history of the radiation (associated to a scale $\mu_i = k_{T,i}$), this corresponds to the following expression for a Final State Radiation (FSR):

$$\mathcal{W}_{\text{new}}^{\text{FSR}} = \frac{\alpha_s(\Delta * \mu_i)}{\alpha_s(\mu_i)} * \mathcal{W}_{\text{orig}}, \quad (52)$$

and to the following expression for Initial State Radiation (ISR), associated to a scale μ_i and fraction of energy x_i :

$$\mathcal{W}_{\text{new}}^{\text{ISR}} = \frac{\alpha_s(\Delta * \mu_i)}{\alpha_s(\mu_i)} \frac{\frac{f_a(x_i, \Delta * \mu_i)}{f_b(x_i, \Delta * \mu_{i+1})}}{\frac{f_a(x_i, \mu_i)}{f_b(x_i, \mu_{i+1})}} * \mathcal{W}_{\text{orig}}, \quad (53)$$

where μ_{i+1} is the scale of the next step in the (initial state) history of radiation.

SYSCALC can include the weight associated to different merging scales in the MLM matching/merging mechanism (for output of the `pythia6` package or `pythia-pgs` package).

2005 In that case, the parton shower does not veto any event according to the MLM
algorithm, although in the output file the scale of the first emission is retained. Hav-
ing this information, SYSCALC can test each value of the specified matching scales
under the `matchscale` parameter block. SYSCALC will then test for each of the
values specified in the parameter `matchscale` if the event passes the MLM criteria
2010 or not. If it does not, then a zero weight is associated to the event, while if it does,
then a weight 1 is kept. As a reminder, those weights are the equivalent of having
a (approximate) Sudakov form-factor and removing at the same time the double
counting between the events belonging to different multiplicities.

2015 Finally, we give an example of the SYSCALC output which follows the LHEF v3
format. The following block appears in the header of the output file:

```
<header>
  <initrwgt>
    <weightgroup type="Central scale variation" combine="envelope">
2020   <weight id="1"> mur=0.5 muf=0.5 </weight>
      <weight id="2"> mur=1 muf=0.5 </weight>
      <weight id="3"> mur=2 muf=0.5 </weight>
      <weight id="4"> mur=0.5 muf=1 </weight>
      <weight id="5"> mur=1 muf=1 </weight>
2025   <weight id="6"> mur=2 muf=1 </weight>
      <weight id="7"> mur=0.5 muf=2 </weight>
      <weight id="8"> mur=1 muf=2 </weight>
      <weight id="9"> mur=2 muf=2 </weight>
    </weightgroup>
2030   <weightgroup type="Emission scale variation" combine="envelope">
      <weight id="10"> alpsfact=0.5</weight>
      <weight id="11"> alpsfact=1</weight>
      <weight id="12"> alpsfact=2</weight>
    </weightgroup>
2035   <weightgroup type="CT10nlo.LHgrid" combine="hessian">
      <weight id="13">Member 0</weight>
      <weight id="14">Member 1</weight>
      <weight id="15">Member 2</weight>
      <weight id="16">Member 3</weight>
```

```

2040      ...
      <weight id="65">Member 52</weight>
    </weightgroup>
  </initrwgt>
</header>

```

2045 For each event, the weights are then written as follows:

```

<rwgt>
  <wgt id="1">83214.7</wgt>
  <wgt id="2">61460</wgt>
  <wgt id="3">47241.9</wgt>
2050  <wgt id="4">101374</wgt>
      ...
  <wgt id="64">34893.5</wgt>
  <wgt id="65">41277</wgt>
</rwgt>

```

2055 7. Conclusions

The ATLAS/CMS Dark Matter Forum concluded its work in June 2015. Its mandate was focused on identifying a prioritized, compact set of simplified model benchmarks to be used for the design of the early Run-2 LHC searches for $\cancel{E}_T + X$ final states. Its participants included many of the experimenters from both collaborations that are involved in these searches, as well as many of the theorists working actively on these models. This report has documented this basis set of models, as well as studies of the kinematically-distinct regions of the parameter space of the models, to aid the design of the searches. Table 6.1 summarizes the state of the art of the calculations, event generators, and tools that are available to the two LHC collaborations to simulate these models at the start of Run-2. It also describes some that are known to be under development as the report was finalized.

This document primarily presents studies related to simplified models. The presentation of results for EFT benchmark models is also discussed. The studies contained in this report are meant to highlight the use of EFTs as a benchmark that is

Benchmark models for ATLAS and CMS Run-2 DM searches				
vector/axial vector mediator, s-channel (Sec. 2.1)				
Signature	State of the art calculation and tools	Implementation	References	
$\text{jet} + \cancel{E}_T$	NLO+PS (POWHEG, SVN r3059)	[113, 114]	[59, 43, 67, 68, 69]	
	NLO+PS (<i>DMsimp</i> UFO + MadGRAPH5_AMC@NLO v2.3.0)	[111]	[73, 74, 75]	
	NLO (MCFM v7.0)	Upon request	[112, 41]	
$W/Z/\gamma + \cancel{E}_T$	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[124]	[73, 74, 75]	
	NLO+PS (<i>DMsimp</i> UFO + MadGRAPH5_AMC@NLO v2.3.0)	[111]	[73, 74, 75]	
scalar/pseudoscalar mediator, s-channel (Sec. 2.2)				
Signature	State of the art calculation and tools	Implementation	References	
$\text{jet} + \cancel{E}_T$	LO+PS, top loop (POWHEG, r3059)	[115, 116]	[59, 43, 67, 68, 69]	
	LO+PS, top loop (<i>DMsimp</i> UFO + MadGRAPH5_AMC@NLO v2.3.0)	[111]	[73, 151, 74, 75]	
	LO, top loop (MCFM v7.0)	Upon request	[112, 41]	
$W/Z/\gamma + \cancel{E}_T$	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)		[73, 74, 75]	
	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[125]	[73, 74, 75]	
$t\bar{t}, b\bar{b} + \cancel{E}_T$	NLO+PS (<i>DMsimp</i> UFO + MadGRAPH5_AMC@NLO v2.3.0)	[111]	[73, 74, 75]	
scalar mediator, t-channel (Sec. 2.3)				
Signature	State of the art calculation and tools	Implementation	References	
$\text{jet}(s) + \cancel{E}_T$ (2-quark gens.)	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[122]	[76, 73, 74, 75]	
$\text{jet}(s) + \cancel{E}_T$ (3-quark gens.)	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[123]	[46, 73, 74, 75]	
$W/Z/\gamma + \cancel{E}_T$	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	TBC	[46, 73, 74, 75]	
$b + \cancel{E}_T$	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[126]	[85, 86, 73, 74, 75]	
Specific simplified models with EW bosons (Sec. 3.1)				
Signature and model	State of the art calculation and tools	Implementation	References	
Higgs + \cancel{E}_T , vector med.	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[135]	[48, 96, 73, 74, 75]	
Higgs + \cancel{E}_T , scalar med.	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[135]	[48, 96, 73, 74, 75]	
Higgs + \cancel{E}_T , 2HDM	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[136]	[96, 73, 74, 75]	
Contact interaction operators with EW bosons (Sec. 3.1)				
Signature and model	State of the art calculation and tools	Implementation	References	
$W/Z/\gamma + \cancel{E}_T$, dim-7	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[139]	[97, 45, 98, 96, 73, 74, 75]	
Higgs + \cancel{E}_T , dim-4/dim-5	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[152]	[48, 47, 96, 73, 74, 75]	
Higgs + \cancel{E}_T , dim-8	LO+PS (UFO + MadGRAPH5_AMC@NLO v2.2.3)	[135]	[48, 47, 96, 73, 74, 75]	

Table 6.1: Summary table for available benchmark models considered within the works of this Forum. The results in this document have been obtained with the implementations in bold.

complementary to simplified models, and to demonstrate how that collider results could be presented a function of the fraction of events that are valid within the contact interaction approximation.

A number of points remain to be developed beyond the scope of this Forum, in
2075 order to fully benefit from LHC searches in the global quest for Dark Matter. First
and foremost, to accomodate the urgent need of a basis set of simplified models,
this work has made many grounding assumptions, as stated in the introduction. De-
partures from these assumptions have not been fully explored. As a consequence,
the list of models and implementations employed by the ATLAS and CMS collabora-
2080 tions for early LHC Run-2 searches is not meant to exhaust the range of possibilities
for mediating processes, let alone cover all plausible models of collider dark matter
production. Rather, it is hoped that others will continue the systematic exploration
of the most generic possibilities for collider dark matter production, building upon
the framework used in this report just as this report has relied heavily on the work of
2085 many others. This also applies to models that exist in literature but do not have an
implementation yet: we hope that this work will further encourage the theory and
generator community to improve the implementation of new models as well as the
precision of the calculations of existing ones. The role of constraints on the media-
tor particles from direct past and present collider searches should also be developed
2090 further.

Furthermore, we see the need for broader discussion on the comparison of ex-
perimental results amongst collider and non-collider searches for particle dark mat-
ter. This point will have to be addressed before the presentation of Run-2 results:
The uncertainties in the comparisons between experiments should be discussed and
2095 conveyed, so that the different results can be placed in their correct context, and so
we can collectively build a fair and comprehensive picture of our understanding of
particle dark matter.

Acknowledgements

The authors would like to thank Daniel Whiteson for helping in the review of
2100 this document. This research was supported by the Munich Institute for Astro- and
Particle Physics (MIAPP) of the DFG cluster of excellence "Origin and Structure of

the Universe". The authors would like to express a special thanks to the Mainz Institute for Theoretical Physics (MITP) for its hospitality and support. P. Pani wishes to thank the support of the Computing Infrastructure of Nikhef. In addition we
2105 thank CERN and acknowledge the support of: DOE and NSF (USA); SER, SNSF and Cantons of Bern and Geneva, Switzerland; MINECO, Spain; European Union; IN2P3-CNRS, CEA-DSM/IRFU, France; NSERC, NRC, and CFI, Canada; INFN, Italy; FNRS and FWO, Belgium; STFC, United Kingdom; ARC, Australia; GRICES and FCT, Portugal; SRC, Sweden; BMBF, DFG, HGF, MPG and AvH Foundation, Germany;
2110 Austrian Academy of Sciences, Austria; MOE and UM, Malaysia; National Research Foundation and Ministry of Education, Science and Technology, Korea; MEXT and JSPS, Japan; FOM and NWO, Netherlands; CNPq and FAPESP, Brazil; Atomic Energy Commission, Pakistan; DNRE, DNSRC and Lundbeck Foundation, Denmark; Thailand Center of Excellence in Physics, Institute for the Promotion of Teaching
2115 Science and Technology, Special Task Force for Activating Research and National Science and Technology Development Agency, Thailand; MOST, Taiwan.

Appendix A. Appendix: Additional models for Dark Matter searches

Appendix A.1. Models with a single top-quark + \cancel{E}_T

Many different theories predict final states with a single top and associated missing transverse momentum (monotop), some of them including dark matter candidates. A simplified model encompassing the processes leading to this phenomenology is described in Refs. [153, 154, 155], and is adopted as one of the benchmarks for Run 2 LHC searches.

The simplified model is constructed by imposing that the model Lagrangian respects the electroweak $SU(2)_L \times U(1)_Y$ gauge symmetry and by requiring minimality in terms of new states to supplement to the Standard Model fields. As a result, two monotop production mechanisms are possible. In the first case, the monotop system is constituted by an invisible (or long-lived with respect to detector distances) fermion χ and a top quark. It is produced as shown in the diagram of A.72 (a)
2130 where a colored resonance φ lying in the triplet representation of $SU(3)_C$ decays

into a top quark and a χ particle. In the second production mode, the monotop state is made of a top quark and a vector state V connected to a hidden sector so that it could decay invisibly into, e.g., a pair of dark matter particles as studied in [155]. The production proceeds via flavor-changing neutral interactions of the
2135 top quark with a quark of the first or second generation and the invisible V boson (see the diagrams of A.72 (b) and (c)).

Resonant production. In this case, a colored 2/3-charged scalar (φ) is produced and decays into a top quark and a spin-1/2 invisible particle, χ . The dynamics of the new sector is described by the following Lagrangian:

$$\mathcal{L} = \left[\varphi \bar{d}^c \left[a_{SR}^q + b_{SR}^q \gamma_5 \right] d + \varphi \bar{u} \left[a_{SR}^{1/2} + b_{SR}^{1/2} \gamma_5 \right] \chi + \text{h.c.} \right], \quad (\text{A.1})$$

where u (d) stands for any up-type (down-type) quark, the notation SR refers to the monotop production mechanism via a scalar resonance and all flavor and color indices are understood for clarity.

2140 In the notation of [154], the couplings of the new colored fields to down-type quarks are embedded into the 3×3 antisymmetric matrices a_{SR}^q (scalar couplings) and b_{SR}^q (pseudoscalar couplings) while those to the new fermion χ and one single up-type quark are given by the three-component vectors $a_{SR}^{1/2}$ and $b_{SR}^{1/2}$ in flavor space.

Under the form of Eq. (A.1), the Lagrangian is the one introduced in the original monotop search proposal [153]. It has been used by the CMS collaboration for Run I analyses after neglecting all pseudoscalar components of the couplings and adding the vector resonance case for which minimality requirements are difficult to accommodate [18]. In contrast, the study of Ref. [155] has imposed electroweak gauge invariance and required minimality. This enforces all new couplings to be right-handed so that

$$a_{SR}^{1/2} = b_{SR}^{1/2} = \frac{1}{2} y_s^* \quad \text{and} \quad a_{SR}^q = b_{SR}^q = \frac{1}{2} \lambda_s, \quad (\text{A.2})$$

2145 where the objects y_s and λ_s are a tridimensional vector and a 3×3 matrix in flavor space respectively. This class of scenarios is the one that has been adopted by the

ATLAS collaboration for its Run I monotop searches [12] and will be considered by both collaborations for Run II analyses.

The resulting model can be likened to the MSSM with an R -parity violating of a top squark to the Standard Model down-type quarks and an R -parity conserving interaction of a top quark and a top-squark to a neutralino.

Non-Resonant production. For non-resonant monotop production, the monotop state is produced via flavor-changing neutral interactions of the top quark, a lighter up-type quark and a new invisible vector particle V . This is the only case considered, as having a new scalar would involve in particular a mixing with the SM Higgs boson and therefore a larger number of free parameters. The Lagrangian describing the dynamics of this non-resonant monotop production case is:

$$\mathcal{L} = \left[V_\mu \bar{u} \gamma^\mu \left[a_{FC}^1 + b_{FC}^1 \gamma_5 \right] u + \text{h.c.} \right], \quad (\text{A.3})$$

where the flavor and color indices are again understood for clarity. The strength of the interactions among these two states and a pair of up-type quarks is modeled via two 3×3 matrices in flavor space a_{FC}^1 for the vector couplings and b_{FC}^1 for the axial vector couplings, the FC subscript referring to the flavor-changing neutral monotop production mode and the (1) superscript to the vectorial nature of the invisible particle.

As for the resonant case, the Lagrangian of Eq. (A.3) is the one that has been used by CMS after reintroducing the scalar option for the invisible state and neglecting all pseudoscalar interactions [18]. As already mentioned, a simplified setup motivated by gauge invariance and minimality has been preferred so that, as shown in Ref. [155], we impose all interactions to involve right-handed quarks only¹⁹,

$$a_{FC}^1 = b_{FC}^1 = \frac{1}{2} a_R \quad (\text{A.4})$$

where a_R denotes a 3×3 matrix in flavor space. This implies the vector field to be an $SU(2)_L$ singlet.

¹⁹Ref. [?] , which became public only after the conclusion of this Forum, introduces both left- and right-handed couplings.

Model parameters and assumptions. The models considered as benchmarks for the first LHC searches contain further assumptions in terms of the flavor structure of the model with respect to the Lagrangians of the previous subsection. In order to have an observable monotop signature at the LHC, the Lagrangians introduced above must include not too small couplings of the new particles to first and second generation quarks. For simplicity, we assumed that only channels enhanced by parton density effects will be considered, so that we fix

$$\begin{aligned} (a_R)_{13} &= (a_R)_{31} = a , \\ (\lambda_s)_{12} &= -(\lambda_s)_{21} = \lambda \quad \text{and} \quad (y_s)_3 = y , \end{aligned} \tag{A.5}$$

2160 all other elements of the matrices and vectors above being set to zero.

Implementation. In order to allow one for the Monte Carlo simulation of events relevant for the monotop production cases described above, we consider the Lagrangian

$$\mathcal{L} = \left[a V_\mu \bar{u} \gamma^\mu P_R t + \lambda \varphi \bar{d}^c P_R s + y \varphi \bar{\chi} P_R t + \text{h.c.} , \right] \tag{A.6}$$

where P_R stands for the right-handed chirality projector and the new physics couplings are defined by the three parameters a , λ and y . We additionally include a coupling of the invisible vector boson V to a dark sector (represented by a fermion ψ) whose strength can be controlled through a parameter g_{DM} ,

$$\mathcal{L} = g_{DM} V_\mu \bar{\psi} \gamma^\mu \psi . \tag{A.7}$$

This ensures the option to make the V -boson effectively invisible by tuning g_{DM} respectively to a . We implement the entire model in the FEYNRULES package [74] so that the model can be exported to a UFO library [75] to be linked to MADGRAPH5_AMC@NLO [73] for event generation, following the approach outlined in [156].

Appendix A.2. Parameter scan

Under all the assumptions of the previous sections, the parameter space of the resonant model is defined by four quantities, namely the mass of the new scalar

field φ , the mass of the invisible fermion χ and the strengths of the interactions
 2170 of the scalar resonance with the monotop system y and with down-type quarks λ .
 One of both coupling parameters could however be traded with the width of the
 resonance.

The parameter space of the non-resonant model is defined by two parameters,
 namely the mass of the invisible state V and its flavor-changing neutral coupling to
 2175 the up-type quarks a_R .

In the case of the non-resonant model, the invisible vector is connected to a hid-
 den sector that could be, in its simplest form, parameterized by a new fermion [155].
 This has effects on the width of the invisible V state.

A consensus between the ATLAS and CMS collaborations has been reached in the
 2180 case of non-resonant monotop production. The results have been described above.
 In contrast, discussions in the context of resonant monotop production are still on-
 going. The related parameter space contains four parameters and must thus be
 further simplified for practical purposes. Several options are possible and a choice
 necessitates additional studies that will be achieved in a near future.

2185 It has been verified that the kinematics do not depend on the width of the in-
 visible state in the case where this width is at most 10% of the V -mass. This is
 illustrated in Fig. A.73, where we show the transverse-momentum spectra of the
 V -boson when it decays into a top-up final state and for different V -boson masses.
 The results are independent of the visible or invisible decay modes as we are only
 2190 concerned with the kinematic properties of the invisible state.

Appendix A.3. Single Top Model implementation

Card files for MADGRAPH5_AMC@NLO are provided on the Forum SVN reposi-
 tory [157] and correspond to the Lagrangian that has been implemented in FEYN-
 RULES. Each coupling constant of the model can be set via the block COUPX of
 2195 the parameter card. Its entries 1, 2 and 3 respectively correspond to the monotop-
 relevant parameters a , λ and y , while the width (and in particular the invisible
 partial width) of the V -boson can be tuned via the g_{DM} parameter to given in the
 entry 10 of the COUPX block.

The masses of the particles are set in the MASS block of the parameter card,
 2200 the PDG codes of the new states being 32 (the vector state V), 1000006 (the φ
 colored resonance), 1000022 (the invisible fermion χ) and 1000023 (the fermion
 ψ connecting the V state to the dark sector). The width of the new vector has to
 be computed from all open tree-level decays (after fixing g_{DM} to a large value and
 setting the relevant entry to Auto in the DECAY block of the parameter card), while
 2205 the way to calculate the width of the resonance ϕ is under discussion by both the
 ATLAS and CMS collaborations. The chi and psi fermions are taken stable so that
 their width vanishes.

Appendix A.4. Further $W+\cancel{E}_T$ models with possible cross-section enhancements

As pointed out in Ref. [65], the mono- W signature can probe the iso-spin vi-
 olating interactions of dark matter with quarks. The relevant operator after the
 electroweak symmetry breaking is

$$\frac{1}{\Lambda^2} \bar{\chi} \gamma_\mu \chi (\bar{u}_L \gamma^\mu u_L + \xi \bar{d}_L \gamma^\mu d_L). \quad (\text{A.8})$$

Here, we only keep the left-handed quarks because the right-handed quarks do not
 radiate a W -gauge boson from the weak interaction. As the LHC constrains the
 cutoff to higher values, it is also important to know the corresponding operators
 before the electroweak symmetry. At the dimension-six level, the following operator

$$\frac{c_6}{\Lambda^2} \bar{\chi} \gamma_\mu \chi \bar{Q}_L \gamma^\mu Q_L \quad (\text{A.9})$$

conserves iso-spin and provides us $\xi = 1$ [65]. At the dimension-eight level, new
 operators appear to induce iso-spin violation and can be

$$\frac{c_8^d}{\Lambda^4} \bar{\chi} \gamma_\mu \chi (H \bar{Q}_L) \gamma^\mu (Q_L H^\dagger) + \frac{c_8^u}{\Lambda^4} \bar{\chi} \gamma_\mu \chi (\tilde{H} \bar{Q}_L) \gamma^\mu (Q_L \tilde{H}^\dagger). \quad (\text{A.10})$$

After inputting the vacuum expectation value of the Higgs field, we have

$$\xi = \frac{c_6 + c_8^d v_{EW}^2 / 2\Lambda^2}{c_6 + c_8^u v_{EW}^2 / 2\Lambda^2}. \quad (\text{A.11})$$

For a nonzero c_6 and $v_{EW} \ll \Lambda$, the iso-spin violation effects are suppressed. On the
 2210 other hand, the values of c_6 , c_8^d and c_8^u depend on the UV-models.

There is one possible UV-model to obtain a zero value for c_6 and non-zero values for c_8^d and c_8^u . One can have the dark matter and the SM Higgs field charged under a new $U(1)'$ symmetry. There is a small mass mixing between SM Z -boson and the new Z' with a mixing angle of $\mathcal{O}(v_{EW}^2/M_{Z'}^2)$. After integrating out Z' , one has different effective dark matter couplings to u_L and d_L fields, which are proportional to their couplings to the Z boson. For this model, we have $c_6 = 0$ and

$$\xi = \frac{-\frac{1}{2} + \frac{1}{3} \sin^2 \theta_W}{\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W} \approx -2.7 \quad (\text{A.12})$$

and order of unity.

Appendix A.5. Simplified model corresponding to dimension-5 EFT operator

As an example of a simplified model corresponding to the dimension-5 EFT operator described in Section 3.2, we consider a Higgs portal with a scalar mediator. Models of this kind are among the most concise versions of simplified models that produce couplings of Dark Matter to pairs of gauge-bosons. Scalar fields may couple directly to pairs of electroweak gauge bosons, but must carry part of the electroweak vacuum expectation value. One may thus consider a simple model where Dark Matter couples to a scalar singlet mediator, which mixes with the fields in the Higgs sector.

$$L \subset \frac{1}{2} m_s S^2 + \lambda S^2 |H|^2 + \lambda' S |H|^2 + y S \chi \bar{\chi} \quad (\text{A.13})$$

Where H is a field in the Higgs sector that contains part of the electroweak vacuum expectation value, S is a heavy scalar singlet and χ is a Dark Matter field. There is then an s -channel diagram where DM pairs couple to the singlet field S , which then mixes with a Higgs-sector field, and couples to W and Z bosons. This diagram contains 2 insertions of EW symmetry breaking fields, corresponding in form to the effective dimension-5 operator in Section 3.2.1.

Appendix A.6. Inert two-Higgs Doublet Model (IDM)

For most of the simplified models included in this report, the mass of the mediator and couplings/width are non-trivial parameters of the model. In these scenarios, we remain agnostic about the theory behind the dark matter sector and try to parameterize it in simple terms.

We have not addressed how to extend the simplified models to realistic and
 2225 viable models which are consistent with the symmetries of the Standard Model.
 Simplified models often violate gauge invariance which is a crucial principle for
 building a consistent BSM model which incorporates SM together with new physics.
 For example, with a new heavy gauge vector boson mediating DM interactions,
 one needs not just the dark matter and its mediator, but also a mechanism which
 2230 provides mass to this mediator in a gauge invariant way.

Considering both the simplified model and other elements necessary for a con-
 sistent theory is a next logical step. The authors of [158] term these Minimal Con-
 sistent Dark Matter (MCDM) models. MCDM models are at the same time still
 toy models that can be easily incorporated into a bigger BSM model and explored
 2235 via complementary constraints from collider and direct/indirect DM search experi-
 ments as well as relic density constraints.

The idea of an inert Two-Higgs Doublet Model (IDM) was introduced more than
 30 years ago in Ref [159]. The IDM was first proposed as a Dark Matter model in
 Ref. [160] and its phenomenology further studied in Refs. [161, 162, 163, 164, 165,
 14, 166, 167, 168, 169, 170, 158]. It is an extension of the SM with a second scalar
 doublet ϕ_2 with no direct coupling to fermions. This doublet has a discrete Z_2
 symmetry, under which ϕ_2 is odd and all the other fields are even. The Lagrangian
 of the odd sector is,

$$\mathcal{L} = \frac{1}{2}(D_\mu \phi_2)^2 - V(\phi_1, \phi_2) \quad (\text{A.14})$$

with the potential V containing mass terms and $\phi_1 - \phi_2$ interactions:

$$\begin{aligned} V = & -m_1^2(\phi_1^\dagger \phi_1) - m_2^2(\phi_2^\dagger \phi_2) + \lambda_1(\phi_1^\dagger \phi_1)^2 + \lambda_2(\phi_2^\dagger \phi_2)^2 \\ & + \lambda_3(\phi_2^\dagger \phi_2)(\phi_1^\dagger \phi_1) + \lambda_4(\phi_2^\dagger \phi_1)(\phi_1^\dagger \phi_2) \\ & + \frac{\lambda_5}{2}[(\phi_1^\dagger \phi_2)^2 + (\phi_2^\dagger \phi_1)^2], \end{aligned} \quad (\text{A.15})$$

where ϕ_1 and ϕ_2 are SM and inert Higgs doublets respectively carrying the same
 hypercharge. These doublets can be parameterized as

$$\phi_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + H \end{pmatrix} \quad \phi_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2}h^+ \\ h_1 + ih_2 \end{pmatrix} \quad (\text{A.16})$$


In addition to the SM, the IDM introduces four more degrees of freedom coming from the inert doublet in the form of a Z_2 -odd charged scalar h^\pm and two neutral Z_2 -odd scalars h_1 and h_2 . The lightest neutral scalar, h_1 is identified as the dark matter candidate. Aspects of the IDM collider phenomenology have been studied in [171, 172, 173, 158, 160, 174, 175, 176, 177, 178, 179, 180, 181, 182]. Its LHC signatures include dileptons [176, 182], trileptons [177] and multileptons [178] along with missing transverse energy, modifications of the Higgs branching ratios [179, 180, 170], as well as \cancel{E}_T + jet, Z, and Higgs and \cancel{E}_T + VBF signals (see Figs. A.74–A.79).

Based on the various LHC search channels, DM phenomenology issues and theoretical considerations, numerous works have proposed benchmark scenarios for the IDM, see e.g. [178, 170] while a FeynRules implementation (including MadGraph, CalcHEP and micrOMEGAs model files) was provided in [178]. An updated analysis of the parameter space has recently been performed in Ref. [158].


The authors suggested to study mono-X signatures that are relevant to model-independent collider DM searches, and evaluated their rates presented below. They have implemented and cross-checked the IDM model into CalcHEP and micrOMEGAs, with an implementation publicly available on the HEPMDB database, including loop-induced HHG and $\gamma\gamma H$ models. They propose an additional set of benchmark points, mostly inspired by mono-X and VBF searches (Table. A.9). Though the overall parameter space of IDM is 5-dimensional, once all relevant constraints are applied the parameter space relevant to a specific LHC signature typically reduces to 1-2 dimensional. In the mono-jet case, one can use two separate simplified models, a $gg \rightarrow h_1 h_1 + g$ process (via Higgs mediator) and a $qq \rightarrow h_1 h_2 + g$ ($gq \rightarrow h_1 h_2 + q$) process (through a Z-boson mediator) to capture the physics relevant to the search. The cross sections for the various mono-X and VBF signatures produced by this model are displayed in Fig. A.80.

BM	1	2	3	4	5
M_{h_1} (GeV)	48	53	70	82	120
M_{h_2} (GeV)	55	189	77	89	140
$M_{h_{\pm}}$ (GeV)	130	182	200	150	200
λ_2	0.8	1.0	1.1	0.9	1.0
λ_{345}	-0.010	-0.024	+0.022	-0.090	-0.100
Ωh^2	3.4×10^{-2}	8.1×10^{-2}	9.63×10^{-2}	1.5×10^{-2}	2.1×10^{-3}
σ_{SI} (pb)	2.3×10^{-10}	7.9×10^{-10}	5.1×10^{-10}	4.5×10^{-10}	2.6×10^{-9}
σ_{LHC} (fb)	1.7×10^2	7.7×10^2	4.3×10^{-2}	1.2×10^{-1}	2.3×10^{-2}


Table A.9: Five benchmarks for IDM in $(M_{h_1}, M_{h_2}, M_{h_{\pm}}, \lambda_2, \lambda_{345})$ parameter space. We also present the corresponding relic density (Ωh^2), the spin-independent cross section for DM scattering on the proton (σ_{SI}), and the LHC cross section at 13 TeV for mono-jet process $pp \rightarrow h_1, h_1 + jet$ for $p_T^{jet} > 100$ GeV cut (σ_{LHC}).




(a) $1 \rightarrow 2$ jets



(b) $2 \rightarrow 3$ jets



(c) $3 \rightarrow 4$ jets



(d) $4 \rightarrow 5$ jets

Figure 64: Distributions of differential jet rates $\frac{dN_{i \rightarrow j}}{d \log_{10}(k_{\text{cut}})}$ for EFT D5 sample with CKKW-L matching scale at 30, 50, 70, 80 and 90 GeV. A zoom of the region around the matching scale values is shown on right.






Figure 65: Missing transverse momentum distributions for EFT D5 sample with CKKW-L matching scale at 80 GeV. Individual contributions from the 0-, 1- and 2-parton emission samples are shown.



(a) No jet multiplicity cut



(b) $N_{\text{jet}} \leq 3$

Figure 66: Missing transverse momentum distributions for EFT D5 sample with CKKW-L matching scale at 80 GeV produced with maximum 1 (black), 2 (red) and 3 (blue) partons emitted at the generator level. The ratios are shown with respect to the latter sample.




Figure 67: Multiplicity of jets with $p_T > 30$ GeV and $|\eta| < 2.8$ for EFT D5 sample with CKKW-L matching scale at 80 GeV produced with maximum 1 (black), 2 (red) and 3 (blue) partons emitted at the generator level. The ratios are shown with respect to the latter sample. The leading jet p_T is required to be larger than 250 GeV.




Figure 68: Comparison of the jet multiplicity (left) and angular correction $\Delta R(j_1, j_2)$ (right) for the $\text{DM} + b\bar{b}$ scalar model generated in the 4-flavor and 5-scheme. The samples are generated for $m_\chi = 1$ GeV and $m_\phi = 10$ GeV.







Figure 69: 95% CL lower limits on the scale of the interaction of the D5 operator at 14 TeV, after the two truncation procedures. The procedure from [144] outlined in Section 5.1.2 is shown in blue, while the procedure from Refs. [143, 7], described in Section 5.1.1 is shown in red. Only the region between the dashed and the solid lines is excluded. Even though the intersection between the two lines is not shown in this plot, it should be noted that no limit can be set anymore for sufficiently low couplings, whatever truncation method is used.




(a) \cancel{E}_T




(b) Center of mass energy E_{cm}



(c) Mediator transverse momentum




(d) Leading DM transverse momentum




(e) DM transverse momentum (sub-leading)


Figure 70: Comparison of the kinematic distributions at 14 TeV between a narrow s -channel mediator and the corresponding D5 contact operator, at generator level for a jet+ \cancel{E}_T signature.




(a) \cancel{E}_T , D5 operator



(b) Invariant mass of the two WIMPs $m_{\chi\chi}$, D5 operator



(c) \cancel{E}_T , simplified model



(d) Invariant mass of the two WIMPs $m_{\chi\chi}$, simplified model

Figure 71: Comparison of the kinematic distributions for a narrow s -channel mediator, at generator level for a jet+ \cancel{E}_T signature, for varying DM masses.




Figure A.72: Feynman diagrams of leading order processes leading to monotop events: production of a colored scalar resonance φ decaying into a top quark and a spin-1/2 fermion χ (a), s - (b) and t -channel (c) non resonant production of a top quark in association with a spin-1 boson V decaying invisibly.




Figure A.73: Distributions of the transverse momentum of the V boson in the case of the process $pp \rightarrow tV \rightarrow t(t\bar{u} + \text{c.c.})$. We have imposed that the V -boson is produced on-shell and have chosen its mass to be $m_V = 200, 600$ and 1000 GeV (left, central and right panels). We have considered three possible cases for the total width of the V -boson, which has been fixed to 0.61%, 0.1% and 10% of the mass.




Figure A.74: Feynman diagrams for $gg \rightarrow h_1 h_1 + g$ process contributing to mono-jet signature, adapted from [158].




Figure A.75: Feynman diagrams for $q\bar{q} \rightarrow h_1 h_2 + g$ ($gq \rightarrow h_1 h_2 + q$) process contributing to mono-jet signature, adapted from [158].




Figure A.76: Feynman diagrams for $q\bar{q} \rightarrow h_1 h_1 + Z$ process contributing to mono-Z signature, adapted from [158].




Figure A.77: Feynman diagrams for $gg \rightarrow h_1 h_1 + H$ process contributing to mono-Higgs signature, adapted from [158].




Figure A.78: Feynman diagrams for $q\bar{q} \rightarrow h_1 h_2 + H$ process contributing to mono-Higgs signature, adapted from [158].




Figure A.79: Diagrams for $qq \rightarrow qq h_1 h_1$ DM production in vector boson fusion process, adapted from [158].




Figure A.80: LHC cross section at 13 TeV for various signatures, from [158].

2265 **Appendix B. Appendix: Presentation of experimental results for reinterpretation**

When collider searches present results with the recommended benchmarks, we suggest the following:

- 2270 • Provide limits in collider language, on fundamental parameters of the interaction: the couplings and masses of particles in simplified model.
- Translate limits to non-collider language, for a range of assumptions, in order to convey a rough idea of the range of possibilities. The details of this point are left for work beyond the scope of this Forum.
- 2275 • Provide all necessary material for theorists to reinterpret simplified model results as building blocks for more complete models (e.g. signal cutflows, acceptances, etc). This point is detailed further in this appendix.
- Provide model-independent results in terms of limits on cross-section times efficiency times acceptance of new phenomena for all cases, but especially when EFTs are employed as benchmarks. This recommendation has been
2280 issued before: see Ref. [183] for detailed suggestions.
- Provide easily usable and clearly labeled results in a digitized format, e.g. [184] entries, ROOT histograms and macros or tables available on analysis public pages.

2285 This appendix describes further considerations for reinterpretation and reimplementation of the analyses, as well as for the use of simplified model results directly given by the collaborations.

Appendix B.1. Reinterpretation of analyses

In the case of reinterpretation for models different than those provided by the experimental collaborations, the information needed primarily includes expected
2290 and observed exclusion lines along with their $\pm 1\sigma$ uncertainty, expected and observed upper limits in case of simplified models, efficiency maps and kinematic

distributions as reported in the analysis. If the kinematics of the new model to be tested in the reinterpretation is similar to that of the original model provided by the collaboration, it will be straight-forward to rescale the results provided to match
2295 the new model cross-section using this information.

Appendix B.2. Reimplementation of analyses

One of the important developments in recent years is an active development of software codes [83, 82, 185, 186, 187, 188] necessary for recasting analyses. The aim of these codes is to provide a public library of LHC analyses that have been reim-
2300 plemented and validated, often by the collaborations themselves. Such libraries can then be used to analyze validity of a BSM scenario in a systematic and effective manner. The availability of public libraries further facilitates a unified framework and can lead to an organized and central structure to preserve LHC information long term. The reimplementation of an analysis consists of several stages. Typically,
2305 the analysis note is used as a basis for the implementation of the preselection and event selection cuts in the user analysis code within the recasting frameworks. Signal events are generated, and passed through a parameterized detector simulation using software such as Delphes or PGS [101, 189]. The reconstructed objects are then analyzed using the code written in the previous step, and the results in terms
2310 of number of events are passed through a statistical analysis framework to compare with the backgrounds provided by the collaborations.

In order to be able to effectively use such codes, it is important to get a complete set of information from the collaborations.

For what concerns the generation of the models, it is desirable to have the fol-
2315 lowing items as used by the collaborations:

- Monte Carlo generators: Monte Carlo generators along with the exact versions used to produce the event files should be listed.
- Production cross sections: The order of production cross sections (e.g. LO,NLO,NLL) as well as the codes which were used to compute them should be provided.

- 2320 Tables of reference cross sections for several values of particle masses are useful as well.
- Process Generation: Details of the generated process, detailing number of additional partons generated.
 - LHE files: selected LHE files (detailing at least a few events if not the entire file) corresponding to the benchmarks listed in the analysis could also be made available in order to cross check process generation. Experimental collaborations may generate events on-the-fly without saving the intermediate LHE file; we advocate that the cross-check of process generation is straightforward if this information is present, so we encourage the generation of a few selected benchmark points allowing for a LHE file to be saved. Special attention should be paid to list the parameters which change the production cross section or kinematics of the process e.g. mixing angles.
 - Process cards: Process cards including PDF choices, details of matching algorithms and scales and details of process generation. If process cards are not available, the above items should be clearly identified.
 - Model files: For models which are not already implemented in MADGRAPH5_AMC@NLO, the availability of the corresponding model files in the UFO format [75] is highly desired. This format details the exact notation used in the model and hence sets up a complete framework. In case MADGRAPH5_AMC@NLO is not used, enough information should be provided in order to clearly identify the underlying model used for interpretations and reproduce the generation.

The ATLAS/CMS Dark Matter Forum provides most of the information needed within its SVN repository [56] and on a dedicated HEPData [57] page dedicated to the results in this report.

- 2345 Efficiency maps and relevant kinematic distributions as reported in the analysis should be provided, in a digitized format with clearly specified units. If selection criteria cannot be easily simulated through parameterized detector simulation, the collaborations should provide the efficiency of such cuts. Overall reconstruction

and identification efficiencies of physics objects are given as an input to the detector simulation software. It is thus very useful to get parametrized efficiencies for reconstructed objects (as a function of the rapidity η and/or transverse momentum p_T), along with the working points at which they were evaluated (e.g. loose, tight selection). Object definitions should be clearly identifiable. Digitized kinematic distributions are often necessary for the validation of the analysis so that the results from the collaboration are obtained, and so are tables containing the events passing each of the cuts.

The availability of digitized data and backgrounds is one of the primary requirements for fast and efficient recasting. Platforms such as HepData [184] can be used as a centralized repository; alternatively, analysis public pages and tables can be used for dissemination of results. Both data and Standard Model backgrounds should be provided in the form of binned histogram that can be interpolated if needed.

A detailed description of the likelihood used in order to derive the limits from the comparison of data to signal plus background should be given. This can be inferred from the analysis documentation itself, however direct availability of the limit setting code as a workspace in RooStats or HistFitter [190] is highly desirable.

Finally, the collaborations can also provide an analysis code directly implemented in one of the public recasting codes detailed above. Such codes can be published via INSPIRE [191] in order to track versioning and citations.

Appendix B.3. Simplified model interpretations

Dark Matter searches at the LHC will include simplified model interpretations in their search results. These interpretations are simple and can be used for a survey of viability of parameter space. Codes such as [192, 193, 194] can make use of the simplified model results given in the form of 95% Confidence Level (CLs) upper limit or efficiency maps in order to test Beyond the Standard Model parameter space. As mentioned above, it will thus be extremely useful if the results are given in a digitized form that is easily usable by the theory community.

The parameter space of these models should be clearly specified. For example, for a simplified model containing dark matter mass m_χ , mediator mass M_{med} and couplings g_χ, g_q it will be very useful to have upper limits on the product of couplings $\sqrt{g_\chi g_q}$ or cross section times branching ratio as a function of m_χ, M_{med} . Limits on visible cross sections of the simplified models considered for interpretations should be made available.

The usage of simplified model results relies on interpolating between upper limit values. In order to facilitate the interpolation, regions where large variation of upper limits is observed should contain denser grid, if a uniform grid over the entire plane is not possible. For simplified model involving more than three parameters (two masses and product of couplings), slices of upper limits in the additional dimensions will be necessary for reinterpretation.

As already mentioned in the introduction to this Chapter, acceptance and efficiency maps for all the signal regions involved in the analysis should be made available. These results are not only useful for model testing using simplified models but also to validate implementation of the analysis. Information about the most sensitive signal regions as a function of new particle masses is also useful in order to determine the validity of approximate limit setting procedures commonly used by theorists.

Appendix C. Appendix: Additional details and studies within the Forum

Further information for baryonic Z' Model

Cross-section scaling

The dependence of the cross section of the $pp \rightarrow H\chi\bar{\chi} + X$ process on $g_{hZ'Z'}$ is shown in Figure C.81. The curves have been fit to second-order polynomials, where y is the cross-section and x is the coupling $g_{hZ'Z'}$.

For $m_{\text{med}} = 100$ GeV, the fit function is

$$y = -0.12 - 3.4 \times 10^{-3}x + 2.7 \times 10^{-4}x^2. \quad (\text{C.1})$$

For $M_{\text{med}} = 1$ TeV, the fit function is is:

$$y = 0.0012 - 2.4 \times 10^{-7} x + 1.5 \times 10^{-7} x^2. \quad (\text{C.2})$$




Figure C.81: Cross section of the $pp \rightarrow H\chi\bar{\chi}$ process as a function of $g_{hZ'Z'}$ for $m_{Z'} = 100$ GeV (left) and $m_{Z'} = 1$ TeV (right). The fit functions are shown in the text.

References

- 2405 [1] A. Birkedal, K. Matchev, M. Perelstein, Dark matter at colliders: A Model independent approach, Phys.Rev. D70 (2004) 077701. [arXiv:hep-ph/0403004](#), doi:10.1103/PhysRevD.70.077701.
- [2] J. L. Feng, S. Su, F. Takayama, Lower limit on dark matter production at the large hadron collider, Phys.Rev.Lett. 96 (2006) 151802. [arXiv:hep-ph/0503117](#), doi:10.1103/PhysRevLett.96.151802.
- 2410 [3] F. J. Petriello, S. Quackenbush, K. M. Zurek, The Invisible Z' at the CERN LHC, Phys.Rev. D77 (2008) 115020. [arXiv:0803.4005](#), doi:10.1103/PhysRevD.77.115020.
- [4] M. Beltran, D. Hooper, E. W. Kolb, Z. A. Krusberg, T. M. Tait, Maverick dark matter at colliders, JHEP 1009 (2010) 037. [arXiv:1002.4137](#), doi:10.1007/JHEP09(2010)037.
- 2415

- [5] Y. Bai, P. J. Fox, R. Harnik, The Tevatron at the frontier of dark matter direct detection, JHEP 1012 (2010) 048. [arXiv:1005.3797](#), [doi:10.1007/JHEP12\(2010\)048](#).
- 2420 [6] T. Aaltonen, et al., A search for dark matter in events with one jet and missing transverse energy in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys.Rev.Lett. 108 (2012) 211804. [arXiv:1203.0742](#), [doi:10.1103/PhysRevLett.108.211804](#).
- 2425 [7] ATLAS Collaboration, Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector Submitted to Eur.Phys.J. C. [arXiv:1502.01518](#).
- 2430 [8] ATLAS Collaboration, Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, Phys.Rev. D91 (2015) 012008. [arXiv:1411.1559](#), [doi:10.1103/PhysRevD.91.012008](#).
- 2435 [9] ATLAS Collaboration, Search for new particles in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, JHEP 1409 (2014) 037. [arXiv:1407.7494](#), [doi:10.1007/JHEP09\(2014\)037](#).
- [10] ATLAS Collaboration, Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector, Phys.Rev. D90 (2014) 012004. [arXiv:1404.0051](#), [doi:10.1103/PhysRevD.90.012004](#).
- 2440 [11] ATLAS Collaboration, Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, Phys.Rev.Lett. 112 (2014) 041802. [arXiv:1309.4017](#), [doi:10.1103/PhysRevLett.112.041802](#).

- [12] ATLAS Collaboration, Search for invisible particles produced in association
 2445 with single-top-quarks in proton-proton collisions at $\sqrt{s} = 8$ TeV with the
 ATLAS detector, Eur.Phys.J. C75 (2015) 79. arXiv:1410.5404, doi:10.
 1140/epjc/s10052-014-3233-4.
- [13] ATLAS Collaboration, Search for dark matter in events with heavy quarks
 and missing transverse momentum in pp collisions with the ATLAS detec-
 2450 tor, Eur.Phys.J. C75 (2015) 92. arXiv:1410.4031, doi:10.1140/epjc/
 s10052-015-3306-z.
- [14] ATLAS Collaboration, Sensitivity to WIMP dark matter in the final states
 containing jets and missing transverse momentum with the ATLAS detector
 at 14 TeV LHC, Tech. Rep. ATL-PHYS-PUB-2014-007, CERN, Geneva (Jun
 2455 2014).
 URL <http://cds.cern.ch/record/1708859>
- [15] CMS Collaboration, Search for dark matter, extra dimensions, and un-
 particles in monojet events in proton-proton collisions at $\sqrt{s} = 8$ TeV,
 Eur.Phys.J. C75 (2015) 235. arXiv:1408.3583, doi:10.1140/epjc/
 2460 s10052-015-3451-4.
- [16] CMS Collaboration, Search for new phenomena in monophoton final states
 in proton-proton collisions at $\sqrt{s} = 8$ TeV Submitted to Phys.Lett.B. arXiv:
 1410.8812.
- [17] CMS Collaboration, Search for physics beyond the standard model in final
 2465 states with a lepton and missing transverse energy in proton-proton collisions
 at $\sqrt{s} = 8$ TeV, Phys.Rev. D91 (2015) 092005. arXiv:1408.2745, doi:
 10.1103/PhysRevD.91.092005.
- [18] CMS Collaboration, Search for monotop signatures in proton-proton colli-
 sions at $\sqrt{s} = 8$ TeV, Phys.Rev.Lett. 114 (2015) 101801. arXiv:1410.1149,
 2470 doi:10.1103/PhysRevLett.114.101801.

- [19] CMS Collaboration, Search for the production of dark matter in association with top-quark pairs in the single-lepton final state in proton-proton collisions at $\sqrt{s} = 8$ TeV Submitted to JHEP. [arXiv:1504.03198](#).
- 2475 [20] CMS Collaboration, Search for the production of dark matter in association with top quark pairs in the di-lepton final state in pp collisions at $\sqrt{s} = 8$ TeV CMS-PAS-B2G-13-004.
URL <http://cds.cern.ch/record/1697173>
- 2480 [21] CMS Collaboration, Search for dark matter direct production using razor variables in events with two or more jets in pp collisions at 8 TeV CMS-PAS-EXO-14-004.
URL <http://cds.cern.ch/record/2002861>
- [22] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. Tait, et al., Constraints on light Majorana dark matter from colliders, Phys.Lett. B695 (2011) 185–188. [arXiv:1005.1286](#), doi:10.1016/j.physletb.2010.11.009.
- 2485 [23] J. Goodman, M. Ibe, A. Rajaraman, W. Shepherd, T. M. Tait, et al., Constraints on dark matter from colliders, Phys.Rev. D82 (2010) 116010. [arXiv:1008.1783](#), doi:10.1103/PhysRevD.82.116010.
- 2490 [24] I. M. Shoemaker, L. Vecchi, Unitarity and monojet bounds on models for DAMA, CoGeNT, and CRESST-II, Phys.Rev. D86 (2012) 015023. [arXiv:1112.5457](#), doi:10.1103/PhysRevD.86.015023.
- [25] J. Kopp, Collider limits on dark matter [arXiv:1105.3248](#).
- [26] P. J. Fox, R. Harnik, J. Kopp, Y. Tsai, LEP shines light on dark matter, Phys.Rev. D84 (2011) 014028. [arXiv:1103.0240](#), doi:10.1103/PhysRevD.84.014028.
- 2495 [27] P. J. Fox, R. Harnik, J. Kopp, Y. Tsai, Missing energy signatures of dark matter at the LHC, Phys.Rev. D85 (2012) 056011. [arXiv:1109.4398](#), doi:10.1103/PhysRevD.85.056011.

- 2500 [28] G. Busoni, A. De Simone, E. Morgante, A. Riotto, On the validity of the effective field theory for dark matter searches at the LHC, *Phys.Lett. B* 728 (2014) 412–421. [arXiv:1307.2253](#), [doi:10.1016/j.physletb.2013.11.069](#).
- [29] J. Alwall, P. Schuster, N. Toro, Simplified models for a first characterization of new physics at the LHC, *Phys.Rev. D* 79 (2009) 075020. [arXiv:0810.3921](#), [doi:10.1103/PhysRevD.79.075020](#).
- 2505 [30] J. Goodman, W. Shepherd, LHC bounds on UV-complete models of dark matter [arXiv:1111.2359](#).
- [31] D. Alves, et al., Simplified models for LHC new physics searches, *J.Phys. G* 39 (2012) 105005. [arXiv:1105.2838](#), [doi:10.1088/0954-3899/39/10/105005](#).
- 2510 [32] H. An, X. Ji, L.-T. Wang, Light dark matter and Z' dark force at colliders, *JHEP* 1207 (2012) 182. [arXiv:1202.2894](#), [doi:10.1007/JHEP07\(2012\)182](#).
- [33] H. An, R. Huo, L.-T. Wang, Searching for low mass dark portal at the LHC, *Phys.Dark Univ.* 2 (2013) 50–57. [arXiv:1212.2221](#), [doi:10.1016/j.dark.2013.03.002](#).
- 2515 [34] A. DiFranzo, K. I. Nagao, A. Rajaraman, T. M. P. Tait, Simplified models for dark matter interacting with quarks, *JHEP* 1311 (2013) 014. [arXiv:1308.2679](#), [doi:10.1007/JHEP11\(2013\)014](#), [doi:10.1007/JHEP01\(2014\)162](#).
- 2520 [35] O. Buchmuller, M. J. Dolan, C. McCabe, Beyond effective field theory for dark matter searches at the LHC, *JHEP* 1401 (2014) 025. [arXiv:1308.6799](#), [doi:10.1007/JHEP01\(2014\)025](#).
- [36] Y. Bai, J. Berger, Fermion Portal dark matter, *JHEP* 1311 (2013) 171. [arXiv:1308.0612](#), [doi:10.1007/JHEP11\(2013\)171](#).

- 2525 [37] Y. Bai, J. Berger, Lepton Portal dark matter, JHEP 1408 (2014) 153. [arXiv:1402.6696](#), doi:10.1007/JHEP08(2014)153.
- [38] H. An, L.-T. Wang, H. Zhang, Dark matter with t -channel mediator: a simple step beyond contact interaction, Phys.Rev. D89 (2014) 115014. [arXiv:1308.0592](#), doi:10.1103/PhysRevD.89.115014.
- 2530 [39] J. Abdallah, et al., Simplified models for dark matter and missing energy searches at the LHC [arXiv:1409.2893](#).
- [40] S. Malik, C. McCabe, H. Araujo, A. Belyaev, C. Boehm, et al., Interplay and characterization of dark matter searches at colliders and in direct detection experiments [arXiv:1409.4075](#).
- 2535 [41] P. Harris, V. V. Khoze, M. Spannowsky, C. Williams, Constraining dark sectors at colliders: Beyond the effective theory approach, Phys.Rev. D91 (2015) 055009. [arXiv:1411.0535](#), doi:10.1103/PhysRevD.91.055009.
- [42] M. R. Buckley, D. Feld, D. Goncalves, Scalar simplified models for dark matter, Phys.Rev. D91 (2015) 015017. [arXiv:1410.6497](#), doi:10.1103/PhysRevD.91.015017.
- 2540 [43] U. Haisch, E. Re, Simplified dark matter top-quark interactions at the LHC, JHEP 1506 (2015) 078. [arXiv:1503.00691](#), doi:10.1007/JHEP06(2015)078.
- [44] Y. Bai, T. M. Tait, Searches with mono-leptons, Phys.Lett. B723 (2013) 384–387. [arXiv:1208.4361](#), doi:10.1016/j.physletb.2013.05.057.
- 2545 [45] L. M. Carpenter, A. Nelson, C. Shimmin, T. M. Tait, D. Whiteson, Collider searches for dark matter in events with a Z boson and missing energy, Phys.Rev. D87 (2013) 074005. [arXiv:1212.3352](#), doi:10.1103/PhysRevD.87.074005.
- 2550 [46] N. F. Bell, J. B. Dent, A. J. Galea, T. D. Jacques, L. M. Krauss, et al., Searching for dark matter at the LHC with a mono-Z, Phys.Rev. D86 (2012) 096011. [arXiv:1209.0231](#), doi:10.1103/PhysRevD.86.096011.

- [47] A. A. Petrov, W. Shepherd, Searching for dark matter at LHC with mono-Higgs production, Phys.Lett. B730 (2014) 178–183. [arXiv:1311.1511](#),
2555 [doi:10.1016/j.physletb.2014.01.051](#).
- [48] L. Carpenter, A. DiFranzo, M. Mulhearn, C. Shimmin, S. Tulin, et al., Mono-Higgs-boson: A new collider probe of dark matter, Phys.Rev. D89 (2014) 075017. [arXiv:1312.2592](#), [doi:10.1103/PhysRevD.89.075017](#).
- [49] J. Abdallah, H. Araujo, A. Arbey, A. Ashkenazi, A. Belyaev, et al., Simplified
2560 models for dark matter searches at the LHCSubmitted to Phys.Dark Univ. [arXiv:1506.03116](#).
- [50] D. Bauer, J. Buckley, M. Cahill-Rowley, R. Cotta, A. Drlica-Wagner, et al., Dark matter in the coming decade: Complementary paths to discovery and beyond, Phys.Dark Univ. 7-8 (2013) 16–23. [arXiv:1305.1605](#), [doi:10.1016/j.dark.2015.04.001](#).
2565
- [51] R. S. Chivukula, H. Georgi, Composite Technicolor Standard Model, Phys.Lett. B188 (1987) 99. [doi:10.1016/0370-2693\(87\)90713-1](#).
- [52] L. Hall, L. Randall, Weak scale effective supersymmetry, Phys.Rev.Lett. 65 (1990) 2939–2942. [doi:10.1103/PhysRevLett.65.2939](#).
- [53] A. Buras, P. Gambino, M. Gorbahn, S. Jager, L. Silvestrini, Universal unitarity
2570 triangle and physics beyond the Standard Model, Phys.Lett. B500 (2001) 161–167. [arXiv:hep-ph/0007085](#), [doi:10.1016/S0370-2693\(01\)00061-2](#).
- [54] G. D’Ambrosio, G. Giudice, G. Isidori, A. Strumia, Minimal Flavor Violation: An effective field theory approach, Nucl.Phys. B645 (2002) 155–187.
2575 [arXiv:hep-ph/0207036](#), [doi:10.1016/S0550-3213\(02\)00836-2](#).
- [55] P. Agrawal, M. Blanke, K. Gemmler, Flavored dark matter beyond Minimal Flavor Violation, JHEP 1410 (2014) 72. [arXiv:1405.6709](#), [doi:10.1007/JHEP10\(2014\)072](#).

- 2580 [56] SVN repository for model, input and parameter cards for models consid-
ered in the ATLAS/CMS Dark Matter Forum studies, <https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/>, [Online; accessed 13-June-2015] (2015).
- [57] The Durham HepData project, atlas/cms dark matter forum repository,
2585 <http://hepdata.cedar.ac.uk/view/ins1381178>, [Online; in preparation] (2015).
- [58] O. Buchmueller, M. J. Dolan, S. A. Malik, C. McCabe, Characterising dark
matter searches at colliders and direct detection experiments: Vector me-
diators, JHEP 1501 (2015) 037. [arXiv:1407.8257](#), doi:10.1007/
2590 JHEP01(2015)037.
- [59] U. Haisch, F. Kahlhoefer, E. Re, QCD effects in mono-jet searches for dark
matter, JHEP 1312 (2013) 007. [arXiv:1310.4491](#), doi:10.1007/
JHEP12(2013)007.
- [60] T. Sjöstrand, S. Mrenna, P. Z. Skands, A brief introduction to PYTHIA 8.1,
2595 Comput.Phys.Commun. 178 (2008) 852–867. [arXiv:0710.3820](#), doi:
10.1016/j.cpc.2008.01.036.
- [61] Y. Bai, J. Bourbeau, T. Lin, Dark matter searches with a mono- Z' jet [arXiv:1504.01395](#).
- [62] M. Chala, F. Kahlhoefer, M. McCullough, G. Nardini, K. Schmidt-Hoberg,
2600 Constraining dark sectors with monojets and dijets [arXiv:1503.05916](#).
- [63] N. Zhou, D. Berge, D. Whiteson, Mono-everything: combined limits on
dark matter production at colliders from multiple final states, Phys.Rev.
D87 (2013) 095013. [arXiv:1302.3619](#), doi:10.1103/PhysRevD.87.
095013.
- 2605 [64] Y. Gershtein, F. Petriello, S. Quackenbush, K. M. Zurek, Discovering hid-
den sectors with mono-photon Z' searches, Phys.Rev. D78 (2008) 095002.
[arXiv:0809.2849](#), doi:10.1103/PhysRevD.78.095002.

- [65] N. F. Bell, Y. Cai, J. B. Dent, R. K. Leane, T. J. Weiler, Dark matter at the LHC: EFTs and gauge invariance *arXiv:1503.07874*.
- 2610 [66] K. Hamaguchi, S. P. Liew, T. Moroi, Y. Yamamoto, Isospin-violating dark matter with colored mediators, *JHEP* 1405 (2014) 086. *arXiv:1403.0324*, doi:10.1007/JHEP05(2014)086.
- [67] S. Alioli, P. Nason, C. Oleari, E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX, *JHEP* 2615 1006 (2010) 043. *arXiv:1002.2581*, doi:10.1007/JHEP06(2010)043.
- [68] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, *JHEP* 0411 (2004) 040. *arXiv:hep-ph/0409146*, doi:10.1088/1126-6708/2004/11/040.
- 2620 [69] S. Frixione, P. Nason, C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method, *JHEP* 0711 (2007) 070, * Temporary entry *. *arXiv:0709.2092*, doi:10.1088/1126-6708/2007/11/070.
- [70] C. Arina, E. Del Nobile, P. Panci, Dark matter with pseudoscalar-mediated 2625 interactions explains the DAMA signal and the Galactic Center excess, *Phys.Rev.Lett.* 114 (2015) 011301. *arXiv:1406.5542*, doi:10.1103/PhysRevLett.114.011301.
- [71] U. Haisch, A. Hibbs, E. Re, Determining the structure of dark-matter couplings at the LHC, *Phys.Rev. D* 89 (2014) 034009. *arXiv:1311.7131*, 2630 doi:10.1103/PhysRevD.89.034009.
- [72] H. M. Lee, M. Park, V. Sanz, Interplay between Fermi gamma-ray lines and collider searches, *JHEP* 1303 (2013) 052. *arXiv:1212.5647*, doi:10.1007/JHEP03(2013)052.
- 2635 [73] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, et al., The automated computation of tree-level and next-to-leading order differential cross sec-

- tions, and their matching to parton shower simulations, JHEP 1407 (2014) 079. [arXiv:1405.0301](#), [doi:10.1007/JHEP07\(2014\)079](#).
- [74] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, B. Fuks, FeynRules 2.0 - a complete toolbox for tree-level phenomenology, Comput.Phys.Commun. 185 (2014) 2250–2300. [arXiv:1310.1921](#).
- [75] C. Degrande, C. Duhr, B. Fuks, D. Grellscheid, O. Mattelaer, et al., UFO - the Universal FeynRules Output, Comput.Phys.Commun. 183 (2012) 1201–1214. [arXiv:1108.2040](#), [doi:10.1016/j.cpc.2012.01.022](#).
- [76] M. Papucci, A. Vichi, K. M. Zurek, Monojet versus the rest of the world I: t -channel models, JHEP 1411 (2014) 024. [arXiv:1402.2285](#), [doi:10.1007/JHEP11\(2014\)024](#).
- [77] S. Chang, R. Edezhath, J. Hutchinson, M. Luty, Effective WIMPs, Phys.Rev. D89 (2014) 015011. [arXiv:1307.8120](#), [doi:10.1103/PhysRevD.89.015011](#).
- [78] G. Busoni, A. De Simone, T. Jacques, E. Morgante, A. Riotto, On the validity of the effective field theory for dark matter searches at the LHC part III: Analysis for the t -channel, JCAP 1409 (2014) 022. [arXiv:1405.3101](#), [doi:10.1088/1475-7516/2014/09/022](#).
- [79] F. Maltoni, A. Martini, K. Mawatari, B. Oehl, Signals of a superlight gravitino at the LHC, JHEP 1504 (2015) 021. [arXiv:1502.01637](#), [doi:10.1007/JHEP04\(2015\)021](#).
- [80] P. de Aquino, F. Maltoni, K. Mawatari, B. Oehl, Light gravitino production in association with gluinos at the LHC, JHEP 1210 (2012) 008. [arXiv:1206.7098](#), [doi:10.1007/JHEP10\(2012\)008](#).
- [81] J. Alwall, S. de Visscher, F. Maltoni, QCD radiation in the production of heavy colored particles at the LHC, JHEP 0902 (2009) 017. [arXiv:0810.5350](#), [doi:10.1088/1126-6708/2009/02/017](#).

- [82] E. Conte, B. Dumont, B. Fuks, C. Wymant, Designing and recasting LHC analyses with MadAnalysis 5, Eur.Phys.J. C74 (2014) 3103. arXiv:1405.3982, doi:10.1140/epjc/s10052-014-3103-0.
- [83] B. Dumont, B. Fuks, S. Kraml, S. Bein, G. Chalons, et al., Toward a public analysis database for LHC new physics searches using MADANALYSIS 5, Eur.Phys.J. C75 (2015) 56. arXiv:1407.3278, doi:10.1140/epjc/s10052-014-3242-3.
- [84] E. Conte, B. Fuks, G. Serret, MadAnalysis 5, a user-friendly framework for collider phenomenology, Comput.Phys.Commun. 184 (2013) 222–256. arXiv:1206.1599, doi:10.1016/j.cpc.2012.09.009.
- [85] T. Lin, E. W. Kolb, L.-T. Wang, Probing dark matter couplings to top and bottom quarks at the LHC, Phys.Rev. D88 (2013) 063510. arXiv:1303.6638, doi:10.1103/PhysRevD.88.063510.
- [86] P. Agrawal, B. Batell, D. Hooper, T. Lin, Flavored dark matter and the Galactic Center gamma-ray excess, Phys.Rev. D90 (2014) 063512. arXiv:1404.1373, doi:10.1103/PhysRevD.90.063512.
- [87] T. Daylan, D. P. Finkbeiner, D. Hooper, T. Linden, S. K. N. Portillo, et al., The characterization of the gamma-ray signal from the central Milky Way: A compelling case for annihilating dark matter arXiv:1402.6703.
- [88] F. Calore, I. Cholis, C. Weniger, Background model systematics for the Fermi GeV excess, JCAP 1503 (2015) 038. arXiv:1409.0042, doi:10.1088/1475-7516/2015/03/038.
- [89] A. Kumar, S. Tulin, Top-flavored dark matter and the forward-backward asymmetry, Phys.Rev. D87 (9) (2013) 095006. arXiv:1303.0332, doi:10.1103/PhysRevD.87.095006.
- [90] B. Batell, T. Lin, L.-T. Wang, Flavored Dark Matter and R-parity violation, JHEP 1401 (2014) 075. arXiv:1309.4462, doi:10.1007/JHEP01(2014)075.

- [91] C. Kilic, M. D. Klimek, J.-H. Yu, Signatures of top flavored dark matter, Phys.Rev. D91 (5) (2015) 054036. [arXiv:1501.02202](#), [doi:10.1103/PhysRevD.91.054036](#).
- 2695 [92] H. M. Lee, M. Park, V. Sanz, Gravity-mediated (or composite) Dark Matter, Eur.Phys.J. C74 (2014) 2715. [arXiv:1306.4107](#), [doi:10.1140/epjc/s10052-014-2715-8](#).
- [93] H. M. Lee, M. Park, V. Sanz, Gravity-mediated (or composite) dark matter confronts astrophysical data, JHEP 1405 (2014) 063. [arXiv:1401.5301](#), [doi:10.1007/JHEP05\(2014\)063](#).
- 2700 [94] D. Y. Bardin, S. M. Bilenky, W. Beenakker, F. A. Berends, W. L. van Neerven, S. C. Van der Marck, G. Burgers, W. F. L. Hollik, T. Riemann, M. Sachwitz, Z line shape, in: Workshop on Z Physics at LEP1 : General Meetings, vol. 1 : Standard Physics, 1989, p. 43 p.
URL <http://cds.cern.ch/record/199969>
- 2705 [95] A. Nelson, L. M. Carpenter, R. Cotta, A. Johnstone, D. Whiteson, Confronting the Fermi Line with LHC data: an effective theory of dark matter interaction with photons, Phys.Rev. D89 (2014) 056011. [arXiv:1307.5064](#), [doi:10.1103/PhysRevD.89.056011](#).
- 2710 [96] A. Berlin, T. Lin, L.-T. Wang, Mono-Higgs detection of dark matter at the LHC, JHEP 1406 (2014) 078. [arXiv:1402.7074](#), [doi:10.1007/JHEP06\(2014\)078](#).
- [97] R. Cotta, J. Hewett, M. Le, T. Rizzo, Bounds on dark matter interactions with electroweak gauge bosons, Phys.Rev. D88 (2013) 116009. [arXiv:1210.0525](#), [doi:10.1103/PhysRevD.88.116009](#).
- 2715 [98] A. Crivellin, U. Haisch, A. Hibbs, LHC constraints on gauge boson couplings to dark matter [arXiv:1501.00907](#).

- [99] D. O’Connell, M. J. Ramsey-Musolf, M. B. Wise, Minimal extension of the Standard Model scalar sector, *Phys.Rev. D* 75 (2007) 037701. [arXiv: hep-ph/0611014](#), [doi:10.1103/PhysRevD.75.037701](#).
- 2720 [100] N. Craig, J. Galloway, S. Thomas, Searching for signs of the Second Higgs Doublet [arXiv:1305.2424](#).
- [101] J. de Favereau, et al., DELPHES 3, a modular framework for fast simulation of a generic collider experiment, *JHEP* 1402 (2014) 057. [arXiv:1307.6346](#), [doi:10.1007/JHEP02\(2014\)057](#).
- 2725 [102] A. Djouadi, A. Falkowski, Y. Mambrini, J. Quevillon, Direct detection of Higgs-Portal dark matter at the LHC, *Eur.Phys.J. C* 73 (2013) 2455. [arXiv:1205.3169](#), [doi:10.1140/epjc/s10052-013-2455-1](#).
- [103] N. Weiner, I. Yavin, How dark are Majorana WIMPs? signals from MiDM and Rayleigh dark matter, *Phys.Rev. D* 86 (2012) 075021. [arXiv:1206.2910](#),
2730 [doi:10.1103/PhysRevD.86.075021](#).
- [104] N. Weiner, I. Yavin, UV completions of magnetic inelastic and Rayleigh dark matter for the Fermi Line(s), *Phys.Rev. D* 87 (2013) 023523. [arXiv:1209.1093](#), [doi:10.1103/PhysRevD.87.023523](#).
- [105] J. Liu, B. Shuve, N. Weiner, I. Yavin, Looking for new charged states at the
2735 LHC: Signatures of magnetic and Rayleigh dark matter, *JHEP* 1307 (2013) 144. [arXiv:1303.4404](#), [doi:10.1007/JHEP07\(2013\)144](#).
- [106] U. Haisch, F. Kahlhoefer, J. Unwin, The impact of heavy-quark loops on LHC dark matter searches, *JHEP* 1307 (2013) 125. [arXiv:1208.4605](#), [doi:10.1007/JHEP07\(2013\)125](#).
- 2740 [107] M. Bahr, S. Gieseke, M. Gigg, D. Grellscheid, K. Hamilton, et al., Herwig++ Physics and Manual, *Eur.Phys.J. C* 58 (2008) 639–707. [arXiv:0803.0883](#), [doi:10.1140/epjc/s10052-008-0798-9](#).

- [108] G. Corcella, I. Knowles, G. Marchesini, S. Moretti, K. Odagiri, et al., HERWIG 6.5 release note [arXiv:hep-ph/0210213](https://arxiv.org/abs/hep-ph/0210213).
- 2745 [109] G. Corcella, et al., HERWIG 6: An event generator for hadron emission reactions with interfering gluons (including supersymmetric processes), JHEP 0101 (2001) 010. [arXiv:hep-ph/0011363](https://arxiv.org/abs/hep-ph/0011363), doi:10.1088/1126-6708/2001/01/010.
- 2750 [110] G. Marchesini, B. Webber, G. Abbiendi, I. Knowles, M. Seymour, L. Stanco, {HERWIG} 5.1 - a monte carlo event generator for simulating hadron emission reactions with interfering gluons, Computer Physics Communications 67 (3) (1992) 465 – 508. doi:[http://dx.doi.org/10.1016/0010-4655\(92\)90055-4](http://dx.doi.org/10.1016/0010-4655(92)90055-4).
URL <http://www.sciencedirect.com/science/article/pii/0010465592900554>
- 2755 [111] M. Backovi?, M. Krämer, F. Maltoni, A. Martini, K. Mawatari, M. Pellen, Higher-order QCD predictions for dark matter production at the LHC in simplified models with s-channel mediators [arXiv:1508.05327](https://arxiv.org/abs/1508.05327).
- [112] P. J. Fox, C. Williams, Next-to-leading order predictions for dark matter production at hadron colliders, Phys.Rev. D87 (2013) 054030. [arXiv:1211.6390](https://arxiv.org/abs/1211.6390), doi:10.1103/PhysRevD.87.054030.
- 2760 [113] SVN repository for POWHEG input card for model with s-channel exchange of axial vector mediator, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_DMA/, [Online; accessed 24-April-2015] (2015).
- 2765 [114] SVN repository for POWHEG input card for model with s-channel exchange of vector mediator, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_DMV/, [Online; accessed 24-April-2015] (2015).

- 2770 [115] SVN repository for POWHEG input card for model with s -channel exchange of scalar mediator, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_DMS_tLoop/, [Online; accessed 24-April-2015] (2015).
- [116] SVN repository for POWHEG input card for model with s -channel exchange
2775 of pseudo-scalar mediator, coupling to the quarks through a top loop, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_DMP_tloop/, [Online; accessed 24-April-2015] (2015).
- [117] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, et al., An introduction to PYTHIA 8.2, *Comput.Phys.Commun.* 191 (2015) 159–177.
2780 arXiv:1410.3012, doi:10.1016/j.cpc.2015.01.024.
- [118] S. Hoeche, S. Kuttimalai, S. Schumann, F. Siegert, Beyond Standard Model calculations with Sherpa, *Eur.Phys.J. C* 75 (3) (2015) 135. arXiv:1412.6478, doi:10.1140/epjc/s10052-015-3338-4.
- [119] R. Frederix, S. Frixione, Merging meets matching in MC@NLO, *JHEP* 1212
2785 (2012) 061. arXiv:1209.6215, doi:10.1007/JHEP12(2012)061.
- [120] J. Alwall, et al., Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions, *Eur.Phys.J. C* 53 (2008) 473–500. arXiv:0706.2569, doi:10.1140/epjc/s10052-007-0490-5.
- 2790 [121] L. Lonnblad, S. Prestel, Matching tree-level matrix elements with interleaved showers, *JHEP* 1203 (2012) 019. arXiv:1109.4829, doi:10.1007/JHEP03(2012)019.
- [122] SVN repository for Madgraph inputs with t -channel exchange of colored scalar mediator, couplings to light quarks only, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_tChannel/contributed_by_PapucciVichiZurek/, [Online; accessed 27-April-2015] (2015).
2795

- [123] SVN repository for Madgraph inputs with t -channel exchange of colored scalar mediator, couplings to all quark generations,
2800 https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/Monojet_tChannel/contributed_by_Amelia_Brennan/, [Online; accessed 27-April-2015] (2015).
- [124] SVN repository for Madgraph input cards for model with s -channel exchange of vector mediator, for electroweak boson final states, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/EW_DMV/,
2805 [Online; accessed 15-May-2015] (2015).
- [125] SVN repository for Madgraph inputs for model with s -channel exchange of pseudo-scalar mediator, produced in association with top quarks, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/HF_S+PS/,
2810 [Online; accessed 24-April-2015] (2015).
- [126] SVN repository for Madgraph inputs for simplified model with a colored scalar mediator coupling to DM and b -quarks, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/HF_S+PS/, [Online; accessed 24-April-2015] (2015).
- 2815 [127] J. M. Campbell, R. K. Ellis, F. Maltoni, S. Willenbrock, Associated production of a Z boson and a single heavy quark jet, Phys.Rev. D69 (2004) 074021. arXiv:hep-ph/0312024, doi:10.1103/PhysRevD.69.074021.
- [128] F. Maltoni, T. McElmurry, S. Willenbrock, Inclusive production of a Higgs or Z boson in association with heavy quarks, Phys.Rev. D72 (2005) 074024.
2820 arXiv:hep-ph/0505014, doi:10.1103/PhysRevD.72.074024.
- [129] J. M. Campbell, R. K. Ellis, F. Maltoni, S. Willenbrock, Production of a Z boson and two jets with one heavy-quark tag, Phys.Rev. D73 (2006) 054007. arXiv:hep-ph/0510362, doi:10.1103/PhysRevD.77.019903, 10.1103/PhysRevD.73.054007.

- 2825 [130] CMS Collaboration, Measurement of the production cross sections for a Z boson and one or more b jets in pp collisions at $\sqrt{s} = 7$ TeV, JHEP 1406 (2014) 120. [arXiv:1402.1521](#), [doi:10.1007/JHEP06\(2014\)120](#).
- [131] CMS Collaboration, Measurement of the cross section and angular correlations for associated production of a Z boson with b hadrons in pp collisions at $\sqrt{s} = 7$ TeV, JHEP 1312 (2013) 039. [arXiv:1310.1349](#), [doi:10.1007/JHEP12\(2013\)039](#).
2830
- [132] CMS Collaboration, Search for H/A decaying into Z+A/H, with Z to ll and A/H to fermion pair (2015).
- [133] R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, R. Pittau, et al., W and Z/γ^* boson production in association with a bottom-antibottom pair, JHEP 1109
2835 (2011) 061. [arXiv:1106.6019](#), [doi:10.1007/JHEP09\(2011\)061](#).
- [134] M. Wiesemann, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, et al., Higgs production in association with bottom quarks, JHEP 1502 (2015) 132. [arXiv:1409.5301](#), [doi:10.1007/JHEP02\(2015\)132](#).
- 2840 [135] SVN repository for Madgraph inputs for vector and scalar mediator models leading to a mono-Higgs signature, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/EW_Higgs_all/, [Online; accessed 24-April-2015] (2015).
- [136] SVN repository for Madgraph inputs for 2HDM model leading to a mono-Higgs signature, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/EW_Higgs_2HDM/, [Online; accessed 12-May-2015] (2015).
2845
- [137] B. Hespel, D. Lopez-Val, E. Vryonidou, Higgs pair production via gluon fusion in the Two-Higgs-Doublet Model, JHEP 1409 (2014) 124. [arXiv:1407.0281](#), [doi:10.1007/JHEP09\(2014\)124](#).
2850

- [138] U. Baur, E. N. Glover, Higgs boson production at large transverse momentum in hadronic collisions, Nucl.Phys. B339 (1990) 38–66. doi:10.1016/0550-3213(90)90532-I.
- 2855 [139] SVN repository for Madgraph inputs for dimension-7 EFT models with direct DM-EW boson couplings, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/EW_Fermion_D7/contributed_by_Renjie_Wang/, [Online; accessed 24-April-2015] (2015).
- 2860 [140] S. Nussinov, Technoc cosmology: could a technibaryon excess provide a 'natural' missing mass candidate?, Phys.Lett. B165 (1985) 55. doi:10.1016/0370-2693(85)90689-6.
- [141] D. B. Kaplan, A single explanation for both the baryon and dark matter densities, Phys.Rev.Lett. 68 (1992) 741–743. doi:10.1103/PhysRevLett.68.741.
- 2865 [142] T. Banks, J.-F. Fortin, S. Thomas, Direct detection of dark matter electromagnetic dipole momentsarXiv:1007.5515.
- [143] G. Busoni, A. De Simone, J. Gramling, E. Morgante, A. Riotto, On the validity of the effective field theory for dark matter searches at the LHC, part II: Complete analysis for the s -channel, JCAP 1406 (2014) 060. arXiv:1402.1275, doi:10.1088/1475-7516/2014/06/060.
- 2870 [144] D. Racco, A. Wulzer, F. Zwirner, Robust collider limits on heavy-mediator dark matter, JHEP 1505 (2015) 009. arXiv:1502.04701, doi:10.1007/JHEP05(2015)009.
- 2875 [145] M. Endo, Y. Yamamoto, Unitarity bounds on dark matter effective interactions at LHC, JHEP 1406 (2014) 126. arXiv:1403.6610, doi:10.1007/JHEP06(2014)126.
- [146] PDF4LHC: Recommendation for LHC cross section calculations, <http://www.hep.ucl.ac.uk/pdf4lh/>, [Online; accessed 13-June-2015] (2015).

- 2880 [147] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, T. Stelzer, MadGraph 5 : Going beyond, JHEP 1106 (2011) 128. [arXiv:1106.0522](#), [doi:10.1007/JHEP06\(2011\)128](#).
- [148] T. Sjöstrand, S. Mrenna, P. Z. Skands, PYTHIA 6.4 physics and manual, JHEP 0605 (2006) 026. [arXiv:hep-ph/0603175](#), [doi:10.1088/1126-6708/2006/05/026](#).
- 2885 [149] M. Whalley, D. Bourilkov, R. Group, The Les Houches accord PDFs (LHAPDF) and LHAGLUE [arXiv:hep-ph/0508110](#).
- [150] M. L. Mangano, M. Moretti, F. Piccinini, M. Treccani, Matching matrix elements and shower evolution for top-quark production in hadronic collisions, JHEP 0701 (2007) 013. [arXiv:hep-ph/0611129](#), [doi:10.1088/1126-6708/2007/01/013](#).
- 2890 [151] V. Hirschi, R. Frederix, S. Frixione, M. V. Garzelli, F. Maltoni, et al., Automation of one-loop QCD corrections, JHEP 1105 (2011) 044. [arXiv:1103.0621](#), [doi:10.1007/JHEP05\(2011\)044](#).
- [152] SVN repository for Madgraph inputs for mono-Higgs EFT models, dimension 4 and 5, https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/EW_Higgs_D4D5/, [Online; accessed 04-May-2015] (2015).
- 2895 [153] J. Andrea, B. Fuks, F. Maltoni, Monotops at the LHC, Phys.Rev. D84 (2011) 074025. [arXiv:1106.6199](#), [doi:10.1103/PhysRevD.84.074025](#).
- [154] J.-L. Agram, J. Andrea, M. Buttignol, E. Conte, B. Fuks, Monotop phenomenology at the Large Hadron Collider, Phys.Rev. D89 (2014) 014028. [arXiv:1311.6478](#), [doi:10.1103/PhysRevD.89.014028](#).
- 2900 [155] I. Boucheneb, G. Cacciapaglia, A. Deandrea, B. Fuks, Revisiting monotop production at the LHC, JHEP 1501 (2015) 017. [arXiv:1407.7529](#), [doi:10.1007/JHEP01\(2015\)017](#).

- 2905 [156] N. D. Christensen, P. de Aquino, C. Degrande, C. Duhr, B. Fuks, et al., A comprehensive approach to new physics simulations, *Eur.Phys.J. C* 71 (2011) 1541. [arXiv:0906.2474](#), [doi:10.1140/epjc/s10052-011-1541-5](#).
- [157] SVN repository for Madgraph inputs for mono-top models, [https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/HF_](https://svnweb.cern.ch/cern/wsvn/LHCDMF/trunk/models/HF_SingleTop/)
2910 [SingleTop/](#), [Online; accessed 27-April-2015] (2015).
- [158] A. Belyaev, G. Cacciapaglia, I. Ivanov, F. Rojas, M. Thomas To appear.
- [159] N. G. Deshpande, E. Ma, Pattern of symmetry breaking with two Higgs doublets, *Phys.Rev. D* 18 (1978) 2574. [doi:10.1103/PhysRevD.18.2574](#).
- [160] R. Barbieri, L. J. Hall, V. S. Rychkov, Improved naturalness with a heavy
2915 Higgs: An alternative road to LHC physics, *Phys. Rev. D* 74 (2006) 015007. [arXiv:hep-ph/0603188](#), [doi:10.1103/PhysRevD.74.015007](#).
- [161] L. Lopez Honorez, E. Nezri, J. F. Oliver, M. H. Tytgat, The Inert Doublet Model: An archetype for dark matter, *JCAP* 0702 (2007) 028. [arXiv:hep-ph/0612275](#), [doi:10.1088/1475-7516/2007/02/028](#).
- 2920 [162] T. Hambye, F.-S. Ling, L. Lopez Honorez, J. Rocher, Scalar Multiplet dark matter, *JHEP* 0907 (2009) 090. [arXiv:0903.4010](#), [doi:10.1007/JHEP05\(2010\)066](#), [10.1088/1126-6708/2009/07/090](#).
- [163] L. Lopez Honorez, C. E. Yaguna, A new viable region of the Inert Doublet Model, *JCAP* 1101 (2011) 002. [arXiv:1011.1411](#), [doi:10.1088/1475-7516/2011/01/002](#).
2925
- [164] M. Gustafsson, E. Lundström, L. Bergström, J. Edsjö, Significant gamma lines from Inert Higgs Dark Matter, *Phys. Rev. Lett.* 99 (2007) 041301. [arXiv:astro-ph/0703512](#), [doi:10.1103/PhysRevLett.99.041301](#).
- [165] E. M. Dolle, S. Su, The Inert Dark Matter, *Phys.Rev. D* 80 (2009) 055012.
2930 [arXiv:0906.1609](#), [doi:10.1103/PhysRevD.80.055012](#).

- [166] P. Agrawal, E. M. Dolle, C. A. Krenke, Signals of Inert Doublet Dark Matter in neutrino telescopes, Phys. Rev. D79 (2009) 015015. arXiv:0811.1798, doi:10.1103/PhysRevD.79.015015.
- 2935 [167] S. Andreas, M. H. Tytgat, Q. Swillens, Neutrinos from Inert Doublet Dark Matter, JCAP 0904 (2009) 004. arXiv:0901.1750, doi:10.1088/1475-7516/2009/04/004.
- [168] E. Nezri, M. H. Tytgat, G. Vertongen, e^+ and anti-p from Inert Doublet model dark matter, JCAP 0904 (2009) 014. arXiv:0901.2556, doi:10.1088/1475-7516/2009/04/014.
- 2940 [169] C. Garcia-Cely, A. Ibarra, Novel gamma-ray spectral features in the Inert Doublet Model, JCAP 1309 (2013) 025. arXiv:1306.4681, doi:10.1088/1475-7516/2013/09/025.
- [170] A. Goudelis, B. Herrmann, O. Stål, Dark matter in the Inert Doublet Model after the discovery of a Higgs-like boson at the LHC, JHEP 1309 (2013) 106. 2945 arXiv:1303.3010, doi:10.1007/JHEP09(2013)106.
- [171] C. Burgess, M. Pospelov, T. ter Veldhuis, The Minimal model of nonbaryonic dark matter: A singlet scalar, Nucl.Phys. B619 (2001) 709–728. arXiv: hep-ph/0011335, doi:10.1016/S0550-3213(01)00513-2.
- 2950 [172] S. Andreas, T. Hambye, M. H. Tytgat, WIMP dark matter, Higgs exchange and DAMA, JCAP 0810 (2008) 034. arXiv:0808.0255, doi:10.1088/1475-7516/2008/10/034.
- [173] A. Arhrib, Y.-L. S. Tsai, Q. Yuan, T.-C. Yuan, An updated analysis of Inert Higgs Doublet Model in light of the recent results from LUX, PLANCK, AMS-02 and LHC, JCAP 1406 (2014) 030. arXiv:1310.0358, doi:10.1088/ 2955 1475-7516/2014/06/030.
- [174] E. Lundström, M. Gustafsson, J. Edsjö, The Inert Doublet Model and LEP II limits, Phys. Rev. D79 (2009) 035013. arXiv:0810.3924, doi:10.1103/PhysRevD.79.035013.

- [175] Q.-H. Cao, E. Ma, G. Rajasekaran, Observing the Dark Scalar Doublet
 2960 and its impact on the Standard-Model Higgs boson at colliders, Phys. Rev.
 D76 (2007) 095011. [arXiv:0708.2939](#), doi:10.1103/PhysRevD.76.
 095011.
- [176] E. Dolle, X. Miao, S. Su, B. Thomas, Dilepton signals in the Inert Dou-
 2965 blet Model, Phys. Rev. D81 (2010) 035003. [arXiv:0909.3094](#), doi:
 10.1103/PhysRevD.81.035003.
- [177] X. Miao, S. Su, B. Thomas, Trilepton signals in the Inert Doublet Model,
 Phys. Rev. D82 (2010) 035009. [arXiv:1005.0090](#), doi:10.1103/
 PhysRevD.82.035009.
- [178] M. Gustafsson, S. Rydbeck, L. Lopez-Honorez, E. Lundstrom, Status of the
 2970 Inert Doublet Model and the role of multileptons at the LHC, Phys.Rev.
 D86 (2012) 075019. [arXiv:1206.6316](#), doi:10.1103/PhysRevD.86.
 075019.
- [179] A. Arhrib, R. Benbrik, N. Gaur, $H \rightarrow \gamma\gamma$ in Inert Higgs Doublet Model,
 Phys. Rev. D85 (2012) 095021. [arXiv:1201.2644](#), doi:10.1103/
 2975 PhysRevD.85.095021.
- [180] B. Swiezewska, M. Krawczyk, Diphoton rate in the inert doublet model with
 a 125 GeV Higgs boson, Phys.Rev. D88 (3) (2013) 035019. [arXiv:1212.](#)
 4100, doi:10.1103/PhysRevD.88.035019.
- [181] A. Goudelis, B. Herrmann, O. Stål, Dark matter in the Inert Doublet Model
 2980 after the discovery of a higgs-like boson at the LHC, JHEP 1309 (2013) 106.
[arXiv:1303.3010](#), doi:10.1007/JHEP09(2013)106.
- [182] G. Belanger, B. Dumont, A. Goudelis, B. Herrmann, S. Kraml, et al., Dilepton
 constraints in the Inert Doublet Model from Run 1 of the LHC [arXiv:1503.](#)
 07367.
- 2985 [183] S. Kraml, B. Allanach, M. Mangano, H. Prosper, S. Sekmen, et al., Searches
 for new physics: Les Houches recommendations for the presentation of LHC

- results, Eur.Phys.J. C72 (2012) 1976. [arXiv:1203.2489](#), [doi:10.1140/epjc/s10052-012-1976-3](#).
- [184] The Durham HepData project, <http://hepdata.cedar.ac.uk/abouthepdata>, [Online; accessed 13-June-2015] (2015).
2990
- [185] J. S. Kim, D. Schmeier, J. Tattersall, K. Rolbiecki, A framework to create customised LHC analyses within CheckMATE [arXiv:1503.01123](#).
- [186] K. Cranmer, I. Yavin, RECAST: Extending the impact of existing analyses, JHEP 1104 (2011) 038. [arXiv:1010.2506](#), [doi:10.1007/JHEP04\(2011\)038](#).
2995
- [187] I.-W. Kim, M. Papucci, K. Sakurai, A. Weiler, ATOM: Automated Testing Of Models, to appear (2015).
- [188] D. Barducci, A. Belyaev, M. Buchkremer, G. Cacciapaglia, A. Deandrea, et al., Framework for model independent analyses of multiple extra quark scenarios, JHEP 1412 (2014) 080. [arXiv:1405.0737](#), [doi:10.1007/JHEP12\(2014\)080](#).
3000
- [189] PGS: simulation of a generic high-energy physics collider detector, <http://www.physics.ucdavis.edu/~conway/research/software/pgs/pgs4-general.htm>, [Online; accessed 13-June-2015] (2015).
3005
- [190] M. Baak, G. Besjes, D. C  rte, A. Koutsman, J. Lorenz, et al., HistFitter software framework for statistical data analysis, Eur.Phys.J. C75 (2015) 153. [arXiv:1410.1280](#), [doi:10.1140/epjc/s10052-015-3327-7](#).
- [191] INSPIRE: High-energy physics literature database, <http://inspirehep.net>, [Online; accessed 13-June-2015] (2015).
3010
- [192] S. Kraml, S. Kulkarni, U. Laa, A. Lessa, W. Magerl, D. Proschofsky, W. Waltenberger, SModelS: a tool for interpreting simplified-model results from the

LHC and its application to supersymmetry, Eur.Phys.J. C74 (2014) 2868.
arXiv:1312.4175, doi:10.1140/epjc/s10052-014-2868-5.

3015 [193] S. Kraml, S. Kulkarni, U. Laa, A. Lessa, V. Magerl, et al., SModelS v1.0: a
short user guidearXiv:1412.1745.

[194] M. Papucci, K. Sakurai, A. Weiler, L. Zeune, Fastlim: a fast LHC limit calculator, Eur.Phys.J. C74 (2014) 3163. arXiv:1402.0492, doi:10.1140/epjc/s10052-014-3163-1.