

Propositional Logic

Dave Dubin and Jodi Schneider

October 10, 2016

Concepts from earlier in the semester

- Propositions are the bearers of truth values: the kinds of things that can be true or false.

Concepts from earlier in the semester

- Propositions are the bearers of truth values: the kinds of things that can be true or false.
- States of affairs are the parts of reality responsible for making propositions true or false.

- We represent propositions with lower case letters (typically p , q , and r).

- We represent propositions with lower case letters (typically p , q , and r).
- A set of proposition letters generates a set of states of affairs: ways the world might be.

- We represent propositions with lower case letters (typically p , q , and r).
- A set of proposition letters generates a set of states of affairs: ways the world might be.
- n proposition letters generate 2^n states of affairs.

- We represent propositions with lower case letters (typically p , q , and r).
- A set of proposition letters generates a set of states of affairs: ways the world might be.
- n proposition letters generate 2^n states of affairs.
- $\{pqr, pq\bar{r}, p\bar{q}r, p\bar{q}\bar{r}, \bar{p}qr, \bar{p}q\bar{r}, \bar{p}\bar{q}r, \bar{p}\bar{q}\bar{r}\}$

Logical Operators

Table 1: 2.15 from van Benthem, et al.

Symbol	In natural language	Technical name
\neg	not	negation
\wedge	and	conjunction
\vee	or	disjunction
\rightarrow	if ... then	implication
\leftrightarrow	if and only if	equivalence

- A sentence constructed from proposition letters and operators is true or false in each state of affairs.

- A sentence constructed from proposition letters and operators is true or false in each state of affairs.
- Consider, for example: $(\neg p \vee q) \rightarrow r$

- A sentence constructed from proposition letters and operators is true or false in each state of affairs.
- Consider, for example: $(\neg p \vee q) \rightarrow r$
- The sentence is mapped to a truth value via the following tables

Semantics of the operators

ϕ	$\neg\phi$
0	1
1	0

Table 3: 2.18 from van Benthem, et al.

ϕ	ψ	$\phi \wedge \psi$	$\phi \vee \psi$	$\phi \rightarrow \psi$	$\phi \leftrightarrow \psi$
0	0	0	0	1	1
0	1	0	1	1	0
1	0	0	1	0	0
1	1	1	1	1	1

Exercise 2.3

You are given the information that p -or- q and $(\text{not-}p)$ -or- r . What can you conclude about q and r ? What is the strongest valid conclusion you can draw? (A statement is stronger than another statement if it rules out more possibilities.)

Drawing truth tables for expressions

p	q	r	$((\sim p \vee q) \rightarrow r)$				
1	1	1	0	1	1	1	1
1	1	0	0	1	1	0	0
1	0	1	0	1	0	1	1
1	0	0	0	1	0	1	0
0	1	1	1	0	1	1	1
0	1	0	1	0	1	0	0
0	0	1	1	0	1	1	1
0	0	0	1	0	1	0	0

Drawing truth tables for expressions

p	q	r	$((\sim p \vee q) \rightarrow r)$				
1	1	1	0	1	1	1	1
1	1	0	0	1	1	0	0
1	0	1	0	1	0	1	1
1	0	0	0	1	0	1	0
0	1	1	1	0	1	1	1
0	1	0	1	0	1	0	0
0	0	1	1	0	1	1	1
0	0	0	1	0	1	0	0

Drawing truth tables for expressions

p	q	r	$((\sim p \vee q) \rightarrow r)$				
1	1	1	0	1	1	1	1
1	1	0	0	1	1	0	0
1	0	1	0	1	0	1	1
1	0	0	0	1	0	1	0
0	1	1	1	0	1	1	1
0	1	0	1	0	1	0	0
0	0	1	1	0	1	1	1
0	0	0	1	0	1	0	0

Drawing truth tables for expressions

p	q	r	$((\sim p \vee q) \rightarrow r)$				
1	1	1	0	1	1	1	1
1	1	0	0	1	1	0	0
1	0	1	0	1	0	1	1
1	0	0	0	1	0	1	0
0	1	1	1	0	1	1	1
0	1	0	1	0	1	0	0
0	0	1	1	0	1	1	1
0	0	0	1	0	1	0	0

Drawing truth tables for expressions

p	q	r	$((\sim p \vee q) \rightarrow r)$				
1	1	1	0	1	1	1	1
1	1	0	0	1	1	0	0
1	0	1	0	1	0	1	1
1	0	0	0	1	0	1	0
0	1	1	1	0	1	1	1
0	1	0	1	0	1	0	0
0	0	1	1	0	1	1	1
0	0	0	1	0	1	0	0

Drawing truth tables for expressions

$(\sim p \vee q) \rightarrow r$

0	0	0
0	0	1
0	1	0
0	1	1
1	0	0
1	0	1
1	1	0
1	1	1

Drawing truth tables for expressions

$(\sim p \vee q) \rightarrow r$

10	0	0
10	0	1
10	1	0
10	1	1
01	0	0
01	0	1
01	1	0
01	1	1

Drawing truth tables for expressions

$(\sim p \vee q) \rightarrow r$

10 1 0 0

10 1 0 1

10 1 1 0

10 1 1 1

01 0 0 0

01 0 0 1

01 1 1 0

01 1 1 1

Drawing truth tables for expressions

$(\sim p \vee q) \rightarrow r$

10 1 0 0 0

10 1 0 1 1

10 1 1 0 0

10 1 1 1 1

01 0 0 1 0

01 0 0 1 1

01 1 1 0 0

01 1 1 1 1

Grammar of propositional logic

Let P be a set of proposition letters and let $p \in P$.

The following expression defines the recursive grammar for a logical expression ϕ in Backus–Naur Form:

$$\phi ::= p \mid \neg\phi \mid (\phi \wedge \phi) \mid (\phi \vee \phi) \mid (\phi \rightarrow \phi) \mid (\phi \leftrightarrow \phi)$$

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r

Grammatically *incorrect* expressions would include:

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$

Grammatically *incorrect* expressions would include:

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$
- $(s \leftrightarrow o)$

Grammatically *incorrect* expressions would include:

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$
- $(s \leftrightarrow o)$
- $(\neg(s \leftrightarrow \neg\neg\neg o) \rightarrow (q \wedge q))$

Grammatically *incorrect* expressions would include:

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$
- $(s \leftrightarrow o)$
- $(\neg(s \leftrightarrow \neg\neg\neg o) \rightarrow (q \wedge q))$

Grammatically *incorrect* expressions would include:

- $\neg \vee p$

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$
- $(s \leftrightarrow o)$
- $(\neg(s \leftrightarrow \neg\neg\neg o) \rightarrow (q \wedge q))$

Grammatically *incorrect* expressions would include:

- $\neg \vee p$
- $\vee)p\neg$

How many correct expressions are consistent with the last one?

Syntactically conforming expressions

Let $P = \{o, q, r, s\}$

Examples of grammatically conforming expressions include:

- r
- $\neg q$
- $(s \leftrightarrow o)$
- $(\neg(s \leftrightarrow \neg\neg\neg o) \rightarrow (q \wedge q))$

Grammatically *incorrect* expressions would include:

- $\neg \vee p$
- $\vee)p\neg$
- $\neg p \vee q \rightarrow r$

How many correct expressions are consistent with the last one?

These conforming expressions are all consistent with $\neg p \vee q \rightarrow r$

- $((\neg p \vee q) \rightarrow r)$

These conforming expressions are all consistent with $\neg p \vee q \rightarrow r$

- $((\neg p \vee q) \rightarrow r)$
- $(\neg(p \vee q) \rightarrow r)$

These conforming expressions are all consistent with $\neg p \vee q \rightarrow r$

- $((\neg p \vee q) \rightarrow r)$
- $(\neg(p \vee q) \rightarrow r)$
- $\neg((p \vee q) \rightarrow r)$

These conforming expressions are all consistent with $\neg p \vee q \rightarrow r$

- $((\neg p \vee q) \rightarrow r)$
- $(\neg(p \vee q) \rightarrow r)$
- $\neg((p \vee q) \rightarrow r)$
- $(\neg p \vee (q \rightarrow r))$

These conforming expressions are all consistent with $\neg p \vee q \rightarrow r$

- $((\neg p \vee q) \rightarrow r)$
- $(\neg(p \vee q) \rightarrow r)$
- $\neg((p \vee q) \rightarrow r)$
- $(\neg p \vee (q \rightarrow r))$
- $\neg(p \vee (q \rightarrow r))$

- Semantics is the relationship of a language to the part of the world that we're modeling.

- Semantics is the relationship of a language to the part of the world that we're modeling.
- Valuations are functions from expressions to truth values.

- Semantics is the relationship of a language to the part of the world that we're modeling.
- Valuations are functions from expressions to truth values.
- “ $V(\phi) = 1$ ” means the formula (or sentence) ϕ is true in the state of affairs represented by the function V . “ $V(\phi) = 0$ ” means that ϕ is false in the state of affairs represented by the function V .

More on semantics

- Semantics is the relationship of a language to the part of the world that we're modeling.
- Valuations are functions from expressions to truth values.
- “ $V(\phi) = 1$ ” means the formula (or sentence) ϕ is true in the state of affairs represented by the function V . “ $V(\phi) = 0$ ” means that ϕ is false in the state of affairs represented by the function V .
- For “ $V(\phi) = 1$ ” we also write “ $V \models \phi$ ” read as “ V is a model of ϕ ” or “ V satisfies ϕ .”

More on semantics

- Semantics is the relationship of a language to the part of the world that we're modeling.
- Valuations are functions from expressions to truth values.
- “ $V(\phi) = 1$ ” means the formula (or sentence) ϕ is true in the state of affairs represented by the function V . “ $V(\phi) = 0$ ” means that ϕ is false in the state of affairs represented by the function V .
- For “ $V(\phi) = 1$ ” we also write “ $V \models \phi$ ” read as “ V is a model of ϕ ” or “ V satisfies ϕ .”
- If V doesn't satisfy ϕ we write “ $V \not\models \phi$ ”. In other words $V(\phi) = 0$.

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.
- A statement ϕ is logically false if it is false in every state of affairs generated by its propositional variables.

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.
- A statement ϕ is logically false if it is false in every state of affairs generated by its propositional variables.
- If a statement ϕ is neither logically true or logically false then it is contingent.

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.
- A statement ϕ is logically false if it is false in every state of affairs generated by its propositional variables.
- If a statement ϕ is neither logically true or logically false then it is contingent.
- Examples:

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.
- A statement ϕ is logically false if it is false in every state of affairs generated by its propositional variables.
- If a statement ϕ is neither logically true or logically false then it is contingent.
- Examples:
 - ① $(q \vee \neg q)$ is logically true.

Logical truth and logical falsity.

- A statement ϕ is logically true if it is true in every state of affairs generated by its propositional variables.
- A statement ϕ is logically false if it is false in every state of affairs generated by its propositional variables.
- If a statement ϕ is neither logically true or logically false then it is contingent.
- Examples:
 - ① $(q \vee \neg q)$ is logically true.
 - ② $(q \wedge \neg q)$ is logically false.

- A set of propositional logic statements is consistent if at least one state of affairs satisfies every statement in the set.

- A set of propositional logic statements is consistent if at least one state of affairs satisfies every statement in the set.
- A set of propositional logic statements is inconsistent if no state of affairs satisfies every statement in the set.

- A conclusion is *valid* with respect to a set of premises if the conclusion is true in every situation where the premises are true (van Benthem, et al, page 2-4).

- A conclusion is *valid* with respect to a set of premises if the conclusion is true in every situation where the premises are true (van Benthem, et al, page 2-4).
- One can validly infer a conclusion ϕ from a set of premises P if the negation of ϕ is inconsistent with the set of statements P .

(From van Benthem, et al., chapter 2)

- ① Computing a truth value for a formula takes linear time.

(From van Bentham, et al., chapter 2)

- 1 Computing a truth value for a formula takes linear time.
- 2 Computing a truth table for validity takes exponential time.

(From van Benthem, et al., chapter 2)

- ① Computing a truth value for a formula takes linear time.
- ② Computing a truth table for validity takes exponential time.
- ③ The problem of testing for validity in propositional logic is decidable: there exists a mechanical method that computes the answer, at least in principle.