exo. Even of the density function, & (x, y) = \( 2 e^{(x,y)} if 0 < y < x Q) Toil a density function? (Q) compute MD X and MD Y. 1) properly 1. Yx E X(a), y E Y(a), g(x,y) >0. positivity It is always justive due to e bury always justing for \$ ZER  $\frac{\text{projecting 2.}}{R^2} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} (x, y) \, dy \, dx = 1 \quad \text{way 2=>} \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{1}{2} e^{-(x+y)} \, dy \, dx = 1$   $x = 0 \quad y = 0$ =20-4, [-0-2] = 2 e-8 (e-8 + e-0) = 2.0-8 e-4  $\int_{2e^{-2y}}^{+\infty} dy = \left[ -e^{-2y} \right]^{+\infty} = e^{0} - e^{-\infty} = 1$ way? \( \int \frac{1}{2} = \frac{1}{2} = \frac{1}{2} \\ \frac{1}{2} = \fra  $\int_{x=0}^{+\infty} 2e^{-x} = 2e^{-2x} dx = 2\int_{e^{-x}}^{+\infty} e^{-2x} dx$   $= 2\left[\frac{1}{2}e^{-2x} - e^{-x}\right]^{+\infty}$   $= 2\left[\frac{1}{2}e^{-2x} - e^{-x}\right]^{+\infty}$   $= 2e^{-x} - 2e^{-2x}$  $=2.\left(\frac{1}{2}.0-0\right)-\left(\frac{1}{2}-1\right)$ =2.(-(-1))

12611	$MD.X = \int_{y=0}^{x} 2e^{-(x+y)} dy = 2e^{-x} \int_{y=0}^{x} e^{-y} dy$
	2-2 [ -47x
	$=2e^{-x}\left[-e^{-y}\right]^{x}$ $=2e^{-x}\left[-e^{-y}\right]^{x}$ $=2e^{-x}\cdot\left(e^{-x}-e^{-x}\right)$ $=2e^{-x}\cdot\left(e^{-x}-e^{-x}\right)$ $=2e^{-x}\cdot\left(e^{-x}-e^{-x}\right)$ $=2e^{-x}\cdot\left(e^{-x}-e^{-x}\right)$
	= 2 e . (e - e 2) (0 otherwin
23	pt00
(site	$MD.Y = \int_{\infty}^{+\infty} \frac{1}{2} e^{-(x+y)} dx = 2e^{-y} \int_{\infty}^{+\infty} e^{-x} dx$
day clas	
	=2e-4. [-e-x] = 2e-4. (e°-e-)
1 Lat 16	=20-9
	MD. Y = {2 e 4 y \ [0; x]
	20 otherwise
	12 C 4 C 4 C 4 C 4 C 4 C 4 C 4 C 4 C 4 C
	10 m
	1 00 0 0 00 1 00 00 00 00 00 00 00 00 00
	1 2 C 1 day = 1 2 C 1 C 2 C 2 C 2 C 2 C 2 C 2 C 2 C 2
	(mil 1 ) changed do s> 2 200 de 2 2 2 2 2 2
100	8.4 az 3
- 1	
	1
	1
	1
	1
	1
	1 2 6 2 2 6 2 4 4 4 5 2 6 2 6 2 6 4 4 4 5 6 2 6 2 6 2 6 2 6 2 6 2 6 2 6 2 6 2 6