ChemCritic 文献证据卡

PMC10731675 | 氢胺化 / C – N 偶联 | 采集状态 imported_unchecked

标题	Hydroamination of alkynes catalyzed by NHC-Gold(I) complexes: the non-monotonic effect of substituted arylamines on the catalyst activity
作者	Sirignano M., D ’ Amato A., Costabile C., Mariconda A., Crispini A., Scarpelli F. 等 7 人
出处	Frontiers in Chemistry 2023
DOI	10.3389/fchem.2023.1260726
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC10731675/
许可	https://creativecommons.org/licenses/by/4.0/This is an open-access article distributed under the terms of the Creative C

摘要（原文节选）

Imines are valuable key compounds for synthesizing several nitrogen-containing molecules used in biological and industrial fields. They have been obtained, as highly regioselective Markovnikov products, by reacting several alkynes with arylamines in the presence of three new N-Heterocyclic carbene gold(I) complexes (3b, 4b, and 6b) together with the known 1-2b and 7b gold complexes as well as silver complexes 1-2a. Gold(I) complexes were investigated by means of NMR, mass spectroscopy, elemental analysis, and X-ray ⋯

数据表 · TABLE 2（4 行，含数值行 4 行）

Screening, promoted by 7b, of the silver salt co-catalyst for the hydroamination reaction of phenylacetylene with aniline.

Entry a	Co-catalyst	Yield (%) b
1	AgSbF6	36
2	AgPF6	29
3	AgNO3	15
4	AgOAc	n.d

脚注（逐字）：a Reaction conditions: 1.0 mmol aniline, 1.5 mmol phenylacetylene, 1% mol 7b, 2% mol Ag salt, 90 ° C oil-bath, 16h, dry CH3CN (1 mL). b Yields are averaged from two runs and determined by 1H-NMR, analysis through internal standard.

正文对应句（逐字）：As depicted in Table 5, all substrates generated the expected imines. Quantitative yields are obtained with aniline and methyl substituted anilines (entries 1 – 4, Table 5). Yields decrease when isopropyl substituted anilines and 2-naphthylamine are used (Entries 5 – 8, Table 5), as well as with 4-metho

数据表 · TABLE 3（6 行，含数值行 6 行）

Screening of the solvent in the reaction of hydroamination promoted by 2b.

Entry a	Solvent	Yield (%) b
1	CH3CN	70
2	1,4-Dioxane	20
3	Dimethylsulfoxide	13
4	C2H2Cl4	45
5	Toluene	50
6	Ethylene Carbonate	35

脚注（逐字）：a Reaction conditions: 1.0 mmol aniline, 1.5 mmol phenylacetylene, 1% mol 2b, 2% mol Ag salt, 90 ° C oil-bath, 16h, solvent (1 mL). b Yields are averaged from two runs and determined by 1H-NMR, analysis through internal standard.

正文对应句（逐字）：As depicted in Table 5, all substrates generated the expected imines. Quantitative yields are obtained with aniline and methyl substituted anilines (entries 1 – 4, Table 5). Yields decrease when isopropyl substituted anilines and 2-naphthylamine are used (Entries 5 – 8, Table 5), as well as with 4-metho

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。