ChemCritic 文献证据卡

PMC11110061 | 烯化（Wittig 等） | 采集状态 imported_unchecked

标题	Flow Chemistry for Synthesis of 2-(C-Glycosyl)acetates from Pyranoses via Tandem Wittig and Michael Reactions
作者	Bennett J., Murphy P.
出处	Organic Process Research & Development 2024
DOI	10.1021/acs.oprd.3c00414
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC11110061/
许可	https://creativecommons.org/licenses/by/4.0/Permits the broadest form of re-use including for commercial purposes, provi

摘要（原文节选）

C-Glycosyl compounds (C-glycosides) are a class of saccharide derivatives with improved stability over their O-linked counterparts. This paper reports the synthesis of several trans-2-(C-glycosyl)acetates via a tandem Wittig – Michael reaction from pyranoses (cyclic hemiacetals) using continuous flow processing, which gave improvements compared to reactions conducted in round-bottom flasks. Products were isolated in yields of >60% from reactions of benzyl-protected xylopyranoses, glucopyranoses, and galactopyranos ⋯

数据表 · Table 2（4 行，含数值行 4 行）

Isolation of Michael Addition Product 7b and Wittig Intermediate 11 in Absence of Base

					isolated yields (%)	
entry	equiv of Ph3PCHCO2Et	temperature (° C)	time (min)	flow rate (mL/min)	7b	11
1	3	220	50	0.1	35	14
2	3	230	60	0.083	30	12
3	5	210	60	0.083	42	10

正文对应句（逐字）：Pyranoses, which are hemiacetals with a latent aldehyde, can undergo the Wittig or Horner – Wadsworth – Emmons (HWE) reaction, and when these reactions can be used in conjunction with conjugate addition (Michael reaction), 2-(glycosyl)acetates can be formed (Scheme 1).19,21 The tandem Wittig – Michael pro

数据表 · Table 3（5 行，含数值行 4 行）

DBU-Improved Formation of C-Glycosyl Compounds from 3 and 8

		reactor 1	reactor 2			products (yields, %)		
entry	reactant	temperature (° C)	time (min)	temperature (° C)	time (min)	flow rate per pump (mL/min)	BP (bar)	β
1	8	210	40	145	40	0.125	16	10 (45)
2	8	180	45	130	45	0.11	8	10 (55)
3	3	200	65	145	65	0.077	8	7b (63)
4	3	210	40	145	40	0.125	16	7b (67)

正文对应句（逐字）：Pyranoses, which are hemiacetals with a latent aldehyde, can undergo the Wittig or Horner – Wadsworth – Emmons (HWE) reaction, and when these reactions can be used in conjunction with conjugate addition (Michael reaction), 2-(glycosyl)acetates can be formed (Scheme 1).19,21 The tandem Wittig – Michael pro

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。