ChemCritic 文献证据卡

PMC11199998 | Suzuki – Miyaura 偶联 | 采集状态 imported_unchecked

标题	Palladium‐Catalyzed Atroposelective Suzuki – Miyaura Coupling to Construct Axially Chiral Tetra‐Substituted α‐Boryl Styrenes
作者	Li X., Kong L., Yin S., Zhou H., Lin A., Yao H. 等 7 人
出处	Advanced Science 2024
DOI	10.1002/advs.202309706
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC11199998/
许可	https://creativecommons.org/licenses/by/4.0/This is an open access article under the terms of the http://creativecommons

摘要（原文节选）

AbstractPalladium‐catalyzed Suzuki – Miyaura (SM) coupling is a valuable method for forming C—C bonds, including those between aryl moieties. However, achieving atroposelective synthesis of axially chiral styrenes via SM coupling remains challenging. In this study, a palladium‐catalyzed atroposelective Suzuki – Miyaura coupling between gem‐diborylalkenes and aryl halides is presented. Using the monophosphine ligand Me‐BI‐DIME (L2), a range of axially chiral tetra‐substituted acyclic styrenes with high yields and excell …

数据表 · Table 1（12 行，含数值行 12 行）

Optimization of the reaction conditions.

Entry a)	[Pd]	Ligand	Base	Yield [%]	Ee [%]
1	Pd(OAc)2	L1	K3PO4	10	15
2	Pd(OAc)2	L1	KOH	28	15
3	Pd(OAc)2	L1	t BuOK	35	14
4	Pd(OAc)2	L1	t BuONa	64	16
5	Pd(OAc)2	L2	t BuONa	82	59
6	Pd(OAc)2	L3	t BuONa	30	38
7	Pd(OAc)2	L4	t BuONa	40	60
8 b)	Pd(OAc)2	L2	t BuONa	85	19
9 c)	Pd(OAc)2	L2	t BuONa	96	49
10	PdCl2	L2	t BuONa	72	57
11	Pd(dba)2	L2	t BuONa	88	57
12 d)	Pd(dba)2	L2	t BuONa	90	86

脚注（逐字）：a) Reaction conditions: 1a (0.1 mmol, 1.0 equiv), 2a (0.12 mmol, 1.2 equiv), [Pd] (10 mol %), L (15 mol %) and base (3.0 equiv in 0.2 mL H2O) in 1.0 mL toluene at 30 °C under argon atmosphere for 48 h. Yields were determined by 1H NMR with 1,3,5‐trimethoxybenzene as the internal standard. The enanti

正文对应句（逐字）：We initiated our studies with 1‐iodo‐2‐methoxynaphthalene (1a) and gem‐diborylalkene 2a as the model substrates to establish an effective catalytic system. The representative results are summarized in Table 1 . First, a variety of bases were examined in the presence of 10 mol % Pd(OAc)2, 15 mol % BI

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。