ChemCritic 文献证据卡

PMC11720983 | 氢胺化 / C – N 偶联 | 采集状态 imported_unchecked

标题	Electrochemical α -C(sp3) – H/N – H Cross-Coupling of Isochromans and Azoles
作者	Li G., Yan B., Wu L., Li Y., Hao X., Gong M. 等 8 人
出处	Molecules 2024
DOI	10.3390/molecules30010004
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC11720983/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

Isochroman and azole moieties are both present in a wide variety of biologically active molecules. Their efficient combination under mild reaction conditions is beneficial for obtaining small-molecule drug candidates. In this paper, we describe electrochemical α -C(sp3) – H/N – H cross-coupling reactions between isochromans and azoles, yielding products in moderate to excellent yields. This protocol does not require any catalysts or exogenous oxidants and can be performed at room temperature under air. Control experimen ⋯

数据表 · Table 1（16 行，含数值行 16 行）

Optimization of the reaction conditions a.

Entry	Anode/Cathode	Electrolyte	Additive	Yield (%) b
1	RVC/Ni	n-Bu4NBF4	-	trace
2	RVC/Ni	n-Bu4NBF4	TfOH	39
3	RVC/Ni	n-Bu4NBF4	AcOH	2
4 c	RVC/Ni	n-Bu4NBF4	TfOH	73
5 c	RVC/Ni	n-Bu4NClO4	TfOH	54
6 c	RVC/Ni	n-Bu4NPF6	TfOH	88
7 c	RVC/Ni	n-Bu4NOAc	TfOH	39
8 c	C/Ni	n-Bu4NPF6	TfOH	73
9 c	RVC/Pt	n-Bu4NPF6	TfOH	71
10 c,d	RVC/Ni	n-Bu4NPF6	TfOH	50
11 c,e,f	RVC/Ni	n-Bu4NPF6	TfOH	72
12 c,g	RVC/Ni	n-Bu4NPF6	TfOH	88
13 c,h	RVC/Ni	n-Bu4NPF6	TfOH	69
14 c,f	RVC/Ni	n-Bu4NPF6	TfOH	84
15 c	RVC/Ni	n-Bu4NPF6	-	70
16 c,i	RVC/Ni	n-Bu4NPF6	TfOH	50

脚注（逐字）：a Reaction conditions: 1a (0.3 mmol), 2a (0.45 mmol, 1.5 equiv.), I = 10 mA, electrolyte (0.1 M), additive (0.45 mmol), MeCN (6 mL), at room temperature for 3 h. b Isolated yield. c 1a (1.2 mmol), 2a (0.3 mmol). d EtOH was used as solvent. e 7.5 mA. f 5 h. g 12.5 mA. h 2.5 h. i Ar.

正文对应句（逐字）：We started our investigation by employing isochroman (1a) and indazole (2a) as template substrates (Table 1). The reaction showed minimal conversion in an undivided RVC/Ni cell at 10 mA without any additives (entry 1). Pleasingly, the introduction of trifluoromethanesulfonic acid (TfOH) significantl

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。