ChemCritic 文献证据卡

PMC12147128 | Miyaura 硼化 / C – H 硼化 | 采集状态 imported_unchecked

标题	An Adaptive Palladium Single-Atom Catalyst Enabling Reactivity Switching between Borylation and C – C Coupling
作者	Saptal V., Saetta C., Laufenb ö ck A., Sterrer M., Kwon I., Lucotti A. 等 12 人
出处	Journal of the American Chemical Society 2025
DOI	10.1021/jacs.4c17943
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC12147128/
许可	https://creativecommons.org/licenses/by/4.0/This article is licensed under CC-BY 4.0

摘要（原文节选）

The development of single-atom catalysts (SACs) with site-specific and tunable catalytic functionalities remains a highly desirable yet challenging goal in catalysis. In this study, we report a SAC featuring anisotropic coordination cavities synthesized via a one-step polymerization of 2,6-diaminopyridine and cyanuric chloride. These cavities provide a robust framework for anchoring isolated Pd single atoms with exceptional stability. The unique broken symmetry of the catalyst ’ s local structure enables precise cont ⋯

数据表 · 1（24 行，含数值行 24 行）

Optimization of the Reaction Conditions for the Miyaura Borylation Reaction

entry	solvent	base	temperature (° C)	time (h)	yield (%)
1	toluene	NaOAc	90	12	20
2	dioxane	NaOAc	90	12	17
3	DMF	NaOAc	90	12	48
4	water	NaOAc	90	12	42
5	water – ethanol	NaOAc	90	12	38
6	toluene – ethanol	NaOAc	90	12	24
7	dioxane – ethanol	NaOAc	90	12	68
8	ethanol	NaOAc	90	12	76
9	methanol	NaOAc	90	12	82
10	methanol	K2CO3	90	12	24
11	methanol	Na2CO3	90	12	17
12	methanol	KOAc	90	12	91
13	methanol	KOAc (3)	90	9	94
14	methanol	KOAc (1.5)	90	9	83
15	methanol	KOAc (1)	90	9	46
16	methanol	no base	80	9	21
17	methanol	KOAc	90	7	90
18	methanol	KOAc	80	7	90
19	methanol	KOAc	50	7	44
20	methanol	KOAc	80	5	84
21	methanol	KOAc	80	3	69
22	methanol	KOAc	80	9	5
23	methanol	KOAc	80	7	60
24	methanol	KOAc	80	7	51

脚注（逐字）：aUnless indicated otherwise, the reaction conditions were iodobenzene (1 mmol), B2Pin2 (1.3 mmol), base (2 mmol), and solvent (5 mL).bIsolated yields.cReaction performed in the absence of Pd1@TRIDAP catalyst.dReaction using Pd@C3N4.eReaction using Pd/C (4.64 mg of 10 wt.% Pd/C).

正文对应句（逐字）：The as-synthesized Pd1@TRIDAP was employed in the Miyaura borylation of aryl halides to synthesize arylboranates, which serve as key building blocks in the construction of complex molecular architectures for drug discovery and development. The optimization of the reaction conditions commenced with i

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。